

Lecture Notes on Core Physics

Will Lancer

`will.m.lancer@gmail.com`

These are notes on basic physics, covering classical mechanics, classical field theory, statistical mechanics, and quantum mechanics. Resources used: **(fill in later)**

- Classical mechanics:
- Classical field theory:
- Statistical mechanics:
- Quantum mechanics:

Remark 0.1 (Notes on notation). A list of notation I will use:

- $\bar{d} = d/2\pi$. $\oint^D(x) = (2\pi)^D \delta^D(x)$.
- My standard “volume” will be Ω . I’ll denote its boundary using the boundary operator ∂ , so my standard surface element is $\partial\Omega$.
- I use the mostly-plus metric, so $\eta_{\mu\nu} = \text{diag}(-1, 1, \dots, 1)$. I’ll also work in D space-time dimensions frequently, so $(D - 1)$ spatial dimensions plus one time dimension.
- We work in natural units almost everywhere, with the notable exception of non-relativistic electromagnetism.

Contents

1	Classical Mechanics	3
1.1	Newtonian mechanics	3
1.2	Configuration space	4
1.3	Examples of Lagrangians	5
1.4	Rigid bodies	6
1.5	Phase space	6
2	Classical Electrodynamics	7
2.1	Mathematical identities	7
2.2	Generalities	7
2.3	Conductor electrostatics	8
2.4	Dielectric electrostatics	13
2.5	Magnetostatics	13
2.6	Electrodynamics	14
2.7	Retarded potentials	14
2.8	Waves	14
3	Statistical Mechanics	15
3.1	The microscopic picture	15
4	Quantum Mechanics	17
4.1	Assumptions	17
4.2	Perturbation theory	17

1 Classical Mechanics

1.1 Newtonian mechanics

All of Newtonian mechanics is predicated upon the fact that we live in Galilean space. Galilean space is the triple $G = (\mathbb{A}^4, t, g)$, where $\mathbb{A}$ is an affine space, t is a time function, and g is the Euclidean metric. t and g do exactly what you think they do. Note that g is restricted to act on the space components of our vectors, which we take to be four-vectors at the outset.

It's an experimental observation that symmetry exists. A symmetry is defined as an action on a mechanical system that doesn't change any observables of that system. The set of transformations that we can perform on Galilean space to give us another Galilean space forms a group, which we call the Galilean group,

$$\mathcal{G} = \text{span}\{\mathbf{a}, \mathbf{V}, R_{ab}, \tau\}.$$

It has ten generators, as you can see. Experimentally, we can observe that a system being symmetric under certain generators of the Galilean group implies that that system has a conserved quantity associated with that generator. While we will have a systematic way to define these later (Noether's theorem), right now we can just look at experiments and come up with the following definitions (particle definitions, not extended bodies).

Symmetry	Conserved quantity
Translational symmetry	Linear momentum: $\mathbf{p} := m\mathbf{v}$
Rotational symmetry	Angular momentum: $\boldsymbol{\ell} := \mathbf{r} \times \mathbf{p}$
Time-translation symmetry	Energy: $E := T + V$

Here, $T := (1/2)m|\dot{\mathbf{r}}|^2$, and V is the potential energy, which is calculated on a system by system basis. Note that it makes intuitive sense that the only scalar conserved quantity is associated with time, which is a scalar, as opposed to translations or rotations, which are inherently associated with vectors.

We may experimentally observe that symmetries are broken. To do one of these very difficult experiments in the comfort of your own home, take a pen and drop it onto the floor. It does not have translational invariance; this symmetry was *broken* by *something*. We now define those somethings:

Symmetry	Conserved quantity	Symmetry breaker
Translational symmetry	Linear momentum: $\mathbf{p} := m\mathbf{v}$	Force: $\mathbf{F} := d\mathbf{p}/dt$
Rotational symmetry	Angular momentum: $\boldsymbol{\ell} := \mathbf{r} \times \mathbf{p}$	Torque: $\boldsymbol{\tau} = d\boldsymbol{\ell}/dt$
Time-translation symmetry	Energy: $E := T + V$	Power: $P = dE/dt$

Again, we see energy is the odd one out here; this is because the practical way of changing the energy of a system is by doing *work* on it, which is defined as the one-form $dW = -F dx$. Yet we define the symmetry breaker as power. This suggests a fundamental fault in the definitions we use, which we get into now.

Remark 1.1 (Why is energy so weird?). Why does energy not follow the nice schema we have laid out for our conserved quantities? After all, the definitions of force and torque are quite

intuitive, and then out flops power. What? I forgot about power after high school; why is that there? (**maybe something to do with entropy**)

1.2 Configuration space

We would like to catalog where things are; a **configuration space** allows us to do this.

Definition 1.1 (Configuration space)

A **configuration space** is the space of all possible positions of our system.

Example 1.1 (Examples of configuration spaces)

Some examples:

- The standard configuration space: $\mathbb{R}^3$. This corresponds to one particle in three dimensions, or three particles in one dimension, or what have you.
- An important example: $\mathbb{R}^6$. This is the configuration space corresponding to two particles moving in three dimensions. Note that spaces like these show that the wave function isn't a field like most people think it is.
- The circle: S^1 . This corresponds to a particle that is constrained to move on, you guessed it, a circle!
- The circle²: $S^1 \times S^1$. Physically, this corresponds to *two* things constrained to move on circles. An example of this is the double pendulum.

Remark 1.2 (The intuition behind this). Intuitively, the configuration space of your system partitions a dimension for each degree of freedom. Look at the previous examples for confirmation of this heuristic (**(fact?)**).

Ok, so configuration space seems like a natural place to try and define things on (it tells us where our particles are, and predicting the positions of particles is the goal of classical mechanics). We can define some sort of tensor function on this space to tell us the dynamics of our system.

The most natural tensors are going to be scalars and vectors; this is just because we have those nice conserved quantities from before, and they are all scalars and vectors. Let's study the simplest case first to get some constraints on this function.

Remark 1.3. The simplest case is the most symmetric one with the fewest components: so a scalar function with Galilean symmetry.

Idea 1.1 (Scalar function on configuration space)

Call this scalar function L . Denote a point in configuration space by $\mathbf{q}$, and its time derivative by $\dot{\mathbf{q}}$. Then the most general L is some function of powers of these quantities, $L = L(\mathbf{q}, \dot{\mathbf{q}}, |\dot{\mathbf{q}}|^2, \dots, t)$. Note that we implemented time-dependence somewhat ad hoc; this is

because we want to talk about dynamics, which implicitly refers to time.

We now implement our symmetry considerations onto L . Note that by translational and rotational invariance, L can only be a function of $|\mathbf{q} - \mathbf{q}'|$, and furthermore, only a function of $|\mathbf{q}|^{2n}$ and $|\dot{\mathbf{q}}|^{2n}$. By time-translation invariance, $L \neq L(t)$. The most natural rotationally invariant objects we can make are $\mathbf{q} \cdot \mathbf{q} = |\mathbf{q}|^2$ and $\dot{\mathbf{q}} \cdot \dot{\mathbf{q}} = |\dot{\mathbf{q}}|^2$.

So the form of L is

$$L(|\mathbf{q}|^2, |\dot{\mathbf{q}}|^2, \dots, |\mathbf{q}|^{2n}, |\dot{\mathbf{q}}|^{2n}, |\mathbf{q} - \mathbf{q}'|^n) = \frac{1}{2} A_{mn} \dot{q}_m \dot{q}_n - \frac{1}{2} B_{mn} q_m q_n + (\text{other stuff}),$$

where A_{mn} and B_{mn} are matrices.

So we've constrained L as much as we can (**I think**). We are now going to invoke the action principle to constrain L more.

Idea 1.2 (The action principle)

It's an experimental observation that the following quantity, when minimized, produces the correct equations of motion for dynamical systems:

$$S[\mathbf{q}] = \int dt L.$$

This quantity is called the **action**.

Let's continue to constrain L . Add a constant shift in position or velocity to L . We want this to contribute a total derivative to S , as this should be a symmetry of L . Doing this, we can see that the only way for this to be a symmetry is if (**BLAH**)

1.3 Examples of Lagrangians

Most important example: oscillations

The ideas behind oscillations are *linearity* and *time-translation invariance*. As we'll see in this section, these considerations give rise to the idea of **normal modes**, which are just the choice of coordinates that makes these ideas most manifest.

Integrable systems

Let's define the term I used in the title of this section.

Definition 1.2 (Integrability)

A system is said to be **integrable** if it has as many first integrals as it has degrees of freedom.

Example 1.2 (Examples of integrable systems)

Here are some common examples of integrable systems.

- (**orbits**)

- (any system with one symmetry and one DOF)
- (something else idk)

1.4 Rigid bodies

(need to re-learn this subject; i'm excited to do that, though! really looking forward to it. tong for a first pass, then goldbart, then landau, then arnol'd? something like that)

1.5 Phase space

Generalities

Regardless of how we derive the Hamiltonian, we first must define the canonical momentum $\mathbf{p} = \partial_{\dot{\mathbf{q}}} L$. (comment on this—what's deep about the canonical momentum? how can it be motivated? maybe in relation to the ele but eh, feels weak) Then, we can define the **Hamiltonian** of a system as the Legendre transform of the Lagrangian for that system,

$$H := \mathbf{p} \cdot \dot{\mathbf{q}} - L|_{\mathbf{p}=\partial_{\dot{\mathbf{q}}} L}.$$

todos:

- cover legendre transform in an idea box; put it before the expression for the lagrangian; use Arnol'd to do this
- motivate the canonical momentum; why is it useful? i don't really get it
- two-form picture of Hamiltonian mechanics

Poisson Brackets

Coordinate Transformations

Symplectic Mechanics

2 Classical Electrodynamics

2.1 Mathematical identities

We'll be using index notation throughout these notes, so most vector identities are useless to put here. We will simply record those identities that are hard to remember.

Differential identities

$$\nabla_{\text{sph}}^2 = \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2} \underbrace{\left(\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\phi^2 \right)}_{\nabla_\Omega^2}$$

$$\nabla_{\text{cyl}}^2 = \frac{1}{r} \partial_r (r \partial_r) + \frac{1}{r^2} \partial_\phi^2 + \partial_z^2$$

Integral identities

Special functions

2.2 Generalities

The action for electromagnetism is

$$S[A] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + j^\mu A_\mu \right),$$

where $F_{\mu\nu} := 2\partial_{[\mu} A_{\nu]}$, and j^μ our current. Varying this action gives the equations of motion

$$\frac{\delta S[A]}{\delta A_\nu} = 0 \implies \boxed{\partial_\mu F^{\mu\nu} = -j^\nu}.$$

There are also two natural constraints that arise from this construction: the first comes from the anti-symmetry of F ,

$$\partial_\nu \partial_\mu F^{\mu\nu} = -\partial_\nu j^\nu = 0 \implies \boxed{\partial_\mu j^\mu = 0}.$$

Thus, to couple the matter sector of electromagnetism with our $U(1)$ gauge field, the current must be locally conserved; you can also view this as a consequence of demanding that $S[A]$ is gauge invariant¹ Additionally, from the definition of $F_{\mu\nu}$, we have that

$$\boxed{\epsilon^{\mu\nu\rho\sigma} \partial_\nu F_{\rho\sigma} = 0}.$$

One can see this is true as follows: write out $\partial_\nu F_{\rho\sigma} = \partial_\nu \partial_\rho A_\sigma - \partial_\nu \partial_\sigma A_\rho$ and then contract with ϵ , renaming indices in the second term:

$$\begin{aligned} \epsilon^{\mu\nu\rho\sigma} \partial_\nu F_{\rho\sigma} &= \epsilon^{\mu\nu\rho\sigma} (\partial_\nu \partial_\rho A_\sigma - \partial_\nu \partial_\sigma A_\rho) \\ &= \epsilon^{\mu\nu\rho\sigma} \partial_\nu \partial_\rho A_\sigma - \epsilon^{\mu\nu\sigma\rho} \partial_\nu \partial_\rho A_\sigma \\ &= \epsilon^{\mu\nu\rho\sigma} \partial_\nu \partial_\rho A_\sigma + \epsilon^{\mu\nu\rho\sigma} \partial_\nu \partial_\rho A_\sigma \\ &= 2\epsilon^{\mu\nu\rho\sigma} \partial_\nu \partial_\rho A_\sigma = 2\epsilon^{\mu[\nu\rho]\sigma} \partial_{(\nu} \partial_{\rho)} A_\sigma = 0. \end{aligned}$$

¹You can trick yourself into thinking that the conservation of j^μ arises due to some sort of “Noether’s theorem” applied to gauge redundancy—this is, of course, false. The conservation of current in electromagnetism is a *consistency condition* for matter to couple with electromagnetism, not a conservation law in the sense of Noether.

Together, these two equations constitute **Maxwell's equations**:

$$\boxed{\partial_\mu F^{\mu\nu} = -j^\nu,} \quad \boxed{\epsilon^{\mu\nu\rho\sigma} \partial_\nu F_{\rho\sigma} = 0.}$$

Besides the equations of motion, we'd like to consider the symmetries of classical electromagnetism. Generically, classical electromagnetism has gauge redundancy and translational symmetry, with the latter giving rise to the stress-energy-momentum tensor:

$$T^\mu{}_\nu$$

(add stress tensor to this discussion)

2.3 Conductor electrostatics

The equations of motion in electrostatics are

$$\begin{aligned} \nabla \cdot \mathbf{E} &= \frac{\rho}{\epsilon_0} & \nabla \times \mathbf{E} &= \mathbf{0} \\ \nabla^2 \Phi &= -\frac{\rho}{\epsilon_0} & \mathbf{F} &= q\mathbf{E}. \end{aligned}$$

Since we're dealing with statics, we should think of these as equations of constraint, not dynamics. The first EOM tells you exactly how $\mathbf{E}$ must be sourced, the second is a constraint on $\mathbf{E}$'s components, and the last tells you how we relate the field picture to the measurable quantity of force. Poisson's equation is just a way of rewriting Gauss' law by taking the curl-free condition into account.

Remark 2.1 (On $\nabla \times \mathbf{E} = \mathbf{0} \implies \mathbf{E} = -\nabla\Phi$). We often conclude that the curl-free condition gives a *global* scalar potential Φ such that $\mathbf{E} = -\nabla\Phi$. This isn't quite right. While we may always define a local Φ that satisfies such a condition, a global potential function requires the domain we're working on to be simply connected, so $H^1(M) = 0$, where M is our domain, and H^1 is the first de Rham cohomology group (lest things become multivalued). This is intuitively obvious, as the statement $\nabla \times \mathbf{E} = \mathbf{0}$ is really just the statement that $dE = 0$, so if $H^1(M) \neq 0$, then Poincaré's lemma doesn't hold, so E isn't necessarily $-d\Phi$ for some Φ .

Field properties

In addition to the equations of motion, we should naturally consider the symmetries—and thus conserved currents—of electromagnetism. Recall the stress-energy-momentum tensor for electromagnetism:

$$T^\mu{}_\nu = (\text{put here})$$

(need to learn more about this before writing this section)

- read Griffiths ch. 7
- landau...
- tong...

Inhomogeneous solutions

We now treat potential problems. The only way we solve Poisson's equation in practice is through Green's functions².

Idea 2.1 (Green's functions)

Let $\mathcal{L}$ be a linear differential operator, $\psi(x)$ a field (x being a parameter), and $S(x)$ a source term. If we would like to solve

$$(\mathcal{L}\psi)(x) = S(x),$$

then we may consider the following alternative problem:

$$(\mathcal{L}G)(x, x') = \delta(x - x'). \quad (1)$$

We call G a **Green's function**. The reason it's useful is because we may recover ψ by integrating G over our source; instead of solving the (probably intractably difficult) PDE each time for a new source term $S(x)$, we may just input it into our general integral solution.

Claim. $\psi(x) = \int_{\Omega} dx' G(x, x') S(x')$.

Proof. Apply $\mathcal{L}$ to the RHS.

$$\begin{aligned} (\mathcal{L}\psi)(x) &= \mathcal{L} \int_{\Omega} dx' G(x, x') S(x') \\ &= \int_{\Omega} dx' (\mathcal{L}G)(x, x') S(x') \\ &= \int_{\Omega} dx' \delta(x - x') S(x') \\ &= S(x). \end{aligned}$$

□

In words, a Green's function tells you about the response of a linear system to a δ -function source; heuristically, it's a weighting factor for a bunch of δ -function sources that each contribute to your solution ψ . The physical analogy here is δ -functions $\leftrightarrow$ point charges, so G basically just counts a bunch of tiny point charges that all contribute to the "potential", ψ .

Some comments on Green's functions, both in generality and in the context of electrostatics:

- Green's functions are, *a priori*, not unique. This is because we may add homogeneous solutions to G and still satisfy (1). This indeterminacy is solved by implementing boundary conditions on G . We should expect G to be ill-defined before implementing boundary conditions, as we're still solving a PDE, and boundary conditions are necessary to specify unique solutions to PDEs. Before we specify boundary conditions, solving (1) actually solves for an entire equivalence class of Green's functions for the operator $\mathcal{L}$. More concretely, if Λ obeys $(\mathcal{L}\Lambda)(x, x') = 0$, then

$$G(x, x') \sim G(x, x') + \Lambda(x, x'),$$

²See Littlejohn's notes on Green's functions [here](#).

where we define equivalence by $(\mathcal{L}G) = (\mathcal{L}G')$, where $G' = G + \Lambda$.

- The way to think about specifying boundary conditions on G is that you're adding "homogeneous solution" to G until the boundary conditions fit. In the bulk, this homogeneous solution does nothing; only when you get to the boundary does it start helping G become unique. The physical interpretation of Λ in electrostatics is some system of charge extraneous to the domain in question; it then follows that $\nabla^2 \Lambda = 0$, but clearly Λ would affect the boundary conditions on our domain.
- Note that we are free to choose the boundary conditions on G as long as the boundary conditions on Φ are satisfied. Physically, this is just because we only care about what's happening inside of our domain, so adding homogeneous solution (extraneous charges) doesn't change anything that we care about. We will use this freedom in our choice to make things as easy as possible for ourselves. **(think about this)**
- The physical interpretation of boundary conditions on Φ and G in electrostatics is some potential and/or charge distribution on the boundary of our domain. **(?)**

Applying this to electrostatics gives some nice results. We first quickly mention Green's theorem, which you can get from the divergence theorem by considering $\mathbf{A} = \psi \nabla \Phi$.

$$\int_{\Omega} d^3 \mathbf{x}' (\psi \nabla^2 \Phi - \Phi \nabla^2 \psi) = \int_{\partial \Omega} dS' \left(\psi \frac{\partial \Phi}{\partial n'} - \Phi \frac{\partial \psi}{\partial n'} \right).$$

For electrostatics, the Green's function problem is:

$$(\nabla^2 G)(\mathbf{x}, \mathbf{x}') = -4\pi \delta(\mathbf{x} - \mathbf{x}'),$$

Implementing scattering boundary conditions on G (i.e. $G \rightarrow 0$ as $|\mathbf{x}| \rightarrow \infty$) gives the unique solution of

$$G_{FF}(\mathbf{x}, \mathbf{x}') = \frac{1}{|\mathbf{x} - \mathbf{x}'|}.$$

This is called the **free-field Green's function**. You can find this via Fourier transform. Integrating this result gives the potential for a localized charge distribution:

$$\Phi(\mathbf{x}) = \frac{1}{4\pi\epsilon_0} \int_{\Omega} d^3 \mathbf{x}' \frac{\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}. \quad (2)$$

If we have prescribed boundary conditions on our domain (physical interpretation: some potential or charge configuration on conducting surfaces), we may use Green's theorem to get

$$\begin{aligned} \Phi_D(\mathbf{x}) &= \frac{1}{4\pi\epsilon_0} \int_{\Omega} d^3 \mathbf{x}' \rho(\mathbf{x}') G_D(\mathbf{x}, \mathbf{x}') - \frac{1}{4\pi} \oint_{\partial \Omega} dS' \Phi(\mathbf{x}') \frac{\partial G_D}{\partial n'}, \\ \Phi_N(\mathbf{x}) &= \langle \Phi \rangle_{\partial \Omega} + \frac{1}{4\pi\epsilon_0} \int_{\Omega} d^3 \mathbf{x}' \rho(\mathbf{x}') G_N(\mathbf{x}, \mathbf{x}') + \frac{1}{4\pi} \oint_{\partial \Omega} dS' G_N \frac{\partial \Phi}{\partial n'}. \end{aligned}$$

Note the inevitable fact that both of these solutions have the form of "free-field" + "boundary terms".

The above remark tells us about an interesting and useful idea: expanding G in the eigenfunctions of $-\nabla^2$. **(fix all of the below after learning about eigenfunction expansions more broadly)**

This is actually quite useful, as it allows us to construct an *approximate* solution to the oftentimes intractable expression (2). **(think about this)** The eigenfunctions of ∇_{sph}^2 and ∇_{cyl}^2 are respectively spherical harmonics and Bessel functions. Expanding G_{FF} in each coordinate system gives³

$$\begin{aligned}\frac{1}{|\mathbf{x} - \mathbf{x}'|} &\simeq \frac{1}{r} + \mathbf{x} \cdot \frac{\mathbf{x}'}{r^3} + \frac{1}{2} x_a x_b \frac{3x'_a x'_b - (r')^2 \delta_{ab}}{r^5} + \mathcal{O}((r')^3/r^4) \\ \frac{1}{|\mathbf{x} - \mathbf{x}'|} &= \sum_{\ell=0}^{\infty} \frac{4\pi}{2\ell+1} \left(\frac{r_{<}^{\ell}}{r_{>}^{\ell+1}} \right) \sum_{m=-\ell}^{\ell} Y_{\ell m}(\Omega) Y_{\ell m}^*(\Omega') = \sum_{\ell=0}^{\infty} \left(\frac{r_{<}^{\ell}}{r_{>}^{\ell+1}} \right) P_{\ell}(\cos \gamma) \\ \frac{1}{|\mathbf{x} - \mathbf{x}'|} &= \frac{2}{\pi} \sum_{m=-\infty}^{\infty} \int_0^{\infty} dq e^{im(\phi - \phi')} \cos(q(z - z')) I_m(q\rho_{<}) K_m(q\rho_{>})\end{aligned}$$

(fix indices on the first expression) Note that the expansion in spherical harmonics uses the addition theorem in its second equality. **(derive the spherical one using the method of direct integration)**

Idea 2.2 (Addition theorem of spherical harmonics)

Place $\mathbf{x}'$ on the z -axis, so $\theta' = 0$. Then we get that

$$\frac{1}{|\mathbf{x} - \mathbf{x}'|} = \sum_{\ell=0}^{\infty} \left(\frac{r_{<}^{\ell}}{r_{>}^{\ell+1}} \right) P_{\ell}(\cos \theta).$$

Note that the LHS is rotationally invariant, so the RHS must be as well. For this to occur, interpret θ as the angle between $\mathbf{x}$ and $\mathbf{x}'$; call it γ . We have that $\hat{\mathbf{x}} \cdot \hat{\mathbf{x}}' = \cos \gamma$, where $\cos \gamma = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi')$. So we have

$$\frac{1}{|\mathbf{x} - \mathbf{x}'|} = \sum_{\ell=0}^{\infty} \left(\frac{r_{<}^{\ell}}{r_{>}^{\ell+1}} \right) P_{\ell}(\cos \gamma).$$

This result always holds, not just for the times when $\mathbf{x}'$ is along the z -axis. Comparing this to the normal eigenfunction expansion in spherical harmonics gives the **addition theorem for spherical harmonics**:

$$P_{\ell}(\cos \gamma) = \frac{4\pi}{2\ell+1} \sum_{m=-\ell}^{\ell} Y_{\ell m}(\Omega) Y_{\ell m}^*(\Omega').$$

We are physicists, so it's uncommon that we take more than the leading term in any sort of approximation. Thus, it's useful to give names to the first few terms in the more common expansions;

³For the procedure of deriving the second two, see the first LJ lecture [here](#), or page 257 of Zangwill. This is called the **method of direct integration**.

in the Cartesian expansion, we write

$$\mathbf{p} := \int d^3\mathbf{x}' \rho(\mathbf{x}') \mathbf{x}', \quad \mathbb{Q}_{ab} := \int d^3\mathbf{x}' \rho(\mathbf{x}') \left(\frac{3x'_a x'_b - (r')^2 \delta_{ab}}{r'^5} \right).$$

These are respectively called the electric **dipole** and **quadrupole** moments. Whether a potential goes like its dipole or quadrupole moment can oftentimes be determined by inspection: if there is reflection symmetry in a given direction, you know that the dipole moment in that direction is zero. **(wb for quadrupole? give examples)** Inserting the definitions of the dipole and quadrupole moments into the expression (2) gives

$$\Phi(\mathbf{x}) \simeq \frac{1}{4\pi\epsilon_0} \left(\frac{Q}{r} + \frac{\mathbf{x} \cdot \mathbf{p}}{r^3} + \frac{x_a x_b \mathbb{Q}^{ab}}{r^5} + \mathcal{O}((r')^3) \right)$$

We can also define spherical moments: assume that $r' < r$. Then, we have that

$$q_{\ell m} := \int d^3\mathbf{x}' \rho(\mathbf{x}') (r')^\ell Y_{\ell m}^*(\Omega')$$

Inserting this definition into (2) gives

$$\begin{aligned} \Phi(\mathbf{x}) &= \frac{1}{4\pi\epsilon_0} \sum_{\ell=0}^{\infty} \frac{4\pi}{2\ell+1} \sum_{m=-\ell}^{\ell} \frac{q_{\ell m}}{r^{\ell+1}} \\ &\simeq \frac{1}{4\pi\epsilon_0} \left(4\pi \frac{q_{00}}{r} + \frac{4\pi}{3} \frac{q_{1,-1} + q_{10} + q_{11}}{r^2} + \mathcal{O}((r')^3) \right). \end{aligned}$$

(write out the first few spherical moments)

Example 2.1

(do an example or two here; talk about method of images as well)

Method of images

Homogeneous solutions

We now consider **Laplace's equation**, the homogeneous counterpart to Poisson's equation.

Idea 2.3 (Laplace's equation)

Laplace's equation is the homogeneous version of Poisson's equation, i.e.

$$\nabla^2 \Phi = 0.$$

There are two main ways to solve this:

- Separation of variables. The standard way of finding eigenfunctions.
- Fourier transform. More standard in field theory at large (QFT), but still quite useful (more useful for finding Green's functions imo).

It should be noted that both of these methods are just ways of finding eigenfunction expansions

of Φ . We may separate variables in any coordinate system we feel like. The most common are Cartesian, spherical, and cylindrical. The solutions for each are listed below:

- **Cartesian:** the solutions all go like products of exponentials (complex and real). **(finish)**

(need to get a lot more clear on what an eigenfunction expansion is—I really don't know what it is)

Idea 2.4 (Eigenfunction expansion)

(talk about more general eigenfunction expansion here; delete the previous section in the Laplace's equation idea)

2.4 Dielectric electrostatics

Remark 2.2 (The Wilsonian view of dielectrics). This section uses a lot of ideas common in the Wilsonian view of QFT, like effective field theories, coarse-graining, and organizing by scale. It's quite beautiful.

We start by defining what a dielectric is physically⁴. To do this, we first have to consider what a conductor is.

Idea 2.5 (Dielectrics and conductors)

Let a charge distribution have length scale L . Consider the length scale at which the *intrinsic* charges of a body can move. “Intrinsic” just means the charge the body has by virtue of existing, so its electrons and protons.

For **conductors**, the length scale at which charges can move is L . This aligns with our intuition for conductors, as we know that charges can rearrange on them however they like.

For **dielectrics**, the length scale at which charges can move is less than L . That's the distinction between conductors and dielectrics: the length scale of the movement of their charges.

2.5 Magnetostatics

(Purcell derivation???)

Magnetostatics is basically just electrostatics upon the following identifications:

Dot products $\rightarrow$ Cross products
 Charge densities $\rightarrow$ Current densities
 Scalar potential $\rightarrow$ Vector potential

⁴The deep considerations of this section are due to Paul Goldbart's incredible lecturing on the subject of dielectrics. I have expanded on what he said, but the core ideas of an “effective field theory” approach to dielectrics is all him.

Great. The core equations of magnetostatics are

$$\begin{aligned}\nabla \cdot \mathbf{B} &= 0 & \nabla \times \mathbf{B} &= \mu_0 \mathbf{J} \\ \nabla^2 \mathbf{A} &= -\mu_0 \mathbf{J} & \mathbf{F} &= q\mathbf{v} \times \mathbf{B}.\end{aligned}$$

You can see that the source shows up in the “curl sector” of the Maxwell’s equation for the magnetic field instead of the “divergence sector” like it did for the electric field. This fundamental asymmetry produces everything different in magnetostatics, more or less.

The first ME gives a constraint on $\mathbf{B}$, and tells us that we may introduce $\mathbf{A}$, the vector potential. The second ME tells us how the $\mathbf{B}$ field is sourced—notice the qualitative difference between the sourcing of $\mathbf{E}$ and $\mathbf{B}$: $\mathbf{E}$ is related to its source via a divergence, while $\mathbf{B}$ is related to its source via a curl. Combining these two equations and choosing the Coulomb gauge gives our Poisson equation, and the Lorentz force law tells us how the magnetic field couples to charged matter.

Vacuum

We are going to be quick here, because a lot of this is review from electrostatics. Solving Poisson’s equation via Green’s function methods gives

$$G_{FF}(\mathbf{x}, \mathbf{x}') = \frac{1}{|\mathbf{x} - \mathbf{x}'|},$$

so our vector potential is

$$\mathbf{A} = \frac{\mu_0}{4\pi} \int d^3\mathbf{x}' \frac{\mathbf{J}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}.$$

The approximation schemes for the vector potential are significantly harder, as we’re now dealing with a vector instead of a scalar. **(derive the results for the dirichlet and neumann bc on vector potential)**

(in approx schemes, talk about multi-pole expansion etc)

(talk about vector spherical harmonics here)

2.6 Electrodynamics

So far we’ve assumed that our fields are not time-dependent. This is not a realistic assumption for most purposes, so now we introduce time-dependence. The good thing is that our previous techniques mostly work with some modifications.

2.7 Retarded potentials

We now solve for the Green’s function **(finish, ts is trivial)**

2.8 Waves

3 Statistical Mechanics

The fundamental idea behind statistical mechanics is that it is mechanics, treated statistically. Statistical mechanics aims to make predictions about the bulk properties of systems with a large number of degrees of freedom. “Large” may generally be defined as when the number of particles is on the order of Avogadro’s number, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$, where the mol^{-1} in the expression for N_A comes from the fact that we want to write things like $N = nN_A$.

Remark 3.1 (Avogadro’s number). Historically, Avogadro’s number was defined as the number of atoms in 12 g of carbon-12. Here is a less stupid derivation of it: just comparing length scales.

An atom is on the order of 10^{-10} m , and a centimeter 10^{-2} m . Considering the number of atoms in 1 cm^3 gives

$$\# \text{ atoms} \sim \frac{10^{-6}}{10^{-30}} = 10^{24} \text{ atoms}$$

which is about the same OOM as Avogadro’s number. Another derivation comes through considering that the scale of atomic mass is 10^{-26} kg , while the gram is 10^{-3} kg , and thus there are roughly 10^{23} atoms per gram of mass.

While we may define thermodynamic quantities when the number of particles is on the order of Avogadro’s number, many results in thermodynamics only hold in the **thermodynamic limit**; this is when you send both the particle number and volume of the system to infinity, but keep the density constant. This limit is actually quite deep—its effects will become most noticable when we talk about phase transitions later in these notes, as we need this limit to make sense of even basic ideas in the subject.

3.1 The microscopic picture

The fundamental object in statistical mechanics is the *density operator*, ρ . We recall its definition here. If there is a probability p_i that a system is in state $|\psi_i\rangle$, the density operator for that system is given by

$$\rho = \sum_i p_i |\psi_i\rangle \langle \psi_i|.$$

For a system in a pure state $|\psi\rangle$, clearly $\rho = |\psi\rangle \langle \psi|$. It can be shown that a density operator generically satisfies $\text{tr} \rho = 1$ and $\rho \geq 0$; this makes sense, as we’d like to think about ρ as a probability distribution of sorts. While ρ is the fundamental object of statistical mechanics—as the world is fundamentally quantum mechanical—treating the microscopic dynamics as *classical* is oftentimes an almost perfect approximation to the quantum mechanical theory. This is, of course, due to the renormalization group—different microscopic theories oftentimes give way to the same effective theory. Thus, we are also interested the classical Hamiltonian of our system, H .

The Hamiltonian of a system is a smooth function on phase space that gives the dynamics of the system when differentiated. Concretely, if we have a system with N degrees of freedom, the phase space is generically $2N$ -dimensional, and the Hamiltonian is a function $H = H(\mathbf{p}, \mathbf{q})$ on this

space. The state of the system evolves due to *Hamilton's equations*,

$$\dot{\mathbf{q}} = \frac{\partial H}{\partial \mathbf{p}}, \quad \dot{\mathbf{p}} = -\frac{\partial H}{\partial \mathbf{q}}.$$

(talk about symplectic two form here, as well as Liouville's theorem)

Along with these fundamental objects, we must make a fundamental assumption called the **ergodic hypothesis**.

Idea 3.1 (The Ergodic Hypothesis)

The ergodic hypothesis states that over long times, a system explores all possible accessible microstates at an energy E . Thus, time averages equal ensemble averages. (fix this “definition” after reading the paper)

Remark 3.2 (Is the ergodic hypothesis true?). Yes and no. For most systems, yes, the ergodic hypothesis seems to hold just fine. But there are classes of systems where it does not hold. We can further delineate between these two classes by observing *how* they break ergodicity. There are those systems whose time scales for exploration are just incredibly long (e.g. longer than the time to the end of the universe), and thus practically break the hypothesis: this is **weak breaking**. Then there are those systems who will *never* explore all of the states available to them, regardless of the time they have to access the states: this is **strong breaking**.

Examples of weak breaking may be found in spin glasses and generic finite systems, where the latter case is an example of the Poincare recurrence theorem. To recall from the notes on classical mechanics, the Poincare recurrence theorem roughly states that chaotic systems return arbitrarily close to their initial state. This practically restricts the volume of phase space that the system explores, thus breaking ergodicity. Spin glasses are just complicated systems with lots of potential barriers, thus making it really difficult to explore microstates; the time scale for exploration becomes absurdly long as the system gets larger, thus weakly breaking ergodicity.

The examples of strong breaking are much more interesting and deep. The quintessential example of strong breaking is an *integrable* system. That is, a Hamiltonian system with as many conserved quantities as degrees of freedom. (expand)

Example 3.1

(give concrete example of where it fails)

We now talk about the three different “ensembles” of thermodynamics. We start with the microcanonical ensemble. A priori, the distinguishing factor delineating the three ensembles from each other is the variables that are assumed to be held constant for that ensemble. In the microcanonical ensemble, energy, volume, and particle number are held constant.

(at some point talk about special relativistic thermodynamics)

4 Quantum Mechanics

4.1 Assumptions

(insert intermediate stuff)

4.2 Perturbation theory

Hamiltonian

We first cover bound-state perturbation theory. Only studying bound states simplifies things, as continuous spectra require more complicated techniques (i.e. TD-perturbation theory). We follow the Brillouin-Wigner (BW) derivation for bound-state perturbation theory, showing equivalence to Rayleigh-Schrodinger perturbation theory (RSPT) along the way (as the latter is generally more immediately useful).




Figure 1: Algebraic and geometric pictures associated with perturbation theory.

It is often true that useful physical systems take the form of a known system plus a small perturbation; indeed, every prediction we see from colliders like the LHC tacitly assume this statement. So we should have a formalism to deal with these kinds of problems; this is the study of **perturbation theory**. We cover the quantum mechanical version here⁵.

Consider a Hamiltonian H , and assume that it may be written as

$$H = H_0 + \lambda H_1 = H_0 + \tilde{H}_1.$$

where $\lambda \in [0, 1]$ interpolates between the free system (H_0) and the full system (H). Another way of looking at this is brought to us by Wilsonian QFT; we may think of λH_1 as being an operator $\tilde{H}_1 = g\mathcal{O}$, where $\mathcal{O}$ is composed of a 's and $a^\dagger$'s. Then λ is defined through $g = \Lambda^{d-\Delta_{\mathcal{O}}} \lambda$, with $\Delta_{\mathcal{O}} := [\mathcal{O}]$ being the mass dimension of $\mathcal{O}$, and we may proceed as usual. As a concrete example, consider the anharmonic oscillator—in this case, $H_0 = p^2/2m + (1/2)m\omega^2 q^2$ and $\tilde{H}_1 = gq^4/4!$. We have that $\lambda = L^{d-\Delta_{\mathcal{O}}} g$, where L is the characteristic length scale of our system, and d the dimension.

We now study the geometry of this problem. Introduce

$$\mathcal{H}_n = \{\text{unperturbed eigenspace}\} \quad \mathcal{H}_n^\perp = \{\text{perturbed eigenspace}\}.$$

These spaces are orthogonal, as they're eigenspaces of different eigenvalues. Label the kets in the spaces respectively by $\{|n\alpha\rangle\}$ and $\{|k\alpha\rangle\}$, where α is a label used to resolve degeneracies. These

⁵This idea is also found in classical mechanics and quantum field theory, and more generally in analysis.

states are orthonormal, $\langle n\alpha|k\beta\rangle = \delta_n^k \delta_\alpha^\beta$. Note that kets correspond to upper indices and bras to lower indices. Introduce commuting projectors onto the eigenspaces

$$P := \sum_{\alpha} |n\alpha\rangle\langle n\alpha|, \quad Q := \sum_{\substack{k \neq n \\ \alpha}} |k\alpha\rangle\langle k\alpha|.$$

Then $|\psi\rangle = P|\psi\rangle + Q|\psi\rangle$. $P|\psi\rangle$ is known by assumption, and $Q|\psi\rangle$ is not. This is the goal of perturbation theory: solve for the small hard part $Q|\psi\rangle$ in terms of big easy part $P|\psi\rangle$. The full Hamiltonian is assumed to satisfy

$$H|\psi\rangle = (H_0 + \lambda H_1)|\psi\rangle = E|\psi\rangle, \quad (3)$$

where E and $|\psi\rangle$ are assumed to be λ -dependent. As $\lambda \rightarrow 0$, $E \rightarrow \epsilon_n$ and $|\psi\rangle \rightarrow |n\alpha\rangle \in \mathcal{H}_n$. This second statement shows the difficulty of degenerate perturbation theory; since $\mathcal{H}_n$ is degenerate, the “tracing back” of $|\psi\rangle$ as λ turns off is not unique, and this is also true in reverse—as we turn on λ , $|\psi\rangle$ generically splits into one of many energy eigenstates, the number of which determined by $\dim \mathcal{H}_n$.

As stated before, we may write $|\psi\rangle = P|\psi\rangle + Q|\psi\rangle$, where the perturbation is entirely found in $Q|\psi\rangle$. Rewriting eq. (3) gives

$$(E - H_0)|\psi\rangle = \lambda H_1|\psi\rangle. \quad (4)$$

So, we’d like to invert $E - H_0$. Write $E - H_0$ as a function of the energy eigenstates,

$$(E - H_0)^{-1} = \sum_{k, \alpha} \frac{|k\alpha\rangle\langle k\alpha|}{E - \epsilon_k}.$$

This expression has a zero eigenvalue at $E = \epsilon_n$, and so it is meaningless as it stands. To resolve this, introduce

$$R = \sum_{\substack{k \neq n \\ \alpha}} \frac{|k\alpha\rangle\langle k\alpha|}{E - \epsilon_k}.$$

R satisfies $RQ = QR = R$, and $[R, P] = 0$. We also note that $R(E - H_0) = (E - H_0)R = Q$, and so $R|_{\mathcal{H}_n^\perp} = \text{id}_{\mathcal{H}_n^\perp}$, and $Q|\psi\rangle = \lambda R H_1 |\psi\rangle$ (which we can see by multiplying (4) through by R). So,

$$|\psi\rangle = P|\psi\rangle + Q|\psi\rangle = P|\psi\rangle + \lambda R H_1 |\psi\rangle.$$

Solving for $|\psi\rangle$ and expanding gives

$$(1 - \lambda R H_1)|\psi\rangle = P|\psi\rangle \implies |\psi\rangle = \sum_{n=0}^{\infty} \lambda^n (R H_1)^n P|\psi\rangle. \quad (5)$$

So we’ve accomplished our goal of writing $|\psi\rangle$ in terms of the easy part $P|\psi\rangle$. We now explore the consequences of this equation with a few general examples.

Example 4.1 (Non-degenerate perturbation theory)

Assume $\dim \mathcal{H}_n = 1$ —call this state $|n\rangle$. Note that $\langle\psi|n\rangle = 1$ by assumption of $\langle n|n\rangle = 1$. P takes an especially simple form in this case, $P = |n\rangle\langle n|$. Referencing eq. (5), we have

$$\begin{aligned} |\psi\rangle &= |n\rangle + \lambda R H_1 |n\rangle + \lambda^2 (R H_1)^2 |n\rangle + \cdots \\ &= |n\rangle + \sum_{\substack{k \neq n \\ \alpha}} |k\alpha\rangle \frac{\langle k\alpha | H_1 | n \rangle}{E - \epsilon_k} + \cdots \end{aligned}$$

The energies are given by bra-ing through eq. (4) by $\langle n|$,

$$\langle n | E - H_0 | \psi \rangle = E - \epsilon_n = \lambda \langle n | H_1 | \psi \rangle,$$

so

$$E = \epsilon_n + \lambda \langle n | H_1 | n \rangle + \lambda^2 \langle n | H_1 R H_1 | n \rangle + \cdots$$

Substituting the zeroth-order approximation for E ($E = \epsilon_n$) gives

$$|\psi\rangle = |n\rangle + \lambda \sum_{\substack{k \neq n \\ \alpha}} |k\alpha\rangle \frac{\langle k\alpha | H_1 | n \rangle}{\epsilon_n - \epsilon_k} + \cdots$$

which, along with the equation for $|\psi\rangle$, agrees with RSPT to first order. We remark upon a general fact here: the λ -expansion for E is always “one-behind” the expansion for $|\psi\rangle$. Concretely, if we know E to order λ^n , then we know $|\psi\rangle$ to order λ^{n+1} . This is because E only shows up in $|\psi\rangle$ *starting at* order λ , so $|\psi\rangle$ is always “one-ahead” of $E(\lambda)$.

In RSPT, we simply identify things slightly differently. $|n\rangle \rightarrow |n_0\rangle$, and $\lambda^k (R H_1)^k |n\rangle \rightarrow |n_k\rangle$. Each of the states $\{n_k\}$ is assumed to be orthogonal to $|n_0\rangle$, which follows from the fact that R projects out all $k = n$ states, or from using eq. (4) and noting that $\langle P\psi | Q\psi \rangle = 0$. Everything else follows from this.

Example 4.2 (Degenerate perturbation theory)

We assume that the unperturbed eigenspace is degenerate, and thus $P = \sum_{\alpha} |n\alpha\rangle\langle n\alpha|$. P acting on $|\psi\rangle$ gives some combination of the $|n\alpha\rangle$ ’s, $P|\psi\rangle = \sum_{\alpha} c_{\alpha} |n\alpha\rangle$. The main difference between non-degenerate and degenerate perturbation theory is that we must solve for the state (c_{α}) and the energy E simultaneously. Subbing $|\psi\rangle$ into the energy equation we previously found gives

$$\begin{aligned} \text{State:} \quad & |\psi\rangle = c_{\alpha} |n\alpha\rangle + \lambda R H_1 (c_{\alpha} |n\alpha\rangle) + \cdots \\ \text{Energies:} \quad & (E - \epsilon_n) c_{\alpha} = \left\{ \lambda |n\alpha\rangle H_1 |n\beta\rangle + \lambda^2 \sum_{\substack{k \neq n \\ \gamma}} \frac{\langle n\alpha | H_1 | k\gamma \rangle \langle k\gamma | H_1 | n\beta \rangle}{E - \epsilon_k} + \cdots \right\} c_{\beta}. \end{aligned}$$

From the second equation we can see what the c_{α} ’s are: eigenvectors with respect to the $g \times g$

matrix in the braces (with $g = \dim \mathcal{H}_n$). Their eigenvalues are $(E - \epsilon_n)$. In practice, we truncate the energy series at $\mathcal{O}(\lambda)$.

(show equivalence to RSPT)

Example 4.3 (Nearly degenerate perturbation theory)

Lagrangian