

# 1 Three stochastic epidemic models

We consider three discrete-time, branching-process models for the spread of an infectious disease. Time is measured in days. Let  $I_t \geq 0$  denote the number of new infections (incidence) on day  $t$ , and let  $R_t > 0$  be the effective reproduction number on day  $t$ .

The *serial interval weights*  $w_s > 0$  ( $s = 1, 2, \dots$ ) satisfy  $\sum_{s=1}^{\infty} w_s = 1$  and encode the probability that a secondary infection appears  $s$  days after the primary case. Write

$$F_r = \sum_{s=1}^r w_s$$

for the cumulative serial interval distribution at lag  $r$ , so  $1 - F_r = \sum_{s=r+1}^{\infty} w_s$  is the probability that transmission occurs after day  $r$ .

The first two models share a *dispersion parameter*  $k > 0$  controlling individual-level variation in infectiousness. Small  $k$  corresponds to high heterogeneity (superspreading); as  $k \rightarrow \infty$  both models collapse onto a common Poisson offspring distribution with no dispersion parameter — our third model.

**Remark.** Throughout,  $\text{NB}(\text{mean} = \mu, \text{disp} = k)$  denotes the negative-binomial distribution with probability mass function

$$\mathbb{P}(X = n) = \binom{n+k-1}{n} \left( \frac{k}{k+\mu} \right)^k \left( \frac{\mu}{k+\mu} \right)^n, \quad n = 0, 1, 2, \dots,$$

mean  $\mu$ , variance  $\mu + \mu^2/k$ , and probability generating function (pgf)  $G(s) = (k / (k + \mu(1-s)))^k$ . Similarly,  $\text{Gamma}(\alpha, \beta)$  denotes the Gamma distribution with shape  $\alpha$  and rate  $\beta$ , so density  $f(x) = \beta^\alpha x^{\alpha-1} e^{-\beta x} / \Gamma(\alpha)$ , mean  $\alpha/\beta$ , and variance  $\alpha/\beta^2$ .

## 1.1 SSE model (Superspreading Events)

In the *superspreading-events* (SSE) model, the force of infection on day  $t$  is the weighted sum of past incidence,

$$\lambda_t = \sum_{s=1}^{\infty} w_s I_{t-s}, \tag{1}$$

and new infections are negative-binomially distributed:

$$I_t \mid \lambda_t \sim \text{NB}(\text{mean} = R_t \lambda_t, \text{disp} = k \lambda_t). \tag{2}$$

The conditional mean of  $I_t$  is  $R_t \lambda_t$  and its variance is  $R_t \lambda_t + R_t^2 \lambda_t / k$ , so the NB inflates the Poisson variance by the factor  $1 + R_t \lambda_t / k$ , capturing superspreading at the population level.

## 1.2 SSI model (Superspreading Individuals)

In the *superspreading individuals* (SSI) model, heterogeneity enters at the individual level. Each infected individual has a latent infectivity that is Gamma distributed with mean 1 and variance  $1/k$ .

Each cohort of  $I_t$  infected individuals then has an aggregate infectivity

$$Y_t \mid I_t \sim \text{Gamma}(\text{shape} = k I_t, \text{rate} = k), \tag{3}$$

so  $\mathbb{E}[Y_t \mid I_t] = I_t$  and  $\text{Var}(Y_t \mid I_t) = I_t / k$ . New infections arise as a Poisson process driven by past infectivity,

$$I_t \mid (Y_s)_{s < t} \sim \text{Poisson} \left( R_t \sum_{s=1}^{\infty} w_s Y_{t-s} \right). \tag{4}$$

**Remark.** Marginalising  $Y_t$  from the SSI model recovers a negative-binomial offspring distribution (by the Poisson–Gamma mixture identity; see below), so the two models share the same mean–variance structure at the population level while differing in the *mechanism* generating overdispersion.

### 1.3 Poisson offspring model

In the *Poisson* offspring model the force of infection on day  $t$  is the same weighted sum as in the SSE model,

$$\lambda_t = \sum_{s=1}^{\infty} w_s I_{t-s}, \quad (5)$$

and new infections are Poisson distributed:

$$I_t \mid \lambda_t \sim \text{Poisson}(R_t \lambda_t). \quad (6)$$

The conditional mean and variance of  $I_t$  are both  $R_t \lambda_t$ : the Poisson is the limit of the SSE-style overdispersion vanishing ( $1 + R_t \lambda_t / k \rightarrow 1$  as  $k \rightarrow \infty$ ).

**Remark.** The Poisson model is the common  $k \rightarrow \infty$  limit of both the SSE and the SSI models. In the SSE step the negative-binomial offspring distribution  $\text{NB}(R_t \lambda_t, k \lambda_t)$  collapses to  $\text{Poisson}(R_t \lambda_t)$  as  $k \rightarrow \infty$ ; in the SSI step the Gamma noise on the individual infectivities degenerates to  $Y_t = I_t$  a.s., again leaving  $\text{Poisson}(R_t \lambda_t)$  for the increment. There is no dispersion parameter; the only model parameter is  $R_t$ .

## 2 Extinction probability

Henceforth assume the reproduction number is constant,  $R_t = R_0$ . We study the probability that the epidemic goes extinct, starting from a single infectious case on day 0.

By branching-process theory, the extinction probability  $q$  is the smallest non-negative fixed point of the pgf of the offspring distribution. For the SSE and SSI models the offspring distribution is  $\text{NB}(\text{mean} = R_0, \text{disp} = k)$ , whose pgf is  $G(s) = (k/(k + R_0(1 - s)))^k$ ; setting  $q = G(q)$  gives

$$q = \left(1 + \frac{R_0}{k}(1 - q)\right)^{-k}. \quad (7)$$

For the Poisson model the offspring distribution is  $\text{Poisson}(R_0)$ , whose pgf is  $G(s) = e^{R_0(s-1)}$ ; setting  $q = G(q)$  gives

$$q = e^{R_0(q-1)}. \quad (8)$$

In all three models, if  $R_0 \leq 1$  the unique solution is  $q = 1$  (extinction is certain); if  $R_0 > 1$  there is a unique solution  $q \in (0, 1)$ , so the probability of a major outbreak is  $1 - q > 0$ .

### 2.1 Extinction probability after observing incidence on days 0 to $r$

Suppose we observe incidence  $I_0, I_1, \dots, I_r$  on the first  $r + 1$  days, with  $I_0 \geq 1$ . We compute the updated extinction probability  $q_r$  for the residual epidemic — the probability that the chain triggered by the observed cases eventually dies out.

A standard branching-process argument applies in all three models: each remaining offspring of the observed cases, once born, initiates an independent fresh epidemic governed by the original offspring distribution ( $\text{NB}(R_0, k)$  for SSE/SSI,  $\text{Poisson}(R_0)$  for Poisson) and goes extinct with the standard probability  $q$  (the smallest fixed point of  $G$ ). The conditioning event constrains only the offspring of the observed cases on days  $0, \dots, r$ ; it does not propagate to later generations. Therefore

$$q_r = G_r(q),$$

where  $G_r$  is the pgf of the (conditional) total remaining offspring of the observed cases. The three models differ in the form of  $G_r$ .

### 2.1.1 SSE model

In the SSE model the offspring process decomposes into independent contributions: each of the  $I_s$  individuals infected on day  $s$  generates a number of offspring at lag  $\ell$  that is  $\text{NB}(\text{mean} = R_0 w_\ell, \text{disp} = k w_\ell)$ , independently across individuals, donor days, and lags. By the NB closure property under a common success probability  $p = k w_\ell / (k w_\ell + R_0 w_\ell) = k / (k + R_0)$  (independent of  $\ell$ ), the population-level  $\text{NB}(R_0 \lambda_t, k \lambda_t)$  law of  $I_t$  is recovered by summing all per-individual-per-lag contributions.

Conditioning on the observed counts  $I_0, \dots, I_r$  pins down only those contributions  $(s, \ell)$  with  $s + \ell \leq r$ . Per-individual-per-lag contributions with  $s \in \{0, \dots, r\}$  and  $\ell > r - s$  are independent of the observations and retain their original distributions. The total remaining offspring of the observed cases is therefore

$$\sum_{s=0}^r \sum_{i=1}^{I_s} \sum_{\ell=r-s+1}^{\infty} \text{NB}(\text{mean} = R_0 w_\ell, \text{disp} = k w_\ell) = \text{NB}(\text{mean} = R_0 \Lambda, \text{disp} = k \Lambda), \quad (9)$$

where the pooled effective weight is

$$\Lambda = \sum_{s=0}^r I_s (1 - F_{r-s}). \quad (10)$$

With the convention  $F_0 = 0$ , the  $s = r$  term enters with full weight, since none of its offspring have had a chance to appear yet.

The pgf is  $G_r(s) = (1 + (R_0/k)(1-s))^{-k\Lambda} = G(s)^\Lambda$ , so

$$q_r = G_r(q) = q^\Lambda = q^{\sum_{s=0}^r I_s (1 - F_{r-s})}. \quad (11)$$

The earlier special case (a single case on day 0 followed by  $r$  days without cases) corresponds to  $I_0 = 1$ ,  $I_1 = \dots = I_r = 0$ , giving  $\Lambda = 1 - F_r$  and recovering  $q_r = q^{1 - F_r}$ . In the same direction, as the zero streak extends ( $r \rightarrow \infty$ ) the contributions  $I_s (1 - F_{r-s})$  all vanish and  $q_r \rightarrow 1$ : extinction approaches certainty.

### 2.1.2 SSI model

For the SSI model the joint posterior of  $(Y_0, \dots, Y_{r-1})$  is in general not available in closed form: as soon as  $I_d > 0$  on at least one day  $d \in \{1, \dots, r\}$ , the day- $d$  rate  $R_0 \sum_{s=1}^{d-1} w_{d-s} Y_s$  couples the latents. We therefore present an MCMC estimator of  $q_r$  here as the canonical SSI workflow. For the special cases of cases on day 0 only, on day 0 and one later day, and on day 0 and two later days, closed forms are however available; these are deferred to § 4 below.

**MCMC posterior.** We sample  $(Y_0, \dots, Y_{r-1})$  from their joint posterior given  $(I_0, \dots, I_r)$  using a Hamiltonian Monte Carlo sampler (implemented in PyMC). Note that  $Y_r$ , the aggregate infectivity of the day- $r$  cohort, has not yet contributed to any observed outcome; its posterior equals its prior,

$$Y_r \sim \text{Gamma}(k I_r, k),$$

and is sampled independently.

**Remaining offspring from observed cases.** Given  $(Y_s)_{s=0}^r$ , the future infections generated by the observed cases arrive as independent Poisson streams: cohort  $s$  contributes at lag  $\ell \geq r - s + 1$  at rate  $R_0 w_\ell Y_s$ . Summing over cohorts and lags gives total remaining offspring

$$I_{\text{rem}} \mid (Y_s)_{s=0}^r \sim \text{Poisson}(R_0 \Lambda), \quad \Lambda = \sum_{s=0}^r Y_s (1 - F_{r-s}), \quad (12)$$

exactly the same pooled-weight formula as in the SSE model.

**Extinction probability and MCMC estimator.** Each remaining offspring initiates an independent fresh epidemic that goes extinct with the standard probability  $q$ . Hence the conditional extinction probability of the residual epidemic, given the latent infectivities, is the pgf of  $\text{Poisson}(R_0\Lambda)$  evaluated at  $q$ :

$$q_r \mid (Y_s)_{s=0}^r = e^{R_0\Lambda(q-1)}. \quad (13)$$

Marginalising over the posterior of  $(Y_s)$  and estimating by a Monte Carlo average over  $M$  MCMC draws,

$$\hat{q}_r = \frac{1}{M} \sum_{m=1}^M e^{R_0\Lambda^{(m)}(q-1)}, \quad \widehat{\text{PMO}} = 1 - \hat{q}_r, \quad (14)$$

where  $\Lambda^{(m)} = \sum_{s=0}^r Y_s^{(m)}(1 - F_{r-s})$ .

**Remark** (Consistency with the day-0-only analytic formula). When  $I_1 = \dots = I_r = 0$  the observations force  $Y_1 = \dots = Y_r = 0$ , so  $\Lambda = Y_0(1 - F_r)$  and the posterior is  $Y_0 \sim \text{Gamma}(kI_0, k + R_0F_r)$ . Taking the expectation of  $e^{R_0Y_0(1-F_r)(q-1)}$  with respect to this posterior via the Gamma moment-generating function gives

$$q_r = \left(1 - \frac{R_0(1 - F_r)(q - 1)}{k + R_0F_r}\right)^{-kI_0} = \left(1 + \frac{R_0(1 - F_r)}{k + R_0F_r}(1 - q)\right)^{-kI_0},$$

which matches the closed-form result (29).

### 2.1.3 Poisson model

The argument of § 2.1.1 carries over verbatim with NB replaced by Poisson. Each of the  $I_s$  individuals infected on day  $s$  generates a  $\text{Poisson}(R_0w_\ell)$  number of offspring at lag  $\ell$ , independently across individuals, donor days, and lags. By Poisson closure under summation the total remaining offspring of the observed cases is

$$\sum_{s=0}^r \sum_{i=1}^{I_s} \sum_{\ell=r-s+1}^{\infty} \text{Poisson}(R_0w_\ell) = \text{Poisson}(R_0\Lambda), \quad (15)$$

with the same pooled effective weight  $\Lambda = \sum_{s=0}^r I_s(1 - F_{r-s})$  as in the SSE model. The pgf is  $G_r(s) = e^{R_0\Lambda(s-1)} = G(s)^\Lambda$ , so

$$q_r = G_r(q) = q^\Lambda, \quad (16)$$

identical in functional form to (11) but with  $q$  the Poisson extinction root (8). In particular, both (11) and (16) are closed-form for any observed history.

## 3 Probability of major outbreak under model uncertainty

So far we have computed the probability of a major outbreak conditional on a chosen model (SSE, SSI, or Poisson). In practice the underlying mechanism generating overdispersion is unknown, and we may wish to combine the candidate models via Bayesian model averaging. Let  $\mathcal{M}$  denote a finite set of candidate models with prior probabilities  $\{\pi_M\}_{M \in \mathcal{M}}$  satisfying  $\sum_{M \in \mathcal{M}} \pi_M = 1$ . Two natural choices are the two-model set  $\mathcal{M} = \{\text{SSE}, \text{SSI}\}$  (a single  $k$ , varying only the noise mechanism) and a wider ensemble including several SSE and SSI specifications with different  $k$  values together with the Poisson limit. Given the observed history  $(I_0, \dots, I_r)$ , Bayes' theorem yields posterior model probabilities

$$\pi_M^* = \frac{\pi_M L_M}{\sum_{M' \in \mathcal{M}} \pi_{M'} L_{M'}}, \quad L_M = \mathbb{P}_M(I_1, \dots, I_r \mid I_0), \quad (17)$$

and the model-averaged probability of a major outbreak is the posterior expectation of the per-model PMOs,

$$\text{PMO}_{\text{avg}} = \sum_{M \in \mathcal{M}} \pi_M^* \text{PMO}_M, \quad (18)$$

where  $\text{PMO}_M = 1 - q_r^M$  as derived above. Specialising to the two-model case  $\mathcal{M} = \{\text{SSE}, \text{SSI}\}$ , (17) and (18) become the familiar pair

$$\pi_{\text{SSE}}^* = \frac{\pi_{\text{SSE}} L_{\text{SSE}}}{\pi_{\text{SSE}} L_{\text{SSE}} + \pi_{\text{SSI}} L_{\text{SSI}}}, \quad \text{PMO}_{\text{avg}} = \pi_{\text{SSE}}^* \text{PMO}_{\text{SSE}} + \pi_{\text{SSI}}^* \text{PMO}_{\text{SSI}}. \quad (19)$$

### 3.1 Likelihood of the observed history

**SSE model.** Conditional on past incidence, the day- $t$  count satisfies  $I_t \mid I_{<t} \sim \text{NB}(\text{mean} = R_0 \lambda_t, \text{disp} = k \lambda_t)$  with  $\lambda_t = \sum_{s=1}^{t-1} w_{t-s} I_s$ , so the joint likelihood factorises into a product of negative-binomial pmfs,

$$L_{\text{SSE}}(I_1, \dots, I_r \mid I_0) = \prod_{t=1}^r \text{NB}(I_t; R_0 \lambda_t, k \lambda_t). \quad (20)$$

This closed form is available for any history.

**Poisson model.** Likewise, the day- $t$  count satisfies  $I_t \mid I_{<t} \sim \text{Poisson}(R_0 \lambda_t)$ , so the joint likelihood factorises into a product of Poisson pmfs,

$$L_{\text{Poisson}}(I_1, \dots, I_r \mid I_0) = \prod_{t=1}^r \text{Poisson}(I_t; R_0 \lambda_t), \quad (21)$$

closed-form for any history. As with (20), the convention is  $\log L = -\infty$  whenever  $\lambda_t = 0$  but  $I_t > 0$ .

**SSI model, closed-form cases.** Closed forms are available in the same three special cases as for  $q_r$  (cases on day 0 only, on day 0 and one later day, on day 0 and two later days); these are derived in § 4 below.

**SSI model, general histories.** The likelihood involves the latent infectivities  $(Y_s)_{s=0}^{r-1}$  and is in general not available in closed form. Unfolding the SSI generative dependence  $I_0 \rightarrow Y_0 \rightarrow I_1 \rightarrow Y_1 \rightarrow \dots \rightarrow I_r$  as a Markov chain, the joint density of latents and observations factorises as

$$p(Y_{0:r-1}, I_{1:r} \mid I_0) = \prod_{t=0}^{r-1} p_Y(Y_t \mid I_t) \cdot \prod_{t=1}^r p_I(I_t \mid Y_{0:t-1}), \quad (22)$$

where  $p_Y(\cdot \mid I_t) = \text{Gamma}(\cdot; k I_t, k)$  is the SSI generative density of  $Y_t$  given the cohort size on day  $t$ , and  $p_I(\cdot \mid Y_{0:t-1}) = \text{Poisson}(\cdot; R_0 \sum_{s=1}^t w_s Y_{t-s})$ . Marginalising the latents,

$$L_{\text{SSI}}(I_{1:r} \mid I_0) = \int \prod_{t=0}^{r-1} p_Y(Y_t \mid I_t) \prod_{t=1}^r p_I(I_t \mid Y_{0:t-1}) dY_{0:r-1}. \quad (23)$$

Days  $s$  with  $I_s = 0$  collapse  $Y_s$  to a point mass at 0, and  $Y_r$  does not enter  $L_{\text{SSI}}$  (it would only contribute to  $I_{r+1}, I_{r+2}, \dots$ ), so the integration runs effectively over  $\mathcal{T} = \{t \in \{0, \dots, r-1\} : I_t > 0\}$ . Three MCMC-based estimators of (23) are derived in § 5.

### 3.2 Simulation-based estimator

For general histories, where the SSI likelihood is unavailable in closed form, we use a rejection-sampling estimator that simultaneously realises the posterior model weighting. We describe it for  $\mathcal{M} = \{\text{SSE}, \text{SSI}\}$ ; it extends to any finite  $\mathcal{M}$  by replacing the Binomial split below with a Multinomial split by prior, and the acceptance-rate argument carries over unchanged. Repeat the following in batches of size  $b$ :

1. Draw the per-batch model split as a Binomial:  $b_{\text{SSE}} \sim \text{Bin}(b, \pi_{\text{SSE}})$  and  $b_{\text{SSI}} = b - b_{\text{SSE}}$ .
2. Run  $b_{\text{SSE}}$  vectorised SSE forward simulations from  $t = 0$  seeded with  $I_0$  index cases, retaining only those whose simulated incidence on days  $1, \dots, r$  exactly matches the observed  $I_1, \dots, I_r$ . Likewise for  $b_{\text{SSI}}$  SSI simulations.
3. For each retained simulation, continue forward until either the single-step incidence reaches the major-outbreak threshold (resolved as “major”) or the most recent  $L = \text{len}(w)$  steps are all zero (resolved as “extinct”).

Continue accumulating batches until a target number of resolved-and-matching simulations is reached. The acceptance rate of model  $M$ 's simulations is  $L_M$  in expectation; with a Binomial split by prior, the expected number of accepted  $M$ -sims is proportional to  $\pi_M L_M$ , so the composition of the accepted pool matches the posterior  $\pi_M^*$ . The pooled estimator

$$\widehat{\text{PMO}}_{\text{avg}} = \frac{N_{\text{SSE}}^{\text{maj}} + N_{\text{SSI}}^{\text{maj}}}{N_{\text{SSE}}^{\text{acc}} + N_{\text{SSI}}^{\text{acc}}} = \hat{\pi}_{\text{SSE}}^* \widehat{\text{PMO}}_{\text{SSE}} + \hat{\pi}_{\text{SSI}}^* \widehat{\text{PMO}}_{\text{SSI}} \quad (24)$$

follows immediately, where  $\hat{\pi}_M^* = N_M^{\text{acc}} / (N_{\text{SSE}}^{\text{acc}} + N_{\text{SSI}}^{\text{acc}})$  is the implied posterior estimate and  $\widehat{\text{PMO}}_M = N_M^{\text{maj}} / N_M^{\text{acc}}$  is the per-model PMO from the accepted  $M$ -sims.

## 4 SSI model: closed forms for histories with cases on a few days

For the SSI model, when non-zero incidence is observed on day 0 and on a small number of later days, the joint posterior of the latent infectivities still factorises (or factorises as a finite mixture) and both the model evidence  $L_{\text{SSI}}$  and the residual extinction probability  $q_r$  admit closed forms. This section treats three such cases in turn: case (i) (§ 4.1), incidence on day 0 only; case (ii) (§ 4.2), incidence on day 0 and one later day; and case (iii) (§ 4.3), incidence on day 0 and two later days.

**Strategy.** Each derivation follows the same five steps:

1. *Conditional likelihood given the latents.* Multiply the per-day Poisson pmfs to obtain  $\mathbb{P}(\text{obs} \mid Y_0, Y_{i_1}, \dots, Y_{i_n})$ .
2. *Joint density of latents and observation.* Multiply by the independent Gamma priors  $Y_s \sim \text{Gamma}(kI_s, k)$  to get  $p(Y_0, Y_{i_1}, \dots, Y_{i_n}, \text{obs})$ , with all normalising constants visible.
3. *Likelihood (model evidence).* Integrate the joint density over the latents using the Gamma normalisation  $\int_0^\infty Y^{\alpha-1} e^{-\beta Y} dY = \Gamma(\alpha) / \beta^\alpha$  in each variable.
4. *Posterior of the latents.* Divide the joint density by the likelihood to read off the posterior; in the cases below this yields independent Gammas (or a finite mixture of independent-Gamma products in case (iii)).
5. *Residual extinction probability and PMO.* The conditional extinction probability given the latents is  $q_r \mid (Y_s) = e^{-R_0(1-q)\Lambda}$  with  $\Lambda = \sum_{s=0}^r Y_s(1-F_{r-s})$  (cf. the SSI MCMC derivation in § SSI model above); marginalising over the posterior via the Gamma MGF  $\mathbb{E}[e^{-tX}] = (1+t/\beta)^{-\alpha}$  for  $X \sim \text{Gamma}(\alpha, \beta)$  gives a closed-form  $q_r$ , and the probability of a major outbreak is  $\text{PMO} = 1 - q_r$ .

A key identity used throughout is that  $I_s = 0$  on day  $s$  forces  $Y_s = 0$  (since  $Y_s \mid I_s \sim \text{Gamma}(kI_s, k)$  collapses to a point mass at 0), so among the latents only those tied to non-zero observed days are non-degenerate. Likewise, the SSE likelihood (20) is closed-form for any history, so each subsection below ends with the model-averaged PMO via Bayes' theorem from the per-model likelihoods and PMOs.

### 4.1 Case (i): $I_0 \geq 1, I_1 = \dots = I_r = 0$

Only  $Y_0$  is non-degenerate; we abbreviate  $y = Y_0$ .

**Step 1: conditional likelihood.** Given  $y$ ,  $I_d$  is independently  $\text{Poisson}(R_0 w_d y)$  for  $d = 1, \dots, r$ , so

$$\mathbb{P}(I_1 = \dots = I_r = 0 \mid y) = \prod_{d=1}^r e^{-R_0 w_d y} = e^{-R_0 F_r y}. \quad (25)$$

**Step 2: joint density.** Multiplying (25) by the prior  $y \sim \text{Gamma}(kI_0, k)$ ,

$$p(y, \text{obs}) = \frac{k^{kI_0}}{\Gamma(kI_0)} y^{kI_0-1} e^{-(k+R_0 F_r)y}. \quad (26)$$

**Step 3: likelihood.** Integrating  $y$  out of (26) via the Gamma normalisation,

$$L_{\text{SSI}}^{(i)} = \frac{k^{kI_0}}{\Gamma(kI_0)} \cdot \frac{\Gamma(kI_0)}{(k + R_0F_r)^{kI_0}} = \left(1 + \frac{R_0F_r}{k}\right)^{-kI_0}. \quad (27)$$

**Step 4: posterior.** Dividing (26) by (27) every constant cancels except the Gamma rate and shape, giving

$$y \mid \text{obs} \sim \text{Gamma}(kI_0, k + R_0F_r). \quad (28)$$

The posterior rate  $k + R_0F_r > k$  reflects the fact that observing no early transmission is evidence that  $y$  is small: the index cohort is likely a low-infectivity one.

**Step 5: residual extinction probability and PMO.** The conditional remaining-offspring rate is  $R_0\Lambda = R_0(1 - F_r)y$ , so the conditional extinction probability is  $q_r \mid y = e^{-R_0(1-q)(1-F_r)y}$ . Marginalising over the posterior (28) via the Gamma MGF,

$$q_r = \mathbb{E}\left[e^{-R_0(1-q)(1-F_r)y}\right] = \left(1 + \frac{R_0(1-F_r)(1-q)}{k + R_0F_r}\right)^{-kI_0}, \quad (29)$$

and  $\text{PMO} = 1 - q_r$ .

**Closed-form SSE counterpart.** Under the SSE model,  $I_t \sim \text{NB}(R_0I_0w_t, kI_0w_t)$  independently across  $t = 1, \dots, r$  when  $I_1 = \dots = I_r = 0$ , so the general likelihood (20) reduces to

$$L_{\text{SSE}} = \prod_{t=1}^r \left(\frac{k}{k + R_0}\right)^{kI_0w_t} = \left(1 + \frac{R_0}{k}\right)^{-kI_0F_r}. \quad (30)$$

The two likelihoods share the same functional shape but with different exponents: the SSE exponent  $kI_0F_r$  is reduced by  $F_r \leq 1$  relative to the SSI exponent  $kI_0$ , reflecting the fact that the SSE total over days  $1, \dots, r$  has dispersion  $kI_0F_r$  whereas the SSI total has dispersion  $kI_0$ . Combined with the SSE PMO  $1 - q^{I_0(1-F_r)}$  from § SSE model and the SSI PMO (29), the model-averaged PMO is

$$\text{PMO}_{\text{avg}} = \pi_{\text{SSE}}^* (1 - q^{I_0(1-F_r)}) + \pi_{\text{SSI}}^* \left[1 - \left(1 + \frac{R_0(1-F_r)(1-q)}{k + R_0F_r}\right)^{-kI_0}\right]. \quad (31)$$

When  $r = 0$ , both likelihoods reduce to 1, the posterior reduces to the prior, and the model-averaged PMO is just the prior-weighted average of the two per-model PMOs.

**Remark** (Consistency with the MCMC formula). When  $I_1 = \dots = I_r = 0$ , the observations force  $Y_1 = \dots = Y_r = 0$ , so  $\Lambda = Y_0(1 - F_r)$  and the general MCMC posterior of § SSI model collapses to (28). Taking the expectation of  $e^{-R_0(1-q)(1-F_r)Y_0}$  with respect to that posterior via the Gamma MGF recovers (29).

## 4.2 Case (ii): $I_0, I_i \geq 1$ for one later day $i$ , all others zero

We now extend to  $I_0 \geq 1$ ,  $I_i \geq 1$  for some  $i \in \{1, \dots, r\}$ , and  $I_d = 0$  for  $d \in \{1, \dots, r\} \setminus \{i\}$ . Only  $Y_0$  and  $Y_i$  are non-degenerate.

**Step 1: conditional likelihood.** Given  $(Y_0, Y_i)$ , the day- $d$  Poisson rate is  $R_0w_dY_0$  for  $d \in \{1, \dots, i\}$  and  $R_0(w_dY_0 + w_{d-i}Y_i)$  for  $d \in \{i+1, \dots, r\}$ . Multiplying the day- $d$  Poisson pmfs,

$$\mathbb{P}(\text{obs} \mid Y_0, Y_i) = \underbrace{\prod_{d=1}^{i-1} e^{-R_0w_dY_0}}_{= e^{-R_0F_{i-1}Y_0}} \cdot \underbrace{\frac{(R_0w_iY_0)^{I_i}}{I_i!} e^{-R_0w_iY_0}}_{\text{day-}i \text{ pmf}} \cdot \underbrace{\prod_{d=i+1}^r e^{-R_0(w_dY_0 + w_{d-i}Y_i)}}_{= e^{-R_0(F_r - F_i)Y_0} e^{-R_0F_{r-i}Y_i}}.$$

The  $Y_0$  exponentials collapse via  $F_{i-1} + w_i + (F_r - F_i) = F_r$ , so

$$\mathbb{P}(\text{obs} \mid Y_0, Y_i) = \frac{(R_0w_iY_0)^{I_i}}{I_i!} e^{-R_0F_rY_0} e^{-R_0F_{r-i}Y_i}. \quad (32)$$

**Step 2: joint density.** Multiplying (32) by the priors  $Y_0 \sim \text{Gamma}(kI_0, k)$  and  $Y_i \sim \text{Gamma}(kI_i, k)$  (independent),

$$p(Y_0, Y_i, \text{obs}) = \underbrace{\frac{(R_0 w_i)^{I_i}}{I_i!} \cdot \frac{k^{kI_0 + kI_i}}{\Gamma(kI_0) \Gamma(kI_i)}}_{\text{constants}} \times \underbrace{Y_0^{kI_0 + I_i - 1} e^{-(k + R_0 F_r) Y_0}}_{\text{Gamma kernel in } Y_0} \cdot \underbrace{Y_i^{kI_i - 1} e^{-(k + R_0 F_{r-i}) Y_i}}_{\text{Gamma kernel in } Y_i}. \quad (33)$$

**Step 3: likelihood.** Integrating each Gamma kernel of (33) via the Gamma normalisation gives

$$L_{\text{SSI}}^{(\text{ii})} = \frac{(R_0 w_i)^{I_i}}{I_i!} \frac{\Gamma(kI_0 + I_i)}{\Gamma(kI_0)} \frac{k^{kI_0}}{(k + R_0 F_r)^{kI_0 + I_i}} \left( \frac{k}{k + R_0 F_{r-i}} \right)^{kI_i}. \quad (34)$$

Setting  $I_i = 0$  collapses the  $\Gamma$ -ratio and the second parenthesis to 1, recovering (27).

**Step 4: posterior.** Dividing (33) by (34), every constant cancels and the two Gamma kernels normalise independently:

$$Y_0 \mid \text{obs} \sim \text{Gamma}(kI_0 + I_i, k + R_0 F_r), \quad Y_i \mid \text{obs} \sim \text{Gamma}(kI_i, k + R_0 F_{r-i}). \quad (35)$$

The shape of  $Y_0$  is enriched by  $I_i$ : each observed offspring of the index cohort acts as a Bayesian data point tilting  $Y_0$  toward higher values. The  $Y_i$  posterior is the day-0-only update (28) applied to the day- $i$  cohort over its residual horizon  $r - i$ ; setting  $i = r$  gives  $F_{r-i} = F_0 = 0$ , so  $Y_r$  collapses to its prior, since the day- $r$  cohort has not had a chance to contribute.

**Step 5: residual extinction probability and PMO.** The remaining-offspring rate is  $R_0 \Lambda$  with  $\Lambda = Y_0(1 - F_r) + Y_i(1 - F_{r-i})$ , so the conditional extinction probability is  $q_r \mid (Y_0, Y_i) = e^{-R_0(1-q)\Lambda}$ . Marginalising over the two independent Gamma factors of (35) via the Gamma MGF gives

$$q_r = \left( 1 + \frac{R_0(1 - F_r)(1 - q)}{k + R_0 F_r} \right)^{-(kI_0 + I_i)} \left( 1 + \frac{R_0(1 - F_{r-i})(1 - q)}{k + R_0 F_{r-i}} \right)^{-kI_i}, \quad (36)$$

and  $\text{PMO} = 1 - q_r$ .

**Remark** (Reduction to case (i)). Setting  $I_i = 0$  in (36) collapses the second factor to 1 and the first factor's exponent to  $-kI_0$ , recovering (29); likewise (34) reduces to (27). As the observation horizon extends ( $r \rightarrow \infty$  with  $i$  fixed), both factors of (36) tend to 1 and  $q_r \rightarrow 1$ .

### 4.3 Case (iii): $I_0, I_i, I_j \geq 1$ for two later days $i < j$ , all others zero

Finally, take  $I_0 \geq 1$ ,  $I_i \geq 1$ ,  $I_j \geq 1$  for some  $1 \leq i < j \leq r$ , and  $I_d = 0$  for  $d \in \{1, \dots, r\} \setminus \{i, j\}$ . Now only  $Y_0, Y_i, Y_j$  are non-degenerate.

**Step 1: conditional likelihood.** The conditional likelihood acquires an extra factor for the day- $j$  Poisson pmf, whose rate  $R_0(w_j Y_0 + w_{j-i} Y_i)$  couples  $Y_0$  and  $Y_i$ . Repeating the telescoping argument of case (ii) for each latent separately,

$$\mathbb{P}(\text{obs} \mid Y_0, Y_i, Y_j) = \frac{(R_0 w_i Y_0)^{I_i}}{I_i!} \frac{(R_0(w_j Y_0 + w_{j-i} Y_i))^{I_j}}{I_j!} e^{-R_0 F_r Y_0} e^{-R_0 F_{r-i} Y_i} e^{-R_0 F_{r-j} Y_j}. \quad (37)$$

Expanding the day- $j$  rate via the binomial theorem,

$$(w_j Y_0 + w_{j-i} Y_i)^{I_j} = \sum_{m=0}^{I_j} \binom{I_j}{m} w_j^{I_j-m} w_{j-i}^m Y_0^{I_j-m} Y_i^m. \quad (38)$$



**Remark** (Latent-partition interpretation). Equivalently, by Poisson superposition we may regard  $I_j$  as the sum of two independent Poissons — one with mean  $R_0 w_j Y_0$  counting day- $j$  infections that originated from the day-0 cohort, and one with mean  $R_0 w_{j-i} Y_i$  counting those from the day- $i$  cohort. Each term  $m$  in (38) is the joint probability that exactly  $I_j - m$  originate from  $Y_0$  and  $m$  from  $Y_i$ .

**Step 2: joint density.** Multiplying (37) by the three independent priors  $\text{Gamma}(kI_0, k)$ ,  $\text{Gamma}(kI_i, k)$ ,  $\text{Gamma}(kI_j, k)$  and inserting (38),

$$p(Y_0, Y_i, Y_j, \text{obs}) = \frac{(R_0 w_i)^{I_i} R_0^{I_j}}{I_i! I_j!} \frac{k^{kI_0+kI_i+kI_j}}{\Gamma(kI_0) \Gamma(kI_i) \Gamma(kI_j)} Y_j^{kI_j-1} e^{-(k+R_0 F_{r-j})Y_j} \\ \times \sum_{m=0}^{I_j} \binom{I_j}{m} w_j^{I_j-m} w_{j-i}^m Y_0^{kI_0+I_i+I_j-m-1} e^{-(k+R_0 F_r)Y_0} Y_i^{kI_i+m-1} e^{-(k+R_0 F_{r-i})Y_i}. \quad (39)$$

This is a constant times a Gamma kernel in  $Y_j$  (the same in every mixture component) times a finite mixture (over  $m \in \{0, \dots, I_j\}$ ) of Gamma-product kernels in  $(Y_0, Y_i)$ .

**Step 3: likelihood.** Integrating (39) via the Gamma normalisation in each variable, define the per-mixture-component constants

$$c_m := \binom{I_j}{m} w_j^{I_j-m} w_{j-i}^m \frac{\Gamma(kI_0 + I_i + I_j - m)}{(k + R_0 F_r)^{kI_0+I_i+I_j-m}} \frac{\Gamma(kI_i + m)}{(k + R_0 F_{r-i})^{kI_i+m}}. \quad (40)$$

Cancelling the  $\Gamma(kI_j)$  factors and grouping  $k^{kI_j}$  with the  $Y_j$ -rate gives

$$L_{\text{SSI}}^{(\text{iii})} = \frac{(R_0 w_i)^{I_i} R_0^{I_j}}{I_i! I_j!} \frac{k^{kI_0+kI_i}}{\Gamma(kI_0) \Gamma(kI_i)} \left( \frac{k}{k + R_0 F_{r-j}} \right)^{kI_j} \sum_{m=0}^{I_j} c_m. \quad (41)$$

Setting  $I_j = 0$  collapses the sum to its  $m = 0$  term and the  $(k/(k + R_0 F_{r-j}))^{kI_j}$  factor to 1, recovering (34).

**Step 4: posterior.** Dividing (39) by (41): the  $Y_j$ -Gamma rate-and-shape pair appears in both numerator and denominator and gives the  $\text{Gamma}(Y_j | kI_j, k + R_0 F_{r-j})$  density. Within the sum, the constants outside cancel; what remains in the  $m$ th term is  $c_m$  (numerator) divided by  $\sum_{m'} c_{m'}$  (denominator), times the two unnormalised  $(Y_0, Y_i)$  Gamma kernels each with their proper normalising constants supplied directly from the  $c_m$  in the denominator. The result is a finite mixture of independent Gamma products,

$$p(Y_0, Y_i, Y_j | \text{obs}) = \text{Gamma}(Y_j | kI_j, k + R_0 F_{r-j}) \\ \times \sum_{m=0}^{I_j} \rho_m \text{Gamma}(Y_0 | kI_0 + I_i + I_j - m, k + R_0 F_r) \text{Gamma}(Y_i | kI_i + m, k + R_0 F_{r-i}), \quad (42)$$

with normalised mixture weights

$$\rho_m = \frac{c_m}{\sum_{m'=0}^{I_j} c_{m'}}. \quad (43)$$

The mixture index  $m$  tracks how many of the  $I_j$  day- $j$  infections are attributed to  $Y_i$  versus  $Y_0$ ; the weights  $\rho_m$  encode which partition is most consistent with the prior infectivity distributions, the serial-interval weights  $w_j$ ,  $w_{j-i}$ , and the dispersion  $k$ .

**Step 5: residual extinction probability and PMO.** The remaining-offspring rate is now  $R_0 \Lambda = R_0[Y_0(1 - F_r) + Y_i(1 - F_{r-i}) + Y_j(1 - F_{r-j})]$ , so the conditional extinction probability is  $q_r | (Y_0, Y_i, Y_j) = e^{-R_0(1-q)\Lambda}$ . Marginalising over the posterior (42),  $Y_j$ 's expectation factors out (its density is independent of  $m$ ); within each mixture component the  $(Y_0, Y_i)$ -densities are independent Gammas whose MGFs evaluate term-by-term, giving

$$q_r = \beta_j^{-kI_j} \sum_{m=0}^{I_j} \rho_m \beta_0^{-(kI_0+I_i+I_j-m)} \beta_i^{-(kI_i+m)}, \quad (44)$$

where, for  $a \in \{0, i, j\}$  (with the convention  $F_{r-0} = F_r$ ),

$$\beta_a = 1 + \frac{R_0(1-q)(1-F_{r-a})}{k + R_0F_{r-a}}.$$

**Remark** (Reductions to simpler cases). When  $I_j \rightarrow 0$  the sum collapses to its  $m = 0$  term ( $\rho_0 = 1$ ) and  $\beta_j^{-kI_j} \rightarrow 1$ : (44) reduces to (36). When also  $I_i \rightarrow 0$ , the exponent on  $\beta_i$  in (36) vanishes and the exponent on  $\beta_0$  reduces from  $-(kI_0 + I_i)$  to  $-kI_0$ , recovering (29). Similarly (41) reduces to (34) and then to (27).

**Computational note.** The  $c_m$  involve  $\Gamma$  factors of size up to  $kI_0 + I_i + I_j$ , so for moderate  $I_j$  a log-domain computation ( $\log \Gamma$ ,  $\text{logsumexp}$ ) is recommended. The number of sum terms is  $I_j + 1$ , typically small for the histories of interest.

## 5 MCMC-based estimators of the SSI marginal likelihood

We describe three estimators of the integral (23). Writing  $\pi(Y) = \prod_{t \in \mathcal{T}} p_Y(Y_t | I_t)$  for the integrand's prior factor restricted to the active days  $\mathcal{T} = \{t \in \{0, \dots, r-1\} : I_t > 0\}$  (a product of independent Gammas, treating the observed cohort sizes  $I_t$  as fixed shape parameters), and  $\mathcal{L}(Y) = \prod_{t=1}^r p_I(I_t | Y_{0:t-1})$  for the integrand's likelihood factor, the target is

$$L_{\text{SSI}} = \int \mathcal{L}(Y) \pi(Y) dY = \mathbb{E}_{Y \sim \pi}[\mathcal{L}(Y)]. \quad (45)$$

The unnormalised posterior (the joint integrand) is  $\tilde{p}(Y) = \mathcal{L}(Y) \pi(Y)$ , and the posterior density of the latents given the observations is  $p_{\text{post}}(Y) = \tilde{p}(Y)/L_{\text{SSI}}$ . The MCMC sampler of § 2.1.2 produces draws  $Y^{p.(j)} \sim p_{\text{post}}$  for  $j = 1, \dots, N_1$ .

### 5.1 Naive Monte Carlo

Equation (45) is itself an expectation under the prior  $\pi$ , so the simplest estimator just averages the conditional likelihood at independent draws from  $\pi$ :

$$\hat{L}_{\text{SSI}}^{\text{naive}} = \frac{1}{N} \sum_{n=1}^N \mathcal{L}(Y^{(n)}), \quad Y^{(n)} \stackrel{\text{iid}}{\sim} \pi. \quad (46)$$

No MCMC trace is needed: drawing  $Y^{(n)}$  requires only independent  $\text{Gamma}(kI_t, k)$  samples for each  $t \in \mathcal{T}$ . In log space the implementation is

$$\log \hat{L}_{\text{SSI}}^{\text{naive}} = \text{logsumexp}_{n=1, \dots, N} \log \mathcal{L}(Y^{(n)}) - \log N.$$

The estimator is unbiased. Its relative variance is  $\text{Var}_{\pi}[\mathcal{L}]/(NL_{\text{SSI}}^2)$ , equivalently  $(\chi^2(p_{\text{post}} \| \pi) + 1)/N - 1/N$ , where  $\chi^2(p \| q) = \int (p/q - 1)^2 q$  is the  $\chi^2$  divergence. The  $\chi^2$  between prior and posterior is small whenever the data are weakly informative about the latents — and remarkably, in the SSI model this remains true even on moderately informative histories, because the prior variance  $I_t/k$  scales automatically with the cohort size  $I_t$  and so retains overlap with the posterior. The estimator becomes unreliable only when the data are very informative (e.g. long histories with substantial incidence at small  $k$ ).

### 5.2 Importance sampling

Importance sampling generalises the naive estimator by sampling from a proposal  $g$  chosen to better cover the posterior:

$$\hat{L}_{\text{SSI}}^{\text{IS}} = \frac{1}{N} \sum_{n=1}^N \frac{\tilde{p}(Y^{(n)})}{g(Y^{(n)})}, \quad Y^{(n)} \stackrel{\text{iid}}{\sim} g. \quad (47)$$

The estimator is unbiased for any proposal  $g$  with  $g(Y) > 0$  wherever  $\tilde{p}(Y) > 0$ ; its relative variance is  $\chi^2(p_{\text{post}} \| g)/N$ , minimised by taking  $g$  as close to  $p_{\text{post}}$  as possible.

We take

$$g(Y) = \prod_{t \in \mathcal{T}} \text{Gamma}(Y_t; \alpha_t, \beta_t), \quad (48)$$

i.e. a product of independent Gammas in the same family as the prior, with shape and rate parameters fitted by method of moments to the per-day marginals of the MCMC posterior trace  $\{Y^{p,(j)}\}_{j=1}^{N_1}$ :

$$\alpha_t = \frac{\bar{Y}_t^2}{S_t^2}, \quad \beta_t = \frac{\bar{Y}_t}{S_t^2}, \quad t \in \mathcal{T}, \quad (49)$$

where  $\bar{Y}_t$  and  $S_t^2$  are the sample mean and (unbiased) sample variance of  $\{Y_t^{p,(j)}\}_j$ . Matching the proposal family to the prior keeps importance weights bounded; matching its marginals to the MCMC posterior makes it close to  $p_{\text{post}}$  marginally.

**Remark** (Caveat: posterior correlations). Equation (48) is a product proposal, while the joint posterior of  $Y_{\mathcal{T}}$  may have non-trivial correlations arising from the shared force-of-infection terms. When such correlations are strong,  $\chi^2(p_{\text{post}} \| g)$  does not shrink to zero in the large-data limit and the IS estimator's variance remains bounded below by that mismatch. Bridge sampling (§ 5.3) is more forgiving of proposal mismatch.

### 5.3 Bridge sampling

Bridge sampling uses draws from *both* the proposal  $g$  and the posterior  $p_{\text{post}}$ , weighted by a bridge function  $\alpha$ . The starting identity (Meng & Wong, 1996) is the algebraic equality

$$L_{\text{SSI}} = \frac{\mathbb{E}_{Y \sim g}[\tilde{p}(Y) \alpha(Y)]}{\mathbb{E}_{Y \sim p_{\text{post}}}[g(Y) \alpha(Y)]}, \quad (50)$$

valid for any  $\alpha$  such that  $0 < \int g(Y) \tilde{p}(Y) \alpha(Y) dY < \infty$ . The identity is verified by writing the numerator as  $\int \tilde{p} g \alpha dY$  and the denominator as  $L_{\text{SSI}}^{-1} \int \tilde{p} g \alpha dY$ . Replacing the expectations by Monte Carlo averages over  $\{Y^{q,(i)}\}_{i=1}^{N_2} \stackrel{\text{iid}}{\sim} g$  and  $\{Y^{p,(j)}\}_{j=1}^{N_1} \sim p_{\text{post}}$  (the MCMC trace) gives the bridge-sampling estimator.

The asymptotic variance of  $\hat{L}_{\text{SSI}}$  depends on the choice of  $\alpha$ . Minimising it (Meng & Wong, 1996) yields the optimal bridge

$$\alpha^*(Y) = \frac{C}{s_1 g(Y) + s_2 \tilde{p}(Y)/L_{\text{SSI}}}, \quad (51)$$

where  $s_1 = N_1/(N_1 + N_2)$ ,  $s_2 = N_2/(N_1 + N_2)$ , and  $C$  is any non-zero constant (any rescaling of  $\alpha$  cancels between numerator and denominator of (50)). The optimal bridge depends on the unknown  $L_{\text{SSI}}$ ; substituting (51) into (50) and replacing the expectations by Monte Carlo averages produces the fixed-point identity

$$L_{\text{SSI}} = \frac{\frac{1}{N_2} \sum_{i=1}^{N_2} \frac{\tilde{p}(Y^{q,(i)})}{s_1 g(Y^{q,(i)}) + s_2 \tilde{p}(Y^{q,(i)})/L_{\text{SSI}}}}{\frac{1}{N_1} \sum_{j=1}^{N_1} \frac{g(Y^{p,(j)})}{s_1 g(Y^{p,(j)}) + s_2 \tilde{p}(Y^{p,(j)})/L_{\text{SSI}}}}}. \quad (52)$$

Solving (52) for  $L_{\text{SSI}}$  requires iterating the right-hand side from an initial guess. Bridge sampling has lower (or equal) asymptotic variance than importance sampling with the same proposal, and the variance bound is governed by a symmetrised divergence rather than  $\chi^2(p_{\text{post}} \| g)$ , so it is significantly more robust to proposal-posterior mismatch.

In practice we work entirely in log space for numerical stability. Define the log importance ratios at the two sample sets,

$$\ell_q^{(i)} = \log \tilde{p}(Y^{q,(i)}) - \log g(Y^{q,(i)}), \quad \ell_p^{(j)} = \log \tilde{p}(Y^{p,(j)}) - \log g(Y^{p,(j)}).$$

Multiplying numerator and denominator of (52) by  $1/g(Y^{q,(i)})$  and  $1/g(Y^{p,(j)})$  respectively and taking logs gives the log-space iteration

$$\begin{aligned} \log L^{(t+1)} = & \log \sum_{i=1, \dots, N_2} \exp \left[ \ell_q^{(i)} - \text{lse}(\log s_1, \log s_2 + \ell_q^{(i)} - \log L^{(t)}) \right] - \log N_2 \\ & - \log \sum_{j=1, \dots, N_1} \exp \left[ -\text{lse}(\log s_1, \log s_2 + \ell_p^{(j)} - \log L^{(t)}) \right] + \log N_1, \quad (53) \end{aligned}$$

where  $\text{lse}(a, b) = \log(e^a + e^b)$ . We use the same Gamma proposal  $g$  as in § 5.2, initialise  $\log L^{(0)}$  at the median of  $\{\ell_q^{(i)}\}_i$ , and iterate (53) until  $\log L^{(t)}$  stops changing appreciably.