

P8123 Final

Farizah Rob, Emily Potts, Harry Wang, William Anderson

2023-12-14

```
library(tidyverse)
library(survey)
library(ggsurvey)
library(tableone)
library(gtsummary)
library(sjPlot)

knitr::opts_chunk$set(
  message = FALSE,
  warning = FALSE
)

theme_set(theme_minimal() + theme(legend.position = "bottom"))

load("../data/nhanes_data.rda") #put files into data folder so everyone can run this
load("../data/nhanes_key.rda")
load("../data/key_guide.rda")
```

Problem statement: Controlling blood pressure (BP) reduces the risk for cardiovascular disease. However, the prevalence of BP control (i.e., systolic BP < 140 and diastolic BP < 90) among US adults with hypertension has decreased since 2013. We invite teams to analyze publicly available data from US adults to help identify potential causes or correlates of worsening BP control among US adults with hypertension over the past decade, as this may allow for development of effective interventions to help control BP and prevent cardiovascular disease.

Svydesign and subset for hypertension patients 2013 onwards

```
dstrat <- svydesign(ids=~svy_psu, strata=~svy_strata, weights=~svy_weight_mec, data=nhanes_data, nest=TRUE)


htsub <- subset(dstrat, svy_subpop_htn==1) #limited to hypertension patients

htsub2 <- subset(htsub, svy_year=="2013-2014" | svy_year=="2015-2016" | svy_year=="2017-2020") #limited to 2013 onwards
```

Some EDA


The **outcome** of interest -> **bp_control_140_90** OR **bp_uncontrolled_140_90**

```
#age -> so for older people, BP control is definitely lower
age <- svyby(~factor(bp_control_140_90), ~factor(demo_age_cat), design=htsub2, svymean, na.rm=TRUE)
barplot(age, legend.text=c("BP uncontrolled", "BP controlled"), col=c("pink", "purple"), main="Proportion of BP control by age category")
```




```
#gender -> not really any difference
gender <- svyby(~factor(bp_control_140_90), ~factor(demo_gender), design=htsub2, svymean, na.rm=TRUE)
barplot(gender, legend.text=c("BP uncontrolled", "BP controlled"), col=c("pink", "purple"), main="Proportion of BP control by gender")
```

Proportion of BP control by gender




```
#race -> for Black people vs non-Black people, BP control is lower
race <- svyby(~factor(bp_control_140_90), ~factor(demo_race_black), design=htsub2, svymean, na.rm=TRUE)
barplot(race,
  legend.text=c("BP uncontrolled", "BP controlled"),
  names.arg=c("Non-Black", "Black"),
  ylim=c(0, 1.0),
  args.legend = list(x = "topright"),
  main="Proportion of BP control by race (non-Black vs Black)")
```

Proportion of BP control by race (non-Black vs Black)




```
#diabetes
diabetes <- svyby(~factor(bp_control_140_90), ~factor(cc_diabetes), design=htsub2, svymean, na.rm=TRUE)
barplot(diabetes,
  legend.text=c("BP uncontrolled", "BP controlled"),
  names.arg=c("No Diabetes", "Diabetes"),
  ylim=c(0, 1.0),
  args.legend = list(x = "topright"),
  main="Proportion of BP control by diabetes")
```

Proportion of BP control by diabetes




```
#ckd
ckd <- svyby(~factor(bp_control_140_90), ~factor(cc_ckd), design=htsub2, svymean, na.rm=TRUE)
barplot(ckd,
  legend.text=c("BP uncontrolled", "BP controlled"),
  names.arg=c("No CKD", "CKD"),
  ylim=c(0, 1.0),
  args.legend = list(x = "topright"),
  main="Proportion of BP control by CKD")
```

Proportion of BP control by CKD



```
#bmi
bmi <- svyby(~factor(bp_control_140_90), ~factor(cc_bmi), design=htsub2, svymean, na.rm=TRUE)
barplot(bmi,
  legend.text=c("BP uncontrolled", "BP controlled"),
  ylim=c(0, 1.0),
  args.legend = list(x = "topright"),
  main="Proportion of BP control by BMI")
```

Proportion of BP control by BMI



#meds recommended -> 100% for those with uncontrolled BP
look at med recommended and meds taken ratio by different subgroups?

```
table(nhanes_data$bp_med_recommended_jnc7, nhanes_data$bp_med_use) #48% of people recommended medication
```

```
##
##           No    Yes
## No  37755      0
## Yes  7062 14690
```

```
svyCreateCatTable(vars= c("bp_med_recommended_jnc7" ,"bp_med_use"),
                  strata="demo_race",
                  data=htsub2) #across all races, the % of med use is lower than % recommended -? this
```

```
##
##           Stratified by demo_race
##           Non-Hispanic White  Non-Hispanic Black
## n  443841370.8  78256419.3
## bp_med_recommended_jnc7 = Yes (%) 156173388.2 (35.2) 33534720.2 (42.9)
## bp_med_use = Yes (%) 111473163.2 (25.1) 23255215.8 (29.7)
##
##           Stratified by demo_race
##           Non-Hispanic Asian  Hispanic
## n  37820092.9  105372525.4
## bp_med_recommended_jnc7 = Yes (%) 11080831.4 (29.3) 26424308.7 (25.1)
## bp_med_use = Yes (%) 6822774.8 (18.0) 15828214.5 (15.0)
```

```
##                                Stratified by demo_race
##                                Other           p       test
##    n                          24220134.3
##    bp_med_recommended_jnc7 = Yes (%) 8029041.0 (33.2) <0.001
##    bp_med_use = Yes (%)             5443707.4 (22.5) <0.001

#for table 1 -> age category, gender, race, BMI, smoking status, diabetes, CKD, cholesterol, recommended
# removed "bp_med_recommended_jnc7", "bp_med_recommended_accaha" for space

# didn't they say something in class about the Rao and Scott being wrong in R? not sure if we should in
# I don't think we should have p-values in the table 1 since we're not running models yet, just showing

reset_gtsummary_theme()
theme_gtsummary_compact()
tab1 <- tbl_svysummary(data = htsub2, by = "bp_control_140_90", include = c("demo_age_cat", "demo_gender",
                                label = list(demo_age_cat ~ "Age",
                                demo_gender ~ "Gender",
                                demo_race ~ "Race",
                                cc_bmi ~ "BMI",
                                cc_ckd ~ "Chronic Kidney Disease",
                                cc_diabetes ~ "Diabetes",
                                cc_cvd_any ~ "Cardiovascular Disease",
                                bp_med_use ~ "BP Medication Use",
                                chol_total ~ "Total Cholesterol",
                                chol_hdl ~ "HDL",
                                chol_ldl ~ "LDL",
                                chol_trig ~ "Triglycerides",
                                chol_med_use ~ "Cholesterol Medication Use"))) %>% add_p() %>% add_
                                italicize_levels()

tab1
```

Table 1: Subject Characteristics (Survey Weighted)

Characteristic	No, N = 110,019,543	Yes, N = 579,490,999	p-value
Age, n (%)			<0.001
18 to 44	18,218,707 (17%)	299,580,713 (52%)	
45 to 64	45,525,230 (41%)	192,746,908 (33%)	
65 to 74	23,816,079 (22%)	58,369,736 (10%)	
75+	22,459,527 (20%)	28,793,643 (5.0%)	
Gender, n (%)			0.050
Men	55,991,881 (51%)	278,426,528 (48%)	
Women	54,027,662 (49%)	301,064,472 (52%)	
Race, n (%)			<0.001
Non-Hispanic White	68,928,072 (63%)	374,913,299 (65%)	
Non-Hispanic Black	18,291,583 (17%)	59,964,836 (10%)	
Non-Hispanic Asian	5,715,421 (5.2%)	32,104,671 (5.5%)	
Hispanic	13,404,152 (12%)	91,968,373 (16%)	
Other	3,680,315 (3.3%)	20,539,820 (3.5%)	
BMI, n (%)			<0.001
<25	23,597,988 (22%)	173,248,879 (30%)	

Characteristic	No, N = 110,019,543	Yes, N = 579,490,999	p-value
25 to <30	33,526,572 (31%)	179,795,985 (32%)	
30 to <35	23,810,864 (22%)	117,994,322 (21%)	
35+	26,600,214 (25%)	97,987,747 (17%)	
Unknown	2,483,906	10,464,066	
Chronic Kidney Disease, n (%)	32,045,575 (29%)	62,111,324 (11%)	<0.001
Diabetes, n (%)	22,552,554 (20%)	55,629,143 (9.6%)	<0.001
Cardiovascular Disease, n (%)	15,856,616 (14%)	40,856,909 (7.1%)	<0.001
BP Medication Use, n (%)	51,549,103 (47%)	111,273,973 (19%)	<0.001
Total Cholesterol, Median (IQR)	192 (163, 222)	183 (158, 211)	<0.001
Unknown	63,100,407	325,591,825	
HDL, Median (IQR)	52 (42, 65)	52 (43, 63)	>0.9
Unknown	63,100,407	325,591,825	
LDL, Median (IQR)	112 (88, 139)	109 (86, 133)	0.013
Unknown	63,221,679	325,731,647	
Triglycerides, Median (IQR)	105 (72, 155)	89 (61, 135)	<0.001
Unknown	63,221,679	325,731,647	
Cholesterol Medication Use, n (%)	16,391,227 (34%)	47,895,434 (19%)	<0.001
Unknown	62,505,218	323,610,642	

Possible Figure 1:

I made a six-panel figure that could serve as our single Figure 1, feel free to change these plots to different things or remove ones we might not need. - Will

```

par(mfrow = c(2, 3))

svyhist(~bp_sys_mean, htsub2, main = "Mean SBP NHANES 2013-2020", xlab = "Mean SBP (mmHg)")

prop <- svymean(~bp_uncontrolled_140_90, htsub2, na.rm=TRUE)

barplot(prop,
        ylim=c(0, 1.0),
        main="BP Control NHANES 2013-2020", names.arg = c("Uncontrolled", "Controlled"),
        ylab = "Proportion")

svyplot(bp_sys_mean~demo_age_years, htsub2, style = "subsample", xlab = "Age (years)", ylab = "Mean SBP")

race <- svyby(~bp_uncontrolled_140_90, ~factor(demo_race), htsub2, svymean, na.rm=TRUE)

barplot(race,
        legend.text=c("Uncontrolled", "Controlled"),
        ylim=c(0, 1.0),
        args.legend = list(x = "topright", bty = "n", inset = c(-0.20, -0.2)),
        main="BP Control by Race",
        ylab = "Proportion",
        names.arg = c("White", "Black", "Asian", "Hispanic", "Other"),
        cex.names = 0.75)

svyboxplot(bp_sys_mean~factor(cc_smoke), htsub2, na.rm=TRUE, xlab = "Smoking Status", ylab = "Mean SBP")

```


```

cvd <- svyby(~bp_uncontrolled_140_90, ~factor(cc_cvd_chd), htsub2, na.rm = T, svymean)


barplot(cvd,
  legend.text=c("Uncontrolled", "Controlled"),
  ylim=c(0, 1.0),
  args.legend = list(x = "topright", bty = "n", inset = c(-0.20, -0.1)),
  main="BP Control by CVD History",
  xlab = "Cardiovascular Disease History",
  ylab = "Proportion")

```


Mean SBP NHANES 2013–202




BP Control NHANES 2013–202




Subsample Bubble Plot




BP Control by Race



SBP per Smoking Status



BP Control by CVD History



Regression models - really don't know which variables to select

using the `bp_uncontrolled_140_90` variable instead of `bp_control` for regression

```

m1 <- svyglm(bp_uncontrolled_140_90 ~ demo_age_cat + demo_race + demo_gender + cc_smoke + cc_bmi + cc_di)

#need to set BMI reference value to normal level rather than underweight

summary(m1)

```

```
##
```

```
## Call:
```

```
## svyglm(formula = bp_uncontrolled_140_90 ~ demo_age_cat + demo_race +
```

```
##      demo_gender + cc_smoke + cc_bmi + cc_diabetes + cc_ckd +
##      cc_cvd_any + chol_trig + chol_ldl + chol_hdl, design = htsub2,
##      family = quasibinomial())
##
## Survey design:
## subset(htsub, svy_year == "2013-2014" | svy_year == "2015-2016" |
##       svy_year == "2017-2020")
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      -4.2784289   0.2847878 -15.023  < 2e-16 ***
## demo_age_cat45 to 64    1.1528233   0.1252890   9.201 5.47e-11 ***
## demo_age_cat65 to 74    1.7627753   0.1598648  11.027 4.28e-13 ***
## demo_age_cat75+        2.4495017   0.1681808  14.565  < 2e-16 ***
## demo_raceNon-Hispanic Black 0.8670093   0.1173396   7.389 1.02e-08 ***
## demo_raceNon-Hispanic Asian 0.3999404   0.1137010   3.517 0.001199 **
## demo_raceHispanic        0.1559047   0.1091321   1.429 0.161743
## demo_raceOther          0.0207119   0.2431672   0.085 0.932594
## demo_genderWomen        -0.3087733   0.1070130  -2.885 0.006566 **
## cc_smokeFormer          0.0729305   0.1250743   0.583 0.563463
## cc_smokeCurrent         0.2004411   0.1685633   1.189 0.242179
## cc_bmi25 to <30        -0.0215815   0.1116762  -0.193 0.847849
## cc_bmi30 to <35         0.2261152   0.1130356   2.000 0.053045 .
## cc_bmi35+              0.7023231   0.1345164   5.221 7.63e-06 ***
## cc_diabetesYes          0.3421597   0.1262108   2.711 0.010214 *
## cc_ckdYes              0.6520772   0.1190915   5.475 3.49e-06 ***
## cc_cvd_anyYes          -0.0273710   0.1208875  -0.226 0.822158
## chol_trig              0.0017308   0.0003819   4.533 6.21e-05 ***
## chol_ldl               0.0052070   0.0012622   4.125 0.000209 ***
## chol_hdl               0.0061692   0.0032180   1.917 0.063190 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for quasibinomial family taken to be 0.9486429)
##
## Number of Fisher Scoring iterations: 5
```

#need to look into subset of high cholesterol and hypertension?

Modeling medication use with bp_uncontrolled_140_90 outcome

Model is limited to survey design subset of hypertension patients from 2013-2020. We selected medication variables concerning self-reported blood pressure medication use (yes/no), the number of antihypertensive medication pills taken per day, the use of vasodilators (yes/no), alpha and beta blockers (yes/no), the use of angiotensin receptor blockers, and the use of statins or other cholesterol medication for cholesterol control (yes/no). These variables were selected based on their prior association with blood pressure levels in existing clinical trials and research studies.

```
meds_binary <- svyglm(bp_uncontrolled_140_90 ~ bp_med_use + bp_med_n_pills + bp_med_vasod + bp_med_alpha
```

Use of blood pressure medications, number of antihypertensive pills (1-4+), and angiotensin receptor blockers, were significantly associated with having uncontrolled blood pressure. The use of vasodilators, statins or

other cholesterol medication, alpha and beta blockers, had non-statistically significant associations with having uncontrolled blood pressure.

Modeling medication use with bp_sys_mean outcome

We now model the continuous outcome of the average systolic blood pressure with the same medication covariates among those with hypertension from 2013-2020.

```
meds_continuous <- svyglm(bp_sys_mean ~ bp_med_use + bp_med_n_pills + bp_med_vasod + bp_med_alpha + bp_med_beta, data = htsub2, family = quasibinomial())
```

The use of blood pressure medications, number of antihypertensive pills (1), and angiotensin receptor blockers were significantly associated with mean systolic blood pressure levels in those with hypertension from 2013-2020. The use of (2-4+) antihypertensive pills per day, vasodilators, alpha and beta blockers, and statins and other cholesterol medications, had non-statistically significant associations with mean systolic blood pressure levels in those with hypertension from 2013-2020.

Nested Models

Level One Demographic Characteristics.

```
model_1 <- svyglm(bp_uncontrolled_140_90 ~ demo_age_cat + demo_race + demo_gender, #demographics
                  design = htsub2, family=quasibinomial())
```

```
summary(model_1)
```

```
##
## Call:
## svyglm(formula = bp_uncontrolled_140_90 ~ demo_age_cat + demo_race +
##       demo_gender, design = htsub2, family = quasibinomial())
##
## Survey design:
## subset(htsub, svy_year == "2013-2014" | svy_year == "2015-2016" |
##       svy_year == "2017-2020")
##
## Coefficients:
##               Estimate Std. Error t value Pr(>|t|)
## (Intercept)      -2.92319    0.09320  -31.364 < 2e-16 ***
## demo_age_cat45 to 64    1.41571    0.07074   20.014 < 2e-16 ***
## demo_age_cat65 to 74    2.01652    0.07848   25.696 < 2e-16 ***
## demo_age_cat75+        2.70181    0.10722   25.200 < 2e-16 ***
## demo_raceNon-Hispanic Black  0.86309    0.07301   11.822 1.1e-15 ***
## demo_raceNon-Hispanic Asian  0.26243    0.07708    3.405 0.00136 **
## demo_raceHispanic        0.23525    0.07650    3.075 0.00350 **
## demo_raceOther          0.27350    0.17971    1.522 0.13474
## demo_genderWomen        -0.21386    0.06224   -3.436 0.00124 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for quasibinomial family taken to be 0.9818051)
##
## Number of Fisher Scoring iterations: 5
```

Level Two Health-related Characteristics

```
model_2 <- svyglm(bp_uncontrolled_140_90 ~ demo_age_cat + demo_race + demo_gender + #demographics
                  cc_smoke + cc_bmi+ cc_diabetes + cc_ckd + cc_cvd_any + chol_trig + chol_ldl + chol_hdl,
                  design = htsub2, family=quasibinomial())
summary(model_2)
```

```
##
## Call:
## svyglm(formula = bp_uncontrolled_140_90 ~ demo_age_cat + demo_race +
##       demo_gender + cc_smoke + cc_bmi + cc_diabetes + cc_ckd +
##       cc_cvd_any + chol_trig + chol_ldl + chol_hdl, design = htsub2,
##       family = quasibinomial())
##
## Survey design:
## subset(htsub, svy_year == "2013-2014" | svy_year == "2015-2016" |
##       svy_year == "2017-2020")
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      -4.2784289   0.2847878  -15.023   < 2e-16 ***
## demo_age_cat45 to 64    1.1528233   0.1252890    9.201 5.47e-11 ***
## demo_age_cat65 to 74    1.7627753   0.1598648   11.027 4.28e-13 ***
## demo_age_cat75+        2.4495017   0.1681808   14.565   < 2e-16 ***
## demo_raceNon-Hispanic Black  0.8670093   0.1173396    7.389 1.02e-08 ***
## demo_raceNon-Hispanic Asian  0.3999404   0.1137010    3.517 0.001199 **
## demo_raceHispanic        0.1559047   0.1091321    1.429 0.161743
## demo_raceOther          0.0207119   0.2431672    0.085 0.932594
## demo_genderWomen        -0.3087733   0.1070130   -2.885 0.006566 **
## cc_smokeFormer          0.0729305   0.1250743    0.583 0.563463
## cc_smokeCurrent         0.2004411   0.1685633    1.189 0.242179
## cc_bmi25 to <30        -0.0215815   0.1116762   -0.193 0.847849
## cc_bmi30 to <35         0.2261152   0.1130356    2.000 0.053045 .
## cc_bmi35+              0.7023231   0.1345164    5.221 7.63e-06 ***
## cc_diabetesYes         0.3421597   0.1262108    2.711 0.010214 *
## cc_ckdYes              0.6520772   0.1190915    5.475 3.49e-06 ***
## cc_cvd_anyYes         -0.0273710   0.1208875   -0.226 0.822158
## chol_trig              0.0017308   0.0003819    4.533 6.21e-05 ***
## chol_ldl               0.0052070   0.0012622    4.125 0.000209 ***
## chol_hdl               0.0061692   0.0032180    1.917 0.063190 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for quasibinomial family taken to be 0.9486429)
##
## Number of Fisher Scoring iterations: 5
```

Level Three Medication Use

```
model_3 <- svyglm(bp_uncontrolled_140_90 ~ demo_age_cat + demo_race + demo_gender + #demographics
                  cc_smoke + cc_bmi+ cc_diabetes + cc_ckd + cc_cvd_any + chol_trig + chol_ldl + chol_hdl,
```

```

        bp_med_use + bp_med_n_pills + bp_med_vasod + bp_med_alpha + #medication
        bp_med_beta + bp_med_angioten + chol_med_statin + chol_med_use, #medication
design = htsub2, family=quasibinomial())
summary(model_3)

```

```

##
## Call:
## svyglm(formula = bp_uncontrolled_140_90 ~ demo_age_cat + demo_race +
##      demo_gender + cc_smoke + cc_bmi + cc_diabetes + cc_ckd +
##      cc_cvd_any + chol_trig + chol_ldl + chol_hdl + bp_med_use +
##      bp_med_n_pills + bp_med_vasod + bp_med_alpha + bp_med_beta +
##      bp_med_angioten + chol_med_statin + chol_med_use, design = htsub2,
##      family = quasibinomial())
##
## Survey design:
## subset(htsub, svy_year == "2013-2014" | svy_year == "2015-2016" |
##      svy_year == "2017-2020")
##
## Coefficients:
##
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      -4.2354978   0.2697846 -15.700 1.86e-14 ***
## demo_age_cat45 to 64    1.0690468   0.1288047   8.300 1.19e-08 ***
## demo_age_cat65 to 74    1.6111371   0.1594032  10.107 2.58e-10 ***
## demo_age_cat75+        2.2844522   0.1762558  12.961 1.35e-12 ***
## demo_raceNon-Hispanic Black  0.8434587   0.1182782   7.131 1.79e-07 ***
## demo_raceNon-Hispanic Asian  0.3873648   0.1161263   3.336 0.002659 **
## demo_raceHispanic        0.1973600   0.1073986   1.838 0.078030 .
## demo_raceOther          0.0066420   0.2505540   0.027 0.979062
## demo_genderWomen        -0.3324135   0.1055494  -3.149 0.004206 **
## cc_smokeFormer          0.0752089   0.1281457   0.587 0.562533
## cc_smokeCurrent         0.2019008   0.1687557   1.196 0.242761
## cc_bmi25 to <30        -0.0631372   0.1149384  -0.549 0.587664
## cc_bmi30 to <35         0.1786980   0.1134249   1.575 0.127718
## cc_bmi35+              0.5917701   0.1431533   4.134 0.000351 ***
## cc_diabetesYes          0.3255830   0.1337452   2.434 0.022397 *
## cc_ckdYes              0.6201295   0.1265869   4.899 4.84e-05 ***
## cc_cvd_anyYes          -0.0996443   0.1392883  -0.715 0.481002
## chol_trig              0.0016698   0.0003928   4.252 0.000259 ***
## chol_ldl               0.0050071   0.0012360   4.051 0.000434 ***
## chol_hdl               0.0063284   0.0032239   1.963 0.060870 .
## bp_med_useYes          0.5344699   0.2079053   2.571 0.016488 *
## bp_med_n_pillsOne       -0.1874577   0.2794390  -0.671 0.508476
## bp_med_n_pillsTwo       -0.2084591   0.2981528  -0.699 0.490901
## bp_med_n_pillsThree     -0.4789832   0.3351850  -1.429 0.165378
## bp_med_n_pillsFour or more -0.4355646   0.3967429  -1.098 0.282731
## bp_med_vasodYes         0.0736384   0.3720228   0.198 0.844692
## bp_med_alphaYes         0.1738527   0.3837901   0.453 0.654464
## bp_med_betaYes          0.2530225   0.1310938   1.930 0.065016 .
## bp_med_angiotenYes      0.3140413   0.1492004   2.105 0.045525 *
## chol_med_statinYes      0.0018895   0.3706140   0.005 0.995973
## chol_med_useYes        -0.1105545   0.3425864  -0.323 0.749602
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```


```
##
## (Dispersion parameter for quasibinomial family taken to be 0.941334)
##
## Number of Fisher Scoring iterations: 5
```

I added the nested models based on three levels. Should we also make figures based on the significant variables from these models? If these models look good, I will proceed making the table two - Harry

Figure 2: Forest plot of estimates from models 1-3

This plots estimates and 95% CI for odds ratios from models 1-3 and can be our figure 2 - Will

```
plot_models(model_1, model_2, model_3, transform = "exp", legend.title = "", show.p = T, grid = T, show
```



Farizah's code for plotting forest plot of model 3, I don't think it's outputting correctly:

```
#age
reg_age <- matrix(c(1.0690468, 1.6111371, 2.2844522, 0.1288047, 0.1594032, 0.1762558), byrow=FALSE, ncol=6)

colnames(reg_age) <- c("log_or", "se")

reg_age <- reg_age %>% as_tibble()
reg_age$age <- c("45 to 64", "65 to 74", "75+" )
reg_age$l95 <- reg_age$log_or-1.96*reg_age$se
reg_age$u95 <- reg_age$log_or+1.96*reg_age$se
```

```

age_plot <- ggplot(reg_age, aes(x = log_or, y = age)) + geom_point() + geom_pointrange(aes(xmin = 195,
#race

reg_race <- matrix(c(0.8434587, 0.3873648, 0.1973600, 0.0066420, 0.1182782, 0.1161263, 0.1073986, 0.250
colnames(reg_race) <- c("log_or", "se")

reg_race <- reg_race %>% as_tibble()
reg_race$race <- c("Non-Hispanic Black", "Non-Hispanic Asian", "Hispanic", "Other" )
reg_race$l95 <- reg_race$log_or-1.96*reg_race$se
reg_race$u95 <- reg_race$log_or+1.96*reg_race$se

race_plot <- ggplot(reg_race, aes(x = log_or, y = race)) + geom_point() + geom_pointrange(aes(xmin = 195
#health


reg_health <- matrix(c(0.5917701, 0.3255830, 0.6201295, 0.0016698, 0.0050071, 0.5344699, 0.3140413, 0.1
colnames(reg_health) <- c("log_or", "se")

reg_health <- reg_health %>% as_tibble()
reg_health$Variable <- c("BMI 35+", "Diabetes", "CKD", "Cholesterol (Trig)", "Cholesterol (LDL)", "Takin

reg_health$l95 <- reg_health$log_or-1.96*reg_health$se
reg_health$u95 <- reg_health$log_or+1.96*reg_health$se

health_plot <- ggplot(reg_health, aes(x = log_or, y = Variable)) + geom_point() + geom_pointrange(aes(x
health_plot

```

```
library(ggpubr)
ggarrange(age_plot, race_plot, health_plot, ncol =3, nrow=1, labels = c("Age (ref = 18-44)", "Race (ref = White)", "Health (ref = No disease)"))
```

