

The image shows a close-up of a rock surface covered with numerous small, green, cup-shaped fungi. Each cup has a dark brown interior and a green, ribbed exterior. The rock surface is dark and textured, with some green moss visible in the background.

# Chapter 3

## Diversity of Fungi of Colombia

Ester Gaya, Viviana Motato-Vásquez & Robert Lücking

*Rhytidhysterium columbiense*  
[Robert Lücking]



# Chapter 3

## Diversity of Fungi of Colombia

Ester Gaya<sup>1</sup>, Viviana Motato-Vásquez<sup>2,3</sup> & Robert Lücking<sup>4\*</sup>

<sup>1</sup> Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3DS, UK.

<sup>2</sup> Grupo de Investigación en Biología de Plantas y Microorganismos (BPM), Departamento de Biología, Facultad de Ciencias Naturales y Exactas, Universidad del Valle, Calle 13 # 100-00, Cali, Valle del Cauca, Colombia.

<sup>3</sup> Calima, Fundación para la Investigación de la Biodiversidad y Conservación en el Trópico, Cali, Colombia.

<sup>4</sup> Botanischer Garten und Botanisches Museum, Freie Universität Berlin, Königin-Luise-Straße 6–8, 14195 Berlin, Germany.

\*Corresponding author: r.luecking@bgbm.org

**Keywords:** animals, Colombian biodiversity, insects, global estimate, plants.

### ABSTRACT

This chapter provides a brief overview of the diversity of fungi of Colombia, both in comparison with other organisms and in a global context, including an assessment of the current state of knowledge and an estimate of the actual species richness. The current checklist of fungi of Colombia, including lichenised and fungus-like organisms unrelated to the true *Fungi*, comprises 7,241 species. Assessments of biodiversity-rich countries are usually based on plants and vertebrates, but here we provide an expanded perspective, including fungi and selected invertebrates. The documented diversity of fungi of Colombia is lower than that of plants, vertebrates, and insects, which we attribute to the lack of rigorous taxonomic studies and systematic inventories. Colombia ranks second in plant and vertebrate diversity in the world, trailing only Brazil. However, in terms of known fungal diversity, Colombia is not among the top ten countries and even far behind smaller temperate countries, such as France, Italy, the UK, Germany, and Japan, again highlighting the need for much additional work. Estimates of the existing species richness of fungi of Colombia oscillate between 27,430 and 380,000 species, depending on the extrapolation method. Although these numbers may seem exaggerated, currently reported numbers for the United States already exceed 45,000 and estimates for Mexico predict up to 260,000 species. To catalogue the diversity of the Colombian funga fully, a thorough assessment is needed, including molecular studies of presumably known taxa that may include morphologically cryptic species and effective detection methods for ecologically hidden fungi.


### RESUMEN

Este capítulo proporciona una breve introducción a la diversidad de hongos de Colombia, tanto en comparación con otros organismos como en un contexto global, incluyendo una evaluación del estado actual del conocimiento y una estimación de la riqueza real de las especies. El listado actual de los hongos de Colombia comprende 7.241 especies, incluyendo especies liquenizadas y organismos similares a los hongos pero no relacionados con los hongos verdaderos, tales como los oomicetes y los mixomicetes. Si bien las evaluaciones de los países ricos en biodiversidad generalmente se basan en plantas y vertebrados, aquí brindamos una perspectiva ampliada, la cual incluye los hongos e también invertebrados seleccionados. La diversidad documentada de hongos de Colombia es menor que la de plantas (26,150 especies), vertebrados (7,607) e insectos (68,000), lo que atribuimos a la falta de estudios taxonómicos rigurosos e inventarios sistemáticos. Colombia ocupa el segundo lugar en diversidad de plantas y vertebrados en el mundo, solo detrás de Brasil. Sin embargo, en términos de diversidad fúngica conocida, Colombia no se encuentra entre los diez primeros países del mundo e incluso muy por detrás de algunos países templados más pequeños, como Francia (24,840 especies), Italia (22,700), el Reino Unido (20,000), Alemania (15,295) y Japón (14,000), lo que también destaca la necesidad de mucho trabajo adicional. Las estimaciones de la riqueza de especies existentes de hongos de Colombia oscilan entre 27,430 y 380,000 especies, según el método de extrapolación. Si bien estos números pueden parecer exagerados, los números reportados actualmente para los Estados Unidos ya superan las 45,000 y las estimaciones para México predicen hasta 260,000 especies. Para catalogar completamente la diversidad de los hongos de Colombia, se necesita inventarios exhaustivos, los cuales incluyen estudios moleculares de taxones supuestamente conocidos que pueden incluir especies morfológicamente crípticas, tanto como métodos efectivos de detección de hongos ecológicamente escondidos, a través del metabarcoding ambiental.

INTRODUCTION

Fungi represent one of the three major eukaryotic life forms on our planet, besides plants and animals (Burki *et al.*, 2019). Mycology, the study of fungi, was traditionally treated as a branch of botany, but fungi are more closely related to animals than plants (Figure 1). To further complicate matters, fungi are not a natural group, with fungal life forms convergently evolving several times among eukaryotes. We can even find fungus-like bacteria in the Actinomycetales (Ventura *et al.*, 2007). Although nearly 99% of all fungi are representatives of the kingdom *Fungi*, fungus-like organisms are also found in various other clades (Beakes

& Thines, 2017; Burki *et al.*, 2019; James *et al.*, 2020). These include the *Oomycota* (egg fungi or water moulds), *Hyphochytridiomycota*, and *Labyrinthulomycetes* (slime nets), which form separate lineages within the *Heterokontophyta* (part of the SAR supergroup and related to brown algae and diatoms), as well as the *Phytomyxea* within the *Rhizaria* (also part of the SAR supergroup). Other fungus-like organisms are represented by the *Myxogastria* or *Myxomycetes* (plasmodial slime moulds) within the *Amoebozoa*, the *Acrasida* (cellular slime moulds) within the *Discoba*, and the *Fonticulida* (another group of cellular slime moulds) within the *Cristidiscoidea*, a clade basally related to the *Fungi* and together with the



**FIGURE 1.** Simplified *Tree of Life* highlighting the position of the true *Fungi* (dark brown) and other fungus-like organisms (light brown) relative to animals (*Animalia*) (blue) and true plants (*Viridiplantae*) (dark green) and plant-like organisms (light green) (compiled after Burki *et al.*, 2019 and James *et al.*, 2020, as well as various other sources for specific clades). Unranked major groups above kingdom are indicated to the right.

latter forming the *Holomycota* (Figure 1). The true fungi are named *Fungi* (as a scientific name written in upper case and italics) to distinguish them from other fungus-like organisms, whereas the non-scientific term “fungi” also encompasses all unrelated fungus-like organisms (as a vernacular name written in lower case and standard font; Thines *et al.*, 2020).

The convergent evolution of the fungal lifestyle within the eukaryotes is comparable to the situation in plants (Burki *et al.*, 2019; James *et al.*, 2020): the true plants are grouped in the kingdom *Plantae* (corresponding to the unranked *Viridiplantae*), but other multicellular plant-like organisms are found in the related red (*Rhodophyta*) and brown algae (*Phaeophyceae*), the latter classified within the *Heterokontophyta* and thus more closely related to the *Oomycota*, *Hyphochytridiomycota*, and *Labyrinthulomycetes* (Figure 1). Fungi are sessile (like plants) and heterotrophic organisms that are incapable of photosynthesis (like animals), with various strategies to obtain carbohydrates. These strategies range from the decomposition of dead organic material (saprotrophs) to attacking living organisms (pathogens, parasites, or carnivores) to forming different types of mutualistic symbioses with phototrophic plants, algae, and cyanobacteria (mycorrhizae and lichens) (Dix & Webster, 1995; Mueller *et al.*, 2004; Piepenbring, 2015; Watkinson *et al.*, 2016). Endophytic and endolichenic fungi are ubiquitous as asymptomatic inhabitants of plants, plant-like organisms, and lichens (Rodríguez *et al.*, 2009; U'Ren *et al.*, 2012). They are often classified as symbionts (e.g., in the FUNGuild database; Nguyen *et al.*, 2016), but their biological roles are diverse and often unknown, and they should only be considered symbionts if there is actual evidence for this (Arnold & Lewis, 2005; Bolívar-Anillo *et al.*, 2016; Bastias *et al.*, 2021). Indeed, the diversity of potential life forms of endophytic and endolichenic fungi has been dubbed the “endophytic continuum” (Schulz & Boyle, 2005).

Although mostly invisible (“ecologically hidden”), fungi are crucial components of most ecosystems on earth, being particularly involved in nutrient cycling (Dix & Webster, 1995; Watkinson *et al.*, 2016). In addition, they have played, and continue to play, important roles in ancient and modern cultures (Guzmán *et al.*, 2003; Dugan, 2008). These include their uses as food, in industrial, biotechnological, and pharmaceutical applications, and in biomonitoring and mycoremediation (Nimis *et al.*, 2002; Rhodes, 2014; Hyde *et al.*, 2019; see also Chapter 13).

Fungi also hold several records among living organisms. For instance, species of *Armillaria* (honey mushrooms) form several of the largest known single organisms on earth, with biomasses corresponding to up to three blue whales (Smith *et al.*, 1992; Schmitt & Tatum, 2008; Anderson *et al.*, 2018). The widespread *Schizophyllum commune* (split gill) and *Coprinopsis cinerea* (gray shag), both model fungi, have been shown to develop more than 20,000 mating types or “sexes” (Kothe, 1999; Casselton & Kües, 2007). The lichenised fungus *Buellia frigida* in dry valleys of Antarctica forms thalli that are among the oldest documented organisms on earth, in part estimated at over 6,500 years old (Green *et al.*, 2012).

In terms of known species, *Fungi* currently rank third amongst the three large kingdoms, with 150,000 species (Lücking *et al.*, 2021a), after *Animalia* (animals), with 1.66 million known species (Zhang, 2013), and *Viridiplantae* (plants), with at least 350,000 known species (Christenhusz & Byng, 2016; The Plant List, 2021). *Fungi* are clearly the least well-known of these three kingdoms, given that estimates predict a much higher number of species, between 1.5 and 6.3 million (Blackwell, 2011; Hawksworth & Lücking, 2017; Baldrian *et al.*, 2021). These estimates would place *Fungi* as the second-largest kingdom after animals, including estimates for unknown arthropods (see below), but far ahead of plants. Predictive estimates of global fungal diversity are based on diverse considerations, including a mean global fungi:plant species ratio of between 6:1 and 10:1, but also taking into account accumulating evidence of hidden diversity among presumably known species, with a mean ratio of over 10:1, as well as a wealth of data emerging from environmental sequencing studies (Baldrian *et al.*, 2021; Lücking *et al.*, 2021a). Cataloguing this unknown diversity is the major challenge for mycology in terms of time and resources and how to deal with fungi only known from sequence data (Lücking *et al.*, 2021a). Much of this unknown fungal diversity is expected to be found in tropical regions, such as Colombia.


### DIVERSITY OF FUNGI OF COLOMBIA IN THE TREE OF LIFE

The ColFungi project (Gaya *et al.*, 2021; <https://colfungi.org>) represents the first compilation of data on all groups of fungi of Colombia into a single resource, including nomenclatural and taxonomic checks. This project resulted in a total of 7,241 accepted species of fungi for Colombia, encompassing a high diversity of fungal life forms, including mushrooms and lichen-formers, plant- and animal-pathogenic microfungi, moulds, yeasts, among many others (Figure 2). By itself, the number of 7,241 species is difficult to evaluate, but when put into a taxonomic and geographic context, it illustrates the current state of knowledge of the Colombian fungi.

Country- or region-based comparisons of biodiversity are often presented in the context of global conservation approaches, but these listings are usually limited to plants and vertebrates (mammals, birds, reptiles, amphibians, and fish; e.g., Butler, 2016, 2019), whereas fungi are rarely included in such comparisons. Colombia is currently ranked second among the ten most biodiversity-rich countries in plant and vertebrate species richness, trailing Brazil but ahead of Indonesia, China, Mexico, Peru, Australia, India, Ecuador, and Venezuela (Butler, 2016, 2019). According to these listings, Colombia has 51,220 plant species, 2,053 fish, 1,878 birds, 771 amphibians, 601 reptiles, and 442 mammals, and updated numbers for all vertebrate groups are consistently higher (Table 1). Notably, with 7,241 known species, fungi would rank second in this enumeration for Colombia.

Still, this ranking would be misleading: plants and fungi essentially represent entire kingdoms, whereas, in terms of





**FIGURE 2.** Examples of fungi of Colombia representing a diversity of lifestyles and phenotypes of spore-bearing structures. **A** *Campanella caesia*. **B** *Cordyceps nidus*. **C** *Laetiporus sulphureus*. **D** *Candida* sp. **E** *Auricularia mesenterica*. **F** *Tolypocladium capitatum*. **G** *Laccaria laccata*. **H** *Boletus* sp. **I** *Yoshimuriella peltigera*. **J** *Scleroderma flavidum*. **K** *Panaeolus semiovatus*. **L** *Cyathus striatus*. **M** *Phillipsia domingensis*. **N** *Rhytidhysterium columbiense*. **O** *Cerioporus scutellatus*. **P** *Hericium erinaceus*. **Q** *Hydnopolyporus fimbriatus*. **R** *Leotia lubrica*. **S** *Gaeastrum pectinatum*. **T** *Cora elephas*. **U** *Rhodotorula* sp. **V** *Calostoma cinnabarinum*. **W** *Trametes versicolor*. **X** *Collybia plectophylla*. **Y** *Chlorociboria aeruginascens*. **Z** *Phallus indusiatus*. Photographs: A, B, C, F, G, I, J, N, O, Q, T, W, Y: Robert Lücking. D, U: Mauricio Ramírez. E, L, R, S: Aída Vasco-Palacios. H, V: Bibiana Moncada. K, M, P, Z: Ana Esperanza Franco-Molano. X: Ana Cristina Bolaños.

**TABLE 1.** Updated totals for the number of known species of selected groups of organisms in Colombia. The global totals exclude fossil taxa.

Group	Butler (2016, 2019)	Updated figures	Global total	Proportion of updated figures vs global total	Reference(s) for updated figures
Animals	—	—	1,530,000	—	Zhang (2013); this paper
Vertebrates	5,745	7,607	57,102	13.2%	This paper
Fish	2,053	3,700	32,000	11.6%	Mejia & Acero (2002); Maldonado-Ocampo <i>et al.</i> (2008); Nelson <i>et al.</i> (2016); DoNascimento <i>et al.</i> (2018)
Amphibia	771	836	8,348	10.0%	Acosta-Galvis (2019); Frost (2021)
Reptiles	601	634	11,440	5.5%	Uetz <i>et al.</i> (2021)
Birds	1,878	1,909	9,159	20.8%	Barrowclough <i>et al.</i> (2016); Avendaño <i>et al.</i> (2017)
Mammals	442	528	6,451	8.2%	Burgin <i>et al.</i> (2018); Mammal Diversity Database (2021)
Invertebrates					
Insects	—	68,000	1,000,000	6.8%	This paper
Papilionoidea	—	3,642	18,768	19.4%	Garwood <i>et al.</i> (2021)
Formicidae	—	1,100	13,809	8.0%	Fernández <i>et al.</i> (2019); Bolton (2020)
Staphylinidae	—	800	48,000	1.7%	Newton <i>et al.</i> (2005)
Cicadellidae	—	679	22,000	3.1%	Freytag & Sharkey (2002)
Odonata	—	335	6,000	5.6%	Pérez-Gutiérrez & Palacino-Rodríguez (2011)
Blattaria	—	133	4,330	3.1%	Vélez (2008)
Arachnida	—	1,546	57,126	2.7%	Perafán <i>et al.</i> (2013)
Plants	51,220	26,150	350,000	7.5%	Bernal <i>et al.</i> (2016a, b, 2019); Diazgranados <i>et al.</i> (2021); The Plant List (2021)
Vascular plants	—	24,500	316,000	7.8%	Bernal <i>et al.</i> (2016a, b, 2019); Diazgranados <i>et al.</i> (2021); The Plant List (2021)
Bryophytes	—	1,650	34,500	4.8%	Bernal <i>et al.</i> (2016a, b, 2019); Diazgranados <i>et al.</i> (2021); The Plant List (2021)
Fungi	—	7,241	150,000	4.8%	Gaya <i>et al.</i> (2021); Lücking <i>et al.</i> (2021a); this paper
Ascomycota	—	4,564	93,632	4.9%	Gaya <i>et al.</i> (2021); Kirk (pers. comm. July 2021); this paper
Basidiomycota	—	2,318	50,460	4.7%	Gaya <i>et al.</i> (2021); Kirk (pers. comm. July 2021); this paper

species richness, vertebrates are only a small fraction of the animal kingdom, the diversity of which is concentrated among arthropods (Stork, 2018). Indeed, the known insect diversity of Colombia alone should easily surpass that of vertebrates and likely even plants. For instance, Newton *et al.*, (2005) listed nearly 800 species only of *Staphylinidae* (rove beetles) for the country, out of a total of 48,000 known globally, and Vélez (2008) enumerated 133 species of *Blattaria* (cockroaches), compared to 4,330 worldwide (Table 1). Pérez-Gutiérrez & Palacino-Rodríguez (2011) reported 335 species of *Odonata* (dragonflies), out of a global total of around 6,000. For *Formicidae* (ants), Fernández *et al.*, (2019) listed 1,100 species for Colombia, compared to over 13,800

globally (Bolton, 2020). The most comprehensive and most recent checklist of Colombian butterflies (*Papilionoidea*) compiled data on 3,642 species out of a global number of 18,768 (Garwood *et al.*, 2021). According to Stork (2018), about one million insect species have been named, which means that globally, *Staphylinidae* corresponds to 4.8%, *Papilionoidea* 1.9%, *Formicidae* 1.4%, *Odonata* 0.6%, and *Blattaria* a little over 0.4% of that number. Extrapolating from these proportions in the case of Colombia implies that a total of between 17,000 and 194,000 insect species are likely known from the country, with a median of 43,350 and a mean of 68,000 (Table 1). Although the exact number has not yet been compiled, this extrapolation is far ahead of the



7,241 fungi and even ahead of the actual known number of plant species (see below).

Compared to the 51,220 species of plants reported for Colombia (Butler, 2016, 2019), the number of 7,241 fungal species also appears relatively low, with a fungus:plant species ratio of 1:7. However, the figure of 51,220 plant species goes back to outdated sources, being recently corrected to about 26,150 species in the *Catálogo de Plantas y Líquenes de Colombia* (Bernal et al., 2016a, b, 2019) and the most recent compilation of ColPlantA (Diazgranados et al., 2021). Based on the corrected number, Colombia's fungus:plant species ratio currently amounts to about 1:3.6, still much lower than ratios of between 6:1 and 10:1 obtained from well-studied extra-tropical areas (Hawksworth & Lücking, 2017). In terms of global diversity, Colombia harbours 4.8% of the 150,000 known fungal species and 7.5% of the 350,000 known plant species (Table 1). While these percentages are in the same order of magnitude, they do not consider the much more limited state of knowledge of global fungal diversity.

Fungi and fungus-like organisms are currently arranged into 64 classes (Species Fungorum 2021), of which 36 have been reported from Colombia. The remaining 28 classes are species-poor, with between one and 321 species globally. These remaining classes potentially occur in Colombia but are challenging to detect, as they largely represent ecologically hidden microfungi. Among the 36 classes present in Colombia, 11 are overrepresented compared to global proportions, by a factor of up to 6.5 (Figure 3). The seven most overrepresented classes are *Glomeromycetes* (factor 6.5), *Coniocybomycetes* (3.9), *Mortierellomycetes* (4.3), *Schizosaccharomycetes* (4.0), *Lecanoromycetes* (3.8), *Candelariomycetes* (3.8), and *Arthoniomycetes* (3.6). Four of these seven classes have low species numbers in Colombia, between one and 26, so the observed deviations are not meaningful, but the three remaining classes are represented by higher species numbers: *Lecanoromycetes* (2,167), *Arthoniomycetes* (189) and *Glomeromycetes* (106). The substantial overrepresentation of *Lecanoromycetes* and *Arthoniomycetes* is because these mostly lichen-forming fungi are much better studied than any other larger group of fungi in Colombia (Gaya et al., 2021), so the observed deviations are based on study bias. On the other hand, five important classes that are highly diverse in tropical regions but substantially underrepresented and understudied in Colombia are *Dothideomycetes*, *Sordariomycetes*, *Leotiomycetes*, *Pezizomycetes*, and *Laboulbeniomycetes* (Figure 3).

### DIVERSITY OF FUNGI OF COLOMBIA IN A GLOBAL CONTEXT

Knowledge on organismic diversity is unevenly distributed across the planet, and even the absolute state of knowledge is opposed to the estimated diversity in a region. This scenario is largely because there are more specialists per area working in regions with fewer species per area for almost any group of organisms, particularly fungi (Hawksworth, 2001; Piepenbring et al., 2018). As


a result, in biodiversity-rich countries, such as Colombia, even though the number of fungal species is expected to be higher than in extra-tropical areas, the absolute number of mycological studies is much lower. For instance, a search in the *Web of Science* [<https://www.webofscience.com/wos/woscc/basic-search>], using the string <(ALL=(Fungi)) AND ALL=(Country)>, retrieved around 16,000 results between 1900 and 2020 for the U.K. (England, Northern Ireland, Scotland, Wales), but only 1,155 for Colombia. Unfortunately, reliable estimates of known fungal diversity are difficult to obtain for most countries due to the lack of comprehensive databases or checklists, exceptions being lichenised fungi and macrofungi (Feuerer & Hawksworth, 2007; Gaya et al., 2021). Also, the nature and detail of existing compilations may differ markedly between countries, making comparisons challenging. Nevertheless, we attempted to compile existing data to compare Colombia with selected countries within and outside the tropics (see Gaya et al., 2021; here updated).

The highest number of fungi, including all lifestyles, has been reported for the United States (Table 2), with approximately 45,000 species (Perlmutter & Weakley, 2018; Bates et al., 2018; Esslinger, 2019). However, this impressive number is tentative, as, for non-lichenised fungi, current names and synonyms have not been fully sorted out. After India, with 29,000 species, France remarkably ranks third, with 24,840 species, followed by China (23,927) and Italy (22,700) (Roux, 2012; Onofri et al., 2013; Medardi, 2006; Venturella et al., 2011; Nimis, 2016; Institute of Microbiology, Chinese Academy of Sciences, 2018; Gargominy et al., 2020; Wang et al., 2020; Wei, 2021). We had originally reported 20,500 species for Brazil (Gaya et al., 2021), but a reanalysis of the underlying data, including accepted names vs synonyms, led to a correction, with now 13,950 species (see Table 2). This number is still substantially higher than the 5,719 species reported by Maia et al., (2015), the latter likely an underestimate based on an incomplete list of selected records for particular groups.

Among the countries assessed here, Colombia ranks seventh for lichenised fungi (see Chapter 6). In comparison, the overall number and particularly the number of non-lichenised fungi is comparatively low, with Colombia currently taking 13th place. In terms of species density and species per  $\log_{10}(\text{area})$ , Colombia also ranks comparatively low, particularly when compared to its richness in other groups of organisms, leading us to the question: how many species of fungi can we expect in Colombia?

### HOW MANY SPECIES OF FUNGI ARE THERE IN COLOMBIA?

Predictive estimates of species richness are naturally speculative, especially for ecologically largely hidden organisms and in groups where taxon concepts have not been fully resolved. Several approaches exist to predict species numbers on the basis of known species and other data. One possible approach is the assumption that Colombia's species per  $\log_{10}(\text{area})$  estimate should be



**FIGURE 3.** Comparison of the proportional phylogenetic diversity of *Fungi* and fungus-like organisms at the class level between Colombia and the global fungi. The percentages indicate the number of species per class compared to the corresponding total. Selected classes that are over- or underrepresented in Colombia are marked in dark grey, and the corresponding percentages are indicated. For detailed data, see Suppl. File S1.



**TABLE 2.** Comparison of numbers of known fungal species between selected countries. The number of 16,900 macrofungi for the United States is an estimate based on Bates *et al.* (2018). The corrected total for Brazil (13,950; as compared to 20,500 reported in Gaya *et al.*, 2021) is based on the accepted number of fungal names for Brazil on SpeciesLink (approximately 11,330), minus the number of lichenised species in that database (approximately 1,690), plus the updated number of lichenised species based on Aptroot (2021).

Country / Territory	All fungi	Lichenised fungi	Macro-fungi	Density [species / log <sub>10</sub> (km <sup>2</sup> )	Reference(s)
United States	45,000	4,341	16,900	6,578	Perlmutter & Weakley (2018); Bates <i>et al.</i> (2018); Esslinger (2019)
India	27,000	2,714	—	4,450	Sarbhoj <i>et al.</i> (1996); Singh & Sinha (2010); Manoharachary & Nagaraju (2016); Sinha <i>et al.</i> (2018)
France	24,840	2,917	—	4,276	Roux (2012); Gargominy <i>et al.</i> (2020)
China	23,927	3,050	7,138	3,427	Institute of Microbiology, Chinese Academy of Sciences (2018); Wang <i>et al.</i> 2020; Wei (2021)
Italy	22,700	2,704	4,198	4,143	Onofri <i>et al.</i> (2013); Medardi (2006); Venturella <i>et al.</i> (2011); Nimis (2016); Checklist of Italian Fungi (2021)
United Kingdom	20,000	2,000	3,100	3,343	Legon & Henrici (2005); Smith <i>et al.</i> (2009); Dines & McCarthy (2014); Clubbe <i>et al.</i> (2020)
Germany	15,295	1,946	6,120	2,754	Wirth <i>et al.</i> (2011); DGfM (2021)
Japan	14,000	1,906	4,160	2,510	Hosoya (2006); Katumoto (2009); Ohmura & Kashiwadani (2018)
Brazil	13,950	4,310	2,300	2,013	Lewinsohn & Prado (2005); Maia <i>et al.</i> (2015); Aptroot (2021); SpeciesLink (2021)
Australia	12,870	4,003	4,000	1,869	May <i>et al.</i> (2004); McCarthy (2020)
Canada	12,800	2,262	4,500	1,829	Perlmutter & Weakley (2018); Bates <i>et al.</i> (2018); Esslinger (2019)
Venezuela	8,300	1,801	—	1,392	Marcano <i>et al.</i> (1996); Itturiaga (2000); Hernández-M. (2021)
Mexico	7,632	2,722	2,900	1,212	Guzmán (1998a, b); Cifuentes (2008); Aguirre-Acosta <i>et al.</i> (2014); Herrera-Campos <i>et al.</i> (2014); Bates <i>et al.</i> (2018)
Colombia	7,241	2,559	1,239	1,210	Vasco-Palacios & Franco-Molano (2013); Gaya <i>et al.</i> (2021); Lücking <i>et al.</i> (2021b); this paper
Thailand	7,300	1,292	2,100	1,278	Hywel-Jones & Boonpratuang (2001); Chandrasrikul <i>et al.</i> (2011); Buaruang <i>et al.</i> (2017); Hyde <i>et al.</i> (2018); Index Fungorum (2021)
Malaysia	3,804	—	—	689	Chua <i>et al.</i> (2012)
Costa Rica	3,500	1,740	2,000	743	Halling & Mueller (2005); Mueller <i>et al.</i> (2007)
Bolivia	2,234	1,353	500	370	Piepenbring (2004); Rodríguez de Flakus <i>et al.</i> (2016); Melgarejo-Estrada <i>et al.</i> (2020a, b)
Panama	1,807	325	300	370	Piepenbring (2007)

comparable to that of a well-studied temperate country, such as the United States. Colombia encompasses an area of 1.143 million km<sup>2</sup>, compared to 9.834 million km<sup>2</sup> for the United States. The latter has 6,578 fungal species per log<sub>10</sub>(km<sup>2</sup>), so with log<sub>10</sub>(1,143,000) = 6.058, one would expect 6,578 x 6.058 = 39,850 species for Colombia (Table 3). This figure is about 5.5 times higher than the currently known number. Given that lichen-forming fungi have been estimated at nearly 5,000 species for Colombia

(see Chapter 6), this would result in approximately 35,000 species of non-lichenised fungi. Since about 5,000 species of non-lichenised fungi are known from Colombia (Gaya *et al.*, 2021), nearly 30,000 still need to be catalogued. However, the actual number could be even higher because tropical ecosystems on average support higher small-scale species richness than temperate areas, which means that patterns found in the United States or other largely temperate countries cannot be directly transferred.

**TABLE 3.** Various predictions of the actual species richness of the Colombian fungi using different approaches.

Prediction	Approach
27,430	Adjusted according to a proportion of well-studied class <i>Lecanoromycetes</i>
39,850	Species versus $\log_{10}(\text{area})$ relationship using the USA as a benchmark
105,600 –182,400	Colombian proportion of global estimated richness, using known fungi as a benchmark (4.8%) and 2.2–3.8 million estimated fungal species globally (Hawksworth & Lücking, 2017)
156,900 –261,500	Fungi:plant ratio between 6:1 and 10:1
165,000 –285,000	Colombian proportion of global estimated richness, using known plants as a benchmark (7.5%) and 2.2–3.8 million estimated fungal species globally (Hawksworth & Lücking, 2017)
220,000 –380,000	Colombian proportion of global estimated richness, using known amphibia as a benchmark (10.0%) and 2.2–3.8 million estimated fungal species globally (Hawksworth & Lücking, 2017)

A similar prediction is achieved when considering the phylogenetic diversity of fungi of Colombia at the class level (Figure 3). As outlined above, the largest overrepresented class is *Lecanoromycetes* (30.3% in Colombia versus 7.9% globally), which includes most of the lichenised fungi, a group particularly well-studied in Colombia. If we assume that the global proportion of *Lecanoromycetes* is relatively constant across terrestrial ecosystems dominated by woody plants, then Colombia would be expected to harbour at least 27,430 species of fungi to render the current number of 2,167 *Lecanoromycetes* at a proportion of 7.9%.


Another approach is to look at proportional richness per organism group in Colombia relative to the known global diversity of the group. The observed values range between 4.3% for insects and 20.8% for birds, with others in between, e.g., 7.5% for plants, 8.2% for mammals, and 10.0% for amphibia (Table 1). Relying on well-studied groups and taking into account how species in particular groups can successfully disperse, that is, how sizable average species ranges probably are, for fungi, one could assume values similar to those of plants or amphibia (or butterflies), that is, 7.5%–10.0% (–19.4%). The global estimate of between 2.2 and 3.8 million species (Hawksworth & Lücking, 2017) would give an estimate of (165,000–)220,000–285,000(–380,000) species for Colombia when applying the 7.5%–10.0% range (the 19.4% range probably being too high). Even taking the currently observed 4.8% for known Colombian vs

globally known fungi as a basis, the estimate would range between 105,600 and 182,400 species, still much higher than the estimate based on  $\log_{10}(\text{area})$  extrapolation (Table 3). Such estimates may seem unrealistic, but figures in the same order of magnitude, namely (100,000–)200,000(–260,000) species of fungi, have been proposed for Mexico (Guzmán 1998a, b; Aguirre-Acosta *et al.*, 2014).

The fungus:plant species ratio is another commonly used measure for predictive species richness estimates (Martin, 1951; Hawksworth, 1991; Hawksworth & Lücking, 2017). Empirical data for this measure vary widely, ranging from 1.8:1 to 19.1:1, depending on the ecosystem and sampling methods. Thus, while plants within a given area can be sampled quite accurately, fungal sampling restricted to macroscopically visible fungi and within a limited time frame will naturally and often grossly underestimate fungal diversity, as evidenced by a study performed in Panama (Piepenbring *et al.*, 2012). The ratio also depends on the group of fungi. For example, rust fungi (*Pucciniomycetes*) show lower empirical fungus:plant species ratios between 1:10 and 1:100, with more realistic ratios ranging between 1:4 and 1:20, resulting in estimates for the diversity of rust fungi alone in Colombia of between 2,500 and nearly 13,000 species (Salazar-Yepes & Alves de Carvalho Júnior, 2013). However, these approaches have largely not been tested using molecular approaches, both in terms of taxon concepts in rust fungi and the detection of species not producing spore-bearing structures. In addition, fungus:plant species ratios vary with latitude, higher ratios in temperate areas going along with substantially lower plant diversity, which requires a downwards correction when extrapolating globally (Tedersoo *et al.*, 2014). Estimates may be further misguided by outdated numbers for plant diversity, such as the earlier - reported exaggerated figure of 51,220 plant species for Colombia (see above). Studies that include more sophisticated sampling methods, such as long-term plots and/or metabarcoding, show overall fungus:plant species ratios of between 6:1 and 10:1. However, such studies have been mostly restricted to temperate ecosystems (Hawksworth & Lücking, 2017). Given the number of 26,150 plant species currently known from Colombia, applying the fungus:plant species ratio approach would thus result in 156,900–261,500 species of fungi for Colombia, a figure comparable to that derived from the proportional richness approach (Table 3).

Although a reliable estimate of fungal diversity in Colombia remains speculative, we can safely assume that the predicted figure is perhaps an order of magnitude higher than the currently number of known species. Apart from the sheer size of that number, the more daunting perspective is the effort required to document this largely unknown diversity, requiring a large task force of experts in the different fungal groups and consequent application of modern methods of biodiversity research. Unfortunately, molecular approaches to catalogue fungi are still limited in Colombia. We performed a structured search in the NCBI GenBank [<https://www.ncbi.nlm.nih.gov/genbank>], using the string <Fungi[organism] AND (5.8S[title] OR ITS1[title] OR ITS2[title] OR “internal





**FIGURE 4.** Number of NCBI GenBank accessions obtained using the fungal ITS barcoding marker (A) and number of biosamples from metabarcoding studies using various markers (B), adjusted for area size [relative to  $\log_{10}(\text{area})$ ] for countries across the Americas.

transcribed spacer”[title]) AND country>, for all countries in the Americas, to assess how many accessions of the fungal ITS barcoding marker exist for each country, and then divided the number by  $\log_{10}(\text{area})$ , to filter the area effect. The United States ranks two orders of magnitude ahead of Colombia, whereas Canada, Mexico, Costa Rica, Ecuador, Brazil, and Argentina rank one order of magnitude ahead (Figure 4A). We further queried the NCBI Sequence Read Archive [<https://www.ncbi.nlm.nih.gov/sra>] using the country name and recording the number of resulting biosamples, also dividing them by  $\log_{10}(\text{area})$  to obtain an estimate of existing metabarcoding studies. Again, the United States ranges two orders of magnitude ahead of Colombia and most other countries except Canada (Figure 4B).

Considering that it took about 200 years to arrive at a list of just 7,241 species of fungi known from Colombia using mostly traditional inventory methods, efforts have to be increased by about two orders of magnitude to catalogue an additional at least 30,000 species within a reasonable time frame of 15–20 years. For this to happen, there needs to be a paradigm shift in the political and societal support for this kind of work in Colombia, and across Latin America and other tropical regions in general.

## CONCLUSIONS

The first assembly of a critical checklist of the Colombian fungi is a milestone in the state of knowledge of fungi in the country. However, compared to other organisms and in

a global context, the inventory of fungi of Colombia is still at its onset in terms of expected numbers. Although a good balance of taxonomic groups has been studied to date in Colombia, including a wide array of macro- and microfungi and fungi representing different lifestyles, the development of mycology is not at a level comparable to that in temperate countries, in which non-professional mycology also plays a critical role in performing inventory work. We believe that mycology in Colombia can only develop further if the focus is not only on applied approaches, but more resources are provided for basic scientific studies, allowing the adoption of modern technologies, such as molecular (meta-)barcoding at a broader scale. In parallel, fostering taxonomic specialist groups for the various taxa is necessary, particularly for taxa that remain grossly understudied. The formation of associations, such as the *Asociación Colombiana de Micología* (ASCOLMI) and the *Grupo Colombiano de Lichenología* (GCOL) can be considered an excellent initiative to achieve this goal.

## Acknowledgements

Paul Kirk is warmly thanked for putting data from *Index Fungorum* and *Species Fungorum* at our disposal and for assistance with obtaining data on the checklist of fungi from China. Kevin Hyde, Gareth Jones, and Gary Perlmutter assisted with obtaining references regarding fungal diversity in Thailand, Malaysia, and North America. Meike Piepenbring and Mauricio Salazar-Yepes provided thoughtful reviews that helped to improve this chapter, and Rafael Felipe de Almeida assisted in linguistic editing.

## References

- Acosta-Galvis AR (2019) *Lista de los Anfibios de Colombia: Referencia en Línea V.09.2019*. Available at: <http://www.batrachia.com>
- Aguirre-Acosta E, Ulloa M, Aguilar S, Cifuentes J, Valenzuela R (2014) Biodiversidad de hongos en México. *Revista Mexicana de Biodiversidad*, 85, 76-81. doi: 10.7550/rmb.33649
- Anderson JB, Bruhn JN, Kasimer D, Wang H, Rodrigue N, Smith ML (2018) Clonal evolution and genome stability in a 2500-year-old fungal individual. *Proceedings of the Royal Society B: Biological Science*, 285, 20182233. doi: 10.1098/rspb.2018.2233
- Aptroot A (2021) The four lichen floras of Brazil. International Association for Lichenology 9th Symposium (IAL 9). Brazil. *IAL9 Program and Abstract Book*. 53p.
- Arnold AE, Lewis LC (2005) Ecology and evolution of fungal endophytes, and their roles against insects. In: Vega FE, Blackwell M (eds), *Insect-Fungal Associations: Ecology and Evolution*. New York: Oxford University Press. Pp.74-96.
- Avendaño JE, Bohórquez CI, Rosselli L, Arzuza-Buelvas D, Estela FA, Cuervo AM, Stiles FG, Renjifo LM (2017) Checklist of the birds of Colombia: a synthesis of the state of knowledge since Hilty and Brown (1986). *Ornitología Colombiana*, 2017(16), eA01-eA11.
- Baldrian P, Větrovský T, Lepinay C, Kohout P (2021) High-throughput sequencing view on the magnitude of global fungal diversity. *Fungal Diversity*, Online First. doi: 10.1007/s13225-021-00472-y
- Barrowclough GF, Cracraft J, Klicka J, Zink RM (2016) How many kinds of birds are there and why does it matter? *PLoS ONE*, 11(11), e0166307. doi: 10.1371/journal.pone.0166307
- Bastías DA, Gianoli E, Gundel PE (2021) Fungal endophytes can eliminate the plant growth-defence trade-off. *New Phytologist*, 230(6), 2105-2113. doi: 10.1111/nph.17335
- Bates ST, Miller AN, Macrofungi Collections and Microfungi Collections Consortia (2018) The protochecklist of North American non-lichenised Fungi. *Mycologia*, 110(6), 1222-1348. doi: 10.1080/00275514.2018.1515 410
- Beakes GW, Thines M (2017) *Hyphochytriomycota* and *Oomycota*. In: Archibald J, Simpson A, Slamovits C (eds). *Handbook of the Protists*. Cham: Springer. Pp. 435-505.
- Bernal R, Gradstein SR, Celis M (2016a) *Catálogo de Plantas y Líquenes de Colombia. Vol. I: Líquenes a Lythraceae*. Bogotá: Instituto de Ciencias Naturales, Universidad Nacional de Colombia. 993p.
- Bernal R, Gradstein SR, Celis M (2016b) *Catálogo de Plantas y Líquenes de Colombia. Vol. II: Magnoliaceae a Zygomycetaceae – Especies Introducidas y Cultivadas*. Bogotá, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. 1556p.
- Bernal R, Gradstein SR, Celis M (2019). *Catálogo de Plantas y Líquenes de Colombia*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá. Available at: <http://catalogoplantasdecolombia.unal.edu.co>
- Blackwell M (2011) The Fungi: 1, 2, 3... 5.1 million species? *American Journal of Botany*, 98(3), 426-438. doi: 10.3732/ajb.1000298
- Bolívar-Anillo HJ, Orozco-Sanchez CJ, Lima GS, dos Santos GF (2016) Endophytic microorganisms isolated of plants grown in Colombia: a short review. *Journal of Microbial and Biochemical Technology*, 8, 1-5. doi: 10.4172/1948-5948.1000335
- Bolton B (2020) *An Online Catalogue of the Ants of the World*. Available at: <https://antcat.org>
- Buaruang K, Boonpragob K, Mongkolsuk P, Sangvichien E, Vongshewarat K, Polyiam W, Rangsiruji A, Saipunkaew W, Naksuwankul K, Kalb J, Parnmen S (2017) A new checklist of lichenized fungi occurring in Thailand. *MycKeys*, 23, 1-91. doi: 10.3897/mycokeys.23.12666
- Burgin CJ, Colella JP, Kahn PL, Upham NS (2018) How many species of mammals are there? *Journal of Mammalogy*, 99(1), 1-14. doi: 10.1093/jmammal/gyx147
- Burki F, Roger AJ, Brown MW, Simpson AG (2019) The new tree of eukaryotes. *Trends in Ecology & Evolution*, 35, 43-55. doi: 10.1016/j.tree.2019.08.008
- Butler R (2016) The Top 10 Most Biodiverse Countries. Available at: <https://news.mongabay.com/2016/05/top-10-biodiverse-countries>
- Butler R (2019) Countries with the Highest Biodiversity. Available at: [https://rainforests.mongabay.com/03highest\\_biodiversity.htm](https://rainforests.mongabay.com/03highest_biodiversity.htm)
- Casseltan LA, Kües U (2007) The origin of multiple mating types in the model mushrooms *Coprinopsis cinerea* and *Schizophyllum commune*. In: Heitman J, Kronstad J, Taylor JW, Casseltan LA (eds). *Sex in fungi: Molecular Determination and Evolutionary Implications*. Washington, DC: American Society of Microbiology Press. Pp. 283-300.
- Chandrasrikul A, Suwanarit P, Sangwanit U, Lumyong S, Payapanon A, Sanoamuang N, Pukahuta C, Petcharat V, Sardud U, Duengkhae K, Klinhom U, Thongkantha S, Thongklam S (2011) *Checklist of Mushrooms (Basidiomycetes) in Thailand*. Bangkok: Office of Natural Resources and Environmental Policy and Planning (ONEP).
- Chua LSL, Lee SS, Lee SS, Alias SA, Jones EGB, Zainuddin N, Chan HT (2012) *Checklist of Fungi of Malaysia*, Issue/No. 132. Selangor, Malaysia: Forest Research Institute Malaysia (FRIM).
- Checklist of Italian Fungi (2021) Available at: <http://dbiodbs.univ.trieste.it/global/mush>
- Christenhusz MJ, Byng JW (2016) The number of known plants species in the world and its annual increase. *Phytotaxa*, 261, 201-217. doi: 10.11646/phytotaxa.261.3.1
- Cifuentes J (2008) Hongos. In: Ocegueda S, Llorente J (eds). *Catálogo Taxonómico de Especies de México en Capital Natural de México, Vol. 1: Conocimiento Actual de la Biodiversidad*. México, DF: Conabio. CD1.
- Clubbe C, Ainsworth AM, Bárrios S, Bensusan K, Brodie J, Cannon P, Chapman T, Copeland AI, Corcoran M, Sanchez MD, David JC, Dines T, Gardiner LM, Hamilton MA, Heller T, Hollingsworth PM, Hutchinson N, Llewelyn T, Forrest LL, McGinn KJ, Miles S, O'Donnell K, Woodfield-Pascoe N, Rich TCG, Rumsey F, Sim J, Smith SR, Spence N, Stanworth A, Stroh P, Taylor I, Trivedi C, Twyford AD, Viruel J, Walker K, Wilbraham J, Woodman J, Fay MF (2020) Current knowledge, status, and future for plant and fungal diversity in Great Britain and the UK Overseas Territories. *Plants, People, Planet*, 2, 557-579. doi: 10.1002/ppp3.10142
- DGfM - Deutsche Gesellschaft für Mykologie (2021) *Datenbank der Pilze Deutschlands. Checkliste. Deutsche Gesellschaft für Mykologie*. Bearbeitet von Dämmrich F, Gminder A, Hardtke HJ, Karasch P, Schmidt M, Wehr K. Available at: <http://www.pilze-deutschland.de/organismen>
- Diazgranados M, Allkin R, Ávila FA, Baker W, Bishop D, Black N, Bystrakova N, Carretero J, Castellanos-Castro C, Cely M, Colville L, Cossu T, Davies L, Díaz A, d'Souza J, García F, Gaya E, Graves E, Green L, Haigh A, Hammond DS, Hendigo P, Herrera A, Hillebrecht W, Howes MJ, Iggulden D, Kersey P, Kor L, Mattana E, Milliken W, Mira M, Moat J, Monro A, Morley J, Murphy V, Nesbitt M, Ondo I, Parker L, Phillips S, Piquer M, Pritchard HW, Reis P, Rojas T, Ruff J, Stone M, Tanimoto M, Torres G, Tovar C, Turner R, Taylor R, Utteridge T, Vargas N, Weech MH, White K, Wilkin P, Williams J, Ulian T (2021) *ColPlantA: Colombian Resources for Plants Made Accessible* (2nd ed.). Richmond: Royal Botanic Gardens, Kew.
- Dines TD, McCarthy B (2014) *Global Strategy for Plant Conservation: A Review of the UK's Progress Towards 2020*. Salisbury: Plantlife.
- Dix NJ, Webster J (1995) *Fungal Ecology*. Dordrecht: Springer.
- Dugan FM (2008) *Fungi in the Ancient World: How Mushrooms, Mildews, Molds, and Yeast Shaped the Early Civilisations of Europe, the Mediterranean, and the Near East*. St. Paul: APS Press.
- Esslinger TL (2019) A cumulative checklist for the lichen-forming, lichenicolous and allied fungi of the continental United States and Canada, version 23. *Opuscula Philolichenum*, 18, 102-378.
- Fernández F, Guerrero RJ, Delsinne TD (2019) *Hormigas de Colombia. 1 Ed*. Bogotá: Universidad Nacional de Colombia.
- Feurerer T, Hawksworth DL (2007) Biodiversity of lichens, including a worldwide analysis of checklist data based on Takhtajan's floristic regions. *Biodiversity and Conservation*, 16(1), 85-98. doi: 10.1007/s10531-006-9142-6
- Freytag PH, Sharkey MJ (2002) A preliminary list of the leafhoppers (Homoptera: Cicadellidae) of Colombia. *Biota Colombiana*, 3(2), 235-283. doi: 10.21068/bc.v3i2.117
- Frost DR (2021) *Amphibian Species of the World: An Online Reference. Version 6.1*. American Museum of Natural History, New York, USA. Available at: <https://amphibiansoftheworld.amnh.org/index.php>
- Gargominy O, Tercerie S, Régner C, Ramage T, Dupont P, Daszkiewicz P, Poncet L (2020) *TAXREF v14, Référentiel Taxonomique pour la France: Méthodologie, Mise en œuvre et Diffusion*. Muséum National d'Histoire naturelle, Paris. Rapport UMS PatriNat (OFB-CNRS-MNHN).
- Garwood K, Huertas B, Ríos-Málaver IC, Jaramillo JG (2021) *Mariposas de Colombia Lista de chequeo/Checklist of Colombian Butterflies (Lepidoptera: Papilionoidea)*. *BioButterfly Database*. V1. 300p. Available at: <http://www.butterflycatalogs.com>



- Gaya E, Vasco-Palacios AM, Vargas-Estupiñan N, Lücking R, Carretero J, Sanjuan T, Moncada B, Allkin B, Bolaños-Rojas AC, Castellanos-Castro C, Coca LF, Corrales A, Cossu T, Davis L, dSouza J, Dufat A, Franco-Molano, AE, Garcia F, Gómez-Montoya N, González-Cuellar FE, Hammond D, Herrera A, Jaramillo-Ciro MM, Lasso-Benavides C, Mira MP, Morley J, Motato-Vásquez V, Niño-Fernandez Y, Ortiz-Moreno ML, Peña-Cañón ER, Ramirez-Castrillón M, Rojas T, Ruff J, Simijaca D, Sipman HJM, Soto-Medina E, Torres G, Torres-Andrade PA, Ulian T, White K, Diazgranados M (2021) *ColFungi: Colombian Resources for Fungi Made Accessible*. Richmond: Royal Botanic Gardens, Kew.
- Green TA, Brabyn L, Beard C, Sancho LG (2012) Extremely low lichen growth rates in Taylor Valley, Dry Valleys, continental Antarctica. *Polar Biology*, 35(4), 535-541. doi: 10.1007/s00300-011-1098-7
- Guzmán GN (1998a) Análisis cualitativo y cuantitativo de la diversidad de los hongos en México (Ensayo sobre el inventario fúngico del país). In: Halffter G (ed). *La Diversidad Biológica de Iberoamérica II*. Pp. 111-175. [Acta Zoológica Mexicana, Nueva Serie Vol. Especial]. Xalapa: CYTED e Instituto de Ecología.
- Guzmán GN (1998b) Inventorying the fungi of Mexico. *Biodiversity and Conservation*, 7, 369-384.
- Guzmán GN, Gómez-Pompa A, Allen MF, Fedick SL, Jiménez-Osorio JJ (2003) Fungi in the Maya Culture: Past, Present and Future. In: Fedick S, Allen M, Jiménez-Osorio J, Gomez-Pompa A (eds). *The Lowland Maya Area. Three Millennia at the Human-Wildland Interface*. New York: Food Products Press. Pp. 315-325.
- Halling RE, Mueller GM (2005) *Common Mushrooms of the Talamanca Mountains, Costa Rica*. Bronx: New York Botanical Garden.
- Hawksworth DL (1991) The fungal dimension of biodiversity: magnitude, significance, and conservation. *Mycological Research*, 95(6), 641-655.
- Hawksworth DL (2001) The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research*, 105(12), 1422-1432.
- Hawksworth DL, Lücking R (2017) Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiology Spectrum*, 5(4), FUNK-0052-2016. doi: 10.1128/microbiolspec.FUNK-0052-2016
- Hernández-M JA (2021) A new look at Venezuelan lichen diversity. International Association for Lichenology 9th Symposium (IAL 9). Brazil (online). *IAL9 Program and Abstract Book*, 77.
- Herrera-Campos MA, Lücking R, Pérez-Pérez RE, Miranda-González R, Sánchez N, Barcenás-Peña A, Carrizosa A, Zambrano A, Ryan BD, Nash III TH (2014) Biodiversidad de líquenes en México. *Revista Mexicana de Biodiversidad*, 85, S82-S99. doi: 10.7550/rmb.37003
- Hosoya T (2006) History and the current status of fungal inventory and databasing in Japan. In: Tomida I (ed). *Proceedings of the 7th and 8th symposia on Collection Building and Natural History Studies in Asia and the Pacific Rim*. [National Museum Monographs 34]. Tokyo: National Science Museum. Pp. 51-61.
- Hyde KD, Norphanphoun C, Chen J, Dissanayake AJ, Doilom M, Hongsan S, Jayawardena RS, Jeewon R, Perera RH, Thongbai B, Wanasinghe DN, Wisitrassameewong K, Tibpromma S, Stadler M (2018) Thailand's amazing diversity: up to 96% of fungi in northern Thailand may be novel. *Fungal Diversity*, 93, 215-239. doi: 10.1007/s13225-018-0415-7
- Hyde KD, Xu JC, Lumyong S, Rapior S, Jeewon R, Lumyong S, Niego AGT, Abeywickrama PD, Aluthmuhandiram JVS, Brahmananage RS, Brooks S, Chaiyasen A, Chethana KWT, Chomnunti P, Chepkirui C, Chuankid B, de Silva NI, Doilom M, Faulds C, Gentekaki E, Gopalan V, Kakumyan P, Harishchandra D, Hemachandran H, Hongsan S, Karunarathna A, Karunarathna SC, Khan S, Kumla J, Jayawardena RS, Liu JK, Liu N, Luangharn T, Macabeo APG, Marasinghe DS, Meeks D, Mortimer PE, Mueller P, Nadir S, Nataraja KN, Nontachaiyapoom S, O'Brien M, Penkhruue W, Phukhamsakda C, Ramanan US, Rathnayaka AR, Sadaba RB, Sandargo B, Samarakoon BC, Tennakoon DS, Siva R, Sriprom W, Suryanarayanan TS, Sujarit K, Suwannarach N, Suwunwong T, Thongbai B, Thongklang N, Wei D, Wijesinghe SN, Winiski J, Yan J, Yasanthika E, Stadler M (2019) The amazing potential of fungi, 50 ways we can exploit fungi industrially. *Fungal Diversity*, 97, 1-136. doi: 10.1007/s13225-019-00430-9
- Hywel-Jones NL, Boonpratuang T (2001) *Updated Preliminary Checklist of Fungi Recorded from Thailand*. BIOTEC Yothi-Mycology Laboratory.
- Index Fungorum (2021) *Index Fungorum*. Available from: <http://www.indexfungorum.org/names/Names.asp>
- Institute of Microbiology, Chinese Academy of Sciences (2018) *Checklist of Fungi in China*. Available from: <http://124.16.146.175:8080/checklist/checklist.html>
- Iturriaga T (2000) *Estado Actual del Conocimiento de la Micobiota en Venezuela* (No. 2). Caracas: Ministerio del Ambiente y de los Recursos Naturales Renovables.
- James TY, Stajich JE, Hittinger CT, Rokas A (2020) Toward a fully resolved fungal tree of life. *Annual Review of Microbiology*, 74, 291-313. doi: 10.1146/annurev-micro-022020-051835
- Katamoto K (2009) *List of Fungi Recorded in Japan*. Tokyo: The Kanto Branch of the Mycological Society of Japan.
- Kothe E (1999) Mating types and pheromone recognition in the homobasidiomycete *Schizophyllum commune*. *Fungal Genetics and Biology*, 27(2-3), 146-152.
- Ligon NW, Henrici A (2005) *Checklist of the British and Irish Basidiomycota*. Richmond: Royal Botanic Gardens, Kew. Available at: <http://www.basidiochecklist.info>
- Lewinsohn TM, Prado PI (2005) How many species are there in Brazil? *Conservation Biology*, 19(3), 619-624.
- Lücking R, Aime MC, Robberts B, Miller AN, Aoki T, Ariyawansa HA, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, Hawksworth DL, Hyde KD, Irinyi L, Jeewon R, Johnston PR, Kirk PM, Malosso E, May TW, Meyer W, Nilsson HR, Öpik M, Robert V, Stadler M, Thines M, Vu D, Yurkov AM, Zhang N, Schoch CL (2021a) Fungal taxonomy and sequence-based nomenclature. *Nature Microbiology*, 6, 540-548. doi: 10.1038/s41564-021-00888-x
- Lücking R, Moncada B, Soto-Medina E, Simijaca D, Sipman HJ (2021b). Actualización nomenclatural y taxonómica del Catálogo de Líquenes de Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 45(174), 147-189. doi: 10.18257/raccefyn.1266
- Maia LC, Carvalho AA, Cavalcanti LH, Gugliotta AM, Drechsler-Santos ER, Santiago ALM, Cáceres ME, Gibertoni T, Aptroot A, Giachini AJ, Soares AMS, Silva ACG, Magnago AC, Goto BT, Lira CRS, Motoya CAS, Pires-Zotarelli CLA, daSilva DKA, Soares DJ, Rezende DHC, Luz EDM, Gumboski EL, Wartchow F, Karstedt F, Freire FM, Coutinho FP, Melo GSN, Sotão HM, Baseia IG, Pereira J, Oliveira JJS, Souza JF, Bezerra JL, Araujo LS, Pfenning H, Gusmão LFP, Neves MA, Capelari M, Jaeger MCW, Palacios MP, Menolli N, Medeiros PS, Friedrich RCS, Chikowski R, Pires RM, Melo RF, Silveira RMB, Urrea-Valencia S, Costez VG, Silva VF (2015) Diversity of Brazilian Fungi. *Rodriguesia* 66(4), 1033-1045. doi: 10.1590/2175-7860201566407
- Maldonado-Ocampo JA, Vari RP, Usma JS (2008) Checklist of the freshwater fishes of Colombia. *Biota Colombiana*, 9(2), 143-237.
- Mammal Diversity Database (2021) *Mammal Diversity Database* (Version 1.5) Zenodo. Available at: <http://doi.org/10.5281/zenodo.4139818>
- Manoharachary C, Nagaraju D (2016) New additions to the fungi of India. *Indian Phytopathology*, 69, 93-96.
- Marciano V, Morales-Méndez A, Sipman H, Calderon L (1996) A first checklist of the lichen-forming fungi of the Venezuelan Andes. *Tropical Bryology*, 12, 193-235.
- Martin GW (1951) The numbers of fungi. *Proceedings of the Iowa Academy of Science*, 58(1), 175-178.
- May TW, Milne J, Wood AE, Shingles S, Jones RH, Neish P (2004) *Interactive Catalogue of Australian Fungi, Version 3.0*. Australian Biological Resources Study, Canberra/Royal Botanic Gardens Melbourne.
- McCarthy PM (2020). *Checklist of the Lichens of Australia and its Island Territories*. Australian Biological Resources Study, Canberra. Version 1 March 2020. Available at: <http://www.anbg.gov.au/abrs/lichenlist/introduction.html>
- Medardi G (2006) *Ascomyceti d'Italia*. A.M.B. Centro Studi Micologici, Associazione Micologica Bresadola, Trento.
- Melgarejo-Estrada E, Suarez ME, Rocabado D, Maillard O, Lechner BE (2020a) Checklist of Bolivian Agaricales: 1. Species with dark and pink spore prints. *Mycotaxon*, 134(4), 739. doi: 10.5248/134.739
- Melgarejo-Estrada E, Rocabado D, Suárez ME, Maillard O, Lechner BE (2020b) Checklist of Bolivian Agaricales: 2. Species with white or pale spore prints. *Mycotaxon*, 135(1), 233. doi: 10.5248/135.233
- Mejia LS, Acero P (2002) *Libro Rojo de Peces Marinos de Colombia*. Bogotá: INVEMAR, Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Ministerio de Medio Ambiente.
- Mueller GM, Foster M, Bills G (2004) *Biodiversity of Fungi. Inventory and Monitoring Methods*. Burlington, MA: Academic Press (Elsevier)

- Mueller GM, Schmit JP, Leacock PR, Buyck B, Cifuentes J, Desjardin DE, Halling RE, Hjortstam K, Iturriaga T, Larsson KH, Lodge DJ (2007) Global diversity and distribution of macrofungi. *Biodiversity and Conservation*, 16(1), 37-48. doi: 10.1007/s10531-006-9108-8
- Do Nascimento C, Herrera-Collazos EE, Maldonado-Ocampo JA (2018) *Lista de Especies de Peces de Agua Dulce de Colombia / Checklist of the Freshwater Fishes of Colombia*. V2.10. Bogotá: Asociación Colombiana de Ictiólogos. Dataset/Checklist. Available at: <http://doi.org/10.15472/numrso>
- Nelson JS, Grande TC, Wilson MV (2016) *Fishes of the World*. Hoboken, NJ: John Wiley & Sons.
- Newton AF, Gutiérrez-Chacón C, Chandler DS (2005) Checklist of the Staphylinidae (Coleoptera) of Colombia. *Biota Colombiana*, 6(1), 1-72.
- Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG (2016) FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, 20, 241-248. doi: 10.1016/j.funeco.2015.06.006
- Nimis PL (2016) *The Lichens of Italy. A Second Annotated Catalogue*. Trieste: EUT Edizioni Università di Trieste.
- Nimis PL, Scheidegger C, Wolseley PA (2002) *Monitoring with Lichens – Monitoring Lichens*. Dordrecht: Springer
- Ohmura Y, Kashiwadani H (2018) *Checklist of Lichens and Allied Fungi of Japan*. Tokyo: National Museum of Nature and Science.
- Onofri S, Bernicchia A, Filipello-Marchisio V, Padovan F, Perini C, Ripa C, Savino E, Venturella G, Vizzini A, Zotti M, Zucconi L (2013) *Checklist of the Macrobasidiomycetes of Italy / Checklist dei Macrobasidiomiceti d'Italia*. Available at: <http://dryades.units.it/macrobasidiomiceti>
- Perafán C, Sabogal A, Moreno-González JA, García-Rincón A, Luna-Sarmiento D, Romero-Ortiz C, Flórez E (2013) Diagnóstico del estado actual de la fauna de arácnidos y de su gestión en Colombia. Simposio 8: Aracnología. In: Rueda-Ramírez D, Torrado-León E, Becerra EH (eds). *Memorias Congreso Colombiano de Entomología*. 40 Congreso Socolen. Socolen, Bogotá, DC: Sociedad Colombiana de Entomología. Pp. 275-335.
- Pérez-Gutiérrez LA, Palacino-Rodríguez F (2011). Updated checklist of the Odonata known from Colombia. *Odonatologica*, 40(3), 1-203.
- Perlmutter GB, Weakley AS (2018) USDA Plants Database Lichen Update, Phase II. Greensboro, NC: NRCS – East National Technology Support Center
- Piepenbring M (2004) 3.2.1. Fungi. In: Ibsch PL, Mérida G (eds). *Biodiversity: the Richness of Bolivia, State of Knowledge and Conservation*. Santa Cruz de la Sierra: Editorial FAN. Pp. 89-94.
- Piepenbring M (2007) Inventorying the fungi of Panama. *Biodiversity and Conservation*, 16(1), 73-84.
- Piepenbring M (2015) *Introduction to Mycology in the Tropics [Introducción a la Micología en los Trópicos]*. St. Paul: APS Press.
- Piepenbring M, Hofmann TA, Unterseher M, Kost G (2012) Species richness of plants and fungi in western Panama: towards a fungal inventory in the tropics. *Biodiversity and Conservation*, 21(9), 2181-2193. doi: 10.1007/s10531-011-0213-y
- Piepenbring M, Lotz-Winter H, Hofmann TA (2018) Incentives and challenges for mycologists in the tropics. *Biosystematics and Ecology Series*, 34, 481-514.
- Rhodes CJ (2014) Mycoremediation (bioremediation with fungi) – growing mushrooms to clean the earth. *Chemical Speciation & Bioavailability*, 26(3), 196-198. doi: 10.3184/095422914X14047407349335
- Rodríguez RJ, White Jr JF, Arnold AE, Redman ARA (2009) Fungal endophytes: diversity and functional roles. *New Phytologist*, 182(2), 314-330. doi: 10.1111/j.1469-8137.2009.02773.x
- Rodríguez de Flakus P, Kukwa M, Etayo J, Lücking R, Meneses RI, Plata ER, Stanton D, Truong C, Vargas R, Flakus A (2016) *Preliminary Catalogue of Lichens and Lichenicolous Fungi from Bolivia*. Available at: <https://bio.botany.pl/lichens-bolivia/en,strona.catalogue.5.htm>
- Roux C (2012) Liste des lichens et champignons lichénicoles de France. *Bulletin de la Société Linnéenne de Provence, N Spécial* 16, 1-220.
- Salazar-Yepes M, Alves de Carvalho Júnior A (2013) *Uredinales* (rust fungi) biota of the Parque Nacional do Itatiaia, Brazil: an analysis of composition, species diversity and altitudinal distribution. *Caldasia* 35(1), 165-176.
- Sarbhoy AK, Agarwal DK, Varshney JL (1996) *Fungi of India 1982-1992*. New Delhi: CBS Publishers and Distributors.
- Schmitt CL, Tatum ML (2008) *The Malheur National Forest: Location of the World's Largest Living Organism (the Humongous Fungus)*. Washington, DC: Forest Service, US Department of Agriculture.
- Singh KP, Sinha GP (2010) *Indian Lichens: An Annotated Checklist*. Kolkata: Botanical Survey of India.
- Sinha GP, Nayaka S, Joseph S (2018) Additions to the checklist of Indian lichens after 2010. *Cryptogam Biodiversity and Assessment, Special Volume*, 2018, 197-206. doi: 10.21756/cab.esp16
- Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA (2009) *The Lichens of Great Britain and Ireland*. London: British Lichen Society.
- Smith ML, Bruhn JM, Anderson JB (1992) The fungus *Armillaria bulbosa* is among the largest and oldest living organisms. *Nature*, 356(6368), 428-431. doi: 10.1038/356428a0
- Species Fungorum (2021) *Species Fungorum*. Available at: <http://www.speciesfungorum.org>
- SpeciesLink (2021) *SpeciesLink*. Available at: <https://specieslink.net>
- Stork NE (2018) How many species of insects and other terrestrial arthropods are there on earth? *Annual Review of Entomology*, 63, 31-45. doi: 10.1146/annurev-ento-020117-043348
- Tedersoo L, Bahram M, Pölme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS, Larsson KH, Kohout P, Hosaka K, Hiiesalu K, Henkel TW, Harend H, Guo L, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen X, Buegger F, Brearley FQ, Bonito G, Anslan S, Abell S, Abarenkov K (2014) Global diversity and geography of soil fungi. *Science*, 346(6213), 1078. doi: 10.1126/science.1256688
- The Plant List (2021) The Plant List. Available at: <http://www.theplantlist.org>
- Thines M, Aoki T, Crous PW, Hyde KD, Lücking R, Malosso E, May TW, Miller AN, Redhead SA, Yurkov AM, Hawksworth DL (2020) Setting scientific names at all taxonomic ranks in italics facilitates their quick recognition in scientific papers. *IMA Fungus*, 11(1), 1-5. doi: 10.1186/s43008-020-00048-6
- Uetz P, Freed P, Aguilar R, Hošek J (2021) *The Reptile Database*. Available at: <http://www.reptile-database.org>
- U'Ren JM, Lutzoni F, Miadlikowska J, Laetsch AD, Arnold AE (2012) Host and geographic structure of endophytic and endolichenic fungi at a continental scale. *American Journal of Botany*, 99(5), 898-914. doi: 10.3732/ajb.1100459
- Vasco-Palacios AM, Franco-Molano AE (2013) Diversity of Colombian macrofungi (Ascomycota–Basidiomycota). *Mycotaxon*, 121(1), 100-158.
- Vélez, A. (2008). Checklist of Colombian cockroaches (Dictyoptera, Blattaria). *Biota Colombiana*, 9(1), 21-38.
- Ventura M, Canchaya C, Tauch A, Chandra G, Fitzgerald GF, Chater KF, van Sinderen D (2007) Genomics of Actinobacteria: tracing the evolutionary history of an ancient phylum. *Microbiology and Molecular Biology Reviews*, 71(3), 495-548.
- Venturella G, Altobelli E, Bernicchia A, Di Piazza S, Donnini D, Gargano ML, Gorjón SP, Granito VM, Lantieri A, Lunghini D, Montemartini A (2011) Fungal biodiversity and *in situ* conservation in Italy. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, 145(4), 950-957.
- Watkinson SC, Boddy L, Money NP (2016) *The Fungi*. Third Edition. Oxford, UK: Academic Press (Elsevier).
- Wang K, Zhao M, Su J, Yang L, Deng H, Wang Y, Wu H, Li Y, Wu H, Wei X, Wei T, Cai L, Yao Y (2020) The use of Checklist of Fungi in China database in the red list assessment of macrofungi in China. *Biodiversity Science*, 28(1), 74-98. doi: 10.17520/biods.2019163
- Wei JC (2021) *The Enumeration of Lichenized Fungi in China*. Beijing: China Forestry Publishing House.
- Wirth V, Hauck M, Brackel WV, Cezanne R, de Bruyn U, Dürhammer O, Eichler M, Gnüchtel A, Litterski B, Otte V, Schiefelbein U (2011) *Checklist of Lichens and Lichenicolous Fungi in Germany*, Version 2. Göttingen: Georg August University.
- Zhang ZQ (2013) Animal biodiversity: An update of classification and diversity in 2013. *Zootaxa*, 3703(1), 5-11. doi: 10.11646/zootaxa.3703.1.3



# Catalogue of Fungi of Colombia

Royal  
Botanic  
Gardens **Kew**

Edited by  
**Rafael F. de Almeida**  
**Robert Lücking**  
**Aída Vasco-Palacios**  
**Ester Gaya**  
**Mauricio Diazgranados**



© The Board of Trustees of the Royal Botanic Gardens, Kew 2022

Text © The Board of Trustees of the Royal Botanic Gardens, Kew, and the Instituto de Investigación de Recursos Biológicos Alexander von Humboldt

Images © The Board of Trustees of the Royal Botanic Gardens, Kew, unless otherwise stated

The authors have asserted their right to be identified as the authors of this work in accordance with the Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form, or by any means, electronic, mechanical, photocopying, recording or otherwise, without written permission of the publisher unless in accordance with the provisions of the Copyright Designs and Patents Act 1988.

Great care has been taken to maintain the accuracy of the information contained in this work. However, neither the publisher nor the authors can be held responsible for any consequences arising from use of the information contained herein. The views expressed in this work are those of the authors and do not necessarily reflect those of the publisher or of the Board of Trustees of the Royal Botanic Gardens, Kew, or Instituto Humboldt.

First published in 2022 by Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, UK, in association with the Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Calle 28A #16-29, Bogotá, Colombia.

[www.kew.org](http://www.kew.org)

ISBN 978-1-84246-790-9

eISBN 978-1-84246-791-6

Distributed on behalf of the Royal Botanic Gardens, Kew in North America by the University of Chicago Press, 1427 East 60<sup>th</sup> Street, Chicago, IL 606037, USA.

British Library Cataloguing in Publication Data

A catalogue record for this book is available from the British Library.

Design: Nicola Thompson, Culver Design

Production Manager: Jo Pillai

Proofreading: Sharon Whitehead

Printed in Great Britain by Halstan & Co Ltd

For information or to purchase all Kew titles please visit [shop.kew.org/kewbooksonline](http://shop.kew.org/kewbooksonline) or email [publishing@kew.org](mailto:publishing@kew.org)

For Instituto Humboldt publications visit <http://repository.humboldt.org.co/>

Kew's mission is to understand and protect plants and fungi, for the wellbeing of people and the future of all life on Earth.

Kew receives approximately one third of its funding from Government through the Department for Environment, Food and Rural Affairs (Defra). All other funding needed to support Kew's vital work comes from members, foundations, donors and commercial activities, including book sales.

Corresponding author: Mauricio Diazgranados, Herbarium, Royal Botanic Gardens Kew, Richmond TW9 3AE, United Kingdom

E-mail: [M.Diazgranados@kew.org](mailto:M.Diazgranados@kew.org)

**Front cover photo credit:** Fungi mosaic made from photographs included in this book, based on the photograph of the species *Cyptotrama asprata* (Basidiomycota) by Nataly Gómez-Montoya. Design cover by R.F. Almeida; **back cover background photo credit:** *Ganoderma* (Mauricio Diazgranados © RBG Kew); **back cover fungi photo credits:** *Lepidostroma calocerum* (Robert Lücking ©), *Tetrapyrgos alba* (Nataly Gómez-Montoya © RBG Kew), *Pleurotus djamor* (Nataly Gómez-Montoya © RBG Kew), *Cordyceps nidus* (Robert Lücking ©), *Dacryopinax spathularia* (Kent Brothers © RBG Kew), *Cyathus striatus* (Aída Marcela Vasco-Palacios ©), *Aseroe rubra* (Nataly Gómez-Montoya © RBG Kew), *Xylaria multiplex* (Robert Lücking ©), *Phyllobaeis imbricata* (Robert Lücking ©), *Cladonia secundana* (Robert Lücking ©)

Funded by:



MINISTERIO DE CIENCIA,  
TECNOLOGÍA E INNOVACIÓN

Implemented by:





# Contents

	Preface	7
	Executive Summary	11
	Foreword	13
	Acknowledgements	15
	List of Contributors	17
Chapter 1	The Useful Plants and Fungi of Colombia (UPFC) project: delivering mycological knowledge to support conservation and sustainable development	19
Chapter 2	Two Centuries of Mycological History in Colombia	33
Chapter 3	Diversity of Fungi of Colombia	45
Chapter 4	Diversity of Basidiomycota in Colombia	49
Chapter 5	Diversity of Non-Lichenised Macro-Ascomycota of Colombia	67
Chapter 6	Diversity, Ecogeography, and Importance of Lichens of Colombia	77
Chapter 7	Diversity of Environmental Yeasts of Colombia: A Systematic Review	93
Chapter 8	Micromycetes of Colombia: Focusing on the Hidden Diversity	105
Chapter 9	Diversity of Rust and Smut Fungi of Colombia	113
Chapter 10	A Critical Assessment of Biogeographic Distribution Patterns of Colombian Fungi	121
Chapter 11	Diversity, Functional Groups, and Community Structure of Fungi of Colombia	139
Chapter 12	Useful Fungi of Colombia	151
Chapter 13	Biotechnology of Fungi of Colombia	165
Chapter 14	Fungal Conservation in Colombia	175
Chapter 15	Fungi in Colombian and International Biological Collections	189
Chapter 16	Annotated Checklist of Fungi of Colombia	209
	Checklist	216
	Laminae	427
	Index of Families	459
	Index of Genera	463
	Alphabetical List of Synonyms	473
	Alphabetical List of Misapplied and Doubtful Names	507



The *Catalogue of Fungi of Colombia* is the first comprehensive listing of the known Colombian fungi. Compiled by a team of Colombian and international mycologists from the Royal Botanic Gardens, Kew, the Humboldt Institute and numerous partner institutions, it consolidates expert-generated information linked and accessible through an online portal (*ColFungi*). The checklist is accompanied by 15 chapters written by specialists, providing perspectives on the state of knowledge on the Colombian fungi, covering a range of topics, from the diversity of the main groups of fungi and the history of mycological studies in this country, to aspects of the biogeography, ecology, biotechnology, conservation, and uses of Colombian fungi and their presence in national and international biological collections. The Catalogue is further enriched by diverse supplementary material, allowing users to explore further open questions and opportunities, to develop new ideas on the use of fungi and their conservation, and to foster social and environmental awareness.



**Rafael F. de Almeida** is a biologist with expertise in Taxonomy and Systematics. He served as research editor for the Useful Plants and Fungi of Colombia (UPFC) project. **Robert Lücking** is a lichenologist at the Botanischer Garten und Botanisches Museum in Berlin, where he works as curator for lichens, fungi, and bryophytes, with emphasis on the fungi of South America. **Aída Vasco-Palacios** is a professor at the Faculty of Microbiology at the University of Antioquia, where she studies fungal biodiversity in Colombia and its biotechnological potential. Also, she is the president of the Colombian Association of Mycologists. **Ester Gaya** is a Senior Research Leader in the Trait Diversity and Function department at the Royal Botanic Gardens, Kew (RBG Kew), where she leads the Comparative Fungal Biology team. **Mauricio Diazgranados** is a research leader in the Ecosystem Stewardship department at RBG Kew. He focuses on investigating how useful plants and fungi can contribute to developing nature-based solutions to tackle environmental and societal challenges. He was also the principal investigator of the UPFC project.

[shop.kew.org/kewbooksonline](https://shop.kew.org/kewbooksonline)

ISBN 978-1-84246-790-9



9 781842 467909