

Name:_____

NetID:_____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined (for n a power of 4) by

$$\begin{aligned}f(1) &= 0 \\f(n) &= 2f(n/4) + n \text{ for } n \geq 4\end{aligned}$$

Your partner has already figured out that

$$f(n) = 2^k f(n/4^k) + n \sum_{p=0}^{k-1} 1/2^p$$

Finish finding the closed form for $f(n)$ assuming that n is a power of 4. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function F defined (for n a power of 2) by

$$\begin{aligned} F(2) &= c \\ F(n) &= F(n/2) + n \text{ for } n \geq 4 \end{aligned}$$

Your partner has already figured out that

$$F(n) = F(n/2^k) + \sum_{i=0}^{k-1} n \frac{1}{2^i}$$

Finish finding the closed form for F . Show your work and simplify your answer.

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function f defined by

$$\begin{aligned} f(0) &= f(1) = 3 \\ f(n) &= 5f(n-2) + d, \text{ for } n \geq 2 \end{aligned}$$

where d is a constant. Express $f(n)$ in terms of $f(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The chromatic number of the
4-dimensional hypercube Q_4

2 ☐3 ☐4 ☐5 ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined (for n a power of 4) by

$$f(1) = 0$$

$$f(n) = 2f(n/4) + n \text{ for } n \geq 4$$

Express $f(n)$ in terms of $f(n/4^{13})$ (assuming n is large enough that this input hasn't reached the base case). Express your answer using a summation and show your work. Do **not** finish the process of finding the closed form for $f(n)$.

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 3) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 3g(n/3) + n \text{ for } n \geq 3 \end{aligned}$$

Express $g(n)$ in terms of $g(n/3^3)$ (where $n \geq 27$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The number of nodes in the
4-dimensional hypercube Q_4

4 ☐16 ☐32 ☐64 ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 4) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 2g(n/4) + n \text{ for } n \geq 4 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 2^k g(n/4^k) + n \sum_{p=0}^{k-1} \frac{1}{2^p}$$

Finish finding the closed form for $g(n)$ assuming that n is a power of 4. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= 1 \\ g(n) &= 4g(n/2) + n^2 \text{ for } n \geq 2 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 4^k g(n/2^k) + kn^2$$

Finish finding the closed form for g . Show your work and simplify your answer.

2. (2 points) Check the (single) box that best characterizes each item.

The Fibonacci numbers can be defined recursively by $F(0) = 0$, $F(1) = 1$, and $F(n) = F(n-1) + F(n-2)$ for all integers ...

$n \geq 0$ ☐

$n \geq 1$ ☐

$n \geq 2$ ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned}g(1) &= 3 \\g(n) &= 4g(n/2) + n \text{ for } n \geq 2\end{aligned}$$

Your partner has already figured out that

$$g(n) = 4^k g(n/2^k) + \sum_{p=0}^{k-1} n2^p$$

Finish finding the closed form for $g(n)$ assuming that n is a power of 2. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name:_____

NetID:_____ Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (8 points) Suppose we have a function g defined by

$$\begin{aligned}g(0) &= g(1) = c \\g(n) &= kg(n-2) + n^2, \text{ for } n \geq 2\end{aligned}$$

where k and c are constants. Express $g(n)$ in terms of $g(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is such that $f(n) = n!$. Give a recursive definition of f

Name: _____

NetID: _____ Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (8 points) Suppose we have a function f defined by

$$\begin{aligned} f(1) &= 5 \\ f(n) &= 3f(n-1) + n^2 \text{ for } n \geq 2 \end{aligned}$$

Express $f(n)$ in terms of $f(n-3)$ (where $n \geq 4$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

2. (2 points) Suppose that G_0 is the graph consisting of a single vertex. Also suppose that the graph G_n consists of a copy of G_{n-1} plus an extra vertex v and edges joining v to each vertex in G_{n-1} . Give a clear picture or precise description of G_4 .

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function F defined (for n a power of 2) by

$$F(2) = 17$$

$$F(n) = 3F(n/2), \text{ for } n \geq 4$$

Use unrolling to find the closed form for F . Show your work and simplify your answer.

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 4g(n/2) + d \text{ for } n \geq 2 \end{aligned}$$

Express $g(n)$ in terms of $g(n/2^3)$ (where $n \geq 8$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

$f(n) = n!$ can be defined recursively by
 $f(0) = 1$, and $f(n+1) = (n+1)f(n)$
 for all integers ...

$n \geq 0$ ☐

$n \geq 1$ ☐

$n \geq 2$ ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 3) by

$$\begin{aligned} g(9) &= 5 \\ g(n) &= 3g(n/3) + n \text{ for } n \geq 27 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 3^k g(n/3^k) + kn$$

Finish finding the closed form for g . Show your work and simplify your answer.

2. (2 points) Suppose that G_0 is the graph consisting of a single vertex. Also suppose that the graph G_n consists of a copy of G_{n-1} plus an extra vertex v and edges joining v to each vertex in G_{n-1} . Give a clear picture or precise description of G_4 .

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function F defined (for n a power of 3) by

$$\begin{aligned} F(1) &= 5 \\ F(n) &= 3F(n/3) + 7 \text{ for } n \geq 3 \end{aligned}$$

Your partner has already figured out that

$$F(n) = 3^k F(n/3^k) + 7 \sum_{p=0}^{k-1} 3^p$$

Finish finding the closed form for F . Show your work and simplify your answer. Recall the following useful closed form (for $r \neq 1$): $\sum_{k=0}^n r^k = \frac{r^{n+1} - 1}{r - 1}$

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 4g(n/2) + n \text{ for } n \geq 2 \end{aligned}$$

Express $g(n)$ in terms of $g(n/2^3)$ (where $n \geq 8$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The diameter of the
4-dimensional hypercube Q_4

1

☐

2

☐

4

☐

16

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function f defined (for n a power of 2) by

$$\begin{aligned} f(1) &= 5 \\ f(n) &= 3f(n/2) + n^2 \text{ for } n \geq 2 \end{aligned}$$

Express $f(n)$ in terms of $f(n/2^3)$ (where $n \geq 8$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The n -dimensional

hypercube Q_n has an Euler circuit.

always ☐ sometimes ☐ never ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined by

$$\begin{aligned} f(0) &= f(1) = 3 \\ f(n) &= 5f(n-2) + d, \text{ for } n \geq 2 \end{aligned}$$

where d is a constant. Your partner has already figured out that

$$f(n) = 5^k f(n-2k) + \sum_{p=0}^{k-1} d5^p$$

Finish finding the closed form for $f(n)$ assuming that n is even. Show your work and simplify your answer. Recall the following useful closed form (for $r \neq 1$): $\sum_{k=0}^n r^k = \frac{r^{n+1} - 1}{r - 1}$

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function f defined by

$$\begin{aligned} f(0) &= f(1) = 3 \\ f(n) &= 5f(n-2) + d, \text{ for } n \geq 2 \end{aligned}$$

where d is a constant. Express $f(n)$ in terms of $f(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

$f(n) = n!$ can be defined recursively
by $f(0) = 1$, and $f(n) = nf(n-1)$
for all integers ...

$n \geq 0$ ☐

$n \geq 1$ ☐

$n \geq 2$ ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 3) by

$$\begin{aligned}g(1) &= c \\g(n) &= 3g(n/3) + n \text{ for } n \geq 3\end{aligned}$$

Express $g(n)$ in terms of $g(n/3^3)$ (where $n \geq 27$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is such that $f(n) = n^2$. Give a recursive definition of f

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 4) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 2g(n/4) + n \text{ for } n \geq 4 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 2^k g(n/4^k) + n \sum_{p=0}^{k-1} \frac{1}{2^p}$$

Finish finding the closed form for $f(n)$ assuming that n is a power of 4. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name:_____

NetID:_____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 4g(n/2) + n \text{ for } n \geq 2 \end{aligned}$$

Your partner has already figured out that

$$g(n) = 4^k g(n/2^k) + n \sum_{p=0}^{k-1} 2^p$$

Finish finding the closed form for $g(n)$ assuming that n is a power of 2. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined by

$$\begin{aligned} g(0) &= g(1) = c \\ g(n) &= kg(n-2) + n^2, \text{ for } n \geq 2 \end{aligned}$$

where k and c are constants. Express $g(n)$ in terms of $g(n-6)$ (where $n \geq 6$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The number of nodes in the
4-dimensional hypercube Q_4

4 ☐16 ☐32 ☐64 ☐

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined (for n a power of 4) by

$$\begin{aligned}f(1) &= 0 \\f(n) &= 2f(n/4) + n \text{ for } n \geq 4\end{aligned}$$

Express $f(n)$ in terms of $f(n/4^{13})$ (assuming n is large enough that this input hasn't reached the base case). Express your answer using a summation and show your work. Do **not** finish the process of finding the closed form for $f(n)$.

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 2) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 4g(n/2) + n^2 \text{ for } n \geq 2 \end{aligned}$$

Express $g(n)$ in terms of $g(n/2^3)$ (where $n \geq 8$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The Fibonacci numbers can be defined recursively by $F(0) = 0$, $F(1) = 1$, and $F(n+1) = F(n) + F(n-1)$ for all integers ...

$n \geq 0$ ☐

$n \geq 1$ ☐

$n \geq 2$ ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function f defined by

$$f(1) = 5$$

$$f(n) = 3f(n-1) + n^2 \text{ for } n \geq 2$$

Express $f(n)$ in terms of $f(n-3)$ (where $n \geq 4$). Show your work and simplify your answer. You do **not** need to find a closed form for $f(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The Fibonacci numbers can be defined

recursively by $F(0) = 0$, $F(1) = 1$, and

$F(n+2) = F(n) + F(n+1)$ for

all integers ...

 $n \geq 0$ ☐ $n \geq 1$ ☐ $n \geq 2$ ☐

Name:_____

NetID:_____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function g defined (for n a power of 2) by

$$g(1) = 3$$

$$g(n) = 4g(n/2) + n \text{ for } n \geq 2$$

Your partner has already figured out that

$$g(n) = 4^k g(n/2^k) + \sum_{p=0}^{k-1} n2^p$$

Finish finding the closed form for $g(n)$ assuming that n is a power of 2. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name:_____

NetID:_____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

(10 points) Suppose we have a function f defined (for n a power of 4) by

$$\begin{aligned} f(1) &= 0 \\ f(n) &= 2f(n/4) + n \text{ for } n \geq 4 \end{aligned}$$

Your partner has already figured out that

$$f(n) = 2^k f(n/4^k) + n \sum_{p=0}^{k-1} 1/2^p$$

Finish finding the closed form for $f(n)$ assuming that n is a power of 4. Show your work and simplify your answer. Recall that $\log_b n = (\log_a n)(\log_b a)$.

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (8 points) Suppose we have a function g defined (for n a power of 4) by

$$\begin{aligned} g(1) &= c \\ g(n) &= 2g(n/4) + n \text{ for } n \geq 4 \end{aligned}$$

Express $g(n)$ in terms of $g(n/4^3)$ (where $n \geq 64$). Show your work and simplify your answer. You do **not** need to find a closed form for $g(n)$.

2. (2 points) Check the (single) box that best characterizes each item.

The chromatic number of the
4-dimensional hypercube Q_4

2 ☐3 ☐4 ☐5 ☐