

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 12-letter strings can be made by rearranging the letters in the word ‘‘apalachicola’’? Show your work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = 2x$ then the set of all even
integers is the ____ of f .

domain
image

co-domain
none of these

$f : \mathbb{N}^2 \rightarrow \mathbb{N}$
 $f(p, q) = pq$

onto not onto not a function

$g : (\mathbb{Z}^+)^2 \rightarrow \mathbb{Z}^+$
 $g(x, y) = \gcd(x, y)$

one-to-one not one-to-one not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there is a true false color that appears on exactly two mailboxes.

$\exists y \in \mathbb{Z}, \forall x \in \mathbb{Z}, y \leq x$

true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Hermione Grainger has 7000 socks in her magically expanding drawer. The socks are colored purple, magenta, and shocking pink. How many socks must she pull out of the drawer before she is guaranteed to have two socks of the same color. Briefly justify your answer.

2. (10 points) Check the (single) box that best characterizes each item.

A function is onto if and only if its image is the same as its co-domain. true false

$g : \mathbb{R} \rightarrow [-1, 1]$
 $g(x) = \sin(x)$ onto not onto not a function

$g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$
 $g(x, y) = (y, 3x)$ one-to-one not one-to-one not a function

Each elf has exactly one gift: charm, strength, or stamina. If there are 10 elves, the pigeonhole principle says that at least three elves have charm. true false

$\exists t \in \mathbb{N}, \forall p \in \mathbb{Z}^+, \gcd(p, t) = p$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) 8 presidential candidates (including Bernie and Hilary) need to line up for a photo. The new editor would like Bernie and Hilary to stand next to each other. How many different ways can we arrange the eight people?

2. (10 points) Check the (single) box that best characterizes each item.

Suppose $f : A \rightarrow B$. For
all $x, y \in A$, if $x = y$, then onto one-to-one neither
 $f(x) = f(y)$.

$g : \mathbb{R} \rightarrow [0, 1]$ onto not onto not a function
 $g(x) = \sin(x)$

$f : \mathbb{R} \rightarrow \mathbb{Z}$ one-to-one not one-to-one not a function
 $f(x) = x$

Each dorm room is given an integer access code between 1 and 10 (inclusive). According to the pigeon-hole principle, if there are 21 dorm rooms, then every access code must be shared by at least two rooms. true false

$\exists y \in \mathbb{Z}, \forall x \in \mathbb{Z}, x - y < 100$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) To make exam grading anonymous and therefore hopefully more fair, each of the 200 students in CS 241 has been assigned a unique 3-character exam code. The character set is $\{\alpha, \beta, \gamma, \delta\}$. Use the Pigeonhole Principle to explain what's wrong with this plan.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = 2x$ then the real numbers is the
_____ of f .

domain
image

co-domain
none of these

$f : \mathbb{Z} \rightarrow \mathbb{Z}$
 $f(x) = x + 3$ (x even), onto not onto not a function
 $f(x) = x - 22$ (x odd)

$g : \mathbb{N}^2 \rightarrow \mathbb{N}$
 $g(x, y) = \gcd(x, y)$ one-to-one not one-to-one not a function

Each elf has exactly one gift: charm, strength, or stamina. If there are 10 elves, there must be at least three elves with the same gift.

true false

$\exists y \in \mathbb{R}^+, \forall x \in \mathbb{R}^+, xy = 1$
 $(\mathbb{R}^+ \text{ is the positive real numbers.})$

true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = p$ and $|B| = q$. How many different functions are there from A to B ?

2. (10 points) Check the (single) box that best characterizes each item.

A function is one-to-one if and only
if each value in the co-domain has
at most one pre-image. true false

$g : \mathbb{R}^2 \rightarrow \mathbb{R}$
 $g(x, y) = \lfloor x \rfloor + y$ onto not onto not a function

$g : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$
 $g(x, y) = (y, 3x)$ one-to-one not one-to-one not a function

Each ACM shirt has one of 6 trendy slogans. I bought
13 ACM shirts. At least three of these shirts must have
the same slogan. true false

$\forall x \in \mathbb{Q}, \exists m, n \in \mathbb{Z}, x = \frac{m}{n}$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) 15 men and 15 women showed up to this week's meeting of the UIUC Swing Dance Society. How many different ways can we form all of them into pairs, each pair containing one man and one woman?

2. (10 points) Check the (single) box that best characterizes each item.

Suppose $f : A \rightarrow B$. For
all $x \in A$, there is a $y \in B$, onto one-to-one neither
 $f(x) = y$.

$g : (\mathbb{Z}^+)^2 \rightarrow \mathbb{Z}^+$
 $g(x, y) = \gcd(x, y)$ onto not onto not a function

$f : \mathbb{N} \rightarrow \mathbb{R}$
 $f(x) = x^2 + 2$ one-to-one not one-to-one not a function

Each ACM shirt has one of 6 trendy slogans. I bought
13 ACM shirts. There is a slogan that appears on at true false
least two shirts.

$\forall x \in \mathbb{Z}, \exists y \in \mathbb{N}, x^2 = y$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = 2$ and $|B| = 3$. How many onto functions are there from A to B ? Briefly justify or show work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = 2x$ then the integers is the _____
of f .

domain
image

co-domain
none of these

$f : \mathbb{N}^2 \rightarrow \mathbb{Z}$
 $f(p, q) = 2^p 3^q$

onto not onto not a function

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = x|x|$

one-to-one not one-to-one not a function

Each elf has exactly one gift: charm, strength, or stamina. If there are 10 elves, the pigeonhole principle says that at least one elf has stamina.

true false

$\forall x \in \mathbb{R}^+, \exists y \in \mathbb{R}^+, xy = 1$

(\mathbb{R}^+ is the positive real numbers.)

true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Prof. Snape is teaching potions to 52 girls and 73 boys. Quiz 1 has integer scores between zero and 100 (inclusive). Assuming no one missed the quiz, what is the probability that two students got the same score? Briefly justify your answer.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : A \rightarrow B$ is one-to-one,
then $|A| \geq |B|$ $|A| \leq |B|$ $|A| = |B|$

$g : \mathbb{N}^2 \rightarrow \mathbb{N}$
 $g(x, y) = \gcd(x, y)$ onto not onto not a function

$g : \mathbb{R} \rightarrow \mathbb{R}^2$
 $g(x) = (x, 3x^2 + 2)$ one-to-one not one-to-one not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there is a true false color that appears on at least two mailboxes.

$\exists m, n \in \mathbb{Z}, \forall x \in \mathbb{Q}, x = \frac{m}{n}$ true false

Name: _____

NetID: _____

Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (5 points) Suppose that $|A| = p$ and $|B| = q$, $p \leq q$. How many different one-to-one functions are there from A to B ?

2. (10 points) Check the (single) box that best characterizes each item.

If a function from \mathbb{R} to \mathbb{R} is increasing,
it must be one-to-one. true false

$f : \mathbb{N} \rightarrow \mathbb{R}$
 $f(x) = x^2 + 2$ onto not onto not a function

$f : \mathbb{N} \rightarrow \mathbb{N}$
 $f(x) = 3 - x$ one-to-one not one-to-one not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, every color appears on at least two mailboxes. true false

$\exists y \in \mathbb{N}, \forall x \in \mathbb{N}, x = xy$ true false

Name: _____

NetID: _____ Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (5 points) Suppose that $|A| = p$, $|B| = q$, $|C| = n$. How many different functions are there from A to $B \times C$?

2. (10 points) Check the (single) box that best characterizes each item.

If a function from \mathbb{R} to \mathbb{R} is strictly increasing, it must be one-to-one. true false

$g : \mathbb{N} \rightarrow \mathbb{Z}$
 $g(x) = |x|$ one-to-one not one-to-one not a function

$g : \mathbb{R} \rightarrow \mathbb{R}$
 $g(x) = \sin(x)$ onto not onto not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there is a color that appears on at least two mailboxes. true false

$\forall x \in \mathbb{Z}, \exists y \in \mathbb{Z}, x \neq y \text{ and } x + y = 0$ true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 7-letter strings can be made by selecting and rearranging letters from the word ‘‘metalworking’’? Show your work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = 2x$ then the set of all even
integers is the _____ of f .

domain co-domain
image none of these

$f : \mathbb{Z} \rightarrow \mathbb{Z}$
 $f(x) = x + 4$ (x even), onto not onto not a function
 $f(x) = x - 22$ (x odd)

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = \lfloor x \rfloor$ one-to-one not one-to-one not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there is a true false color that appears on exactly two mailboxes.

$\exists y \in \mathbb{Z}, \forall x \in \mathbb{Z}, y \leq x$ true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) 10 men and 15 women showed up to this week's meeting of the UIUC Swing Dance Society. How many different ways can they line up (left to right) in front of the stage without any men being next to another man?

2. (10 points) Check the (single) box that best characterizes each item.

A function is onto if and only if its image is the same as its co-domain. true false

$f : \mathbb{Z} \rightarrow \mathbb{Z}$
 $f(x) = x + 3$ (x even), one-to-one not one-to-one not a function
 $f(x) = x - 21$ (x odd)

$g : \mathbb{Z} \rightarrow \mathbb{R}$
 $g(x) = x + 2.137$ onto not onto not a function

Each elf has exactly one gift: charm, strength, or stamina. If there are 10 elves, the pigeonhole principle says that at least three elves have charm. true false

$\exists y \in \mathbb{R}^+, \forall x \in \mathbb{R}^+, xy = 1$
 $(\mathbb{R}^+$ is the positive real numbers.) true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = p$, $|B| = q$, $|C| = n$. How many different functions are there from A to $B \times C$?

2. (10 points) Check the (single) box that best characterizes each item.

A function from \mathbb{R} to \mathbb{R} is strictly increasing if and only if it is one-to-one.

true false

$$f : \mathbb{Z} \rightarrow \mathbb{Z}$$

$f(x) = x + 3$ (x even), onto not onto not a function
 $f(x) = x - 22$ (x odd)

$$g : \mathbb{R} \rightarrow \mathbb{Z}$$

$g(x) = |x|$ one-to-one not one-to-one not a function

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there are two mailboxes with the same color.

true false

$$\exists y \in \mathbb{N}, \forall x \in \mathbb{N}, y \leq x$$

true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = 3$ and $|B| = 3$. How many onto functions are there from A to B ? Briefly justify or show work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that $f(x) = 2x$ then the integers is the _____ domain co-domain
of f . image none of these

$g : \mathbb{Z} \rightarrow \mathbb{Z}$ one-to-one not one-to-one not a function
 $g(x) = |x|$

$g : \mathbb{R} \rightarrow [0, 1]$ onto not onto not a function
 $g(x) = \sin(x)$

Each elf has exactly one gift: charm, strength, or stamina. If there are 10 elves, there must be at least true false

$\exists y \in \mathbb{N}, \forall x \in \mathbb{Z}, x^2 = y$ true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Let n and k be integers. Consider the integer powers of n from n^0 to n^k . Use the Pigeonhole Principle to show that there are two distinct (i.e. not equal) integers i and j , both between 0 and k (inclusive), such that $n^i \equiv n^j \pmod{k}$. (Your solution should be clear but does not need to be very formal.)

2. (10 points) Check the (single) box that best characterizes each item.

If a function is onto, then each value in the co-domain has exactly one pre-image. true false

$g : \mathbb{R} \rightarrow \mathbb{R}^2$
 $g(x) = (x, 3x^2 + 2)$ one-to-one not one-to-one not a function

$f : \mathbb{N} \rightarrow \mathbb{R}$
 $f(x) = x^2 + 2$ onto not onto not a function

If $f : A \rightarrow B$ is one-to-one,
then $|A| \geq |B|$ $|A| \leq |B|$ $|A| = |B|$

$\exists t \in \mathbb{Z}^+, \forall p \in \mathbb{Z}^+, \gcd(p, t) = 1$ true false

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 14-letter strings can be made by rearranging the letters in the word ‘‘classification’’? Show your work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{Z}$ is a function such that $f(x) = -|x|$ then \mathbb{N} is the _____ of f . domain co-domain
image none of these

$f : \mathbb{N}^2 \rightarrow \mathbb{N}$ one-to-one not one-to-one not a function
 $f(p, q) = pq$

$g : \mathbb{Z} \rightarrow \mathbb{Z}$ onto not onto not a function
 $g(x) = |x|$

We painted 12 mailboxes. There were 5 colors to choose from and each mailbox is painted with a single color. By the pigeonhole principle, there is a true false color that appears on at least two mailboxes.

$\exists t \in \mathbb{N}, \forall p \in \mathbb{Z}^+, \gcd(p, t) = p$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = p$ and $|B| = q$, $p \leq q$. How many different one-to-one functions are there from A to B ?

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = |x|$ then \mathbb{N} is the _____ of f .

domain
 image

co-domain

$g : \mathbb{R}^2 \rightarrow \mathbb{R}$
 $g(x, y) = \lfloor x \rfloor + y$

onto not onto not a function

$g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$
 $g(x, y) = (y, 3x)$

one-to-one not one-to-one not a function

Suppose a graph with 12 vertices is colored with
 exactly 5 colors. By the pigeonhole principle, every color appears on at least two vertices.

true false

$\forall x \in \mathbb{Q}, \exists m, n \in \mathbb{Z}, x = \frac{m}{n}$

true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Xin plans to randomly draw a hand of cards from a standard deck of 52 cards (evenly divided among 4 suits). He'd like to be sure the hand includes 3 cards with the same suit. How large must the hand be? Briefly justify your answer.

2. (10 points) Check the (single) box that best characterizes each item.

A function is one-to-one if and only if each value in the domain has exactly one image.

true false

$g : \mathbb{R}^2 \rightarrow \mathbb{R}$
 $g(x, y) = \lfloor x \rfloor + y$

one-to-one not one-to-one not a function

$g : \mathbb{R} \rightarrow [-1, 1]$
 $g(x) = \sin(x)$

onto not onto not a function

$f : \mathbb{N}^2 \rightarrow \mathbb{Z}$
 $f(p, q) = 2^p 3^q$

one-to-one not one-to-one not a function

$\forall x \in \mathbb{Z}^+, \exists y \in \mathbb{Z}^+, xy = 1$

true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that $|A| = 50$ and $B = \{5, 6\}$. How many onto functions are there from A to B ? Briefly justify or show work. (Hint: how many non-onto functions are there?)

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{N} \rightarrow \mathbb{Z}$ is a function such that $f(x) = -|x|$ then \mathbb{N} is the _____ of f .

domain	<input type="checkbox"/>	co-domain	<input type="checkbox"/>
image	<input checked="" type="checkbox"/>		

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = x|x|$

onto	<input type="checkbox"/>	not onto	<input type="checkbox"/>	not a function	<input type="checkbox"/>
------	--------------------------	----------	--------------------------	----------------	--------------------------

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = 7 - \lfloor \frac{x}{3} \rfloor$

one-to-one	<input type="checkbox"/>	not one-to-one	<input type="checkbox"/>	not a function	<input type="checkbox"/>
------------	--------------------------	----------------	--------------------------	----------------	--------------------------

$g : \mathbb{Z} \rightarrow \mathbb{Z}$
 $g(x) = x|x|$

one-to-one	<input type="checkbox"/>	not one-to-one	<input type="checkbox"/>	not a function	<input type="checkbox"/>
------------	--------------------------	----------------	--------------------------	----------------	--------------------------

$\forall x \in \mathbb{Z}, \exists y \in \mathbb{Z}, x - y < 100$

true	<input type="checkbox"/>	false	<input type="checkbox"/>
------	--------------------------	-------	--------------------------

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 10-letter strings can be made by rearranging the characters in the word ‘‘minimalist’’? Show your work.

2. (10 points) Check the (single) box that best characterizes each item.

If a function is onto, then each value in the co-domain has at least one pre-image.

true false

$g : (\mathbb{Z}^+)^2 \rightarrow \mathbb{Z}^+$
 $g(x, y) = \gcd(x, y)$

one-to-one not one-to-one not a function

$g : (\mathbb{Z}^+)^2 \rightarrow \mathbb{Z}^+$
 $g(x, y) = \gcd(x, y)$

onto not onto not a function

$f : \mathbb{R} \rightarrow \mathbb{Z}$
 $f(x) = x$

one-to-one not one-to-one not a function

$\exists m, n \in \mathbb{Z}, \forall x \in \mathbb{Q}, x = \frac{m}{n}$

true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose that A is a set containing $k+1$ (distinct) integers. Use the Pigeonhole Principle to show that there are x and y in A ($x \neq y$) such that $x - y$ is a multiple of k .

2. (10 points) Check the (single) box that best characterizes each item.

A function is one-to-one if and only if each value in the co-domain has true false at most one pre-image.

$g : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ one-to-one not one-to-one not a function

$$g(x, y) = (y, 3x)$$

$g : \mathbb{Z} \rightarrow \mathbb{N}$ one-to-one not one-to-one not a function

$$g(x) = x$$

$g : \mathbb{N}^2 \rightarrow \mathbb{N}$ onto not onto not a function

$$g(x, y) = \gcd(x, y)$$

$\exists y \in \mathbb{Z}, \forall x \in \mathbb{Z}, x - y < 100$ true false

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) How many different 10-letter strings can be made by rearranging the characters in the word ‘‘tattletale’’? Show your work.

2. (10 points) Check the (single) box that best characterizes each item.

If $f : \mathbb{Z} \rightarrow \mathbb{R}$ is a function such that
 $f(x) = 2x$ then the real numbers is the
_____ of f .

domain
image

co-domain

$g : \mathbb{N}^2 \rightarrow \mathbb{N}$
 $g(x, y) = \gcd(x, y)$

one-to-one not one-to-one not a function

$f : \mathbb{N}^2 \rightarrow \mathbb{N}$
 $f(p, q) = pq$

onto not onto not a function

Each dorm room is given an access code between 1 and 10 (inclusive). According to the pigeonhole principle, if there are 21 dorm rooms, then every access code must be shared by at least two rooms.

true false

$\forall m, n \in \mathbb{Z}, \exists x \in \mathbb{Q}, x = \frac{m}{n}$

true false