

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Multiply out and simplify, showing your work.

$$(x^{x-2} + x^2)^2 =$$

2. (10 points) Check the (single) box that best characterizes each item.

Shorthand for the set of integers.

 \mathbb{J} ☐ \mathbb{N} ☐ \mathbb{W} ☐ \mathbb{Z} ☐

If $\sqrt{2}$ is rational,
then -3 is positive.

true ☐false ☐undefined ☐

$$\log_2 3 < \log_3 2$$

true ☐false ☐

$$(p \vee \neg p) \rightarrow q$$

true ☐false ☐depends on q ☐

$$\neg(p \wedge \neg q) \equiv \neg p \vee q$$

true ☐false ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{\log_2(32^3)}{5} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$\sqrt{2} \in \mathbb{Q}$$

true ☐ false ☐

For all positive integers n ,
if $n! < 10$, then $n < 100$.

true ☐ false ☐

$$a^{b^c} = (a^b)^c$$

true ☐ false ☐

$$p \wedge q \equiv \neg(p \rightarrow \neg q)$$

true ☐ false ☐

For any real number x ,
 $\lceil \lfloor x \rfloor \rceil = \lfloor x \rfloor$.

true ☐ false ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\log_2(40) - \log_2(5) =$$

2. (10 points) Check the (single) box that best characterizes each item.

Shorthand for the set of rationals.

 \mathbb{R} ☐
 \mathbb{F} ☐
 \mathbb{Q} ☐
 \mathbb{B} ☐

Assume x is real.

If $x^2 < 0$, then x is even.

 true ☐

 false ☐

 undefined ☐

7 is a rational number

 true ☐

 false ☐

$(p \wedge q) \vee r \equiv (p \vee r) \wedge (q \vee r)$

 true ☐

 false ☐

$\neg(\neg p \rightarrow \neg q) \equiv \neg p \wedge q$

 true ☐

 false ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$x - \frac{x^2 - 2}{x - \sqrt{2}} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$-5 \in \mathbb{Z}$$

true ☐false ☐

$$\exists n \in \mathbb{Z}, n \geq 3 \text{ and } n \leq 3.$$

true ☐false ☐undefined ☐

$$\text{For some real number } x, \\ \lceil x \rceil \leq \lfloor x \rfloor.$$

true ☐false ☐

$$(p \wedge \neg p) \rightarrow q$$

true ☐false ☐depends on q ☐

$$\text{For any real numbers } x \text{ and } y, \\ \text{if } x \leq y, \text{ then } x^2 \leq y^2.$$

true ☐false ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{1}{\left(\frac{1}{2}\right)^4 + \left(-\frac{1}{2}\right)^5} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$0 \in \mathbb{Z}$$

true

☐

false

☐

$$\text{If } \pi < 7, \text{ then } 3 < 1$$

true

☐

false

☐

undefined

☐

$$\text{For any real number } x, 2\lfloor x \rfloor \leq \lfloor 2x \rfloor$$

true

☐

false

☐

$$\neg(p \rightarrow q) \equiv \neg p \rightarrow \neg q$$

true

☐

false

☐

$$\binom{n}{k}$$

$$\frac{n}{k}$$

☐

$$\frac{n!}{k!}$$

☐

$$\frac{k!}{n!}$$

☐

$$\frac{n!}{k!(n-k)!}$$

☐

$$\frac{n!}{k!(k-n)!}$$

☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{1}{\left(\frac{1}{2}\right)^4 + \left(-\frac{1}{2}\right)^6} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$0 \in \mathbb{Z}^+$$

true

☐

false

☐For any integer x ,if x is positive, then x is a real number.

true

☐

false

☐

$$\log_5 7 < 1$$

true

☐

false

☐

$$\exists n \in \mathbb{Z}, \text{ such that } n^2 = 10.$$

true

☐

false

☐

undefined

☐

$$\exists n \in \mathbb{Z}, \text{ such that } n^2 = 1.$$

true

☐

false

☐

undefined

☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\log_3(45x) - \log_3(5x) =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$0 \in \mathbb{R}$$

true

☐

false

☐

$$\text{If } \pi > 7, \text{ then } 3 < 1$$

true

☐

false

☐

undefined

☐

$$0!$$

0

☐

1

☐

-1

☐

undefined

☐

$$p \wedge \neg q \equiv \neg(p \rightarrow q)$$

true

☐

false

☐

$$2 \text{ is in the interval } (0, 2).$$

true

☐

false

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Express $\frac{1}{2}[(2 \log_2 a + \log_2 b) - 5 \log_2 c]$ as a single logarithm.

2. (10 points) Check the (single) box that best characterizes each item.

$\sqrt{-1} \in \mathbb{C}$

true

☐

false

☐

For any real number x ,
if $x > 10$, then $x^2 > 0$.

true

☐

false

☐

For any integer x , $\lfloor x \rfloor = x$.

true

☐

false

☐

$\neg(p \rightarrow q) \equiv \neg q \rightarrow \neg p$

true

☐

false

☐

$5 \in \mathbb{Q}$

true

☐

false

☐

Name: _____

NetID: _____

Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (5 points) Simplify, showing your work.

$$\log_3(45x) - \log_3(5x) =$$

2. (10 points) Check the (single) box that best characterizes each item.

The interval (a, b) contains b .

true

☐

false

☐

For any real number x ,
if $x \geq 10$, then $x > 10$.

true

☐

false

☐

$$3^{\lfloor -1.5 \rfloor} =$$

$$-3 \quad \boxed{}$$

$$\frac{1}{9} \quad \boxed{}$$

$$\frac{1}{3\sqrt{3}} \quad \boxed{}$$

$$\frac{1}{3} \quad \boxed{}$$

$$\log_3 2 \leq \log_2 3$$

true

☐

false

☐

$$\neg(p \wedge \neg q) \equiv \neg p \wedge q$$

true

☐

false

☐

Name: _____

NetID: _____ Lecture: B

Discussion: Friday 11 12 1 2 3 4

1. (5 points) Suppose $\log_k x = 8$. Then $\log_k(x\sqrt{x}) =$

2. (10 points) Check the (single) box that best characterizes each item.

The interval (a, b) contains b .

true

☐

false

☐

If 7 is even,
then 7 is a perfect square.

true

☐

false

☐

undefined

☐

$\exists n \in \mathbb{R}$, such that $n^2 = 10$.

true

☐

false

☐

undefined

☐

$\log_7 5 < 1$

true

☐

false

☐

For any real number x , $2\lceil x \rceil \leq \lceil 2x \rceil$

true

☐

false

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\log_7(\log_2 32 - \log_3 81) =$$

Solution:

Notice that $\log_2 32 = 5$ and $\log_3 81 = 4$.

$$\text{So } \log_7(\log_2 32 - \log_3 81) = \log_7(5 - 4) = \log_7 1 = 0$$

2. (10 points) Check the (single) box that best characterizes each item.

The interval (a, b) contains b .

true

☐

false

☒

For any real number x ,
if $x \geq 10$, then $x > 10$.

true

☐

false

☒

$(p \wedge q) \vee r \equiv (p \vee r) \wedge (q \vee r)$

true

☒

false

☐

$\log_3 2 \leq \log_2 3$

true

☒

false

☐

For any real number x , $2\lceil x \rceil \leq \lceil 2x \rceil$

true

☐

false

☒

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Remembering that log takes only positive inputs, solve $\log_{10} x + \log_{10}(x - 3) = 1$ for x .

Solution: This is equivalent to $\log_{10}(x(x - 3)) = 1$. So $x(x - 3) = 10$. Then we have $x^2 - 3x = 10$, so $x^2 - 3x - 10 = 0$. Factoring this, we get $(x - 5)(x + 2) = 0$. So x is -2 or 5. But -2 is not a legal input to log, so $x = 5$.

2. (10 points) Check the (single) box that best characterizes each item.

$$\binom{n}{k} \quad \frac{n}{k} \quad \frac{n!}{k!} \quad \frac{k!}{n!} \quad \frac{n!}{k!(n-k)!} \quad \frac{n!}{k!(k-n)!}$$

If Mickey Mouse is president of the US, then $\pi = 3$.

true ☒ false ☐ undefined ☐

$\exists n \in \mathbb{R}$, such that $n^2 = 10$.

true ☒ false ☐ undefined ☐

$\log_7 5 < 1$

true ☒ false ☐

For any integers x and y , if $x \leq y$, then $x^2 \leq y^2$.

true ☐ false ☒

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{\log_2(48) - \log_2(3)}{3} =$$

Solution: $\frac{\log_2(48) - \log_2(3)}{3} = \frac{\log_2(16 \cdot 3) - \log_2(3)}{3} = \frac{\log_2(16) + \log_2 3 - \log_2(3)}{3} = \frac{\log_2(16)}{3} = \frac{4}{3}$

2. (10 points) Check the (single) box that best characterizes each item.

$1.5 \in \mathbb{Z}$

true ☐ false ☒

$\pi < 7 \text{ or } \frac{3}{4} > \frac{2}{3}$

true ☒ false ☐

$p \wedge \neg q \equiv \neg(p \rightarrow q)$

true ☒ false ☐

$0! = 1!$

true ☒ false ☐

For any reals x and y ,

$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor.$

true ☐ false ☒

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Suppose $\log_k x = 5$. Then $\log_k(kx^{-3}) =$

Solution: $\log_k(kx^{-3}) = \log_k k + \log_k x^{-3} = 1 + \log_k x^{-3} = 1 + -3 \log_k x = 1 + -3 \cdot 5 = -14$

2. (10 points) Check the (single) box that best characterizes each item.

$0 \in \mathbb{N}$

true

☒

false

☐

If $\pi < 7$, then $3 < 1$

true

☐

false

☒

undefined

☐

$\neg(\neg p \rightarrow \neg q) \equiv \neg p \wedge q$

true

☒

false

☐

For any real number x , $2\lfloor x \rfloor = \lfloor 2x \rfloor$

true

☐

false

☒

$\lfloor -3 \rfloor$

3

☐

-3

☒

4

☐

-4

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\log_4 \frac{16^2}{2^{-3}} =$$

Solution: $\log_4 \frac{16^2}{2^{-3}} = \log_4 2^8 2^3 = \log_4 2^{11} = 11 \log_4 2 = \frac{11}{2}$

2. (10 points) Check the (single) box that best characterizes each item.

Shorthand for the set of integers. \mathbb{Z} ☒ \mathbb{I} ☐ \mathbb{N} ☐ \mathbb{W} ☐

$\sqrt{2} > 0$ or $\sqrt{2} > 3$ true ☒ false ☐

$(p \wedge q) \vee r \equiv (p \wedge r) \vee (q \wedge r)$ true ☐ false ☒

7 is a real number true ☒ false ☐

For any real number x ,
 $\lceil \lfloor x \rfloor \rceil = \lfloor x \rfloor$. true ☒ false ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 9 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\log_2\left(\frac{\sqrt[3]{16}}{8}\right) =$$

Solution: $\log_2\left(\frac{\sqrt[3]{16}}{8}\right) = \log_2(\sqrt[3]{16}) - \log_2 8 = \frac{1}{3} \log_2(16) - 3 = \frac{4}{3} - 3 = -\frac{5}{3}$

2. (10 points) Check the (single) box that best characterizes each item.

$1 \in \mathbb{Z}^+$ true ☒ false ☐

If U. Illinois is in Paris,
then $\pi < 0$. true ☒ false ☐ undefined ☐

$3^2 = 9$ or $4^2 = 15$ true ☒ false ☐

For any integer x ,
 $2\lfloor x \rfloor = \lfloor 2x \rfloor$ true ☒ false ☐

$\lfloor -3.4 \rfloor$ -3 ☐ 3.4 ☐ -4 ☒ undefined ☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$2^{\log_4 n} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$\neg(p \wedge \neg q) \equiv \neg p \wedge q$$

true

☐

false

☐

Shorthand for the set of integers.

 \mathbb{J} ☐ \mathbb{N} ☐ \mathbb{W} ☐ \mathbb{Z} ☐For any reals x and y ,

$$\lfloor x - y \rfloor = \lfloor x \rfloor - \lfloor y \rfloor.$$

true

☐

false

☐For any real number x ,

$$\text{if } |x + 5| \leq 10, \text{ then } |x| \leq 20.$$

true

☐

false

☐

$$\log_2 3 < \log_3 2$$

true

☐

false

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Multiply out and simplify, showing your work.

$$(x^{x-2} + x^2)^2 =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$\sqrt{2} \in \mathbb{Q}$$

true ☐ false ☐

$$\exists n \in \mathbb{Z}, n \geq 3 \text{ and } n \leq 3.$$

true ☐ false ☐ undefined ☐

$$\neg(p \rightarrow q) \equiv \neg q \rightarrow \neg p$$

true ☐ false ☐

$$\lfloor -3.4 \rfloor + \lceil -3.4 \rceil$$

-6 ☐ -6.8 ☐ -7 ☐ -8 ☐

For any real number x ,
if $x > 10$, then $x \geq 10$.

true ☐ false ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{1}{(\frac{1}{2})^4 + (-\frac{1}{2})^5} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$6 \in \mathbb{C}$

true

☐

false

☐

For any real numbers x and y ,
if $x \leq y$, then $x^2 \leq y^2$.

true

☐

false

☐

For any real number x , $\lfloor x \rfloor \leq \lceil x \rceil$.

true

☐

false

☐

$(p \vee \neg p) \rightarrow q$

true

☐

false

☐

depends on q

☐

If U. Illinois is in Paris,
then $\pi < 0$.

true

☐

false

☐

undefined

☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$(\log_2 17)(\log_{17} 64) =$$

2. (10 points) Check the (single) box that best characterizes each item.

$\exists n \in \mathbb{Z}$, such that $n^2 = 10$.

true

☐

false

☐

undefined

☐

7 is a rational number

true

☐

false

☐

For some real number x ,
 $\lceil x \rceil \leq \lfloor x \rfloor$.

true

☐

false

☐

$\sqrt{2} \in \mathbb{R}$

true

☐

false

☐

For any real numbers x and y ,
if $x \leq 6$ and $y \leq 2$, then $x - y \leq 4$.

true

☐

false

☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{1}{\left(\frac{1}{2}\right)^{10} + \left(-\frac{1}{2}\right)^{11}} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$p \wedge q \equiv \neg(p \rightarrow \neg q)$$

true

☐

false

☐For any integer x ,if x is positive, then x is a real number.

true

☐

false

☐For any real numbers x and y ,if $xy \leq 17y$, then $x \leq 17$.

true

☐

false

☐

Shorthand for the set of rationals.

 \mathbb{R} ☐ \mathbb{F} ☐ \mathbb{Q} ☐ \mathbb{B} ☐

$$3^{\lfloor -1.5 \rfloor} =$$

-3

☐ $\frac{1}{9}$ ☐ $\frac{1}{3\sqrt{3}}$ ☐ $\frac{1}{3}$ ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Simplify, showing your work.

$$\frac{(2^3 \times 2^5)^{10}}{512} =$$

2. (10 points) Check the (single) box that best characterizes each item.

$$p \rightarrow q \equiv \neg q \rightarrow \neg p$$

true

☐

false

☐

$$\text{If } \pi > 7, \text{ then } 3 < 1$$

true

☐

false

☐

undefined

☐

$$\lfloor -3.4 \rfloor$$

-3

☐

-4

☐

3.4

☐

undefined

☐The interval $[a, b]$ contains b .

true

☐

false

☐For any real number x , $2\lfloor x \rfloor \leq \lfloor 2x \rfloor$

true

☐

false

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) $\frac{\log_3(81^n)}{\log_3 \frac{1}{27}} =$

2. (10 points) Check the (single) box that best characterizes each item.

$\exists n \in \mathbb{Z}$, such that $n^2 = 1$.

true ☐ false ☐ undefined ☐

2 is in the interval $(0, 2)$.

true ☐ false ☐

$\log_5 7 < 1$

true ☐ false ☐

If $\sqrt{2}$ is rational, then -3 is positive.

true ☐ false ☐ undefined ☐

$\log_7 1$

0 ☐ 1 ☐ 7 ☐ undefined ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) $4^{\log_2(7n)} =$

2. (10 points) Check the (single) box that best characterizes each item.

 $\frac{3}{7}$ is a real number

true

☐

false

☐

$p \rightarrow q \equiv \neg p \rightarrow \neg q$

true

☐

false

☐

$p \vee q \equiv \neg p \rightarrow q$

true

☐

false

☐For any integer x , $\lfloor x \rfloor = x$.

true

☐

false

☐ $\forall x \in \mathbb{R},$
if $x^2 > 100$, then $|x| \geq 10$.

true

☐

false

☐

Name: _____

NetID: _____

Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) $x - \frac{x^2 - 2}{x - \sqrt{2}} =$

2. (10 points) Check the (single) box that best characterizes each item.

$-5 \in \mathbb{N}$

true ☐ false ☐

$0!$

0 ☐ 1 ☐ -1 ☐ undefined ☐

$a^{b^c} = (a^b)^c$

true ☐ false ☐

$(p \wedge \neg p) \rightarrow q$

true ☐ false ☐ depends on q ☐

For any real number x ,
if $x \in \mathbb{N}$, then $x \in \mathbb{Q}$.

true ☐ false ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) $25 \times 2^{-3 \log_2(5)} =$

2. (10 points) Check the (single) box that best characterizes each item.

$\neg(p \rightarrow q) \equiv p \rightarrow \neg q$

true

☐

false

☐For all positive integers n ,
if $n! < 10$, then $n < 100$.

true

☐

false

☐

$\lfloor -3.4 \rfloor$

-3

☐

3.4

☐

-4

☐

undefined

☐

$0 \in \mathbb{N}$

true

☐

false

☐

$3^{\lfloor -1.5 \rfloor} =$

-3

☐ $\frac{1}{9}$ ☐ $\frac{1}{3\sqrt{3}}$ ☐ $\frac{1}{3}$ ☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) Express $\frac{1}{2}[(2 \log_2 a + \log_2 b) - 5 \log_2 c]$ as a single logarithm.

2. (10 points) Check the (single) box that best characterizes each item.

$$\neg(p \rightarrow q) \equiv \neg p \rightarrow \neg q$$

true

☐

false

☐

For any real number x ,
if $x > 10$, then $x^2 > 0$.

true

☐

false

☐

For any integer x , $\lfloor x \rfloor = x$.

true

☐

false

☐

$$0 \in \mathbb{Z}^+$$

true

☐

false

☐

7 is a real number

true

☐

false

☐

Name: _____

NetID: _____ Lecture: A B

Discussion: Thursday Friday 10 11 12 1 2 3 4 5 6

1. (5 points) $\frac{(-1)^{k+2}2^{2k}}{(-2)^{k+1}} =$

2. (10 points) Check the (single) box that best characterizes each item.

$\exists n \in \mathbb{Z}$, such that $n^2 = 10$.

true ☐ false ☐ undefined ☐

$\pi < 7$ or $\frac{3}{4} > \frac{2}{3}$

true ☐ false ☐

If 7 is even, then 7 is a perfect square.

true ☐ false ☐ undefined ☐

$-5 \in \mathbb{Z}$

true ☐ false ☐

For any real number x , $2\lfloor x \rfloor = \lfloor 2x \rfloor$

true ☐ false ☐