

GALAXY ZOO: MORPHOLOGICAL CLASSIFICATIONS FOR 120,000 GALAXIES IN HST LEGACY IMAGING

KYLE W. WILLETT^{1,2}, MELANIE A. GALLOWAY¹, STEVEN P. BAMFORD³, CHRIS J. LINTOTT⁴, KAREN L. MASTERS^{5,6}, CLAUDIA SCARLATA¹, B.D. SIMMONS^{4,7}, MELANIE BECK¹, CAROLIN N. CARDAMONE⁸, EDMOND CHEUNG⁹, EDWARD M. EDMONDSON⁵, LUCY F. FORTSON¹, ROGER L. GRIFFITH^{10,11}, BORIS HÄUSSLER^{4,12,13}, ANNA HAN¹⁴, ROSS HART³, TOM MELVIN⁵, MICHAEL PARRISH¹⁵, KEVIN SCHAWINSKI¹⁶, R.J. SMETHURST⁴, ARFON M. SMITH^{5,15,17}

¹School of Physics and Astronomy, University of Minnesota, USA

²Department of Physics and Astronomy, University of Kentucky, USA

³School of Physics and Astronomy, University of Nottingham, UK

⁴Oxford Astrophysics, University of Oxford, UK

⁵Institute of Cosmology and Gravitation, University of Portsmouth, UK

⁶SEPnet, South East Physics Network, UK

⁷Center for Astrophysics and Space Sciences, UC San Diego, USA

⁸Department of Mathematics and Science, Wheelock College, USA

⁹Kavli Institute for the Physics and Mathematics of the Universe (WPI), Japan

¹⁰Infrared Processing and Analysis Center, California Institute of Technology, USA

¹¹Department of Astronomy and Astrophysics, Pennsylvania State University, USA

¹²Centre for Astrophysics, University of Hertfordshire, UK

¹³European Southern Observatory, Santiago, Chile

¹⁴Department of Astronomy, Yale University, USA

¹⁵Adler Planetarium, USA

¹⁶Institute for Astronomy, ETH Zürich, Switzerland

¹⁷GitHub

ABSTRACT

We present the data release paper for the Galaxy Zoo: Hubble (GZH) project.^a This is the third phase in a large effort to measure reliable, detailed morphologies of galaxies by using crowdsourced visual classifications of color composite images. Images in GZH were selected from various *Hubble Space Telescope* Legacy programs using the Advanced Camera for Surveys, with filters that probe the rest-frame optical emission from galaxies out to $z \sim 1$. The bulk of the sample is selected to have $m_{I814W} < 23.5$ across AEGIS, COSMOS, GEMS, and 2-epoch exposures in GOODS-N and GOODS-S. Full-depth 5-epoch images from the GOODS fields are also included and have $m_{I814W} < 26.8$. The median redshift of the combined samples is $\langle z \rangle = 0.9 \pm 0.6$, with a tail extending out to $z \simeq 4$. The GZH morphological data include measurements of both bulge- and disk-dominated galaxies, details on spiral disk structure that relate to the Hubble type, bar identification, and numerous measurements of clump identification and geometry. This paper also describes a new method for calibrating morphologies for galaxies of different luminosities and at different redshifts by using artificially-redshifted galaxy images as a baseline. The GZH catalog contains both raw and calibrated morphological vote fractions for 119,849 galaxies, providing the largest dataset to date suitable for large-scale studies of galaxy evolution out to $z \sim 1$.

Keywords: galaxies:structure — galaxies:high redshift — galaxies:evolution — methods:data analysis — catalogs

1. INTRODUCTION

^a This publication has been made possible by the participation of more than 200,000 volunteers in the Galaxy Zoo project. Their contributions are individually acknowledged at <http://authors.galaxyzoo.org/authors.html>.

The morphology of galaxies encodes information on the orbital parameters and assembly history of their contents, including gas, dust, stars, and the central black hole. The morphology is also closely related to the local environment of the galaxy, as mutual interactions such as tides, shocks in cluster environments, and direct

mergers can all change the shape of the galaxy’s gravitational potential. For M^* galaxies in the local Universe, this interplay between the physical development of a galaxy and its external appearance typically manifests at the most basic level as the difference between bulge-dominated systems with no/little spiral structure (early-types) and disk-dominated, rotationally-supported galaxies (late-types) frequently exhibiting spiral arms. This dichotomy has been used to explore much of the astrophysics governing galaxy formation and evolution, and has been shown to be closely linked with other galactic properties such as stellar mass, halo mass, bolometric luminosity, black hole activity, effective radius, and the relative ages of the stellar populations.

The advent of larger telescopes in an increasing range of observing wavelengths has revealed that the distribution and properties of galaxy morphology have strongly evolved over the lifetime of the Universe. At redshift $z \simeq 1$ (roughly 6 Gyr after the Big Bang), many galaxies are still in the process of assembling the baryonic mass required to reproduce the mature, coherent structures seen in the present day. This growth occurs in a variety of pathways, including accretion of baryons from large-scale galactic filaments onto halos via streaming, mergers of individual dark matter halos along with their baryons, conversion of gas into stars through gravitational collapse and star formation, etc. The process can also be slowed or even reversed via feedback from stellar winds, supernovae, and active black holes. Each of these processes affects the galaxy morphology in different ways, and so an accurate measurement of the demographics as a function of redshift provides an extremely powerful observational constraint on the physics involved (for recent reviews see [Buta 2013](#); [Conselice 2014](#)).

Theoretical predictions for the morphology of galaxies as a function of redshift are primarily computed within the Λ CDM cosmological framework. Full treatments model gravitational interactions between baryons and dark matter, hydrodynamics of the gas, and baryonic physics related to star formation and evolution. The most advanced simulations now span volumes up to $\sim 100 \text{ Mpc}^3$ while simultaneously resolving the smaller ($< 1 \text{ kpc}$) scales necessary to reproduce the influence of baryonic physics ([Vogelsberger et al. 2014](#); [Schaye et al. 2015](#)). Such simulations predict clustering of galaxies on large scales in a hierarchical assembly model ([Silk & Mamon 2012](#)). The structure of individual galaxies is affected by their merger history ([Toomre & Toomre 1972](#); [Steinmetz & Navarro 2002](#); [Hopkins et al. 2010](#)), local environment (*e.g.*, the morphology-density relation; [Dressler 1980](#)), initial dark halo mass, secular evolution rate, and many other factors. Morphologies of individ-

ual simulated high-mass galaxies at $z \sim 2 - 3$ commonly show kpc-scale “clumpy” structures, with few galaxies that are either smooth or well-ordered spirals; asymmetric galaxies with strong density contrasts dominate simulated populations in the early Universe until at least $z \sim 1$ ([Bell et al. 2012](#); [Genel et al. 2014](#)).

Observational studies of galaxies at high-redshift also display a wide range of morphological types, many of which are rare or absent at $z \sim 0$. These include spheroids and disks (akin to the ellipticals and spirals seen in the local Universe), but also a significant population of massive, more irregular galaxies, including mergers, tadpoles, chains, double-clumps, and clump-clusters ([Elmegreen et al. 2005, 2007](#); [Cameron et al. 2011](#); [Förster Schreiber et al. 2011](#); [Kartaltepe et al. 2015](#)). In contrast, while grand-design spirals have been observed as far back as $z = 2.18$ ([Law et al. 2012a](#)), their spatial density suggests that they are exceedingly rare at these high redshifts, with a very low overall disk fraction ([Mortlock et al. 2013](#)). Current observational data thus strongly suggests that the classical Hubble sequence/tuning fork ([Hubble 1936](#)) is not a suitable framework for characterizing high-redshift morphology.

Space-based observatories, particularly the *Hubble Space Telescope* (*HST*), have been responsible for the bulk of imaging studies of high-redshift galaxies. Observations of fields with very deep imaging (*eg.*, [Williams et al. 1996](#); [Giavalisco et al. 2004](#); [Beckwith et al. 2006](#); [Davis et al. 2007](#); [Scoville et al. 2007](#); [Grogin et al. 2011](#)) give the photometric sensitivity necessary to detect L^* galaxies at $z > 1$, while also providing the angular resolution to distinguish internal structure and characterize the morphology. While these measurements are helped by the fact that the angular diameter distance is relatively flat beyond $z > 1$ in a flat Λ CDM cosmology, the relevant angular scales are only of the order $\sim 5 - 10 \text{ kpc}/''$ ([Wright 2006](#)). *HST* can thus resolve much of the structure for a Milky Way-sized galaxy out to moderately high redshifts (at least distinguishing a disk from a bulge), but is limited for more compact structures. Since the size of galaxies evolves as roughly $r \propto (1 + z)^{-1}$ ([Mao et al. 1998](#); [Law et al. 2012b](#)), the compact sizes of high-redshift galaxies make detailed morphologies a challenge even for *HST* ([Chevance et al. 2012](#)). However, the public availability of more than 10^5 galaxies in archival imaging has generated a data sample with the potential for statistically-robust studies of galaxy demographics and evolution.

One of the major difficulties in studying the morphologies of galaxies lies in the techniques used for measurement. Visual classification by experts has been used for many decades (*eg.*, [Hubble 1926](#); [de Vaucouleurs 1959](#); [Sandage 1961](#); [van den Bergh 1976](#); [Nair & Abraham 2010](#); [Baillard et al. 2011](#); [Kartaltepe et al. 2015](#)). These

methods have the advantage of using the significant processing power of the human brain to identify patterns, but suffer from issues such as lack of scaling to large surveys and potential issues with replicability and calibration (e.g., see Lahav et al. 1995 for a discussion on the extent to which eight expert classifiers agree with each other). Automated measurements, both parametric (Peng et al. 2002; Simard et al. 2011; Lackner & Gunn 2012) and non-parametric (Abraham et al. 2003; Conselice 2003; Lotz et al. 2004; Scarlata et al. 2007; Bamford et al. 2008; Freeman et al. 2013), scale well to very large sample sizes, but do not always fully capture the relevant features, especially for asymmetric galaxies that become increasingly common at high redshifts. The Galaxy Zoo project (Lintott et al. 2008; Fortson et al. 2012) utilizes crowdsourced visual classifications to measure galaxies in color-composite images. The efforts of more than 2×10^5 classifiers allow for multiple independent classifications of each image which are combined and calibrated to give a distribution of vote fractions proportional to the probability of a feature being visible. While crowdsourced data require extensive calibration (Bamford et al. 2009; Willett et al. 2013), they have a proven reliability and have been used for a wide variety of scientific studies (eg, Land et al. 2008; Bamford et al. 2009; Darg et al. 2010; Masters et al. 2011; Skibba et al. 2012; Simmons et al. 2013; Schawinski et al. 2014; Willett et al. 2015).

This paper presents the classifications collected from the Galaxy Zoo Hubble (GZH) project.¹ GZH was the third phase of Galaxy Zoo, following its initial results classifying $\sim 900,000$ Sloan Digital Sky Survey (SDSS) images into primarily early/late types (Lintott et al. 2011) and Galaxy Zoo 2, which covered a subset of $\sim 250,000$ images using a more detailed classification scheme that included bars, spiral arms, and galactic bulges (Willett et al. 2013). GZH used a similarly detailed classification scheme, but focused for the first time on images of high-redshift galaxies taken with *HST*. The Galaxy Zoo: CANDELS project has also classified morphologies of galaxies at high redshift, but using ACS and WFC3 rest-frame infrared imaging (Simmons et al., submitted).

The sample selection and creation of the images used for GZH is described in Section 2. Section 3 describes the GZH interface and the collection of classifications. Section 4 outlines the process used to calibrate and correct the crowdsourced vote fractions for redshift-dependent bias. Section 5 gives the main catalog of results, with several examples of how the data may be

queried in Section 6. Section 7 gives a short overview of the observed morphological demographics and compares them to several other catalogs, with a summary in Section 8.

This paper assumes the WMAP9 cosmological parameters of $(\Omega_m, \Omega_\Lambda, h) = (0.282, 0.718, 0.697)$ (Hinshaw et al. 2013).

2. SAMPLE AND DATA

The GZH project contains images drawn from a number of different dedicated surveys and sample selection criteria. The majority of the data (as implied by the project name) were sourced directly from *HST* Legacy Surveys, all of which primarily used imaging from the Advanced Camera for Surveys (ACS). In addition to *HST* images, the project uses images from SDSS Stripe 82, as well as simulated *HST* images from multiple sources. Below we provide details on each source of imaging and describe the creation process for images shown to classifiers. We also detail the sources of metadata for galaxies in the sample, such as photometry and redshifts.

2.1. *Hubble Legacy Surveys*

Image information from the individual surveys have been combined into a single photometric and morphological database, the Advanced Camera for Surveys General Catalog (ACS-GC; Griffith et al. 2012). A summary of the key parameters is given in Table 1.

The properties of the individual surveys are as follows:

- The All-Wavelength Extended Groth Strip International Survey (AEGIS; Davis et al. 2007) covers a strip centered at $\alpha = 14^{\text{h}}17^{\text{m}}, \delta = +52^\circ30'$. This area of the sky was selected for a deep survey due to a combination of low extinction and low Galactic/zodiacal emission. The ACS images covered 63 separate tiles over a total area of ~ 710 arcmin². The two ACS bands for AEGIS had exposure times of 2300 seconds in F606W (V_{606W}) and 2100 seconds in F814W (I_{814W}). The final mosaic images were dithered to a resolution of 0.03 ''/pixel. For extended objects, the limiting magnitude of sources was 26.23 (AB) in V_{606W} and 25.61 (AB) in I_{814W} .
- The Great Observatories Origins Deep Survey (GOODS; Giavalisco et al. 2004) covered two separate fields in the northern and southern hemispheres: the Hubble Deep Field-North ($\alpha = 12^{\text{h}}36^{\text{m}}, \delta = +62^\circ14'$) and the Chandra Deep Field-South ($\alpha = 03^{\text{h}}32^{\text{m}}, \delta = -27^\circ48'$). The *HST* ACS imaging data from the two fields are referred to as GOODS-N and GOODS-S, respectively. ACS imaging in GOODS fields used 4 filters

¹ <http://zoo3.galaxyzoo.org/>




Figure 1. Transmission curves of the filters used by *HST* Advanced Camera for Surveys (ACS) in wide-field channel mode for the various surveys in GZH. The unfilled black curves show the filters for the Suprime Camera on *Subaru*, which was used to create color gradients in the GZH COSMOS images.

– F435W (B_{435W}), V_{606W} , F775W (i_{775W}), and F850LP (z_{850LP}). The mean exposure times for each epoch varied by band, from 1050–2100 seconds. The B_{435W} images were completed in a single epoch at the beginning of the survey, but the V_{606W} , i_{775W} , and z_{850LP} images were taken in five separate epochs separated by 40–50 days each. Images were dithered to a pixel scale of 0.03 ''/pixel and covered a total area of $\sim 320 \text{ arcmin}^2$ (160 arcmin^2 per north/south field). The 5σ limiting magnitude for extended sources was 25.7 for V_{606W} and 25.0 for i_{775W} .

- The Cosmic Evolution Survey (COSMOS; Scoville et al. 2007) covered an area of $\sim 1.8 \text{ deg}^2$ centered at $\alpha = 10^{\text{h}}00^{\text{m}}$, $\delta = +02^\circ 12'$. Its location near the celestial equator was designed to enable coverage by ground-based telescopes in both the Northern and Southern Hemispheres, in addition to space-based observatories. The ACS data for COSMOS consisted of 1 orbit per pointing with an exposure time of 2028 seconds in I_{814W} ; 590 total pointings were used to cover the entire field. The image resolution was dithered to 0.05 ''/pixel. The 50% completeness magnitude for a galaxy with a half-light radius of 0.50'' in I_{814W} was 24.7 mag.
- The Galaxy Evolution from Morphologies and

SEDs (GEMS; Rix et al. 2004; Caldwell et al. 2008) survey was also centered on the Chandra Deep Field-South. The GEMS data covered $\sim 800 \text{ arcmin}^2$, completely surrounding the area covered by GOOD-S-N. Images from ACS in GEMS had 1 orbit per pointing for a total of 63 pointings. The exposure times were 2160 and 2286 seconds in V_{606W} and z_{850LP} , respectively. The image resolution had a pixel scale of 0.03 ''/pixel. The 5σ limiting magnitude for source detection was 25.7 AB in V_{606W} and 24.2 AB in z_{850LP} .

In the ACS-GC (Griffith et al. 2012), individual galaxies were identified using a combination of SExtractor (Bertin & Arnouts 1996) and the galaxy-profile fitting framework GALAPAGOS (Barden et al. 2012). GZH included all galaxies with $m < 23.5$, where m is in the I_{814W} , z_{850LP} , or i_{775W} for the AEGIS + COSMOS, GEMS + GOOD-S-N (2-epoch), and GOOD-S-N (2-epoch) surveys, respectively. The full-depth GOODS images from both fields included galaxies with $m < 26.8$. This yielded a total of 119,849 images (Table 1).

Images from *HST* legacy surveys were used to create multiple different color and grayscale images using different depths and filter combinations; these are described further in Section 2.4. Additionally, the use of a small set of *HST* images to create simulated images of active galactic nucleus (AGN) host galaxies is described

in Section 2.3.1.

2.2. The Sloan Digital Sky Survey

The project includes images from the SDSS (York et al. 2000; Strauss et al. 2002), in particular those from Data Release 7 (Abazajian et al. 2009). A small number of images were used in the creation of simulated *HST* images, described in Sections 2.3.2 and 4 below; the majority of SDSS images used in GZH were from Stripe 82. The Stripe 82 samples in GZH include single-epoch and co-added images. The single-epoch images provide a local sample for comparison to higher-redshift galaxies (including the new measurements of clumpy structure), while the co-added images allow for analysis of morphological properties as a function of image depth.

Single-epoch images from SDSS Stripe 82 were selected using the criteria from Willett et al. (2013), which required limits of `petroR90_r > 3''` (where `petroR90_r` is the radius containing 90% of the *r*-band Petrosian flux) and a magnitude brighter than $m_r < 17.77$. 21,522 galaxies in SDSS met these criteria. Co-added images from Stripe 82 were selected from the union of galaxies with co-added magnitudes brighter than 17.77 mag, and the galaxies detected in the `stripe-82-single` images and matched to a co-add source. This resulted in a total set of 30,339 images. Of the images in the co-added sample, 5144 (17 percent) were dimmer than the initial magnitude cut of 17.77.

2.3. Simulated *HST* images

To facilitate correction of classifications in the presence of known redshift bias (Section 4), GZH includes two different samples of simulated *HST* images: real *HST* galaxy images (Section 2.1) with nuclear emission added to simulate AGN host galaxies, and lower-redshift galaxy images from SDSS (Section 2.2), artificially redshifted from $0.3 \leq z \leq 1$. Each is described separately below.

2.3.1. Images with simulated nuclear point sources

As active galactic nuclei often have bright, unresolved optical emission, AGN have the potential to mimic or distort the identification of a bulge component. GZH thus includes a set of images designed to measure the effect of AGN on morphological classifications. The presence of an AGN was simulated by modeling the point spread function (PSF) of the telescope and then inserting a bright source near the center of a real galaxy. For each image, the simulated AGN was assigned one of three colors – either blue, red, or flat (white) as seen in the color images – and a range of brightnesses such that $L_{\text{ratio}} \equiv L_{\text{galaxy}}/L_{\text{AGN}}$ is in $(0.2, 1.0, 5.0, 10.0, 50.0)$. Combining these parameters generated 15 images with different simulated AGN for each host, in addition to the original galaxy image.

Two sets of simulated AGN were generated in GZH. The first set (version 1) was assembled from 95 galaxies from GOODS-S imaging and empirical PSFs made by combining stars in the GOODS fields using the PSF creation tools in `daophot`. The second set (version 2) was assembled from 96 galaxies in GOODS-S; this version used simulated PSFs from `TinyTim` (Krist 1993), drizzled using the same procedures as those used in the reduction of the GOODS-S images (Koekemoer et al. 2002, 2003; Giavalisco et al. 2004). The use of these two versions facilitates comparisons between these different PSF creation methods, which are widely used in AGN host galaxy morphology studies (e.g., Sánchez et al. 2004; Simmons & Urry 2008; Pierce et al. 2010; Simmons et al. 2011). Each PSF creation method has advantages and disadvantages: the empirical PSFs better represent the nuances of the PSF in the specific data being used and look more realistic at lower luminosities, but the extended features of the noiseless `TinyTim` PSFs are visually more realistic at higher luminosities.

Images with simulated AGN were classified in the interface in an identical manner and evenly distributed with unaltered images of the galaxies. Volunteers were not explicitly told that the images had been altered during classification, as the goal was to measure the effect on normal classifications using the same technique as closely as possible. Following classification, a classifier could view a page with additional details about each galaxy; where applicable, these pages contained further information regarding image modifications. Simmons et al. (2014) uses similarly-processed images for reliable identification of bulgeless galaxies using SDSS data.

2.3.2. Generating images of artificially-redshifted galaxies

The dimming and resolution effects of redshift can significantly affect galaxy classifications derived from any method. To facilitate corrections due to redshift, we include a sample of 288 galaxies with SDSS imaging that can be transformed to simulate *HST* imaging out to $z = 1$. The SDSS images were redshifted and processed to mimic simulated *HST* imaging parameters using the FERENGI code (Barden et al. 2008).

In addition to the physical parameters of the input images, the FERENGI output depends on assumptions of the global galaxy evolution model. This evolution is a crude mechanism that mimics the brightness increase of galaxies with increasing redshift (e.g., Lilly et al. 1998; Loveday et al. 2012). The effect on the redshifted images is simply an empirical addition to the magnitude of a galaxy of the form $M' = e \times z + M$, where M' is the corrected magnitude, and e is the evolutionary correction in magnitudes ($e = -1$ essentially brightens the entire galaxy by 1 magnitude by $z = 1$). FERENGI was run on the images for values of e starting from $e = 0$

Table 1. Summary of GZH *Hubble Space Telescope* imaging

Survey	Total t_{exp} [sec]	Filters	Resolution ["/pix]	Area [arcmin 2]	N_{galaxies}
AEGIS	2100–2300	V_{606W}, I_{814W}	0.03	710	8507
COSMOS	2028	I_{814W}	0.05	6480	84954
GEMS	2160–2286	V_{606W}, z_{850LP}	0.03	800	9087
GOODS	—	—	—	—	—
<i>GOODS-N 2 epoch</i>	2100–4200	V_{606W}, i_{775W}	0.03	320	2551
<i>GOODS-S 2 epoch</i>	2100–4200	V_{606W}, z_{850LP}	0.03	320	3593
<i>GOODS-N 5 epoch</i>	5100–10500	$B_{435W}, V_{606W}, i_{775W}, z_{850LP}$	0.03	"	6015
<i>GOODS-S 5 epoch</i>	5100–10500	$B_{435W}, V_{606W}, i_{775W}, z_{850LP}$	0.03	"	5142
all <i>HST</i> surveys	—	—	—	8630	119849

and decreasing to $e = -3.5$ in increments of $\Delta e = 0.5$. Figure 2 shows several examples of the effects of “losing” spiral/disc features with increasing redshift for two galaxies with no evolution corrections ($e = 0$); as the signal-to-noise ratio in the images decreases and the galaxies become fainter, the contrast between features like spiral arms goes down and f_{features} drops to the point where both galaxies would have been classified as likely ellipticals.

The final number of FERENGI images produced for each galaxy is ultimately a function of the galaxy’s redshift (since the new images cannot be resampled at better angular resolution than the original SDSS data), as well as the number of e values selected. The use of these images to correct for redshift-dependent bias is described in Section 4.

2.4. Creating color images

The images used for classification in GZH were color-composite JPEGs made from multi-band data. These were created following the method of Lupton et al. (2004), which preserves color information irrespective of intensity. An asinh intensity mapping was applied to enhance the appearance of faint features while avoiding saturating galaxy centers. The relative scalings of the filter bands were chosen while trying to reproduce the color appearance of the SDSS images in previous iterations of Galaxy Zoo.

Many of the Legacy surveys described in Section 2.1 provided *HST* images in only two filters. For these, the shorter-wavelength band was mapped to the blue channel, the longer-wavelength band to the red channel, and the green channel created by taking the arithmetic mean of the two. The bands used in each of the surveys are listed in Table 1. Although four bands were available for the GOODS survey, only two bands were used to create the original 2-epoch images, for consistency with AEGIS and GEMS. The 2-epoch GOODS-N and GOODS-S images were created using different filters — this was a

deliberate choice made so that the GEMS images could be directly compared with the overlapping coverage of GOODS-S (Figure 1).

Only 2-epoch GOODS images were included at the launch of GZH. Deeper, 5-epoch GOODS images were added into GZH later (in March 2015). These images made use of the full four-band data by using the arithmetic mean of B_{435W} and V_{606W} in the blue channel, I_{814W} in the green channel, and z_{850LP} in the red channel.

The COSMOS survey provides only I_{814W} HST imaging. For these galaxies, GZH used “pseudocolor” images created by using the ACS I_{814W} data as an illumination map and ground-based imaging from the *Subaru* telescope in B_J , r^+ , and i^+ filters to provide color information (see Griffith et al. 2012 for further details). This resulted in images with the angular resolution of *HST* (~ 0.05 "/pixel) for the overall intensity, but color gradients at ground-based resolution, with seeing between 0.95" and 1.05" (Taniguchi et al. 2007).

Stripe 82 single-epoch images were taken directly from the DR7 SDSS Skyserver, which combined g' , r' , and i' exposures into the RGB channels (Nieto-Santisteban et al. 2004). The co-added Stripe 82 images were assembled from runs 106 and 206 in DR7 and processed into color composites in the same manner as previous iterations of Galaxy Zoo.

In some cases, we found that attempting to emphasize faint features in the images resulted in the sky noise taking the appearance of brightly colored speckles. This impaired the aesthetics of the images and might have been a distraction to visual classification. To counteract this, a soft-edged object mask was applied to the color images and a desaturation operation performed. This masking procedure was effective in preserving the color balance for galaxies and retained the visibility of faint features, while reducing the color contrast in the sky noise and greatly improving the appearance of the images. This solution was applied to the coadded Stripe 82, COSMOS




Figure 2. Examples of two galaxies which have been run through the FERENGI code to produce simulated *HST* images. The measured value of f_{features} from GZH for the images in each panel are (1) Top row: $f_{\text{features}} = (0.900, 0.625, 0.350, 0.350, 0.225)$ and (2) Bottom row: $f_{\text{features}} = (1.000, 0.875, 0.875, 0.625, 0.375)$.

and 5-epoch GOODS images.

In addition to the primary *HST* legacy imaging sample with colored images described as above, GZH also includes samples of *HST* galaxy images with different color prescriptions. In particular, there are two sets of 3,927 images each, drawn from the COSMOS sample; the first has a dramatically reduced color saturation, and the second has reversed the colors so that the blue and red filters have exchanged places in the RGB image. The ‘faded’ galaxy set facilitates measuring of possible variations in classification due to the presence or absence of color features. The ‘recolored’ set enables an alternative test of potential color biases in classifications.

The simulated AGN host galaxy color images (described in Section 2.3.1) were created using the same prescription as for the GOODS 2-epoch imaging. The artificially redshifted color images (Section 2.3.2) were created using the same prescription as for the AEGIS images.

2.5. Galaxy sample labels

The full GZH sample is composed of 8 different galaxy sub-samples. Throughout this paper and in the published catalogs they are referred to with the following labels:

- **main:** *HST* imaging with RGB colors making use of all available filters with typical saturation and in correct order, as described in Section 2.4. This sample includes AEGIS, COSMOS, GEMS, and full-depth GOODS (North and South) images. (113,705 galaxies)
- **faded:** a subset of COSMOS images with very low color saturation. (3,927 galaxies)
- **recolored:** a subset of COSMOS images with red

and blue channels reversed. Note: this sub-sample uses the same galaxies as the **faded** sub-sample. (3,927 galaxies)

- **goods-shallow:** Images from GOODS-North and GOODS-South, with color from 2 filters and imaged at 2-epoch depth. (6,144 galaxies)
- **stripe-82-single:** Single-epoch images from SDSS Stripe 82. (21,522 galaxies)
- **stripe-82-coadd:** Co-added images from SDSS Stripe 82. Note: this sub-sample includes all galaxies in the **stripe-82-single** sub-sample, with additional sources detected in the deeper imaging. (30,339 galaxies)
- **redshifted:** Simulated *HST* images constructed using SDSS images and artificially processed to redshifts between $0.3 \leq z \leq 1$. (288 original galaxies; 6,624 redshifted images)
- **simulated-agn:** Simulated AGN host galaxies constructed using *HST* images and PSFs. (96 original galaxies; 2,961 images of simulated AGN hosts)

2.6. Galaxy metadata

Photometric data for the bulk of the GZH **main**, **faded**, **recolored**, and **goods-shallow** samples were largely drawn from the tables in Griffith et al. (2012). This included photometric parameters such as the fluxes, magnitudes, radii, ellipticities, position angles, and positions drawn from both SExtractor and GALFIT. All photometric parameters were measured in both bands of the ACS imaging, with the exception of the single-band COSMOS images. Photometric data

for the GOODS 5-epoch imaging, including SExtractor parameters, is from Giavalisco (2012).

Redshifts for the GZH catalog were compiled from a variety of sources. For each galaxy, the primary redshift is in the Z_BEST column of Table 4. The redshift type (spectroscopic: SPEC_Z, photometric: PHOTO_Z, or grism: GRISM_Z) is listed in the column Z_BEST_TYPE, and the source catalog of the redshift is included as Z_BEST_SOURCE.

For galaxies which have published redshifts from multiple sources, the following algorithm was used to select the Z_BEST quantity. A high-quality spectroscopic redshift in the ACS-GC is the primary option, provided in the ACS-GC (Griffith et al. 2012), 3DHST (Momcheva et al. 2015), and MUSYC (Cardamone et al. 2010) catalogs and used in that order. For galaxies with multiple spectroscopic redshifts, more than 98% are consistent ($\Delta z < 0.001$), and so the order of selection made no practical difference. Galaxies with inconsistent spectroscopic redshifts between any pair of catalogs are marked with a flag in Table 4. If no spectroscopic redshifts were available, the 1- σ errors of the photometric (ACS-GC, 3DHST, MUSYC, UltraVISTA; Ilbert et al. 2013) and UltraVISTA grism data were used to select the estimated redshift with the smallest error.

Photometric and spectroscopic data for the stripe-82-single and stripe-82-coadd galaxies were taken from the CasJobs DR7 tables. This included *ugriz* Petrosian magnitudes and fluxes, as well as the relative de Vaucouleurs and exponential fits from the model magnitudes. All redshifts used for SDSS galaxies were spectroscopic. 82.6% of galaxies in the stripe-82-single images and 65.1% of galaxies in the stripe-82-coadd images had a measured DR7 spectroscopic redshift.

The technique for redshift debiasing (Section 4) requires consistent measurements of the galaxy surface brightness. For both the redshifted calibration images and the *HST* images which have their morphologies corrected, we calculate the mean surface brightness μ within the :w effective radius (R_e) as:

$$\mu = m + 2.5 * \log_{10} (2 \times (b/a) \times \pi R_e^2). \quad (1)$$

All photometric parameters are taken from SExtractor. For the FERENGI galaxies, m is MAG_AUTO in the I_{814W} band, (b/a) is the galaxy ellipticity (the profile RMS along the semi-major and -minor axes), and R_e is the 50% FLUX_RADIUS converted into arcsec (Melvin 2016). For the *HST* galaxies in the main sample, the parameters are identical except that we use MAG_BEST instead of MAG_AUTO (Griffith et al. 2012) in either the I_{814W} , i_{775W} , or z_{850LP} bands, depending on the available imaging (Table 1).

3. GZH INTERFACE AND CLASSIFICATIONS

Below we describe the classification structure of GZH, including the software interface and the hierarchical structure of a classification. Section 3.2 describes the process of combining individual classifications into vote fractions for each galaxy.

3.1. Interface and decision tree

Classifications for GZH were made using a web-based interface (Figure 3), similar in design to Galaxy Zoo and Galaxy Zoo 2. The front-end runs on a Ruby on Rails framework with classifications stored in a MySQL backend. Classifiers were shown a randomly-selected color composite image from the GZH sample; the default showed the image with a black sky background, although they had the option to invert the color palette if desired. The questions and responses for morphology appeared on the right side of the image as a panel, including both text and icons. There was no tutorial required for participation, although classifiers could access an extensive “Help” section containing example images and descriptive text for all the morphological labels.

The procedure for classifying an image in GZH followed a hierarchical decision tree (Figure 4). Every classification began with the step of identifying whether the object at the center of the image was a smooth galaxy, a galaxy with a disk or other features, or a star/artifact. Subsequent questions in the tree depended on the previous answer(s) given by the classifier; the decision tree was designed so that every question relevant to the morphology in the process of being identified was answered. Questions that were not answered were implicitly assumed to be absent in the image — for example, if the classifier identified a galaxy as being smooth, they were not asked to count the number of spiral arms. For every task, the classifier chose a single answer before continuing to the next question; they also had the option to restart any classification in-progress.

The GZH decision tree was designed to remain close to that used by GZ2 while taking account of the likely differences in high redshift systems, and has four broad sets of morphologies classified by the users. The first set identified stars or image artifacts (the result of either bad data or incorrect identification as a galaxy by the ACS pipeline); in this case, the classification process ended and no further questions were asked. The second set was smooth galaxies, intended to select ellipticals/early-types; volunteers also indicate the relative axial ratio (roundness) for these galaxies. The third set was for disk/late-type galaxies, which labeled the features necessary to place a galaxy on the standard Hubble tuning fork (bars, spiral arm, strength of the central bulge). The final set, which was new in this




Figure 3. Screenshot of the GZH interface (<http://zoo3.galaxyzoo.org>) at the beginning of a classification, with the classifier ready to select an answer for the first question in the decision tree.

phase of Galaxy Zoo and designed for high-redshift targets, identified objects dominated by clumpy morphologies. Further annotations for clumpy galaxies included assessing the number, arrangement, relative brightness, and location of the clumps within the galaxy. Finally, every classification of a galaxy gave the option of identifying “odd” features within the image; these labels were for relatively rare ($\lesssim 1\%$) phenomena, including dust lanes, gravitational lenses, and mergers.

The number of independent classifications per subject collected by GZH was on average higher than GZ1 or GZ2, due to both the increased complexity of the decision tree and the relative difficulty of classifying images of small and distant galaxies. Images from the **main** AEGIS, GEMS, and GOODS data sets had a median of 122 independent classifications per image. The remainder of the images either had a later activation date (**main** COSMOS, **simulated-agn**) or a lower retirement limit (**stripe-82-single** and **stripe-82-coadd**). Images from these samples had a median of 46–48 classifications per image (Figure 5).

The GZH project was launched on 23 Apr 2010 with the inclusion of the AEGIS, GEMS, GOODS 2-epoch, and SDSS Stripe 82 images. Images from COSMOS and the simulated AGN were activated in Dec 2010, as well

as a small sample of images from AEGIS, GEMS, and GOODS that were previously excluded from the original sample due to cuts on blended and/or saturated objects and subsequently confirmed as classifiable galaxies. The GZH site collected data until its replacement, the fourth phase of Galaxy Zoo² (including data from both the *HST* CANDELS survey and SDSS DR8) began on 10 Sep 2012. Classifications for the GOODS 5-epoch images were separately obtained from Mar-Jun 2015 using the fourth version of the Galaxy Zoo site and the same GZH decision tree. The GZH project had a total of 10,349,357 classifications from 93,898 registered participants.

3.2. Classifier weighting

As a first step to producing a consensus measurement of each galaxy, the votes of individual volunteers who classified galaxies in GZH were combined to make a vote fraction for each response ($f_{response}$) to a question in the decision tree. Votes were subsequently weighted and re-combined in an iterative method similar to that in previous versions of Galaxy Zoo (Land et al. 2008;

² <http://zoo4.galaxyzoo.org>




Figure 4. Flowchart of the questions presented to GZH classifiers, labeled with the corresponding task numbers. Tasks in the decision tree are color-coded by tier level: Gray-colored tasks are 1st-tier questions which are asked in every classification. Tasks colored green, blue, purple, and pink (respectively) are one, two, three, or four steps below branching points in the decision tree. The dashed line between T15 and T03 indicates the unique case where volunteers could label both clumpy and disk-like morphologies for galaxies with clumps arranged in a spiral pattern.




Figure 5. Distribution of the total number of classifications per image for GZH, split by survey.

Willett et al. 2013), using a method chosen to be as egalitarian as possible while also identifying and down-weighting classifiers who frequently disagreed with others. The weighting factor w was 1 for the top 95% of classifiers, as ranked by consistency. For the bottom 5% of classifiers, w was designed to drop smoothly and was effectively zero for the bottom 1% in the distribution function. Since downweighting only occurs for the bottom few percent of classifiers (and an even smaller percentage of the classifications), the overall effect on the GZH dataset was minimal. The method was effective, however, at filtering out contributions from randomly classifying or malicious participants.

Classifications were aggregated and weighted only if the classifier was logged into GZH under their username (which was encouraged, but not required for participation). Classifications by participants who were not logged in were marked as “Anonymous” and receive the same $w = 1$ weighting as the vast majority of logged-in classifiers.

4. CORRECTING FOR REDSHIFT-DEPENDENT CLASSIFICATION BIAS

The previous versions of Galaxy Zoo morphology classifications (Lintott et al. 2011; Willett et al. 2013) were based on observations of galaxies in the SDSS, which have a median redshift of $z < 0.2$. In these cases, it was assumed that there was no cosmological evolution of the morphologies of galaxies and therefore any observed changes in the morphological distribution were

due to a redshift-dependent bias that affects image quality and classification accuracy (*i.e.*, galaxies of a given mass/size will appear smaller and dimmer at higher redshifts). This bias is not unique to the crowdsourcing technique; its dependence in data quality is a potential problem for *both* automated and visual classifications, and must be addressed in order to accurately measure demographics over any significant redshift range.

For both previous releases of Galaxy Zoo morphologies, a correction for redshift-dependent bias was applied based on matching the mean classification fractions at the highest redshifts with those at the lowest redshift. Bamford et al. (2009) and Willett et al. (2013) provide complete descriptions of the process for GZ1 and GZ2, respectively.

In GZH, the redshift range is large enough that cosmological evolution of the types and morphologies of galaxies is expected for the *HST* sample. In addition, the effects of band shifting will change the images even more across these redshift ranges. As a result, the previous methods of correcting for redshift-dependent bias do not work.

In order to test and correct for the effects of redshift, GZH includes a set of calibration images. These are simulated images of a set of nearby galaxies as they would appear observed at a variety of redshifts. The input images are from the SDSS (York et al. 2000; Strauss et al. 2002). The sample consists of 288 well-resolved galaxies at $z < 0.013$. The galaxies spanned a variety of morphologies (as selected by GZ2 classifications, including identifications of spiral structure, ellipticals, mergers, edge-on disks, bulge prominence/shape, and bars) and r -band surface brightnesses. The overall selection spanned the redshift range of SDSS targets and maximized the number of *HST* galaxies at the same surface brightness and redshift (Figure 6). The FERENGI code was used to produce sets of images corresponding to observations of these galaxies out to $z_{\text{sim}} = 1.0$ (see Section 2.3.2). The resulting images were classified in the GZH interface using the standard classification scheme.

4.1. Correcting morphologies for classification bias

The approach used in GZH for correcting the weighted classifications for redshift bias rests on the assumption that the *degree* of bias is a function of the apparent size and brightness of the galaxy. This is controlled by two types of parameters: **intrinsic** properties of the galaxy itself, such as its physical diameter and luminosity, and **extrinsic** properties, such as the distance (redshift) of the galaxy and its relative orientation. The combination of all such parameters forms a high-dimensional space, and it is not clear how to separate this into individual effects. Instead, the method used here employs only two parameters as a simple model of the effect of redshift




Figure 6. Surface brightness as a function of redshift for 3,449 FERENGI images and the 102,548 **main** galaxies with measured μ and z values. The color histogram shows the number of FERENGI images as a function of μ and z_{sim} . White contours show counts for the galaxies in the **main** sample, with the outermost contour starting at $N = 1500$ and separated by intervals of 1500.

brightness on classification: the surface brightness (μ ; intrinsic) and redshift (z ; extrinsic).

We model the debiased vote fraction $f_{\mu,z}$ for an artificially-redshifted galaxy as:

$$f_{\mu,z} = (f_{\mu,z=0.3}) \times e^{\frac{z-z_0}{\zeta}}, \quad (2)$$

where $f_{\mu,z=0.3}$ is the vote fraction at the lowest redshift in the artificially-redshifted sample ($z_0 = 0.3$). Here, ζ is a parameter that controls the rate at which f_{features} decreases with increasing redshift (and that may depend on other galaxy properties). This function bounds the observed vote fractions between $f_{\mu,z=0.3}$ and zero based on two assumptions: 1) the vote fractions for featured galaxies decrease monotonically with increasing redshift, therefore restricting vote fractions for a given galaxy to be less than $f_{\mu,z=0.3}$ (which is almost always true of the data), and 2) the vote fractions cannot be less than zero (which is always true).

Figure 7 shows the change in vote fraction and the best-fit model for a random set of galaxies in the FERENGI sample. The results show that there is a clear dependence on redshift for the observed changes in f_{features} . To examine whether a better fit should involve additional parameters, we examine the global dependence of ζ on the galaxy surface brightness. This is motivated by the fact that brighter galaxies would presumably be easier to identify; Bamford et al. (2009) and Willett et al. (2013) found changes in f_{features} depending on a combination of physical size and absolute magni-

tude. To simplify our model, we subsume these into a single parameter μ .

Figure 8 examines the results of fitting the FERENGI images with Equation 2. This fit only includes galaxies where SExtractor robustly measured the photometry in all images between $0.3 < z_{\text{sim}} < 1.0$ (such that $\sigma_{M_I} < 1.0$ mag; Melvin 2016). Interestingly, the derived correction is only a very weak function of surface brightness. Higher-surface brightness galaxies have on average slightly stronger corrections, possibly because these galaxies have larger f_{features} values at high redshifts. Low surface brightness galaxies are more likely to begin low and remain low; the bounded nature of the dropoff (and variance among the individual voters) means that the average magnitude of ζ will be lower.

Since there is little evidence for any strong systematic dependence of ζ on μ , we do not include any additional parameters in fitting to our calibration model. We fit the data in Figure 8 with a linear function:

$$\log_{10}(\hat{\zeta}) = \zeta_0 + (\zeta_1 \times \mu), \quad (3)$$

where $\hat{\zeta}$ is the correction factor applied to each galaxy as a function of surface brightness. The best-fit parameters to the linear fit (from least-squares optimization) are $\zeta_0 = 0.50$, $\zeta_1 = -0.03$. To make the final debiased correction, the simple exponential form of Equation 2 is modified to bound the debiased vote fractions between f and 1:

$$f_{\text{features,debiased}} = 1 - (1 - f_{\text{features,weighted}})e^{\frac{z-z_0}{\hat{\zeta}}} \quad (4)$$

where $f_{\text{features,weighted}}$ is the weighted vote fraction described in Section 3.2.

4.2. Effects of morphological debiasing

Figure 9 shows the change in f_{features} as a function of their lowest simulated redshift for the 3,449 FERENGI images with robust photometric measurements across the full range of redshifts. For each simulated redshift value z_{sim} at a fixed surface brightness, $f_{\text{features},z}$ is the value measured at that simulated redshift. We plot $f_{\text{features},z}$ against $f_{\text{features},z=0.3}$, the value measured for the same galaxy at $z_{\text{sim}}=0.3$.

The objective is to use these data to predict, for a galaxy with a measured $f_{\text{features},z}$ value, what its f_{features} value *would have been* if it had been viewed at $z = 0.3$. This predicted value is defined as the “debiased” vote fraction $f_{\text{features,debiased}}$, and is calculated by applying a correction to the measured value of f_{features} , determined by the ζ function described in the previous section. A reliable predicted value can be obtained so long as the relationship between $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$ is single-valued; that is, for a given $f_{\text{features},z}$, there is




Figure 7. Behavior of the normalized, weighted vote fractions of features visible in a galaxy (f_{features}) as a function of redshift in the artificial FERENGI images. Galaxies in this plot were randomly selected from a distribution with $e = 0$ and at least three detectable images in redshift bins of $z \geq 0.3$, and are sorted by $f_{\text{features},z=0.3}$. Measured vote fractions (blue solid line) are fit with an exponential function (red dashed line; Equation 2); the best-fit parameter for ζ is given above each plot.

exactly one corresponding value of f_{features} at $z = 0.3$.

Figure 9 shows that the relationship between $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$ is *not* always single-valued; hence, it is not appropriate to correct galaxies that lie in certain regions of surface brightness/redshift/ f_{features} space. These regions tend to have low f_{features} values at high redshift, but a wide range of values at $z = 0.3$. These regions contain two morphological types of galaxies: the first set are genuine ellipticals, which have low values of f_{features} at both high and low redshift. The second group are disks whose features become indistinct at high redshift; hence their f_{features} value at $z = 0.3$ may be quite high, while the value observed at high redshift is very low. This effect is strongest at high z and low μ , where features become nearly impossible to discern in the images (see top right panel of Figure 9).

The criteria for determining whether a region of this

space is single-valued, and therefore correctable, is as follows: In each surface brightness and redshift bin, the relationship between $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$ is modeled by fitting the data with polynomials of degrees $n = 3, 2$, and 1 , and using the best formal fit out of the three as measured by the sum of the residuals. These fits are shown as the dashed black lines in Figure 9. Flat regions of the bins are areas in which there is not a clear single-valued relationship between $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$. We quantify this by measuring the slope of the best-fit polynomial; regions of the bins with a slope less than 0.4 are considered *not* one-to-one, and therefore $f_{\text{features},z}$ cannot be boosted to its $f_{\text{features},z=0.3}$ value. Galaxies in this region are referred to as the *lower limit* sample, because the most stringent correction available is that the weighted f_{features} is a lower limit to the true value. These regions are high-




Figure 8. All fits for the FERENGI galaxies of the vote fraction dropoff parameter ζ for f_{features} as a function of surface brightness. This includes only galaxies with a bounded range on the dropoff ($-10 < \log(\zeta) < 10$) and sufficient points to fit each function (37 original galaxies, each artificially redshifted in eight bins over a range from $0.3 \lesssim z_{\text{sim}} \lesssim 1.0$).

lighted in blue in Figure 9. Uncolored (white) regions of the plot have sufficiently high slopes to consider the relationship as single-valued; galaxies in these regions are considered “correctable”, and only these are used in measuring the parameters for the ζ function (Section 4.1). Only surface brightness/redshift bins with at least 5 galaxies were considered; regions with fewer than 5 galaxies are considered to have “not enough information” to determine the $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$ relationship, colored gray in Figure 9. These galaxies are hereafter referred to as the “NEI” sample.

The unshaded regions in Figure 9 define discrete ranges of redshift, surface brightness, and f_{features} a galaxy must have in order for the debiased correction to be confidently applied to a galaxy in the GZH sample. While the appropriate correctable regions were defined discretely, the true correctable region is assumed to be a smooth function of z , μ , and f_{features} . To define this smooth space, we calculate the shape of the convex hull that encloses the correctable and lower-limit FERENGI galaxies in z - μ - f_{features} space. The boundaries are then adjusted until the contamination from both groups is minimized. The resulting hulls define the correctable and lower-limit regions for categorizing the *HST* galaxies. The results of this method and final categorization of the *HST* sample are in Table 3. Of the galaxies at redshift higher than $z = 0.3$, 17% can be debiased us-

Table 2. Distribution of FERENGI images analyzed in Figure 9. Correctable images have a single-valued relationship between their measured f_{features} values at high and low redshifts (white regions in Figure 9). Images with only a lower-limit on f_{features} have a non single-valued relationship (blue regions). NEI (“not enough information”) images have undetermined relationships due to a lack of data ($N < 5$) in their corresponding z - μ bins (gray regions).

	N	%
Correctable	1,690	49%
Lower-limit	1,678	49%
NEI	81	2%
Total	3,449	100%

ing the ζ method, 27% cannot be debiased since the relationship between $f_{\text{features},z}$ and $f_{\text{features},z=0.3}$ is not monotonic, and 56% cannot be debiased since they either have an unknown redshift or insufficient numbers of FERENGI images in their z - μ value to determine a reliable value of ζ .

For the “lower-limit” galaxies for which a single debiased f_{features} value cannot be confidently assigned, the range of debiased values is estimated using a method visualized in Figure 10. This uses the FERENGI simulated data to analyze the range of intrinsic $f_{\text{features},z=0.3}$ values for any given observed f_{features} value, again as a function of surface brightness and redshift. In each unclassifiable (shaded blue) z - μ bin, the spread of intrinsic values of $f_{\text{features},z=0.3}$ for four ranges of observed f_{features} is computed. The range of intrinsic values is defined as the inner 80% of the data, represented by the orange bars in Figure 10. For any galaxy which cannot be directly debiased by the ζ method, these ranges are used to denote the upper and lower limits on the expected values $f_{\text{features},z=0.3}$ as a function of the observed f_{features} .

Few galaxies in the sample have sufficiently high corrections to completely change them from being confidently “smooth” to “featured” following the correction for redshift bias (Figure 11). As a check, highly boosted galaxies are compared to the expert classifications in CANDELS (Kartaltepe et al. 2015). There are only eight galaxies that were strongly boosted ($f_{\text{features}} < 0.2$ and $f_{\text{features,best}} > 0.5$) in GZH and also appears in the CANDELS expert sample. Of those eight galaxies, Kartaltepe et al. (2015) classify 5 as spheroids, 2 as disks, and 1 as irregular/disky. The $f_{\text{features,best}}$ values for GZH are all between 0.5 and 0.6, making them intermediate disk candidates. One of the spheroids has a hint of extended emission to the south that may have been missed by the CANDELS experts, but almost all images are dim and with very low surface brightnesses in the ACS imaging, making positive identification challenging. Since the surveys use different rest-frame filters and the overlap sample is tiny, though, detailed compar-




Figure 9. Effects of redshift bias in 3,449 images in the FERENGI sample. Each point in a given redshift and surface brightness bin represents a unique galaxy. On the y -axis in each bin is the f_{features} value of the image of that galaxy redshifted to the value corresponding to that redshift bin. On the x -axis is the f_{features} value of the image of the same galaxy redshifted to $z = 0.3$. The dashed black lines represent the best-fit polynomials to the data in each square. The solid black line represents $f_{\text{features},z=f_{\text{features},z=0.3}}$. Regions in which there is a single-valued relationship between f_{features} at high redshift and at $z = 0.3$ are white; those in which there is not are blue, and those with not enough data ($N < 5$) are gray. A larger version of the bin outlined at $z = 1.0$ and $20.3 < \mu < 21.0$ (mag/arcsec 2) is shown in Figure 10.




Figure 10. A larger version of the dark-outlined square in Figure 9, containing FERENGI galaxies that have been artificially redshifted to $z = 1.0$ and have surface brightnesses between $20.3 < \mu < 21.0$ (mag/arcsec 2). The orange bars represent the inner 80th percentiles of the data, the boundaries of which are the lower and upper limits of the debiased values.

isons between the overall morphologies are highly difficult.

4.3. Challenges of debiasing questions beyond “smooth or features”

As with the HST images, each FERENGI subject had a varying number of classifiers answering the various questions in the hierarchical decision tree. Every classifier answers the first question, “*Is the galaxy smooth and rounded, with no sign of a disk?*”; as such the vote fractions f_{smooth} , f_{features} , and f_{artifact} are all computed with the minimum statistical error for any question, with between 40 and 140 responses (see Section 3). The number of participants answering subsequent questions, however, is always equal to or less than the number who answered the preceding question. The average number of responses per task for fourth- or fifth-tier questions (such as spiral arm structure; Tasks 12–14) is only 4 ± 4 for the FERENGI sample. While this distribution is strongly bimodal (reflecting the true morphologies of selected galaxies), the very low absolute numbers of votes introduce very high variance when attempting to calculate a statistical correction.

In the FERENGI data, these numbers severely limit the amount of information that can be extracted for the higher-tier questions. The debiasing technique used (Section 4.1) requires that at least 10 classifiers answer each question for a galaxy with $z_{\text{sim}} = 0.3$ and the corre-

sponding image at higher redshift. This requirement is (by default) met by all galaxies for the smooth/features question. However, this is often *not* met for questions beyond Task 01. On average, $60\% \pm 24\%$ of the galaxies do not have sufficient data to measure a correction, as compared to 2.0% achieved for Task 01 (Table 2). Of the remaining FERENGI galaxies which did meet this requirement, polynomials were fit to the data in each redshift-surface brightness bin, as was done for Task 01. The goodness-of-fit was evaluated in each bin, for each Task, using a normalized χ^2 metric. The mean χ^2 for all higher-order tasks was 0.10 ± 0.04 , significantly larger than the Task 01 value of 0.04. Based on this statistic, the data cannot not be modeled with sufficient confidence for the higher order tasks. An example of the debiasing technique applied to f_{bar} is shown in Section A.2; visual inspection of Figure A5 confirms that the points in each bin are scattered to a degree at which no strong correlation is present. For these reasons, debiased vote fractions are only offered in the GZH catalog for Task 01 (smooth/features).

5. THE GALAXY ZOO: HUBBLE CATALOG

The catalog for GZH includes morphological data for 189,149 images (generated from a total of 145,741 unique galaxies). The full table can be accessed at <http://data.galaxyzoo.org>. The online data also includes a secondary metadata table, which is drawn from a variety of sources detailed in Section 2.

Each image is listed under a unique project ID (eg, AHZ000001); the actual galaxy in the image is identified by the combination of the OBJNO and original survey. For each of the 55 responses in the GZH decision tree, the following classification data is provided: for each question, N_{votes} is the number of classifiers who answered that question. For each unique answer, fraction is the fraction of classifiers who selected that answer ($N_{\text{answer}}/N_{\text{votes}}$), and weighted is the weighted fraction, which takes into account overall consistency (Section 3.2).

The GZH vote fractions can be largely dependent on the resolution of the image. Two otherwise morphologically identical galaxies which differ significantly in redshift, brightness, or size may result in very different vote fractions for any given question, given that many features of a galaxy are difficult to discern in less-resolved images (bars, spiral arms, disk structure, etc). For this reason, caution must be used when taking vote fractions as cut-offs to determine morphological structure; guidelines for careful classification are given in Section 6.

The GZH catalog is corrected for redshift bias only for the first question of the GZH decision tree (Section 4), which asks “*Is the galaxy smooth and round, with no sign of a disk?*” For this question, the catalog pro-




Figure 11. Debiased f_{features} corrected to $z = 0.3$ vs weighted f_{features} for the correctable (left), lower-limit (middle), and “not enough information” (NEI; right) galaxies in the GZH sample.

Table 3. Correctable fractions for the top-level task in GZH, split by survey.

	Correction type	AEGIS	COSMOS	GEMS	GOODS-N		GOODS-S	Total
					5-epoch	5-epoch		
correctable	0	2,908	21,169	2,802	1,459	1,189	29,527	
lower-limit	1	833	5,169	1,021	1,377	1,267	9,667	
no correction needed ($z \leq 0.3$)	2	955	10,870	1,175	415	400	13,815	
not enough information (NEI)	3	2,677	43,058	3,559	2,077	2,184	53,555	
no redshift information	4	1,134	4,688	530	687	102	7,141	
total		8,507	84,954	9,087	6,015	5,142	113,705	

vides the additional parameters `debiased`, `lower_limit`, `upper_limit`, and `best` vote fractions. The `best` fraction for f_{features} is chosen based on the categorization of the galaxy: if it is “correctable”, `best = debiased`; if it is a lower limit, `best = lower_limit`; if neither condition applies, then `best = weighted`.

The debiased and best vote fractions for f_{smooth} are calculated on the criteria that vote fractions for all answers must sum to unity:

$$f_{\text{smooth}} \equiv 1 - f_{\text{features,best}} - f_{\text{artifact}}. \quad (5)$$

In rare cases (1.2% of the `main` sample), this requirement resulted in negative vote fractions for f_{smooth} ; these were cases in which the f_{features} vote fraction was boosted to a high value relative to f_{artifact} . In these cases, the constraint of Equation 5 is met by setting $f_{\text{smooth}}=0.0$ and decreasing $f_{\text{features,best}}$ accordingly. This correction

is typically small, with a median change of f_{features} or f_{smooth} of $\Delta f = 0.04$.

Tables 4–10 also contain flags for Task 01 that identify clean (but not complete) samples of galaxies with high likelihoods of being either smooth, featured, or a star/artifact. These are set as `smooth/featured/artifact_flag = 1` if $f_{\text{smooth/features/artifact,best}} > 0.8$. For a galaxy to be flagged as “smooth”, an additional criterion of `correction_type = 0, 1, or 2` is applied. This accounts for the uncertainty in distinguishing between genuine ellipticals and disks whose features have been washed out due to surface brightness and redshift effects (Section 4).

Data products for GZH are split by the type of image being classified. Each sample in Section 2.5 corresponds to the data in Tables 4–10.

Table 4. GZH morphological classifications for *HST* images from AEGIS, COSMOS, GEMS, and GOODS

Zooniverse ID	Survey ID	Imaging	t01_smooth_or_features_			t01_smooth_or_features_a01_smooth_					
			Correction ¹	N _{votes}	fraction	weighted	debiased	best	lower limit	upper limit	flag
AHZ100002g	10010842	AEGIS	0	127	0.118	0.128	0.085	0.085	0.226	0.226	0
AHZ100002h	10010870	AEGIS	4	127	0.567	0.592	0.927	0.592	—	—	0

Table 4 (*continued*)

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>						flag
			Correction ¹	N _{votes}	fraction	weighted	debiased	best	lower limit	upper limit		
...												
AHZ20004kd	20014731	COSMOS	3	44	0.682	0.675	0.147	0.675	—	—	—	0
AHZ20004ke	20014732	COSMOS	2	45	0.689	0.756	0.893	0.756	—	—	—	0
...												
AHZ400043g	90022729	GEMS	1	121	0.702	0.733	0.487	0.734	0.483	0.800	0.800	0
AHZ4000416	90022735	GEMS	1	127	0.646	0.698	0.508	0.698	0.171	0.727	0.727	0
...												
AGZ0007z47	10014	GOODS-N-FULLDEPTH	1	40	0.475	0.475	0.197	0.475	0.011	0.496	0.496	0
AGZ0007z48	10017	GOODS-N-FULLDEPTH	3	40	0.675	0.675	0.048	0.675	0.168	0.669	0.669	0
...												
AGZ00083jb	8869	GOODS-S-FULLDEPTH	1	40	0.425	0.425	0.109	0.425	0.070	0.548	0.548	0
AGZ00083jc	8878	GOODS-S-FULLDEPTH	0	40	0.205	0.205	0.048	0.048	-0.005	0.287	0.287	0
...												

¹ Flag indicating how the vote fractions for this galaxy were corrected through debiasing (Section 4.2), if possible. 0 = correctable, 1 = lower-limit ($f_{\text{raw}} - f_{\text{adj}}$ single-valued), 2 = uncorrected ($z < 0.3$), 3 = uncorrected (insufficient FERENGI galaxies in this z - μ bin), 4 = uncorrected (no galaxy redshift available).

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 113,705 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 5. GZH morphological classifications for color-faded *HST* images

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>						...
			Correction	N _{votes}	fraction	weighted	debiased	best	lower limit	upper limit	flag	
AHZF000001	20000002	COSMOS	1	48	0.708	0.755	0.228	0.754	0.325	0.829	0	
AHZF000003	20000004	COSMOS	3	49	0.367	0.379	0.100	0.379	0.198	0.198	0	
AHZF000004	20000006	COSMOS	3	49	0.265	0.271	0.010	0.270	—	—	0	
AHZF00000z	20000102	COSMOS	1	44	0.727	0.78	0.233	0.780	0.316	0.820	0	
AHZF000010	20000104	COSMOS	2	53	0.811	0.849	0.904	0.848	—	—	0	
...												

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 3,927 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 6. GZH morphological classifications for color-inverted *HST* images

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>						...
			Correction	N _{votes}	fraction	weighted	debiased	best	lower limit	upper limit	flag	
AHZC000001	20000002	COSMOS	1	168	0.615	0.664	0.160	0.663	0.271	0.775	0	
AHZC000003	20000004	COSMOS	0	235	0.333	0.364	0.002	0.002	0.063	0.063	0	
AHZC000004	20000006	COSMOS	3	316	0.235	0.252	-0.011	0.252	—	—	1	
AHZC00000z	20000102	COSMOS	1	207	0.755	0.757	0.272	0.756	0.249	0.796	0	
AHZC000010	20000104	COSMOS	2	158	0.843	0.882	0.936	0.881	—	—	1	
...												

Table 6 *continued*

Table 6 (*continued*)

Zooniverse ID	Survey ID	Imaging	Correction	<i>N_{votes}</i>	fraction	weighted	debiased	best	lower limit	upper limit	flag	...
---------------	-----------	---------	------------	--------------------------	----------	----------	----------	------	-------------	-------------	------	-----

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 3,927 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 7. GZH morphological classifications for GOODS 2-epoch images

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>					...	
			Correction	<i>N_{votes}</i>	fraction	weighted	debiased	best	lower limit	upper limit	flag	
AHZ3000001	50000000	GOODS-N	0	123	0.390	0.415	0.090	0.090	—	—	0	
AHZ3000002	50000001	GOODS-N	2	126	0.341	0.355	0.356	0.356	0.220	0.279	0	
AHZ3000003	50000005	GOODS-N	1	129	0.760	0.826	0.633	0.825	0.596	0.834	0	
AHZ3000004	50000008	GOODS-N	1	120	0.758	0.787	0.639	0.787	0.658	0.834	0	
AHZ3000005	50000010	GOODS-N	1	123	0.854	0.890	0.611	0.889	0.597	0.914	0	
...												

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 6,144 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 8. GZH morphological classifications for SDSS Stripe 82 single-epoch images

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>					...	
			Correction	<i>N_{votes}</i>	fraction	weighted	debiased	best	flag	...		
AHZ5000001	587730845812064684	SDSS	2	41	0.585	0.595	0.759	0.594	0			
AHZ5000002	587730845812065247	SDSS	2	46	0.609	0.651	0.897	0.651	0			
AHZ5000003	587730845812196092	SDSS	2	51	0.039	0.044	0.067	0.043	0			
AHZ5000004	587730845812196825	SDSS	2	35	0.514	0.605	0.928	0.605	0			
AHZ5000005	587730845812524122	SDSS	2	47	0.766	0.812	1.038	0.810	1			
AHZ5000006	587730845812654984	SDSS	2	42	0.500	0.542	0.680	0.541	0			
AHZ5000007	587730845812655541	SDSS	2	41	0.488	0.526	0.697	0.525	0			
AHZ5000008	587730845812720365	SDSS	2	53	0.792	0.84	1.050	0.839	1			
AHZ5000009	587730845812720640	SDSS	4	43	0.000	0.0	0.000	0.0	0			
AHZ500000a	587730845812720699	SDSS	2	40	0.425	0.478	0.588	0.477	0			
...												

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 21,522 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 9. GZH morphological classifications for SDSS Stripe 82 co-added images

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>			<u>t01_smooth_or_features_a01_smooth_</u>					...
			Correction	<i>N_{votes}</i>	fraction	weighted	debiased	best	flag	...	
AHZ6000001	8647474690312306978	SDSS	4	40	0.275	0.289	0.762	0.289	0		

Table 9 *continued*

Table 9 (*continued*)

Zooniverse ID	Survey ID	Imaging	<u>t01_smooth_or_features_</u>		<u>t01_smooth_or_features_a01_smooth_</u>				...	
			Correction	N_{votes}	fraction	weighted	debiased	best	flag	
AHZ6000002	8647474690312307154	SDSS	2	43	0.605	0.634	0.858	0.635	0	
AHZ6000003	8647474690312307877	SDSS	2	51	0.608	0.627	0.906	0.627	0	
AHZ6000004	8647474690312308301	SDSS	4	52	0.038	0.038	0.723	0.038	0	
AHZ6000005	8647474690312308318	SDSS	2	44	0.614	0.632	0.776	0.631	0	
AHZ6000006	8647474690312308880	SDSS	2	36	0.667	0.683	0.901	0.683	0	
AHZ6000007	8647474690312372644	SDSS	4	48	0.646	0.674	1.145	0.674	0	
AHZ6000008	8647474690312372789	SDSS	4	45	0.489	0.571	0.964	0.570	0	
AHZ6000009	8647474690312372931	SDSS	4	47	0.553	0.587	0.926	0.587	0	
AHZ600000a	8647474690312373190	SDSS	4	47	0.574	0.559	1.008	0.559	0	
...										

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 30,339 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

Table 10. GZH morphological classifications for *HST* images with simulated AGN

Zooniverse ID	Survey ID	Imaging	Correction	Version	L_{ratio}	AGN color ¹	N_{votes}	<u>t01_smooth_or_features_a01_smooth_</u>				
								fraction	weighted	debiased	best	lower limit
AHZ7000001	90024700	GEMS	1	1	0.2	1	42	0.238	0.239	-0.110	0.238	-0.113
AHZ7000002	90024700	GEMS	1	1	1.0	1	51	0.255	0.265	-0.107	0.264	-0.128
AHZ7000003	90024700	GEMS	0	1	5.0	1	47	0.170	0.167	-0.018	-0.018	-0.049
AHZ7000004	90024700	GEMS	0	1	10.0	1	41	0.195	0.195	0.045	0.045	0.044
AHZ7000005	90024700	GEMS	0	1	50.0	1	47	0.170	0.178	0.067	0.067	0.146
...												
AHZ700013m	90024700	GEMS	0	2	0.0	0	35	0.171	0.136	0.011	0.011	0.029
AHZ700013n	90024700	GEMS	1	2	0.2	1	20	0.150	0.158	-0.278	0.049	-0.351
AHZ700013o	90024700	GEMS	1	2	1.0	1	32	0.281	0.300	-0.086	0.281	-0.119
AHZ700013p	90024700	GEMS	0	2	5.0	1	29	0.103	0.115	-0.098	-0.098	-0.152
AHZ700013q	90024700	GEMS	0	2	10.0	1	35	0.171	0.181	0.027	0.027	0.023
AHZ700013r	90024700	GEMS	0	2	50.0	1	34	0.206	0.206	-0.005	-0.005	-0.056
...												

¹ Flag indicating the color of the PSF in the simulated AGN. 0 = no simulated AGN, 1 = blue, 2 = flat, 3 = red.

NOTE—The full version of this table is available in electronic form, as well as at <http://data.galaxyzoo.org>. The complete version includes data for 2,961 galaxies and morphological information for all tasks in the tree. A subset of the information is shown here to illustrate form and content.

6. USING THE CATALOG

The primary purpose of the GZH catalog is to provide a reliable method of selecting galaxies of a desired morphological type. This section provides instructions for creating such pure samples using the vote fractions corresponding to the tasks shown in Figure 4 (eg, Masters et al. 2011; Cheung et al. 2015; Galloway et al. 2015). Increasing the levels of the thresholds can create purer, but not necessarily complete, morphologically-selected

samples. These are useful for selecting galaxies of rare or unique types that merit individual study. Looser cuts can be applied to obtain samples with a higher level of completeness, although the rate of false positives must be closely monitored. We stress that the details of any selection process will vary based on the particular science case; for example, Bamford et al. (2009); Skibba et al. (2009); Smethurst et al. (2015) all demonstrate the advantages of using the vote fractions directly as weights rather than applying discrete thresholds.

To select galaxies of a morphological type identified with a particular task, a cut is placed on the vote fraction for that task (f_{task}), *as well as* the vote fractions for the tasks preceding it, because of the dependency induced by the decision-tree structure. For example, to select barred galaxies, a cut may be placed on f_{bar} such that only galaxies where a high fraction of votes for this task voted for the *bar, yes* answer. This is not the only necessary cut, however, since not all classifiers answer this question; only those who have previously selected “features” in Task 01, “not clumpy” in Task 12, and “not edge-on” in Task 02 will have the opportunity to vote on the bar question, Task 03. To ensure that f_{bar} is well-sampled, cuts on all previous tasks must be applied.

The flexibility of this catalog allows users to set their own selection criteria for vote fraction thresholds to create a morphologically pure sample. Table 11 provides baseline cuts for selecting galaxies of a variety of morphologies. These thresholds are determined by a visual inspection of subsamples. The thresholds used are a combination of 20 votes for the task being considered, as well as limits on the vote fractions for previous response(s) in the decision tree.

We visually analyzed subsamples of 50 galaxies meeting both criteria, as well as a control sample of galaxies which had 20 classifiers vote on the task, but did not meet the threshold cut set for the previous task. The threshold cut was adjusted and new subsamples were inspected until both the original and control samples achieved $> 80\%$ purity.

As an example of how to use Table 11 to create a sample of 3-armed spiral galaxies, we suggest selecting objects with $N_{\text{arm number}} \geq 20$, $f_{\text{features}} > 0.23$, $f_{\text{clumpy,no}} > 0.30$, $f_{\text{edgeon,no}} > 0.25$, and $f_{\text{spiral,yes}} > 0.25$. These cuts define a sample of galaxies of “arm number candidates”; *i.e.*, galaxies for which answering the arm number question makes physical sense and the vote fraction $f_{\text{arm number}}$ is well-sampled. In this example, such galaxies are featured, non-clumpy, non-edge on spirals. At this point a cut can be made on $f_{\text{arm number}=3}$ to select spirals with three arms.

Tasks 03, 04 and 05 have an additional possible pathway; as shown in Figure 4, a classifier might also be shown this question if they select “featured/disk” in Task 01, “clumpy” in Task 12, two or more clumps in Task 16, and “spiral arrangement” in Task 15. After applying the appropriate thresholds for this path, $< 0.5\%$ of the galaxies which have ≥ 20 answers to these questions used this pathway to arrive at these Tasks. None of these images show obvious disk structure, although the clumps within are arranged in a spiral pattern.

This section described how to use Table 11 to select galaxies for which a particular task is reliably sampled. The following two examples extend this and show how

to use the vote fractions to obtain samples of galaxies with a specific morphological type.

6.1. Example 1: Selecting barred galaxies

A sample of barred disk galaxies is created by applying cuts on the previous tasks as listed in Table 11. We identify 11,049 “bar candidates” as disk galaxies for which bar identification can be attempted in GZH. These galaxies were selected by applying the cuts $N_{\text{bar}} \geq 20$, $f_{\text{features}} > 0.23$, $f_{\text{clumpy,no}} > 0.30$, and $f_{\text{edgeon,no}} > 0.25$. These galaxies are featured, non-clumpy, non-edge on galaxies. Of these, a pure sample of 730 barred disks was identified by applying a cut of $f_{\text{bar}} > 0.7$. A sub-sample of 50 galaxies were visually inspected and 94% were found to contain strong bars. A complete sample of strong and weak bars was created by applying a cut of $f_{\text{bar}} > 0.3$. This sample contained 3,218 galaxies, 86% of which were found to contain weak or strong bars through visual inspection.

The resulting bar sample can be used to estimate the redshift evolution of bar fraction; we find a steady decrease of $f_{\text{bar}} \sim 0.32$ at $z = 0.4$ to $f_{\text{bar}} \sim 0.24$ at $z = 1.0$. This decrease agrees with Melvin et al. (2014), although they report a lower overall bar fraction going from $f_{\text{bar}} = 0.22$ at $z = 0.4$ to $f_{\text{bar}} = 0.11$ at $z = 1.0$. The difference in total bar fraction is expected, as this analysis used a looser cut on f_{bar} , there is no luminosity cut, and the use of debiased values for f_{features} increases the total amount of disks in the sample compared to Melvin et al. (2014). However, the results from GZH indicate that if features are reliably identified, the bar is always visible; for galaxies at increasing redshifts, the discs fade first but the galactic bar remains visible (Melvin 2016). As a result, we do *not* recommend using an evolving cut on f_{bar} as a function of redshift for the selection of barred discs.

Another existing study using GZH to select barred galaxies was presented in Cheung et al. (2015). This study used a slightly different bar selection method to demonstrate that AGN hosts show no statistically significant enhancement in bar fraction or average bar likelihood compared to closely-matched inactive galaxies. We note that their technique matched test and control galaxies over the same redshift ranges, which minimizes the possible impact of redshift bias.

6.2. Example 2: Identifying clump multiplicity

Clumps are known to be a characteristic feature of galaxies at high redshift, and there is evidence they play a crucial role in the evolution of modern spirals, particularly in the formation of central bulges (Elmegreen et al. 2005; Elmegreen & Elmegreen 2014; Guo et al. 2015; Behrendt et al. 2016). Simulations show clumps migrate from the outer disk to the galactic center in only

a few orbital periods (Mandelker et al. 2015) and observations show increasing bulge to clump mass and density ratios as the Universe evolves since $z \sim 1.5$ (Elmegreen et al. 2009), suggesting that clumps coalesce over time to form the modern bulges of disk galaxies. GZH includes a “clumpy” path in the decision tree for the purposes of identifying clumps and investigating their evolution with redshift.

For galaxies identified as “clumpy” in GZH, the number of clumps can be determined using Task 16. Table 11 can be used to select 8,444 galaxies measured as “clumpy” using $f_{\text{features}} > 0.23$ and $f_{\text{clumpy, yes}} > 0.80$ to ensure the vote fractions for Task 16 are well-sampled. The clump number can be reasonably identified for 1,112 of the clumpy galaxies; for the remainder, the unique clumps were less distinguishable from each other and the exact number of clumps could not be deduced without careful visual inspection. In the 1,112 which did have distinguishable clumps, there are 61 one-clump galaxies using $f_1 \text{ clump} > 0.50$, 442 two-clump galaxies using $f_2 \text{ clumps} > 0.80$, 275 three-clump galaxies using $f_3 \text{ clumps} > 0.75$, 71 four-clump galaxies using $f_4 \text{ clumps} > 0.70$, and 263 galaxies with more than four clumps using $f_{>4 \text{ clumps}} > 0.70$. Alternatively, these data may be used to create more general samples of clumpy galaxies with few clumps and many clumps. A sample of 989 “few clumps” galaxies can be made using $(f_1 \text{ clump} + f_2 \text{ clumps}) > 0.5$ and 2,910 “many clumps” galaxies using $(f_3 \text{ clumps} + f_4 \text{ clumps} + f_{>4 \text{ clumps}}) > 0.5$.

7. ANALYSIS

7.1. Demographics of morphology

Any analysis of the morphological distribution of galaxies must properly consider morphology with respect to other physical properties of the sample, such as color, mass, size, environment, and redshift. We defer such analyses to later papers, and comment here only on a few broad characteristics of the overall GZH sample.

Figure 12 shows the breakdown of morphologies for the combined surveys in GZH as a flow diagram, where each node represents a task and the flows between the nodes represent the possible responses. The width of the flows is proportional to the number of galaxies associated with that response. Figure 12 uses a simple plurality vote to assign the combined set of labels for each galaxy. This method is likely to undersample some of the categories; visual inspection of images suggests that $f_{\text{features}} \gtrsim 0.3$ identifies disk-dominated or irregular galaxies reliably, for example. This method also does not use the suggested thresholds in Table 11.

Given these caveats, the overall distribution of galaxy types is significantly different than the low-redshift sample classified in SDSS imaging from GZ1 and GZ2. Lin-

Table 11. Suggested thresholds for selecting morphological samples from GZH.

No.	Task	Previous task(s)	Vote fraction threshold $N_{\text{task}} \geq 20$
01	smooth or features	—	—
02	edge on	01,12	$f_{\text{clumpy,no}} > 0.30$
03	bar	01,12,02	$f_{\text{edgeon,no}} > 0.25$
		01,12,16,15	$f_{\text{clumpy spiral}} > 0.65$
04	spiral arms	01,12,02	$f_{\text{edgeon,no}} > 0.25$
		01,12,16,15	$f_{\text{clumpy spiral}} > 0.65$
05	bulge prominence	01,12,02	$f_{\text{edgeon,no}} > 0.25$
		01,12,16,15	$f_{\text{clumpy spiral}} > 0.65$
06	odd yes/no	—	—
07	rounded	01	$f_{\text{smooth}} > 0.70$
08	odd feature	06	$f_{\text{odd, yes}} > 0.50$
09	bulge shape	01,12,02	$f_{\text{edgeon, yes}} > 0.40$
10	arms winding	01,12,02,04	$f_{\text{spiral, yes}} > 0.25$
11	arms number	01,12,02,04	$f_{\text{spiral, yes}} > 0.25$
12	clumpy	01	$f_{\text{features}} > 0.23$
13	bright clump	01,12,16	$f_{\text{one clump}} < 0.40$
14	bright central clump	01,12,16,13	$f_{\text{bright clump, yes}} > 0.50$
15	clump arrangement	01,12,16	$f_{\text{multiple clumps}} > 0.45$
16	clump count	01,12	$f_{\text{clumpy, yes}} > 0.80$
17	clumps symmetrical	01,12	$f_{\text{clumpy, yes}} > 0.80$
18	clumps embedded	01,12	$f_{\text{clumpy, yes}} > 0.80$

tott et al. (2011) found that elliptical galaxies exceeded spiral galaxies by a factor of $\sim 2 : 1$ in their spectroscopic sample if using a plurality vote criterion (although Bamford et al. 2009 show that this strongly depends on the selection method; spirals are the dominant population in a volume-limited sample at $z < 0.088$, for example). The weighted votes for GZH have smooth galaxies outnumbering disks and clumpy galaxies by a factor of $\sim 3 : 1$, however. The fraction of objects identified as stars or artifact is also much higher in the *HST* imaging. By plurality votes, these encompass only $\sim 0.1\%$ of images in SDSS (Willett et al. 2013), but 6% of images in GZH.

Within the sample of galaxies identified as “not smooth”, it is clear that the addition of the clumpy branch is necessary to describe a large fraction of the sample, since disk-dominated galaxies outnumber clumpy morphologies by less than a factor of 2. Disk galaxies are primarily unbarred Melvin et al. (2014) and possessing two visible spiral arms over a flat distribution of pitch angles and bulge prominence. Clumpy galaxies are identified across the full range of clump multiplicities, with the exception of 1-clump galaxies (which would be difficult to differentiate from compact




Figure 12. Demographics of the morphologies for all galaxies in GZH (including both *HST* and SDSS imaging). Each node in each diagram (dark blue horizontal bars of uniform height) represents a task in the tree. The left diagram shows the full decision tree. The right diagram zooms in on the features/clumpy tasks, which are otherwise difficult to see. The paths between tasks represent each possible answer to the task, flowing from top to bottom between their origin and the subsequent task in the tree. Labels are assigned to each galaxy based on the plurality answer for each task. A galaxy is assigned only one label at each node. Widths of the paths are proportional to the number of galaxies assigned to that path. The widths of the nodes are proportional to the number of galaxies for which the question was reliably answered.

spheroids). Roughly half of the galaxies have at least 1 clump identified as the brightest. Clumps are most commonly asymmetrically arranged in clusters and are not often seen embedded in larger structures.

7.2. Comparing GZH morphologies to other catalogs

All of the Legacy surveys included in the GZH imaging have had morphological catalogs previously published; these catalogs have significant differences in the number of galaxies, size and magnitude limits, classification scheme, and the methods used for measuring morphology. These catalogs have been cross-matched to GZH to compare the results. This is not presented as an endorsement of any particular method, but as an exploration of the strengths and weaknesses of the Galaxy Zoo crowdsourced catalogs compared to products made with machine learning, automatic fits, and expert visual classification.

The types and accuracy of morphological classification

strongly depend on the sample and methods being used. In an attempt to make a consistent comparison between different techniques, galaxies are broadly grouped into three categories: bulge-dominated/elliptical/smooth, disk-dominated/spiral, and irregular/clumpy. These categories are compared to two parameters in GZH: $f_{\text{features,best}}$, which identifies disk-dominated and clumpy galaxies, and f_{odd} , which identifies deviations from well-formed spirals or S0s and which constitutes a “catch-all” for the variety of asymmetric morphologies that can constitute an irregular galaxy.

The comparison and analysis of the GZH morphologies is done a set of matched, volume-limited samples at $z < 1.0$ and $m_{I|i|z} < 22.5$ mag. The redshift and magnitude limits for the sample are chosen to match the shared constraints for the shallowest depths (GEMS; [Bundy et al. 2005](#)) and the limits on morphological reliability (COSMOS/ZEST; [Scarlata et al. 2007](#)) in the comparison catalogs. We match all catalogs against GZH using

a positional radius of $0.5''$.

Morphologies for GEMS galaxies were measured by Häufler et al. (2007), who used single-component Sérsic fits to the F850LP imaging. A $1''$ positional match for the GEMS galaxies in Table 9 in Häufler et al. (2007) to the GZH sample gives 8,846 galaxies in both Häufler et al. (2007) and GZH. The primary morphological parameter in the automated catalog is the Sérsic index n defining the radial surface brightness profile. “Elliptical” galaxies are selected by $n > 2.5$ and “spirals” by $n < 2.5$. There is no automatic measurement of irregular or clumpy structure.

AEGIS galaxies were morphologically classified using non-parametric measurements by Lotz et al. (2008). Their method used a combination of the Gini coefficient (G), which measures the relative inequality in pixel brightness, and M_{20} , the second-order moment of the brightest 20% of the light (Lotz et al. 2004). A linear combination of G and M_{20} delineates three broad categories of galaxy morphology: E/S0/Sa (“elliptical”), Sb/Sc/Ir (“spiral”), and mergers (“irregular”). We limit the AEGIS galaxies to those with reliably-measured morphologies and $S/N > 3$ in both V - and I -bands in Lotz et al. (2008).

Galaxies in both of the GOODS fields down to a limit of $z_{AB} = 22.5$ were visually classified by a single expert (R.S. Ellis), inspecting both z -band and composite Viz color images (Bundy et al. 2005). These morphologies are assigned a numerical value based on categories in Brinchmann et al. (1998): the corresponding morphologies used are “elliptical” (classes 0,1,2), “spirals” (classes 3,4,5), and “irregular” (classes 6,7,8).

COSMOS galaxies have multiple published datasets automatically classifying morphology, all using a variation of non-parametric measurements. Cassata et al. (2007) used a combination of concentration (C), asymmetry (A), G , and M_{20} (Cassata et al. 2005) make empirical divisions based on several hundred training images to assign galaxy morphology. A similar method was employed by Tasca (2011), using the same non-parametric indices but with a different method of calculating the Petrosian radius and total light profile, and using a nearest-neighbors method to label morphologies. Scarlata et al. (2007, ZEST) used C , A , G , M_{20} , the galaxy ellipticity (ϵ), and Sérsic index (n) to quantify morphology; a principal component analysis is used to assign galaxies to categories. All three methods use discrete morphological categories of ellipticals, spirals, and peculiar/irregulars. We use the same categories with the exception of assigning S0/bulge-dominated disks (type 2.0) from ZEST into the elliptical category.

Figure 13 shows the proportion of galaxies as split by their automated/expert visual morphologies for each of the six catalogs matched to GZH. Galaxies from every

catalog show a strong correlation between $f_{\text{features,best}}$ and the fraction of galaxies identified as spirals, with a corresponding anti-correlation between $f_{\text{features,best}}$ and the fraction of ellipticals.

The AEGIS sample had the fewest number of matched galaxies (485) with GZH; while the statistical uncertainties are as a result higher, the strongest difference between spirals and early-types is seen at $f_{\text{features,best}} > 0.8$. GOODS galaxies, which are the only comparison sample that also uses visual morphology, shows a very similar trend, with nearly 100% pure samples at the lowest and highest values of $f_{\text{features,best}}$. Galaxies from GEMS (which are only split into two types) also have a strong correlation with $f_{\text{features,best}}$, which is somewhat surprising (but encouraging) considering the known limits of assigning galaxy morphology based only on a single Sérsic parameter.

Visual inspection of images of the exceptions — where both $f_{\text{features,best}}$ and n are high — show that most are obvious spirals but with prominent bulges. This means that the single-component Sérsic fit is likely choosing too small of an effective radius and missing the extended disk structure for a large population of galaxies. This is a known issue with using n as a standalone measure of galaxy morphology – it is relatively common, for example, to measure high n for an object with an exponential disk but a central cusp in the light profile (eg, due to an AGN, cluster, foreground star, noise, etc).

Galaxies from the three COSMOS surveys all have morphologies that are reasonably well-predicted by the GZH based on $f_{\text{features,best}}$. The relation is strongest for early-type galaxies, since the comparable fractions of irregulars and spirals shows that clumpy/asymmetric morphologies are clearly also being captured by the $f_{\text{features,best}}$ parameter. The small dropoff in the fraction of early-types for the Cassata and ZEST samples at $f_{\text{features,best}} < 0.1$ is somewhat puzzling, but could potentially be due to the presence of a small population of extremely compact objects (stars or quasars) that would be classified as artifacts in GZH, explaining the dip in $f_{\text{features,best}}$.

We also emphasize that the various COSMOS catalogs are not consistent; for example, only 64% of the galaxies in Tasca (2011) have the same morphological class as Cassata et al. (2007). The bulk of these are galaxies classified as “spiral” using one technique and “irregular” using another (Cassata et al. 2007 is more likely to classify galaxies as irregular). A detailed comparison of the external catalogs is outside the scope of this paper, but we emphasize only that it is *impossible* for any catalog to be consistent with all three. We interpret the broad agreements between early-types, spirals, and irregulars for all three samples as a validation of the crowdsourced




Figure 13. Distributions of morphological parameters for a volume-limited sample ($z < 1.0, m < 22.5$) of galaxies matched between GZH and six published morphological catalogs, split by survey (AEGIS, COSMOS, GEMS, and GOODS). This plot shows the fraction of overall galaxies for each of the external morphological categories as a function of $f_{\text{features}, \text{best}}$ in GZH. The shaded regions around the binned fractions are confidence intervals calculated for a binomial population (Cameron 2011).

visual classifications provided by GZH.

Figure 14 analyzes the same sets of matched galaxies, but compares the data the f_{odd} parameter in GZH to examine its correlation with irregular morphologies. The relationship in this case is less clear; for the COSMOS catalogs, in particular, early-types, spirals, and irregulars are roughly equally probable at $f_{\text{odd}} > 0.5$. In both AEGIS and GOODS (there is no measurement of irregular structure for GEMS in Häubler et al. 2007), the proportion of irregular galaxies does increase with f_{odd} . However, the actual number of galaxies at $f_{\text{odd}} > 0.5$ is quite small ($N \lesssim 100$) in both samples, and so estimating an accurate fraction is less precise. We also note that this particular question (“*anything odd?*”) has somewhat uncertain phrasing in GZH, and that the design of the tree explicitly separates clumpy from irregular or merging morphologies, which would have similar properties in automatic techniques. A more accurate selection

of irregular morphologies in GZH should use a weighted combination of $f_{\text{features}, \text{best}}$, f_{clumpy} , and f_{odd} .

We repeated all of the analysis comparing morphologies between GZH and external catalogs on the entire catalog, rather than the volume-limited sample ($z < 1.0, m < 22.5$) discussed above. If all redshifts and the original magnitude limit of $m < 23.5$ are used, the correlations between all of the automatic morphological populations and GZH decrease significantly. The only exception are the GEMS galaxies, which are the only sample visually classified (similar to GZH) and had a shallower magnitude cut on their catalog. Based on these comparisons, we suggest that use of the GZH catalog be limited to galaxies with $m < 22.5$; fainter targets at the sensitivity of the *HST* Legacy surveys have been demonstrated to be less reliable, regardless of the method of morphological classification being used.




Figure 14. Same as Figure 13, but comparing external morphologies to f_{odd} in GZH. Measurement of f_{odd} excludes galaxies labeled as “odd” due to lenses or dust lanes (< 1% of the total), which are GZH categories not expected to correlate with larger-scale structural parameters.

7.3. Galaxy Zoo: Hubble and Galaxy Zoo: CANDELS

Finally, there are 7,681 galaxies in the GOODS-S field with morphological classifications in both GZH and the Galaxy Zoo: CANDELS project (Simmons et al., submitted). Since both the sensitivity and filters for the two sets of images are significantly different (and there is no correction for redshift bias applied to GZC), there is no prior reason to expect a perfect correlation between the separate vote fractions for the projects. Briefly, we note that the f_{features} value for GZH is on average higher than GZC; the effect is strongest at $f_{\text{features,GZC}} < 0.3$, for which roughly half the galaxies have $f_{\text{features,GZH}} > 0.5$. However, the correlation between vote fractions is single-valued (although not linear, with a Pearson correlation coefficient of $r = 0.6$), and should be possible to calibrate using a similar approach to that described in Section 4; the correlation between other tasks, such as edge-on galaxies is significantly stronger ($r = 0.9$). So while the raw vote fractions are not directly comparable, the

initial analysis indicates that the broad morphologies are at least consistent.

8. SUMMARY

This paper presents the catalog release for the Galaxy Zoo Hubble project, which uses crowdsourced visual classifications to measure galaxy morphologies. The first two phases of Galaxy Zoo (Lintott et al. 2011; Willett et al. 2013) used images of low-redshift galaxies from SDSS; this is the first result of the project with space-based images of high-redshift targets (in addition to the Galaxy Zoo: CANDELS collaboration; Simmons et al., submitted). The final sample includes classifications for 189,149 images generated from 145,741 unique galaxies (of which 115,402 are in *HST* Legacy imaging).

Galaxies were selected from a brightness-limited sample from multiple Legacy surveys using the Advanced Camera for Surveys on the *Hubble Space Telescope*, including AEGIS, GEMS, GOODS-N, GOODS-S, and

COSMOS. The catalog also includes classifications for SDSS images from Stripe 82 at $z < 0.25$; these images serve both as a low-redshift anchor for cosmological studies and a potential comparison for the different epochs of classification between GZH and Galaxy Zoo 2 (Willett et al. 2013).

The data for the GZH catalogs has been extensively tested and reduced. The dominant effect is a known bias against identifying disk and asymmetric sub-structures at either low resolution or surface brightness. This can be the result either of genuinely small or dim galaxies, or a perceived effect from observing galaxies at further distances (higher redshift). To calibrate this *without* potentially overcorrecting for the genuine morphological evolution of galaxies over cosmic time, the GZH project uses SDSS images of low-redshift galaxies, processes them to appear as if they were at higher redshift, and classifies them through the GZH interface in an identical fashion. The resulting change in f_{features} as a function of z and μ is applied as a multiplicative correction to the top-level vote fractions for $\sim 50\%$ of the GZH galaxies.

Galaxies in GZH show significant changes in the disk/elliptical fraction as a function of redshift, along with an increasing number of galaxies dominated by smaller clumps and presumed to be in the process of building up their baryonic mass through a combination of hierarchical merging and in-situ star formation. While the majority of scientific interpretation is left to future work, this paper confirms the decrease in observed bar fraction with increasing redshift and identifies a new way of selecting clumpy galaxies as a function of clump multiplicity.

The full data tables for the catalogs can be accessed in machine-readable form from both the journal and at <http://data.galaxyzoo.org>. All the code and data tables used to generate this manuscript can be found at <https://github.com/willettk/gzhubble>.

KW, MG, CS, MB, and LF gratefully acknowledge support from the US National Science Foundation Grant AST1413610. KS gratefully acknowledges support from Swiss National Science Foundation Grant PP00P2_138979/1. The development and hosting of Galaxy Zoo Hubble was supported by a grant from the Alfred P. Sloan Foundation. The Zooniverse acknowledges support from a Google Global Impact Award.

We thank Meg Schwamb and the ASIAA for hosting

the “Citizen Science in Astronomy” workshop, 3-7 Mar 2014 in Taipei, Taiwan, at which some of this analysis was done. We thank Jennifer Lotz for sharing her G - M_{20} measurements for the AEGIS sample. We thank Coleman Krawczyk for his assistance in producing Figure 4.

This project made heavy use of the Astropy packages in Python (Astropy Collaboration et al. 2013), the seaborn plotting package (Waskom et al. 2015), astroML (Vanderplas et al. 2012), and TOPCAT (Taylor 2005, 2011). Modified code from Nick Wherry and David Schlegel was used to create the JPG images. Figure 12 was generated with <http://sankeymatic.com/>. Holwerda (2005) provided invaluable assistance in interpreting SExtractor output.

This work is based on (GO-10134, GO-09822, GO-09425.01, GO-09583.01, GO-9500) program observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS website is <http://www.sdss.org>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory and the University of Washington.

REFERENCES

- Abazajian K. N. et al., 2009, ApJS, 182, 543
- Abraham R. G., van den Bergh S., Nair P., 2003, ApJ, 588, 218
- Astropy Collaboration et al., 2013, A&A, 558, A33
- Baillard A. et al., 2011, A&A, 532, A74
- Bamford S. P. et al., 2009, MNRAS, 393, 1324
- Bamford S. P., Rojas A. L., Nichol R. C., Miller C. J., Wasserman L., Genovese C. R., Freeman P. E., 2008, MNRAS, 391, 607

- Barden M., Häußler B., Peng C. Y., McIntosh D. H., Guo Y., 2012, MNRAS, 422, 449
- Barden M., Jahnke K., Häußler B., 2008, ApJS, 175, 105
- Beckwith S. V. W. et al., 2006, AJ, 132, 1729
- Behrendt M., Burkert A., Schartmann M., 2016, ApJL, 819, L2
- Bell E. F. et al., 2012, ApJ, 753, 167
- Bertin E., Arnouts S., 1996, A&AS, 117, 393
- Brinchmann J. et al., 1998, ApJ, 499, 112
- Bundy K., Ellis R. S., Conselice C. J., 2005, ApJ, 625, 621
- Buta R. J., 2013, Galaxy Morphology, Oswalt T. D., Keel W. C., eds., Springer, pp. 1–89
- Caldwell J. A. R. et al., 2008, ApJS, 174, 136
- Cameron E., 2011, PASA, 28, 128
- Cameron E., Carollo C. M., Oesch P. A., Bouwens R. J., Illingworth G. D., Trenti M., Labb   I., Magee D., 2011, ApJ, 743, 146
- Cardamone C. N. et al., 2010, ApJS, 189, 270
- Cassata P. et al., 2005, MNRAS, 357, 903
- Cassata P. et al., 2007, ApJS, 172, 270
- Cheung E. et al., 2015, MNRAS, 447, 510
- Chevance M., Weijmans A.-M., Damjanov I., Abraham R. G., Simard L., van den Bergh S., Caris E., Glazebrook K., 2012, ApJL, 754, L24
- Conselice C. J., 2003, ApJS, 147, 1
- Conselice C. J., 2014, ARA&A, 52, 291
- Darg D. W. et al., 2010, MNRAS, 401, 1552
- Davis M. et al., 2007, ApJL, 660, L1
- de Vaucouleurs G., 1959, Handbuch der Physik, 53, 275
- Dressler A., 1980, ApJ, 236, 351
- Elmegreen D. M., Elmegreen B. G., 2014, ApJ, 781, 11
- Elmegreen D. M., Elmegreen B. G., Ferguson T., Mullan B., 2007, ApJ, 663, 734
- Elmegreen D. M., Elmegreen B. G., Marcus M. T., Shahinyan K., Yau A., Petersen M., 2009, ApJ, 701, 306
- Elmegreen D. M., Elmegreen B. G., Rubin D. S., Schaffer M. A., 2005, ApJ, 631, 85
- F  rster Schreiber N. M., Shapley A. E., Erb D. K., Genzel R., Steidel C. C., Bouch   N., Cresci G., Davies R., 2011, ApJ, 731, 65
- Fortson L. et al., 2012, Galaxy Zoo: Morphological Classification and Citizen Science, Way M. J., Scargle J. D., Ali K. M., Srivastava A. N., eds., CRC Press, Taylor & Francis Group, pp. 213–236
- Freeman P. E., Izbicki R., Lee A. B., Newman J. A., Conselice C. J., Koekemoer A. M., Lotz J. M., Mozena M., 2013, MNRAS, 434, 282
- Galloway M. A. et al., 2015, MNRAS, 448, 3442
- Genel S. et al., 2014, MNRAS, 445, 175
- Giavalisco M., 2012, VizieR Online Data Catalog, 2318
- Giavalisco M. et al., 2004, ApJL, 600, L93
- Griffith R. L. et al., 2012, ApJS, 200, 9
- Grogan N. A. et al., 2011, ApJS, 197, 35
- Guo Y. et al., 2015, ApJ, 800, 39
- H  u  ller B. et al., 2007, ApJS, 172, 615
- Hinshaw G. et al., 2013, ApJS, 208, 19
- Holwerda B. W., 2005, ArXiv Astrophysics e-prints
- Hopkins P. F. et al., 2010, ApJ, 715, 202
- Hubble E. P., 1926, ApJ, 64, 321
- Hubble E. P., 1936, Realm of the Nebulae. Yale University Press
- Ilbert O. et al., 2013, A&A, 556, A55
- Kartaltepe J. S. et al., 2015, ApJS, 221, 11
- Koekemoer A., Fruchter A., Hack W., 2003, Space Telescope European Coordinating Facility Newsletter, Volume 33, p.10, 33, 10
- Koekemoer A. M., Fruchter A. S., Hook R. N., Hack W., 2002, in The 2002 HST Calibration Workshop : Hubble after the Installation of the ACS and the NICMOS Cooling System, Proceedings of a Workshop held at the Space Telescope Science Institute, Baltimore, Maryland, October 17 and 18, 2002. Edited by Santiago Arribas, Anton Koekemoer, and Brad Whitmore. Baltimore, MD: Space Telescope Science Institute, 2002., p.339, pp. 339–+
- Krist J., 1993, in Astronomical Society of the Pacific Conference Series, Vol. 52, Astronomical Data Analysis Software and Systems II, R. J. Hanisch, R. J. V. Brissenden, & J. Barnes, ed., pp. 536–+
- Lackner C. N., Gunn J. E., 2012, MNRAS, 421, 2277
- Lahav O. et al., 1995, Science, 267, 859
- Land K. et al., 2008, MNRAS, 388, 1686
- Law D. R., Shapley A. E., Steidel C. C., Reddy N. A., Christensen C. R., Erb D. K., 2012a, Nature, 487, 338
- Law D. R., Steidel C. C., Shapley A. E., Nagy S. R., Reddy N. A., Erb D. K., 2012b, ApJ, 745, 85
- Lilly S. et al., 1998, ApJ, 500, 75
- Lintott C. et al., 2011, MNRAS, 410, 166
- Lintott C. J. et al., 2008, MNRAS, 389, 1179
- Lotz J. M. et al., 2008, ApJ, 672, 177
- Lotz J. M., Primack J., Madau P., 2004, AJ, 128, 163
- Loveday J. et al., 2012, MNRAS, 420, 1239
- Lupton R., Blanton M. R., Fekete G., Hogg D. W., O'Mullane W., Szalay A., Wherry N., 2004, PASP, 116, 133
- Mandelker N., Dekel A., Ceverino D., DeGraf C., Guo Y., Primack J., 2015, ArXiv e-prints, 1512.08791
- Mao S., Mo H. J., White S. D. M., 1998, MNRAS, 297, L71
- Masters K. L. et al., 2011, MNRAS, 411, 2026
- Melvin T., 2016, PhD thesis, University of Portsmouth
- Melvin T. et al., 2014, MNRAS
- Momcheva I. G. et al., 2015, ArXiv e-prints, 1510.02106
- Mortlock A. et al., 2013, MNRAS, 433, 1185
- Nair P. B., Abraham R. G., 2010, ApJS, 186, 427
- Nieto-Santisteban M. A., Szalay A. S., Gray J., 2004, in Astronomical Society of the Pacific Conference Series, Vol. 314, Astronomical Data Analysis Software and Systems (ADASS) XIII, Ochsenbein F., Allen M. G., Egret D., eds., p. 666
- Peng C. Y., Ho L. C., Impey C. D., Rix H.-W., 2002, AJ, 124, 266
- Pierce C. M. et al., 2010, MNRAS, 405, 718
- Rix H.-W. et al., 2004, ApJS, 152, 163
- S  nchez S. F. et al., 2004, ApJ, 614, 586
- Sandage A., 1961, The Hubble atlas of galaxies. Carnegie Institute of Washington
- Scarlata C. et al., 2007, ApJS, 172, 406
- Schawinski K. et al., 2014, MNRAS, 440, 889
- Schaye J. et al., 2015, MNRAS, 446, 521
- Scoville N. et al., 2007, ApJS, 172, 1
- Silk J., Mamon G. A., 2012, Research in Astronomy and Astrophysics, 12, 917
- Simard L., Mendel J. T., Patton D. R., Ellison S. L., McConnachie A. W., 2011, ApJS, 196, 11
- Simmons B. D. et al., 2013, MNRAS, 429, 2199
- Simmons B. D. et al., 2014, MNRAS, 445, 3466
- Simmons B. D., Urry C. M., 2008, ApJ, 683, 644
- Simmons B. D., Van Duyne J., Urry C. M., Treister E., Koekemoer A. M., Grogan N. A., The GOODS Team, 2011, ApJ, 734, 121
- Skibba R. A. et al., 2009, MNRAS, 399, 966
- Skibba R. A. et al., 2012, MNRAS, 423, 1485
- Smethurst R. J. et al., 2015, MNRAS, 450, 435
- Steinmetz M., Navarro J. F., 2002, NewA, 7, 155
- Strauss M. A. et al., 2002, AJ, 124, 1810
- Taniguchi Y. et al., 2007, ApJS, 172, 9

Table A1. Correctable fractions for the top-level task in GZH in the GOODS shallow-depth (2-epoch) images.

	GOODS-N	GOODS-S	Total
correctable	1,051	730	1,781
lower-limit	131	334	465
no correction needed ($z \leq 0.3$)	267	267	534
NEI	943	2,078	3,021
no redshift information	159	184	343
total	2,551	3,593	6,144

Tasca L. A. M., 2011, VizieR Online Data Catalog, 7265
 Taylor M., 2011, TOPCAT: Tool for OPerations on Catalogues And Tables. Astrophysics Source Code Library
 Taylor M. B., 2005, in Astronomical Society of the Pacific Conference Series, Vol. 347, Astronomical Data Analysis Software and Systems XIV, Shopbell P., Britton M., Ebert R., eds., p. 29
 Toomre A., Toomre J., 1972, ApJ, 178, 623
 van den Bergh S., 1976, ApJ, 206, 883

Vanderplas J., Connolly A., Ivezić Ž., Gray A., 2012, in Conference on Intelligent Data Understanding (CIDU), pp. 47–54
 Vogelsberger M. et al., 2014, MNRAS, 444, 1518
 Waskom M. et al., 2015, seaborn: v0.6.0 (june 2015)
 Willett K. W. et al., 2013, MNRAS, 435, 2835
 Willett K. W. et al., 2015, MNRAS, 449, 820
 Williams R. E. et al., 1996, AJ, 112, 1335
 Wright E. L., 2006, PASP, 118, 1711
 York D. G. et al., 2000, AJ, 120, 1579

APPENDIX

A. GOODS SHALLOW DEPTH DATA

GZH used both 5-epoch and 2-epoch sets of data to construct the GOODS set of images. The 11,157 full depth 5-epoch images are used in the `main` catalog; the classifications for the 6,144 `goods-shallow` images are provided as a supplementary table. This section analyzes the effect of image depth on the ability of the Galaxy Zoo classifiers to identify features or disk structure in the images.

A.1. Comparing shallow and full depth morphologies

Of the 11,157 galaxies in the GOODS-N and GOODS-S full depth sample, 4,461 are also in the shallow-depth sample. Figure A1 shows a strong correlation between f_{features} for both sets of images. The mean change in f_{features} from the shallow to full depth images $f_{\text{features,full}} - f_{\text{features,shallow}} \equiv \Delta f = 0.00$, with a standard deviation of $\sigma = 0.17$. While there is some variance in Δf in the whole sample, the change is usually small and not often significant enough to change a morphological classification. Defining a clean sample of disk galaxies as those with $f_{\text{features,best}} > 0.8$, elliptical galaxies as those with $f_{\text{smooth,best}} < 0.2$, and intermediate as those in between, 75% of the sample would not change morphology. Of the remaining 25% that would change morphology, only 0.3% (10 galaxies) drastically change morphology either from smooth to featured or vice versa, while the rest transition to or from the “intermediate” morphology. Details can be seen in Table A2 and examples of images representing the 9 possible changes (or lack of) in morphology are shown in Figures A2, A3, and A4.

A.2. Debiasing higher-order tasks: f_{bar}

To determine the rate at which the vote fraction for any Task decreases with redshift or surface brightness, data from simulated FERENGI images were modeled with functions in discrete redshift and surface brightness bins (Section 4.2). A cut of $N > 10$ was placed on the number of votes to reduce the error in the vote fractions and ensure they were well-sampled before applying a fit to the data. This requirement was only met for Task 01, which significantly reduced the goodness-of-fit for the model functions as compared with to the smooth/features task. Here we show an example of results obtained for the bar Task. Figure A5 shows that both the low sample rate and the degree of scatter in each bin inhibit fitting a reliable function that predicts $f_{\text{bar},z=0.3}$. Table A6 summarizes the correction types for the FERENGI data; 71% of the galaxies simulated did not meet the GZH requirements for well-sampled vote fractions for f_{bar} , as compared to 2% for f_{features} . Similar results were obtained for all higher-order Tasks in GZH (Section 4.3).





Figure A1. Distribution of f_{features} for the 4,461 GOODS galaxies with both shallow (2-epoch; `goods-shallow`) and full-depth (5-epoch; `main`) images morphologically classified in GZH. For most galaxies, the value of f_{features} is consistent ($\Delta f_{\text{features}} < 0.2$) between depths. Examples of galaxies with sharp changes in f_{features} , as well as those with little to no change are shown in Figures A2–A4.

Table A2. Properties of galaxies whose morphologies changed or stayed the same in the shallow vs full images. Featured here is defined as $f_{\text{features,best}} > 0.8$, intermediate $= 0.2 < f_{\text{features,best}} < 0.8$, smooth $= f_{\text{smooth,best}} < 0.2$.


	N	%	$< \Delta f >$	$< z >$
smooth to smooth	758	17.0	-0.00	0.69
smooth to intermediate	367	8.2	0.18	0.69
smooth to featured	7	0.2	0.76	0.57
intermediate to smooth	214	4.8	-0.18	0.65
intermediate to intermediate	2,303	51.6	0.01	0.78
intermediate to featured	168	3.8	0.19	0.83
featured to smooth	3	0.1	-0.74	0.71
featured to intermediate	337	7.6	-0.18	0.68
featured to featured	301	6.8	-0.05	0.71
Total	4,461	100		

Table A3. Distribution of FERENGI images analyzed in Figure A5. Correctable images had a single-valued relationship between their measured f_{bar} values at high and low redshifts (white regions in Figure A5). Galaxies with a lower-limit on f_{bar} had a non single-valued relationship (blue regions). NEI images had undetermined relationships due to a lack of data ($N < 5$) in their corresponding z - μ bins (gray regions). Only 17% (maximum) of FERENGI galaxies in the sample were considered “correctable”, which is not sufficient to compute a ζ function applicable to the *HST* data.


	N	%
Correctable	664	17%
Lower-limit	483	12%
NEI	2,803	71%
Total	3,950	100%



[a]




[b]




[c]


Figure A2. Galaxies whose shallow images were classified as smooth and full depth images were classified as smooth, intermediate, or featured.



[b]




[b]




[b]


Figure A3. Galaxies whose shallow images were classified as intermediate and full depth images were classified as smooth, intermediate, or featured.



[b]



[b]



[b]

Figure A4. Galaxies whose shallow images were classified as featured and full depth images were classified as smooth, intermediate, or featured.




Figure A5. Similar to Figure 9, showing effects of redshift bias for 3,449 images in the FERENGI sample. Each point in a given redshift and surface brightness bin represents a unique galaxy. The y -axis in each bin is the f_{features} value of the image of that galaxy redshifted to the value corresponding to that redshift bin. The x -axis is the f_{features} value of the image of the same galaxy redshifted to $z = 0.3$. The dashed black lines represent the best-fit polynomials to the data in each square. The solid black line represents $f_{\text{features},z=f_{\text{features},z=0.3}}$. Regions in which there is a single-valued relationship between f_{features} at high redshift and $z = 0.3$ are marked in white; those in which there is not are blue, and those with not enough data ($N < 5$) are gray. A larger version of the bin outlined at $z = 1.0$ and $20.3 < \mu < 21.0$ ($\text{mag}/\text{arcsec}^2$) is shown in Figure A6. The mean normalized χ^2 of the change in vote fraction is 0.08. None of the correlations for higher-order tasks (including bars) were applied to galaxies in the GZH catalog.




Figure A6. A larger version of the dark-outlined square in Figure A5, containing FERENGI galaxies that have been artificially redshifted to $z = 1.0$ and have surface brightnesses between $20.3 < \mu < 21.0$ (mag/arcsec 2). The orange bars represent the inner 80th percentile of the data, the boundaries of which are the lower and upper limits of the debiased values.