

## Review Article

# Operating room management: why, how and by whom?

R. MARJAMAA<sup>1</sup>, A. VAKKURI<sup>1</sup> and O. KIRVELÄ<sup>2</sup>

<sup>1</sup>Department of Anesthesiology and Intensive Care Medicine, Peijas Hospital and <sup>2</sup>Department of Anesthesiology and Intensive Care Medicine, Töölö Hospital, Helsinki University Hospital, Helsinki, Finland

Operating room (OR) is a cost-intensive environment, and it should be managed efficiently. When improving efficiency, shortening case duration by parallel processing, training of the resident surgeons, the choice of anesthetic methods, effective scheduling, and monitoring of the overall OR performance are important. When redesigning the OR processes, changes should be given a clear target and the achieved results monitored and reported to everyone involved. Advanced, reliable, and easy to use information technology solutions for OR management are under development. Pre-operative clinic and functionally designed

facilities support efficiency. OR personnel must be kept motivated by clear management and leadership, supported by superiors.

Accepted for publication 14 December 2007

**Key words:** Operating room; management; parallel processing; resources; personnel.

© 2008 The Authors

Journal compilation © 2008 The Acta Anaesthesiologica Scandinavica Foundation

## Why?

OPERATING rooms (ORs) incur high costs and considerable amount of hospitals revenues. Demand for OR and procedural facilities appears to be increasing due to aging population and developments in surgery. The conventional solution has been to build new facilities. Building them and, of course, staffing them adequately is increasingly expensive. By contrast, committing to increase the productivity of existing facilities seems to be a wiser strategy. One of the key methods here is proper OR management and optimizing the whole process or chain of processes involved in the treatment of a patient.

### Goals for OR management

Clear goals for OR management are essential: improving productivity and efficiency while maintaining high quality of care at all times. Improving efficiency means shorter case durations, rational scheduling of various types of surgery, and minimizing nonoperative time (NOT) (Table 1) by reorganizing OR tasks. This requires motivated personnel and teamwork in every step of the patient care process. If all professionals working in the OR remain interested in developing their

own work, we reach our goal: working smarter, not faster (Tables 2 and 3).

## How?

### Shortening case times

Novel surgical techniques, although enhancing recovery, may prolong operations (2). Case durations (time spent in the OR) are highly variable even between similar procedures (3, 4). Differences

Table 1

Operating room (OR) management glossary [partly adopted from Abouleish et al. (1)].

Raw utilization	The percentage of time a patient is in the OR of the total allocated time for a given service (case durations only)
Adjusted utilization	The percentage of time a patient is in the OR of the total allocated time for a given service plus the time for set up and clean up (= case duration plus turnover time)
Case duration	The time the patient is in the OR
Turnover time	The time from the previous patient out-room time to the succeeding patient in-room time to the same OR
Nonoperative time (NOT)	The time between the end of surgery of the previous patient to the beginning of surgery of the succeeding patient
Tardiness	The delay from the scheduled start time (patient enters OR) to the actual start time of the patient

Table 2

Key factors for operating room efficiency.

Streamlining processes with other units  
 Patient flow and its coordination  
 Timely patient preparation  
 Efficient patient reception  
 Parallel processing by use of induction area/PACU  
 Recovery room/ICU/ward capacity  
 Personnel: number/professional skills/motivation  
 Flexible facilities  
 Patient focused processes  
 Continuous process improvement

PACU, post anesthesia care unit; ICU, intensive care unit.

Table 3

Common problems with operating room management.

Poorly designed processes  
 Change reluctance/friction  
 Lack of motivation (no financial incentives?)  
 Dodging responsibility/placing blame  
 Lack of discipline (physicians!)

arise mainly from the surgeon's work rate (4, 5). The presence or participation of a resident physician prolongs the duration of the surgery up to 70% increasing costs accordingly (6–8). Adequate resident training, possibly with the aid of a simulator (9) and experienced assistance should be provided to the residents starting to operate more independently (10). Even small reductions in operative time can increase OR throughput (11).

Individual differences between anesthesiologists have very little effect on the case length (4), although specializing in certain types of surgery may speed up the anesthesiologists work (12). Teaching a resident seems to delay the anesthesiologist only by 2–3 min (13). Covering more than one room statistically causes a delay of 6 min (14).

Selecting the right type of anesthesia is crucial as it is the second important factor affecting the time spent in OR – right after the surgeon's speed. Anesthesia induction, taking roughly 10% of the time spent in the OR, is most affected by the type of anesthesia, the type of the airway, the need for invasive monitoring, and the American Society of Anesthesiologists (ASA) physical status. ASA class IV doubles (in children) or triples (in adults) the anesthesia preparation time compared with ASA physical status class I (15, 16).

Administering neuraxial blocks may take longer than inducing general anesthesia, yet this appears not to cause inefficiency, because no time is needed for emergence from anesthesia (17). However, per-

forming a brachial plexus block tends to take too much OR time and should therefore be performed in a separate area well in advance of the anticipated start of the surgery (18). Moreover, local anesthesia should be considered whenever possible. Local anesthesia is cost efficient for example for hernia repair (19, 20), especially if administered beforehand in the induction room (21). As for general anesthesia, titrating short-acting anesthetic agents carefully while monitoring the depth of hypnosis (22), may help minimizing anesthesia-controlled time. Choosing a laryngeal mask for airway instead of an endotracheal tube may also save time (23).

The time needed for anesthesia preparation should always be considered, when scheduling a case. Cases requiring invasive hemodynamic monitoring and neuraxial block should not be placed first on the schedule. The first case should be short and predictable in length (24, 25). During the first listed case, anesthesia preparations such as neuraxial blocks can be done either in the induction room, recovery room, or a block room for the second case (17, 26). For inpatients, inserting epidural catheters, central venous catheters, etc. already the day before surgery may speed up the start of the operation the following day.

### *Changing the way of working*

Parallel processing improves OR throughput by shortening NOT. Using separate rooms for anesthesia induction – a common way of working in some countries – shortens NOT and may allow extra cases each day compared with the traditional sequenced induction in the OR (27–30). Emergence from anesthesia can also take place in a separate room (27, 30) while the OR is being cleaned up. However, long procedures with few turnovers per OR seem not to benefit from parallel processing (31).

Using induction rooms may require additional staff, equipment, and facilities and thus have multiple financial implications to consider. It seems, however, that the benefits of the increased throughput, as well as the profits gained, will outweigh the increased costs (28, 29, 32). There are also ways to employ parallel processing without extra costs: redesigning work flow patterns and reassigning tasks of various professionals (33). Obviously, anticipating and preparing for the following case during the ongoing procedure will also help (33, 34).

Considering the fact that the OR is often the most cost-intensive and the most productive unit in the hospital (35), it is important to reduce time running

at idle in the OR. Delays due to unavailability of various professionals, lack of staff, or the patient not being prepared in time can amount up to five working days per month (36). Surgeons have been found to cause delays more frequently than any other profession in the OR environment (37, 38). Moreover, the delays caused by surgeons are significantly longer than the delays caused by any other single reason (16). Substantial improvement can be achieved by increasing communication between the surgeon and the rest of the team, as well as by notifying those often being late (16, 36–38). Sometimes a minor change of the process may help: providing a space and equipment for dictating in the OR (27) or computerizing patient records (33, 34) will keep the surgeons in the vicinity and readily available for the next procedure.

#### *Monitoring OR performance*

The effects from organizational changes can be expected to take considerable amount of time. As improvements from changes may take a year to achieve, monitoring the outcome is warranted. To maintain the achievements, monitoring should be frequent, if not real time (39). Useful efficiency measures include output/input-type of measures (40), NOT (27, 28, 30, 34), OR start times (41), and raw utilization (28, 40). Number of operations, overtime, costs, cancellation rate, complications, surgery durations, and under- and over-utilized time may also be worth measuring (42). Measuring patient waiting time is a key to better customer satisfaction and patient safety (25, 40). Statistical process control may help in interpreting the results (43, 44). It is essential that changes are given a clear target and results are reported to everyone involved in the process.

#### *Technical tools for management*

The OR management information system should support process management in real time. Of those currently used, most are designed primarily for scheduling and do not have visually high-quality displays for ongoing procedures. An ideal system allows tracking of patients and resources, and monitoring and reporting of the OR performance (45). Most of the currently used systems require manual data input. Wireless patient tracking systems can automatically timestamp key events, thereby decreasing the need for manual data input (39).

#### *Focus on the processes*

Well-defined processes enhance mutual understanding of all parties involved in the perioperative care (40). When each person involved has a clear understanding of his responsibilities and duties, the process can run efficiently (33). Multidisciplinary teamwork, parallel processing, careful reorganization, and reallocation of tasks have been shown to diminish the delays and NOT, to speed up turnover times, and to get the first operation of the day to start earlier (37). The process of receiving patients in the morning and the discharge to ward must be carefully planned to avoid 'bottle necks.' Collaboration with other units, timing and sequencing of patient preparation as well as post-operative care, is therefore important.

#### *Preoperative clinic*

A well-functioning pre-operative clinic enhances OR efficiency by reducing the number of cancellations and delays (46). Careful consideration between the risks and expected long-term benefits for the patient is needed to make a rational decision whether surgery is justified in a high-risk case. Also, patients that have been on the waiting list for months may need a re-evaluation. This evaluation requires a considerable amount of experience from the surgeon and the anesthesiologist and should therefore be performed by a specialist. To be of any use, the evaluation has to take place well in advance of the planned surgery to allow necessary changes in medication to be made, additional tests and consultations to be arranged, as well as any corrective measures taken during the evaluation to have enough time to take effect.

#### *Facilities*

The hospital facilities should be designed to support fluent patient flows (40) and to allow their flexible use for a variety of functions (35, 47). Sometimes a trivial structural feature, such as limited elevator capacity, can be a serious rate-limiting factor. Separate and dedicated processes for emergency and urgent surgery may be a necessity in some hospitals (48). This, although ensuring a quick response in an emergency, may cause inefficiency when these resources have to be reserved at all times, even when not needed.

## Personnel

In many countries shortage of anesthesiologists or anesthesia nurses restricts the availability of ORs (49). Sometimes the work flow can be improved by changing the ratio of nurses and doctors: an extra nurse may enable efficient use of induction room with the same number of anesthesiologists. There is also some evidence that financial incentives improve performance (50).

OR work requires high level of professional skills and knowledge, requiring constant updating. Motivation to keep up one's skills is best achieved in an encouraging organizational climate with healthy interdisciplinary relationships (51).

## By whom?

### Management and leadership

Successful management of the OR requires sound organizational structure, good leadership, as well as interdisciplinary collaboration. Mostly for historical reasons, distribution of roles and responsibilities is not always clear in the OR environment. Yet, it is vital that the status and authority of the person in charge is known and acknowledged by everyone in the perioperative process as well as supported by his/her superiors. This way the conflicts can be solved quickly and locally, and resource allocation can be done efficiently (52). Leadership skills and emotional intelligence are equally important as credentials. The specialty or profession of the leader is of secondary importance.

## References

1. Abouleish AE, Hensley SL, Zornow MH et al. Inclusion of turnover time does not influence identification of surgical services that over- and underutilize allocated block time. *Anesth Analg* 2003; **96**: 813–8.
2. Purkayastha S, Tilney HS, Georgiou P et al. Laparoscopic cholecystectomy versus mini-laparotomy cholecystectomy: a meta-analysis of randomised control trials. *Surg Endosc* 2007; **21**: 1294–300.
3. Dexter F, Davis M, Egger Halbeis CB et al. Mean operating room times differ by 50% among hospitals in different countries for laparoscopic cholecystectomy and lung lobectomy. *J Anesth* 2006; **20**: 319–22.
4. Strum DP, Sampson AR, May JH et al. Surgeon and type of anesthesia predict variability in surgical procedure times. *Anesthesiology* 2000; **92**: 1454–66.
5. Abouleish AE, Prough DS, Zornow MH et al. The impact of longer-than-average anesthesia times on the billing of academic anesthesiology departments. *Anesth Analg* 2001; **93**: 1537–43.
6. Babineau TJ, Becker J, Gibbons G et al. The “cost” of operative training for surgical residents. *Arch Surg* 2004; **139**: 366–9.
7. Bridges M, Diamond DL. The financial impact of teaching surgical residents in the operating room. *Am J Surg* 1999; **177**: 28–32.
8. Traverso LW, Koo KP, Hargrave K et al. Standardizing laparoscopic procedure time and determining the effect of patient age/gender and presence or absence of surgical residents during operation. A prospective multicenter trial. *Surg Endosc* 1997; **11**: 226–9.
9. Korndorffer JR Jr, Dunne JB, Sierra R et al. Simulator training for laparoscopic suturing using performance goals translates to the operating room. *J Am Coll Surg* 2005; **201**: 23–9.
10. Koperna T. How long do we need teaching in the operating room? The true costs of achieving surgical routine. *Langenbecks Arch Surg* 2004; **389**: 204–8.
11. Seim AR, Dahl DM, Sandberg WS. Small changes in operative time can yield discrete increases in operating room throughput. *J Endourol* 2007; **21**: 703–8.
12. Kain ZN, Fasulo A, Rimar S. Establishment of a pediatric surgery center: increasing anesthetic efficiency. *J Clin Anesth* 1999; **11**: 540–4.
13. Eappen S, Flanagan H, Bhattacharyya N. Introduction of anesthesia resident trainees to the operating room does not lead to changes in anesthesia-controlled times for efficiency measures. *Anesthesiology* 2004; **101**: 1210–4.
14. Paoletti X, Marty J. Consequences of running more operating theatres than anaesthetists to staff them: a stochastic simulation study. *Br J Anaesth* 2007; **98**: 462–9.
15. Saadat H, Escobar A, Davis EA et al. Task analysis of the preincision period in a pediatric operating suite: an independent observer-based study of 656 cases. *Anesth Analg* 2006; **103**: 928–31.
16. Escobar A, Davis EA, Ehrenwerth J et al. Task analysis of the preincision surgical period: an independent observer-based study of 1558 cases. *Anesth Analg* 2006; **103**: 922–7.
17. Williams BA, Kentor ML, Williams JP et al. Process analysis in outpatient knee surgery: effects of regional and general anesthesia on anesthesia-controlled time. *Anesthesiology* 2000; **93**: 529–38.
18. Chan VW, Peng PW, Kaszas Z et al. A comparative study of general anesthesia, intravenous regional anesthesia, and axillary block for outpatient hand surgery: clinical outcome and cost analysis. *Anesth Analg* 2001; **93**: 1181–4.
19. Nordin P, Zetterstrom H, Gunnarsson U et al. Local, regional, or general anaesthesia in groin hernia repair: multicentre randomised trial. *Lancet* 2003; **362**: 853–8.
20. Song D, Greilich NB, White PF et al. Recovery profiles and costs of anesthesia for outpatient unilateral inguinal herniorrhaphy. *Anesth Analg* 2000; **91**: 876–81.
21. Friedman DM, Sokal SM, Chang Y et al. Increasing operating room efficiency through parallel processing. *Ann Surg* 2006; **243**: 10–4.
22. Vakkuri A, Yli-Hankala A, Sandin R et al. Spectral entropy monitoring is associated with reduced propofol use and faster emergence in propofol-nitrous oxide-alfentanil anesthesia. *Anesthesiology* 2005; **103**: 274–9.
23. Hartmann B, Banzhaf A, Junger A et al. Laryngeal mask airway versus endotracheal tube for outpatient surgery: analysis of anesthesia-controlled time. *J Clin Anesth* 2004; **16**: 195–9.
24. Lebowitz P. Schedule the short procedure first to improve OR efficiency. *Aorn J* 2003; **78**: 651–4, 7–9.

25. Dexter F, Epstein RH, Traub RD et al. Making management decisions on the day of surgery based on operating room efficiency and patient waiting times. *Anesthesiology* 2004; **101**: 1444–53.
26. Armstrong KP, Cherry RA. Brachial plexus anesthesia compared to general anesthesia when a block room is available. *Can J Anaesth* 2004; **51**: 41–4.
27. Sandberg WS, Daily B, Egan M et al. Deliberate perioperative systems design improves operating room throughput. *Anesthesiology* 2005; **103**: 406–18.
28. Torkki PM, Marjamaa RA, Torkki MI et al. Use of anesthesia induction rooms can increase the number of urgent orthopedic cases completed within 7 hours. *Anesthesiology* 2005; **103**: 401–5.
29. Sokolovic E, Biro P, Wyss P et al. Impact of the reduction of anaesthesia turnover time on operating room efficiency. *Eur J Anaesthesiol* 2002; **19**: 560–3.
30. Hanss R, Buttgerit B, Tonner PH et al. Overlapping induction of anesthesia: an analysis of benefits and costs. *Anesthesiology* 2005; **103**: 391–400.
31. Eappen S, Flanagan H, Lithman R et al. The addition of a regional block team to the orthopedic operating rooms does not improve anesthesia-controlled times and turnover time in the setting of long turnover times. *J Clin Anesth* 2007; **19**: 85–91.
32. Stahl JE, Sandberg WS, Daily B et al. Reorganizing patient care and workflow in the operating room: a cost-effectiveness study. *Surgery* 2006; **139**: 717–28.
33. Cendan JC, Good M. Interdisciplinary work flow assessment and redesign decreases operating room turnover time and allows for additional caseload. *Arch Surg* 2006; **141**: 65–9.
34. Harders M, Malangoni MA, Weight S et al. Improving operating room efficiency through process redesign. *Surgery* 2006; **140**: 509–14.
35. Sieber TJ, Leibundgut DL. Operating room management and strategies in Switzerland: results of a survey. *Eur J Anaesthesiol* 2002; **19**: 415–23.
36. Weinbroum AA, Ekstein P, Ezri T. Efficiency of the operating room suite. *Am J Surg* 2003; **185**: 244–50.
37. Overdyk FJ, Harvey SC, Fishman RL et al. Successful strategies for improving operating room efficiency at academic institutions. *Anesth Analg* 1998; **86**: 896–906.
38. Lapierre SD, Batson C, McCaskey S. Improving on-time performance in health care organizations: a case study. *Health Care Manage Sci* 1999; **2**: 27–34.
39. Marjamaa RA, Torkki PM, Torkki MI et al. Time accuracy of a radio frequency identification patient tracking system for recording operating room timestamps. *Anesth Analg* 2006; **102**: 1183–6.
40. Torkki PM, Alho AI, Peltokorpi AV et al. Managing urgent surgery as a process: case study of a trauma center. *Int J Technol Assess Health Care* 2006; **22**: 255–60.
41. Vitez TS, Macario A. Setting performance standards for an anesthesia department. *J Clin Anesth* 1998; **10**: 166–75.
42. Macario A. Are your hospital operating rooms “efficient”? A scoring system with eight performance indicators. *Anesthesiology* 2006; **105**: 237–40.
43. Fasting S, Gisvold SE. Statistical process control methods allow the analysis and improvement of anesthesia care. *Can J Anaesth* 2003; **50**: 767–74.
44. Seim A, Andersen B, Sandberg WS. Statistical process control as a tool for monitoring nonoperative time. *Anesthesiology* 2006; **105**: 370–80.
45. Macario A, Vasanaawala M. Technology and computing in the surgical suite: key features of an or management information system and opportunities for the future. *Anesth Analg* 2002; **95**: 1120–1.
46. Correll DJ, Bader AM, Hull MW et al. Value of preoperative clinic visits in identifying issues with potential impact on operating room efficiency. *Anesthesiology* 2006; **105**: 1254–9.
47. Sokal SM, Craft DL, Chang Y et al. Maximizing operating room and recovery room capacity in an era of constrained resources. *Arch Surg* 2006; **141**: 389–93; discussion 93–5.
48. Bhattacharyya T, Vrahas MS, Morrison SM et al. The value of the dedicated orthopaedic trauma operating room. *J Trauma* 2006; **60**: 1336–40.
49. Vakkuri A, Niskanen M, Meretoja OA et al. Allocation of tasks between anesthesiologists and anesthesia nurses in Finland. *Acta Anaesthesiol Scand* 2006; **50**: 659–63.
50. St Jacques PJ, Patel N, Higgins MS. Improving anesthesiologist performance through profiling and incentives. *J Clin Anesth* 2004; **16**: 523–8.
51. Flin R, Fletcher G, McGeorge P et al. Anaesthetists’ attitudes to teamwork and safety. *Anaesthesia* 2003; **58**: 233–42.
52. Marjamaa RA, Kirvela OA. Who is responsible for operating room management and how do we measure how well we do it? *Acta Anaesthesiol Scand* 2007; **51**: 809–14.

Address:

Anne Vakkuri  
 Department of Anesthesiology and Intensive Care Medicine  
 Peijas Hospital  
 Helsinki University Hospital  
 P.O. Box 900  
 FIN-00029 HUS, Helsinki  
 Finland  
 e-mail anne.vakkuri@hus.fi

Copyright of *Acta Anaesthesiologica Scandinavica* is the property of Blackwell Publishing Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.