SuSiE versus EM SuSiE−mix | iteration 18

P < 1e−08 | mean reads >= 100 | >=1 CS in either model | 324 regions across 27 tissues

A CS−count agreement: four high−disagreement tissues

69.8% same CS count overall | 226 / 324 regions

Rows: SuSiE CSs | Columns: EM CSs | Cells: region counts

Vagina
Disagreement rank 1 | n = 12
Same: 0 (0.0%) | Different: 12

	0	1	2
0	–	0	0
1	10	0	0
2	2	0	0

Heart
Disagreement rank 2 | n = 12
Same: 6 (50.0%) | Different: 6

	0	1	2
0	–	0	0
1	1	6	1
2	0	4	0

Testis
Disagreement rank 3 | n = 12
Same: 6 (50.0%) | Different: 6

	0	1	2
0	–	0	0
1	1	6	1
2	0	4	0

Blood
Disagreement rank 4 | n = 12
Same: 7 (58.3%) | Different: 5

	0	1	2
0	–	0	0
1	1	7	1
2	0	3	0

B How far do leads differ? All tissues

63.5% of matched leads retained | 181 / 285 pairs



Curves exclude 181 identical pairs and 41 extra CSs in model−only regions.

Changed pairs use one−to−one matches; extra CSs use the nearest other−model lead.

C Distance to TSS for the extra leads in B



Same extra CSs as B; 1 missing TSS distances excluded.

Model−only regions excluded from both ECDFs. TSS proximity does not establish accuracy.

D Posterior coding support in the same cohort

All 294 EM CSs: additive 81.8% | recessive 6.2% | dominant 12.0%

Mean component posterior probabilities; additive mass retained.



Each reported EM CS contributes its full component posterior; counts shown at right.

This is posterior support among reported CSs, not the fraction of nonadditive regions.

Same parent cohort in A−D. Table ranking: CS−count disagreement rate, minimum n = 10; B−D use all cohort tissues.

Distance matching is descriptive, not evidence of shared signals or LD. Matching unavailable for 0 selected regions.

SYNTHETIC VALIDATION − NOT GTEx RESULTS