SuSiE versus EM SuSiE−mix | iteration 18

P < 1e−08 | mean reads >= 100 | at least one CS in either model | 9,358 regions across 27 tissues

A CS−count agreement: four tissue extremes

Rows: SuSiE CSs | Columns: EM CSs | Cells: region counts

Skin

Most disagreement | n = 518

Same: 427 (82.4%) | Different: 91

	0	1	2	3	4	5
0	–	0	0	0	0	0
1	24	388	1	0	0	0
2	4	51	33	1	0	0
3	0	1	8	3	0	0
4	0	0	1	0	2	0
5	0	0	0	0	0	1

Muscle

Most disagreement | n = 628

Same: 526 (83.8%) | Different: 102

	0	1	2	3	4
0	–	0	0	0	0
1	28	450	3	0	0
2	2	59	72	0	0
3	0	2	7	4	0
4	0	0	0	1	0

Liver

Most agreement | n = 182

Same: 175 (96.2%) | Different: 7

	0	1	2
0	–	0	0
1	5	168	0
2	1	1	7

Prostate

Most agreement | n = 157

Same: 149 (94.9%) | Different: 8

	0	1	2
0	–	0	0
1	7	140	0
2	0	1	9

B How far do leads differ? All tissues



Excluded here: 9,676 identical pairs; 503 extra CSs without a counterpart model CS

C Lead retention across CS−count groups



D Coding and support of extra EM CSs



Extra = unpaired after identical−lead and distance matching; includes EM−only regions.

Tissue ranking: equal−count rate, minimum n = 100. CS leads compare exact SNP IDs, ignoring coding.

Positional matches do not establish shared signals or LD. Audited exclusions: 0 lead comparisons; 0 distance comparisons.