

CYSTIC FIBROSIS - INHALED ANTIBIOTICS

toll free 855.274.1694 toll free fax 844.306.0200

krogerspecialtypharmacy.com

PATIENT INFO			PRESCRIBER INFO			
Patient Name	lale 🖵 Female	Date of Birth	Prescriber Nar		Supervising MD NPI	
Address Ci	City, State, Zip		DEA#	NPI#	License #	
Phone Al	Allergies		Address		City, State, Zip	
CFR Mutation	Weight □ lbs □ kg				Fax	
PLEASE FAX COPY OF: 📮 PRES	CRIPTION CA	RD FRONT & BA	CK 🗆 CLII	NICAL NOTES	☐ MEDICAL CARD FROM	IT & BACK
		CLINICAL IN	FORMATIC	ON		
Diagnosis: □ E84.0 Cystic Fibrosis with pulmor □ E84.19 Cystic Fibrosis with intesti □ Other: Drug Allergies:	nal manifestations	■ E84.8 Cystic Fib	rosis with other		1 E84.11 Mecoium ileus in Cystic Fik 1 B96.5 pseudomonas (mallei) causi	
MEDICATION		DIRECT	IONS			QTY. REFILLS
Amikacin ☐ 500mg/2ml vial ☐ Sodium chloride 0.9% (3mL) ☐ BD 5mL syringes - Sig: use as directed	☐ Amikacin 250mg UDN BID x 28 days on/off or continuous ☐ Amikacin 500mg UDN BID x 28 days on/off or continuous ☐ <u>Sodium chloride 0.9% (3mL):</u> Add 1 vial (3mL) of sodium chloride 0.9% into neb cup with amikacin mixture and inhale contents via nebulizer BID					
☐ Bethkis 300mg/4mL	Sig: 1 vial via nebulizer BID x 28 days on/off					
Cayston 75mg vial ☐ Altera Nebulizer - Sig: use as directed ☐ Pari Smart mask (ped/kid) Sig: use as directed	Inhale 1 vial via ALTERA nebulizer TID x 28 days on/off ☐ Altera Handset – sig: use as directed					
Ceftazidime ☐ 1 gram vial of Ceftazidime ☐ 2 gram vial of Ceftazidime ☐ Sterile water (10mL) OR ☐ Sodium chloride 0.9% (10mL) ☐ BD 5mL syringes - Sig: use as directed	□ 1 gram BID x 28 days on/off or continuous □ 2 gram BID x 28 days on/off or continuous Reconstitution instructions for: Mix 1 vial of Ceftazidime withmL of sterile water/sodium chloride and give via nebulizer					
Colistimethate 150mg vial ☐ Sterile water for injection (10mL vial) ☐ BD syringes 5mL 22G x 1 ½	Reconstitute 1 vial (150mg) with 8mL water and give 4mL (75mg) via neb B on and 28 days off Sterile water 10mL vial Sig: Draw 8mL to reconstitute 1 vial of c BD syringes 5mL 22G x 1 ½ Sig: Use as directed with Colistimetha		BID x 28 days	water and give 4r on and 28 days of Sterile water 10m Sig: Draw 4mL to r BD syringes 5mL	nL vial reconstitute 1 vial of colistimethate	
Gentamicin ☐ 80mg/2ml vial ☐ BD 5mL syringes Sig: Use as directed	Sig: 80mg via nebulizer BID					
☐ Kitabis Pak (Tobramycin inhalation solution with PARI LC nebulizer)	Sig: 1 vial via nebulizer BID x 28 days on/off					
☐ Tobi podhaler 28mg per cap	Inhale the contents of 4 capsules via podhaler BID x 28 days on/off					
☐ Tobramycin 300mg/5mL	Sig: 1 vial via nebulizer BID x 28 days on/off					
Vancomycin ☐ 500mg vial of Vancomycin ☐ 250mg vial of Vancomycin ☐ Sterile water (10mL) ☐ BD 5mL syringes - Sig: use as directed	250mg BID x 28 days on/off or continuous Reconstitution instructions for: Mix 1 vial of Vancomycin withmL of sterile water and give via nebulizer					
By signing this form and utilizing our services, you are authorizi	ng Kroger Specialty Phari	macy™ and it's employees	o serve as your prior	authorization designated	agent in dealing with medical and prescriptic	n insurance companies.

Prescriber's Signature (no stamps) Substitution Permitted Date Prescriber's Signature (no stamps) Dispense As Written

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged property or exempt from disclosure under applicable law. If you are not the named addresses you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.