



SPECIALTY PHARMACY

DERMATOLOGY

Garden Grove, CA toll free 800.228.3643 toll free fax 866.539.1092

krogerspecialtypharmacy.com

DATE: _____ NEEDS BY DATE: _____ SHIP TO: ☐ PATIENT ☐ OFFICE ☐ OTHER _____

| PATIENT INFO | | PRESCRIBER INFO | |
|-------------------|---|------------------|-----------------|
| Patient Name | | Prescriber Name | |
| Address | | DEA # | NPI # License # |
| City, State, Zip | | Address | |
| Main Phone | Alternate Phone | City, State, Zip | |
| Social Security # | | Phone | Fax |
| Date of Birth | <input type="checkbox"/> Male <input type="checkbox"/> Female | Contact Person | |

PLEASE FAX COPY OF: ☐ PRESCRIPTION CARD FRONT & BACK ☐ CLINICAL NOTES ☐ MEDICAL CARD FRONT & BACK

CLINICAL INFORMATION

 Diagnosis: ☐ L20.9 Atopic Dermatitis ☐ L40.8 Moderate to Severe Plaque Psoriasis ☐ L40.50 Psoriatic Arthritis ☐ L73.2 Hidradenitis Suppurativa - Hurley Stage _____
☐ Other: Dx code _____ Condition _____

Drug Allergies: _____

Location: % BSA: _____ ☐ Hands ☐ Feet ☐ Scalp ☐ Groin ☐ Nails ☐ Other: _____Prior Failed Meds: ☐ Biologics ☐ Cimzia ☐ Cosentyx ☐ Enbrel ☐ Humira ☐ Orencia ☐ Remicade ☐ Rituxan ☐ Simponi ☐ Stelara☐ MTX ☐ Soriatane ☐ CYA Length of Treatment _____ Reason for Discontinuing _____☐ PUVA/UVB Length of Treatment _____ Reason for Discontinuing _____☐ Topicals Length of Treatment _____ Inadequate Response List Specific Names _____☐ Contraindicated Medication _____ Reason _____Does patient have a latex allergy? ☐ Yes ☐ No TB/PPD Test given (or intended to be given before biologic started)? ☐ Yes ☐ No (PLEASE send LAB result)

PRESCRIPTION INFORMATION

QUANTITY

REFILLS

| | | | | |
|---|---|---|--------------------------------|---------------------|
| <input type="checkbox"/> Cosentyx* | 300mg (2x150) <input type="checkbox"/> Pen <input type="checkbox"/> PFS 150mg <input type="checkbox"/> Pen <input type="checkbox"/> PFS | <input type="checkbox"/> Load: Inject <input type="checkbox"/> 300mg or <input type="checkbox"/> 150mg subcutaneously week 0, 1, 2, 3, 4 <input type="checkbox"/> Maintenance: Inject <input type="checkbox"/> 300mg or <input type="checkbox"/> 150mg subcutaneously every 4 weeks | 5 week supply 4 week supply | none |
| <input type="checkbox"/> Dupixent* | 300mg/2 mL PFS w/ shield | <input type="checkbox"/> Load: Inject 600mg (2-300mg injections in different injection sites) on Day 1, then 300mg on Day 15, then 300mg every other week <input type="checkbox"/> Maintenance: Inject 300mg subcutaneously every other week | 4 syringes 2 syringes | none |
| <input type="checkbox"/> Enbrel* | <input type="checkbox"/> 50mg Sureclick <input type="checkbox"/> 50mg Prefilled Syringe <input type="checkbox"/> 25mg Prefilled Syringe <input type="checkbox"/> 25mg Vials Wt: _____ | <input type="checkbox"/> Inject 50mg subcutaneously TWICE a week 72-96 hours apart <input type="checkbox"/> Inject 50mg subcutaneously ONCE a week <input type="checkbox"/> Inject 25mg subcutaneously TWICE a week 72-96 hours apart <input type="checkbox"/> Inject 2-25mg (50mg) on same day TWICE a week 72-96 hours apart <input type="checkbox"/> Inject 0.8mg/kg (_____mg) subcutaneously ONCE a week | 4 week supply | _____ |
| <input type="checkbox"/> Erivedge* | 150mg capsule | Take one capsule by mouth daily | 4 week supply | _____ |
| <input type="checkbox"/> Humira* | <input type="checkbox"/> Psoriasis Starter Kit <input type="checkbox"/> 40mg Pen <input type="checkbox"/> 40mg Prefilled Syringe | <input type="checkbox"/> Inject 2-40mg (80mg) on Day 1, then 40mg on Day 8, then 40mg every other week <input type="checkbox"/> Inject 40mg subcutaneously EVERY OTHER week <input type="checkbox"/> Inject 40mg subcutaneously ONCE a week | Loading Dose 4 week supply | none |
| <input type="checkbox"/> Humira* HS | <input type="checkbox"/> HS Starter Package <input type="checkbox"/> 40mg Pen <input type="checkbox"/> 40mg Prefilled Syringe | 160mg given as <input type="checkbox"/> Four 40mg SubQ day 1 OR <input type="checkbox"/> Two 40mg SubQ days 1 & 2 then week 2 inject 80mg (Two 40mg injections) subcutaneously on day 15 <input type="checkbox"/> Week 4 +: Inject 40mg SQ weekly | Loading Dose 4 week supply | none |
| <input type="checkbox"/> Odomzo* | 200mg capsule | Take one capsule by mouth daily on an empty stomach, 1 hour before or 2 hours after a meal | 30 | _____ |
| <input type="checkbox"/> Otezla* | <input type="checkbox"/> Starter Pack <input type="checkbox"/> 30mg Tablets | <input type="checkbox"/> Titrate: Take 1 tablet on day 1 then twice daily as directed OR date provided _____ <input type="checkbox"/> Maintenance: Take 1 tablet by mouth twice daily. <input type="checkbox"/> Bridge Rx: Take 1 tablet by mouth twice daily; dispensed by OSP | 1 Starter Pack 60 28 | none _____ 12 |
| <input type="checkbox"/> Remicade* Wt: _____ | 100mg Vial | <input type="checkbox"/> Infuse _____mg at week 0, 2, 6 <input type="checkbox"/> Infuse _____mg at every _____ weeks | Loading dose | none |
| <input type="checkbox"/> Siliq* | 210mg Prefilled Syringe | <input type="checkbox"/> Load: Inject 210mg subcutaneously at weeks 0, 1 and 2 and then every 2 weeks <input type="checkbox"/> Maintenance: Inject 210mg subcutaneously every 2 weeks | 4 syringes 2 syringes | none |
| <input type="checkbox"/> Simponi* | 50mg <input type="checkbox"/> SmartJect <input type="checkbox"/> PFS | Inject 50mg subcutaneously once a month as directed | 4 week supply | _____ |
| <input type="checkbox"/> Stelara* Wt: _____ | <input type="checkbox"/> 45mg Prefilled Syringe <input type="checkbox"/> 90mg Prefilled Syringe | Inject 45mg on day 0, then week 4, then every 12 weeks (for Patients ≤ 220 lbs) Inject 90mg on day 0, then week 4, then every 12 weeks (for Patients > 220 lbs) | 1 1 | _____ _____ |
| <input type="checkbox"/> Taltz™ | <input type="checkbox"/> 80mg/mL Autoinjector <input type="checkbox"/> 80mg/mL Prefilled Syringe | <input type="checkbox"/> Load: Inject 160mg (2 – 80mg) subcutaneously on week 0, then inject 80mg week 2 then Inject 80mg every 2 weeks (weeks 4-10) then Inject 80mg at week 12 <input type="checkbox"/> Maintenance Dose: Inject 80mg every 4 weeks | 3 2 1 1 | none 1 none |
| <input type="checkbox"/> Tremfya™ | 100mg/ml Prefilled Syringe | <input type="checkbox"/> Load: Inject 100mg subcutaneously on week 0 and on week 4 <input type="checkbox"/> Maintenance: Inject 100mg subcutaneously every 8 weeks | 1 1 | _____ _____ |
| <input type="checkbox"/> Other | | | | |

By signing this form and utilizing our services, you are authorizing Kroger Specialty Pharmacy™ and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature (no stamps)

Substitution Permitted

Date

Prescriber's Signature (no stamps)

Dispense As Written

Date

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged property or exempt from disclosure under applicable law. If you are not the named addressee you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately. 04866 08-07-2017 GGCA