

NEUROLOGY

Lake Mary, FL toll free

toll free fax

krogerspecialtypharmacy.com

SPECIALTY PHARMACY		DATE: NEEDS BY DATE: SHIP TO: □ PATIENT □ OFFICE □ OTHER			
	PATIEN	IT INFO	PRESCRIBER INFO		
Patient Name			Prescriber Name		
Address			DEA# NPI#	License #	
City, State, Zip			Address		
Main Phone	Alternate Phone		City, State, Zip		
Social Security # Phone			Phone Fax		
Date of Birth		☐ Male ☐ Female	Contact Person		
PLEASE	FAX COPY OF: 📮 I	PRESCRIPTION CARD FRONT & B.	ACK 🗖 CLINICAL NOTES 📮 MEDIC	AL CARD FRONT	& BACK
		CLINICAL IN	NFORMATION		
Drug Allergies: History: •	ls the patient currently on t Will patient stop taking cur How long will the patient v	□ Primary Progressive □ Secondary Pously treated for this condition? therapy? rent therapy before starting new therapy? wait before starting the new therapy?	□ No □ Yes Medication failed □ No □ Yes Medication failed □ Yes □ No		
•.	Are there other medication	ns patient currently taking? Please list:			
		PRESCRIPTION INFO		QUANTITY	REFILLS
□ Avonex®	30mcg □ PFS □ PEN □ Pwo		□ Inject 30mcg intramusculary once weekly □ Other dosing:		
□ Betaseron®	0.3mg Vial	(0.1875mg (0.75ml), Week 7+: 0.25mg (☐ Initial Week 1&2: 0.0625mg (0.25ml), Week 3&4: 0.125mg (0.5ml), Week 5&6: 0.1875mg (0.1875mg (0.75ml), Week 7+: 0.25mg (1ml) SQ every other day ☐ Maintenance Dose: Inject 0.25mg (1ml) SQ every other day		
□ Copaxone®	□ 20mg PFS □ 40mg /mL PFS	☐ Inject 20mg subcutaneously daily☐ Inject 40mg subcutaneously three tir	□ Inject 20mg subcutaneously daily □ Inject 40mg subcutaneously three times a week		
□ Extavia [®]	0.3mg Vial	(0.1875mg (0.75ml), Week 7+: 0.25mg (☐ Initial Week 1&2: 0.0625mg (0.25ml), Week 3&4: 0.125mg (0.5ml), Week 5&6: 0.1875mg (0.1875mg (0.75ml), Week 7+: 0.25mg (1ml) SQ every other day ☐ Maintenance Dose: Inject 0.25mg (1ml) SQ every other day		
□ Gilenya°	0.5mg Capsule	Take 1 capsule PO daily	Take 1 capsule PO daily		
□ Glatopa°	20mg PFS	Inject 20mg SQ daily	Inject 20mg SQ daily		
□ Plegridy°	□ Starter Pack □ PEN □ 125mcg PFS □ 125mcg PEN	PFS Inject 63mcg SQ on day-1, then 94m Inject 125mcg SQ every 14 days	□ Inject 63mcg SQ on day-1, then 94mcg SQ on day -15, then 125mcg SQ on day-29 □ Inject 125mcg SQ every 14 days		
□ Rebif°	☐ Titration Rebidose ☐ Titration Pack ☐ 22mcg/0.5mL Rebidos ☐ 22mcg/0.5mL PFS ☐ 44mcg/0.5mL Rebidos ☐ 44mcg/0.5mL PFS	wk 5+: inject 44mcg SQTIW e ☐ Maintenance Dose: Inject mcg SQ	□ Titration Dose: wk 1&2: inject 8.8mcg SQ TIW; wk 3& 4: inject 22mcg SQ TIW; wk 5+: inject 44mcg SQ TIW □ Maintenance Dose: Inject mcg SQ TIW		
□ Tecfidera®	☐ Titration Starter Pack Caps ☐ 240mg ☐ 120mg		7 days, then take 240mg PO BID thereafter	1kit 1 month supply 1 month supply	None 11 11
□ Epipen° □ Epipen Jr.°		Inject 1 pen into thigh area in case of a	naphylaxis; may repeat	2 pen pack	
□ Other		<u> </u>			<u> </u>
By signing this form	and utilizing our convices, you are a	uthorizing Krager Specialty Pharmacy/M and it's employees	s to some as your prior authorization designated agent in dealing w	ith modical and procerintian i	nsuransa sampanias

Prescriber's Signature (no stamps) Substitution Permitted Date Prescriber's Signature (no stamps) Dispense As Written

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged property or exempt from disclosure under applicable law. If you are not the named addresses you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.