

# DEFINABLE COMBINATORICS AT THE FIRST UNCOUNTABLE CARDINAL

WILLIAM CHAN AND STEPHEN JACKSON

**ABSTRACT.** We work throughout in the theory ZF with the axiom of determinacy, AD. We introduce and prove some club uniformization principles under AD and  $\text{AD}_{\mathbb{R}}$ . Using these principles, we establish continuity results for functions of the form  $\Phi: [\omega_1]^{\omega_1} \rightarrow \omega_1$  and  $\Psi: [\omega_1]^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$ . Specifically, for every function  $\Phi: [\omega_1]^{\omega_1} \rightarrow \omega_1$ , there is a club  $C \subseteq \omega_1$  so that  $\Phi \upharpoonright [C]_*^{\omega_1}$  is a continuous function. This has several consequences such as establishing the cardinal relation  $|[\omega_1]^{<\omega_1}| < |[\omega_1]^{\omega_1}|$  and answering a question of Zapletal by showing that if  $\langle X_\alpha : \alpha < \omega_1 \rangle$  is a collection of subsets of  $[\omega_1]^{\omega_1}$  with the property that  $\bigcup_{\alpha < \omega_1} X_\alpha = [\omega_1]^{\omega_1}$ , then there is an  $\alpha < \omega_1$  so that  $X_\alpha$  and  $[\omega_1]^{\omega_1}$  are in bijection.

We show that under  $\text{AD}_{\mathbb{R}}$  everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization holds which is the following statement: Let  $\text{club}_{\omega_1}$  denote the collection of club subsets of  $\omega_1$ . Suppose  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  is  $\subseteq$ -downward closed in the sense that for all  $\sigma \in [\omega_1]^{<\omega_1}$ , for all clubs  $C \subseteq D$ ,  $R(\sigma, D)$  implies  $R(\sigma, C)$ . Then there is a function  $F: \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  so that for all  $\sigma \in \text{dom}(R)$ ,  $R(\sigma, F(\sigma))$ .

We show that under AD almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization holds which is the statement that for every  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed, there is a club  $C$  and a function  $F: \text{dom}(R) \cap [C]_*^{<\omega_1} \rightarrow \text{club}_{\omega_1}$  so that for all  $\sigma \in \text{dom}(R) \cap [C]_*^{<\omega_1}$ ,  $R(\sigma, F(\sigma))$ .

## 1. INTRODUCTION

The setting throughout this article will be  $\text{ZF} + \text{AD}$ . AD is the axiom of determinacy which asserts that for every two player integer game, one of the two players must have a winning strategy. AD and its various extensions have been shown to be a fruitful and general framework for extending properties of simple subsets of  $\mathbb{R}$  to a much more general class. Within this setting, sets which are surjective images of  $\mathbb{R}$  have a very interesting structure.

The definable properties of  $\mathbb{R}$  and its subsets have long been studied within descriptive set theory. Under determinacy, the first uncountable cardinal,  $\omega_1$ , is a minimal uncountable set much like  $\mathbb{R}$ . AD can distinguish  $\omega_1$  and  $\mathbb{R}$  via bijections:  $\omega_1$  and  $\mathbb{R}$  are incomparable cardinals in the sense that neither can inject into the other. Moreover, under a strengthening of AD called  $\text{AD}^+$ , Woodin's perfect set dichotomy implies that every uncountable set  $X$  which is a surjective image of  $\mathbb{R}$  must contain a copy of  $\mathbb{R}$  or  $\omega_1$ . (See [3] Section 8 or [4].) More generally, [1] showed that in  $L(\mathbb{R}) \models \text{AD}$ , every uncountable set  $X$  must contain a copy of  $\mathbb{R}$  or  $\omega_1$ . Like its companion  $\mathbb{R}$ ,  $\omega_1$  and its subsets deserves a definable analysis.

Note that  $\mathbb{R}$ ,  $\mathcal{P}(\omega)$ , and  $[\omega]^\omega$  (where  $[\omega]^\omega$  is the collection of increasing functions from  $\omega$  into  $\omega$ ) are all in bijection. Let  $[\omega_1]^{\omega_1}$  denote the collection of increasing functions from  $\omega_1$  to  $\omega_1$ .  $[\omega_1]^{\omega_1}$  is in bijection with  $\mathcal{P}(\omega_1)$ . Under AD, the cardinal structure below  $|\mathbb{R}| = |\mathcal{P}(\omega)| = |[\omega]^\omega|$  is fully understood. One motivation for this article was to explore the definable cardinals around  $|\mathcal{P}(\omega_1)| = |[\omega_1]^{\omega_1}|$  under AD. A continuity phenomenon for functions of the form  $\Phi: [\omega_1]^{\omega_1} \rightarrow \omega_1$  will be a useful tool for studying the cardinals below  $\mathcal{P}(\omega_1)$ . The continuity phenomenon will be shown to be a consequence of a choice principle for club subsets of  $\omega_1$  which is fundamentally useful for studying definable combinatorics on  $|[\omega_1]^{\omega_1}| = |\mathcal{P}(\omega_1)|$  under AD.

The continuity phenomenon in a general sense asserts that a local property of the output of a function can be determined by a local behavior of the input. Philosophically, this is motivated by a question of whether it is possible for one to truly use all of a function  $f \in [\omega_1]^{\omega_1}$  in order to assign to  $f$  a countable ordinal.

As motivation, consider the classical case of a function  $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ . As customary in descriptive set theory,  $\mathbb{R}$  denotes  ${}^\omega\omega$  which is the collection of functions from  $\omega$  into  $\omega$ . A priori,  $\Phi$  may need all of  $f \in \mathbb{R}$  even to determine the first bit  $\Phi(f)(0)$  of  $\Phi(f)$ . That is, if  $g$  differs from  $f$  at any place,  $\Phi(f)(0)$  could potentially be different from  $\Phi(g)(0)$ . However, if  $\Phi$  is continuous, then there is a  $j \in \omega$  so that if  $f \upharpoonright j = g \upharpoonright j$ , then  $\Phi(f)(0) = \Phi(g)(0)$ . Thus one can determine the value of  $\Phi(f)(0)$  forever by freezing an appropriate local

---

July 24, 2020. The first author was supported by NSF grant DMS-1703708. The second author was supported by NSF grant DMS-1800323.

behavior of the input  $f$ . Certainly not all functions  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  are continuous. However, under AD, every function is continuous almost everywhere in the sense that there is a comeager set  $C \subseteq \mathbb{R}$  so that  $\Phi \upharpoonright C$  is a continuous function.

Now consider a function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ . First, one needs an appropriate notion of “almost-everywhere”. Let  $\mu$  be the collection of subsets of  $\omega_1$  which contain a club subset of  $\omega_1$ . Solovay showed that  $\mu$  is a normal countably complete measure on  $\omega_1$  under AD. It has the distinction of being the unique normal measure on  $\omega_1$ . Let  $\mu_{\omega_1}$  be the filter on  $[\omega_1]_*^{\omega_1}$  defined by  $X \in \mu_{\omega_1}$  if and only if there is a club  $C \subseteq \omega_1$  so that  $[C]_*^{\omega_1} \subseteq X$ . (If  $A \subseteq \omega_1$ ,  $[A]_*^{\omega_1}$  is the collection of increasing functions from  $\omega_1$  into  $A$  which are of the correct type. See Definition 2.1.) Using the (correct type) strong partition property  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$  of Martin, one can show that  $\mu_{\omega_1}$  is a countably complete measure on  $[\omega_1]_*^{\omega_1}$ . Using  $\mu_{\omega_1}$  as the notion of almost-everywhere is both natural and robust since it allows the strong partition property as a powerful tool in analyzing the continuity phenomenon. (The use of correct type is needed to obtain club homogeneous sets for partitions. One can show  $[\omega_1]^{\omega_1}$  and  $\mathcal{P}(\omega_1)$  are in bijection with  $[\omega_1]_*^{\omega_1}$ . For this reason, this article will prefer  $[\omega_1]_*^{\omega_1}$  over  $\mathcal{P}(\omega_1)$ .)

So the question becomes: For every  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$ , is  $\Phi$  continuous  $\mu_{\omega_1}$ -almost everywhere? Precisely, is there a club  $C \subseteq \omega_1$  so that for all  $f \in [C]_*^{\omega_1}$ , there is an  $\alpha < \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$  with  $f \upharpoonright \alpha = g \upharpoonright \alpha$ ,  $\Phi(f) = \Phi(g)$ ?

There is a great deal of empirical evidence that the continuity property holds. Any function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$  which is of bounded dependence  $\mu_{\omega_1}$ -almost everywhere is continuous  $\mu_{\omega_1}$ -almost everywhere. (This means that there is an  $\epsilon < \omega_1$  and a function  $\Psi : [\omega_1]_*^\epsilon \rightarrow \omega_1$  so that for  $\mu_{\omega_1}$  almost all  $f$ ,  $\Phi(f) = \Psi(f \upharpoonright \epsilon)$ .) The function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$  defined by  $\Phi(f) = \sup_{\alpha < f(0)} f(\alpha)$  does not have bounded dependence, but it is continuous.

One can even attempt to use definability notions to construct a function that ostensibly seems to use the entire sequence to define an output: For instance, let  $\Phi(f) = \omega_1^{L[f]}$ . This function is discussed in Example 4.2 where it is shown that  $\mu_{\omega_1}$ -almost everywhere this function is constant. Thus for  $\mu_{\omega_1}$ -almost all  $f$ ,  $\Phi$  actually uses no information about  $f$  to determine the output  $\Phi(f)$ .

This article will show that the continuity phenomenon holds for every function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ :

**Theorem 4.5.** *Assume ZF + AD. Every function  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is continuous  $\mu_{\omega_1}$ -almost everywhere.*

The continuity property, in its various forms, has interesting mathematical consequences for definable combinatorics under determinacy. The continuity property for function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is an important tool for the study of the Mycielski and Jónsson property for quotient of  $E_0$  in [6] and [2]. Furthermore in [5], a form of the continuity property is established for functions  $\Phi : [\omega_1]_*^\epsilon \rightarrow \omega_1$  where  $\epsilon < \omega_1$  and for functions  $\Phi : [\omega_2]_*^\epsilon \rightarrow \omega_2$  where  $\epsilon < \omega_2$  in order to give a purely descriptive set theoretic proof under AD that  $|[\omega_1]^\omega| < |[\omega_1]^{<\omega_1}|$  and  $|[\omega_2]^\omega| < |[\omega_2]^{<\omega_1}| < |[\omega_2]^{\omega_1}| < |[\omega_2]^{<\omega_2}|$ .

Using the continuity property at  $\omega_1$ , one can give a purely descriptive set theoretic proof of the following cardinality computation:

**Theorem 4.7.** *Assume ZF + AD.  $|[\omega_1]^{<\omega_1}| < |[\omega_1]^{\omega_1}|$ .*

Zapletal also asked the first author the following basic combinatorial question: Assume AD. If one partitions  $[\omega_1]^{\omega_1}$  (or equivalently  $\mathcal{P}(\omega_1)$ ) into  $\omega_1$  many pieces,  $\langle X_\alpha : \alpha < \omega_1 \rangle$ , so that  $X_\alpha \subseteq [\omega_1]^{\omega_1}$  and  $\bigcup_{\alpha < \omega_1} X_\alpha = [\omega_1]^{\omega_1}$ , then must there be a piece  $X_\alpha$  so that  $X_\alpha \approx [\omega_1]^{\omega_1}$ , meaning  $X_\alpha$  is in bijection with  $[\omega_1]^{\omega_1}$ ? A positive answer follows from the continuity property.

**Theorem 4.6.** *Assume ZF + AD. Suppose  $\langle X_\alpha : \alpha < \omega_1 \rangle$  is a sequence of subsets of  $[\omega_1]^{\omega_1}$  so that  $\bigcup_{\alpha < \omega_1} X_\alpha = [\omega_1]^{\omega_1}$ . Then there is an  $\alpha < \omega_1$  so that  $X_\alpha \approx [\omega_1]^{\omega_1}$ .*

A natural question extending Theorem 4.5 is to ask whether every function  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  is continuous  $\mu_{\omega_1}$ -almost everywhere. (Here  ${}^{\omega_1}\omega_1$  refers to the set of all functions  $f : \omega_1 \rightarrow \omega_1$ .) Given such a function  $\Phi$ , one can define  $\Phi_\beta : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  by  $\Phi_\beta(f) = \Phi(f)(\beta)$ . By applying Theorem 4.5 to  $\Phi_\beta$ , there is a club  $C$  so that  $\Phi_\beta \upharpoonright [C]_*^{\omega_1}$  is continuous. Although it is possible to show there is a sequence  $\langle C_\beta : \beta < \omega_1 \rangle$  so that

for all  $\beta < \omega_1$ ,  $\Phi_\beta \upharpoonright [C_\beta]_*^{\omega_1}$  is continuous (see [3] Fact 4.8), it is not clear how to use this sequence to obtain one single club  $C$  which witnesses that the original function  $\Phi : [\omega_1]^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  is continuous on  $[C]_*^{\omega_1}$  since an intersection of  $\omega_1$ -many club subsets of  $\omega_1$  may not be a club. Using ideas similar to the proof of Theorem 4.5 but with more elaborate partitions, the following almost everywhere continuity result can be shown.

**Theorem 5.3** (*With Trang*) *Assume ZF + AD. Every function  $\Phi : [\omega_1]^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  is continuous  $\mu_{\omega_1}$ -almost everywhere.*

The strong partition property for  $\omega_1$  is crucial in the arguments for establishing the continuity property for functions  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ . The second uncountable cardinal  $\omega_2$  fails to have the strong partition property but by a result of Martin and Paris, it does have the weak partition property, that is,  $\omega_2 \rightarrow (\omega_2)_2^\epsilon$  for each  $\epsilon < \omega_2$ . Using an explicit failure of the strong partition property for  $\omega_2$ , Section 6 shows that there is a function  $\Phi : [\omega_2]^{\omega_2} \rightarrow 2$  with no club  $C \subseteq \omega_2$  such that  $\Phi \upharpoonright [C]_*^{\omega_2}$  is continuous.

The main challenge in establishing Theorem 4.5 is to show that a certain natural partition  $P : [\omega_1]_*^{\omega_1} \rightarrow 2$  has a club homogeneous set for the desired side of the partition. As described in the proof of Theorem 4.5, one needs to make choices of club subsets of  $\omega_1$  which is dependent on previous choices of clubs. The axiom of determinacy is incompatible with many consequences of the axiom of choice. A selection principle for subsets of  $\omega_1$  is generally not possible in AD. To perform the construction mentioned above, one would need to prove a club uniformization result.

Let  $\text{club}_{\omega_1}$  denote the club subsets of  $\omega_1$ . In the applications of this paper, one has a relation  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed in the sense that for all  $C \subseteq D$  which are club subsets of  $\omega_1$  and for all  $\sigma$ , if  $R(\sigma, D)$  holds, then  $R(\sigma, C)$  holds.  $[\omega_1]^{<\omega_1}$ -club uniformization is the statement that there is a function  $\Lambda : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  so that for all  $\sigma \in [\omega_1]^{<\omega_1}$ ,  $R(\sigma, \Lambda(\sigma))$ .

For any  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  as above, there is a coded version  $\tilde{R} \subseteq \mathbb{R} \times \mathbb{R} \times \mathbb{R}$  of  $R$ . Theorem 3.7 shows that if  $\tilde{R}$  has a uniformization, then one can use the simple  $\omega_1$ -version of the Kechris-Woodin generic coding function (see [9]) and a category argument to establish that  $R$  has an (everywhere) uniformization. Thus under  $\text{AD}_{\mathbb{R}}$ , (everywhere)  $[\omega_1]^{<\omega_1}$ -club uniformization holds:

**Theorem 3.7.** *Assuming ZF +  $\text{AD}_{\mathbb{R}}$ ,  $[\omega_1]^{<\omega_1}$ -club uniformization holds.*

Under  $\text{AD}_{\mathbb{R}}$ , every relation  $S \subseteq \mathbb{R} \times \mathbb{R}$  can be uniformized. AD cannot prove this full uniformization since  $L(\mathbb{R}) \models \text{AD}$  has a relation on  $\mathbb{R} \times \mathbb{R}$  that cannot be uniformized. However, there is an almost everywhere uniformization result that does hold in AD: for any relation  $S \subseteq \mathbb{R} \times \mathbb{R}$ , there is a comeager  $C \subseteq \mathbb{R}$  and a function  $F : C \rightarrow \mathbb{R}$  which uniformizes  $S$  on  $C$ .

Similarly, AD cannot prove (everywhere)  $[\omega_1]^{<\omega_1}$ -club uniformization since Fact 3.9 shows that it fails in  $L(\mathbb{R}) \models \text{AD}$ . One says that almost-everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization holds if and only if for every relation  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed, there is a club  $C \subseteq \omega_1$  so that  $R \cap ([C]_*^{<\omega_1} \times \text{club}_{\omega_1})$  has a uniformization. By combining the generic coding function, category arguments, the Moschovakis coding lemma, and a fundamental idea of Martin (used in the study of the partition properties on  $\omega_1$ ) where the player with the winning strategy determines the property of the output but the losing player determines the identity of the output, one can prove the following which is one of the main results of this paper:

**Theorem 3.10.** *Assume ZF + AD. Almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization holds.*

Neeman has shown similar uniformization results (such as [10] Theorem 3.9) in  $L(\mathbb{R})$  using inner model theory techniques.

Almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization is used to verify that the partition used in the proof of the continuity property (Theorem 4.5) has a homogeneous club which is homogeneous for the desired side. Moreover, Theorem 3.10 is a powerful general technique for constructing functions  $h \in [\omega_1]_*^{\omega_1}$  which verify that partitions of a certain form are homogeneous for the desired side. The following template illustrates a very typical and simple use of Theorem 3.10:

Suppose  $P : [\omega_1]_*^{\omega_1} \rightarrow 2$  is a partition defined by  $P(f) = 0$  if and only if  $f$  does not have any “errors”. An error is a property of  $f$  so that if  $f$  has an error, then it must be witnessed at a  $\gamma < \omega_1$ . An example

of an error property could be that  $L[f] \models \neg \text{GCH}$ , i.e. the generalized continuum hypothesis fails in  $L[f]$ . For this example, if  $f$  has an error, then by a condensation argument, there is a  $\gamma < \omega_1$  which witnesses this error in the sense that  $L[f] \models 2^\gamma > \gamma^+$ . By the partition relations, there is a club  $D_0 \subseteq \omega_1$  which is homogeneous for  $P$ . Suppose one could show that for all  $\sigma \in [D_0]^{<\omega_1}$ , there is a club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$  such that  $\sup(\sigma) < g(0)$ ,  $\sigma \hat{\ } g$  does not have an error at any  $\gamma$  such that  $\sup(\sigma) \leq \gamma < g(0)$ . Define a relation  $R \subset [D_0]^{<\omega_1} \times \text{club}_{\omega_1}$  by  $R(\sigma, C)$  if and only if  $C$  has the above property with respect to  $\sigma$ .  $R$  is a relation which is  $\subseteq$ -downward closed in the  $\text{club}_{\omega_1}$ -coordinate. By Theorem 3.10, let  $D_1 \subseteq D_0$  and  $\Lambda : \text{dom}(R) \cap [D_1]_*^{<\omega_1} \rightarrow \text{club}_{\omega_1}$  be such that for all  $\sigma \in \text{dom}(R) \cap [D_1]_*^{<\omega_1}$ ,  $R(\sigma, \Lambda(\sigma))$ . Now construct a function  $h \in [D_1]_*^{\omega_1}$  by recursion as follows: Let  $F_0 = D_1 \cap \Lambda(\emptyset)$ . Let  $h(0)$  the  $\omega^{\text{th}}$  element of  $F_0$ . Suppose for some  $\alpha$ ,  $h \restriction \alpha$  and  $F_\beta$ , for all  $\beta < \alpha$ , have been defined. Let  $F_\alpha = \Lambda(h \restriction \alpha) \cap \bigcap_{\beta < \alpha} F_\beta$  and let  $h(\alpha)$  be the  $\omega^{\text{th}}$  element of  $F_\alpha$  larger than  $\sup(h \restriction \alpha)$ . This completes the construction. Note that  $h$  belongs to  $[D_1]_*^{\omega_1}$ . For each  $\alpha < \omega_1$ , let  $\text{drop}(h, \alpha) \in [D_1]_*^{\omega_1}$  be defined by  $\text{drop}(h, \alpha)(\gamma) = h(\alpha + \gamma)$ . For all  $\alpha < \omega_1$ , by construction, one has  $\text{drop}(h, \alpha) \in [F_\alpha]_*^{\omega_1} \subseteq [\Lambda(h \restriction \alpha)]_*^{\omega_1}$  and therefore  $h$  does not have an error at any  $\gamma$  with  $\sup(h \restriction \alpha) \leq \gamma < \text{drop}(h, \alpha)(0) = h(\alpha)$ . Thus  $h$  has no errors at any  $\gamma < \omega_1$ . Because errors must be witnessed at some  $\gamma < \omega_1$ ,  $h$  has no error.  $P(h) = 0$  and therefore  $D_0$  is homogeneous for  $P$  taking value 0.

The almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization of Theorem 3.10 is particularly important for studying the stable theory of the partition measure  $\mu_{\omega_1}$ : Each  $f \in [\omega_1]^{\omega_1}$ ,  $L[f]$  is naturally an  $\mathcal{L} = \{\dot{\in}, \dot{E}\}$  structure. Since  $\mu_{\omega_1}$  is an ultrafilter, for any  $\mathcal{L}$ -sentence, either (1) for  $\mu_{\omega_1}$ -almost all  $f$ ,  $L[f] \models \varphi$  or (2) for  $\mu_{\omega_1}$ -almost all  $f$ ,  $L[f] \models \neg \varphi$ . The  $\omega_1$ -stable theory is  $\mathfrak{T}^{\omega_1}$  which is defined to be the collection of  $\mathcal{L}$ -sentences  $\varphi$  so that for  $\mu_{\omega_1}$ -almost all  $f$ ,  $L[f] \models \varphi$ . One can ask which natural statements of set theory, such as  $\text{GCH}$ , belong to  $\mathfrak{T}^{\omega_1}$ . For instance, in forthcoming work of Chan, Jackson, and Trang, one can show that for any  $\sigma \in [\omega_1]^{<\omega_1}$ , there is a club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$ , for all  $\kappa$  with  $\sup(\sigma) \leq \kappa < g(0)$ ,  $L[\sigma \hat{\ } g] \models 2^\kappa = \kappa^+$ . Thus using the outline above, one has that  $\text{GCH} \in \mathfrak{T}^{\omega_1}$ . Using Theorem 3.10, one can also show that for  $\mu_{\omega_1}$ -almost all  $f$ ,  $L[f] \models (\forall \alpha < \omega_1)(f(\alpha) \text{ is a strongly inaccessible cardinal})$  and  $L[f]$  satisfies  $\Sigma_1^1$ -determinacy. One can also show that for  $\mu_{\omega_1}$ -almost all  $f$ ,  $L[f]$  has a canonical inner model  $L[\bar{v}_f]$  where  $\bar{v}_f$  is an  $\omega_1$ -length sequence of normal measure with discontinuous increasing sequence of critical points  $\bar{\kappa}$  so that  $f$  is generic over  $L[\bar{v}_f]$  for a generalized Prikry forcing  $\bar{\mathbb{P}}_{\bar{v}_f}$ , considered by Fuchs [7]. This can be used to show that for  $\mu_{\omega_1}$ -almost all  $f$ ,  $\Delta_2^1$ -determinacy fails in  $L[f]$ . Welch [11] has investigated similar questions in a different setting.

## 2. BASICS

Throughout the entire paper, assume  $\text{ZF} + \text{AD}$  (but not necessarily  $\text{DC}_{\mathbb{R}}$ ) unless otherwise explicitly stated.

Except for Theorem 2.16 and Theorem 2.17 which were proved or observed by the authors for this paper, the results of this section are well known and due to Martin and Solovay. This section will introduce the necessary notations and results. Although the proofs use a simple and fundamentally important idea of Martin that appears in his arguments for the partition properties, the exposition is quite tedious. A careful presentation is given in [3]. Specifically, see [3] Section 2, 3, and 4 for more details.

**Definition 2.1.** Let  $[\omega_1]^{\omega_1}$  denote the collection of strictly increasing functions  $f : \omega_1 \rightarrow \omega_1$ . A function  $f \in [\omega_1]^{\omega_1}$  has uniform cofinality  $\omega$  if and only if there is a function  $F : \omega_1 \times \omega \rightarrow \omega_1$  so that for all  $\alpha < \omega_1$ , for all  $n \in \omega$ ,  $F(\alpha, n) < F(\alpha, n+1)$  and  $f(\alpha) = \sup\{F(\alpha, n) : n \in \omega\}$ . A function  $f \in [\omega_1]^{\omega_1}$  has correct type if and only if  $f$  has uniform cofinality  $\omega$  and for all  $\alpha < \omega_1$ ,  $f(\alpha) > \sup\{f(\beta) : \beta < \alpha\}$ , that is,  $f$  is discontinuous everywhere. Let  $[\omega_1]_*^{\omega_1}$  denote the subset of  $[\omega_1]^{\omega_1}$  consisting of the functions of correct type.

**Fact 2.2.**  $[\omega_1]^{\omega_1} \approx [\omega_1]_*^{\omega_1}$ .

*Proof.* Let  $A = \{\omega \cdot (\alpha + 1) : \alpha \in \omega_1\}$ . Suppose  $f \in [A]^{\omega_1}$ . Let  $F'(\alpha)$  be the unique  $\beta$  so that  $f(\alpha) = \omega \cdot (\beta + 1)$ . Let  $F : \omega_1 \times \omega \rightarrow \omega_1$  be defined by  $F(\alpha, n) = \omega \cdot F'(\alpha) + n$ . Note that for all  $\alpha$ ,  $f(\alpha) = \sup\{F(\alpha, n) : n \in \omega\}$ . Thus  $f$  has uniform cofinality  $\omega$ . For any  $\alpha$ , for any  $\beta < \alpha$ ,  $f(\beta) = \omega \cdot (F'(\beta) + 1) \leq \omega \cdot F'(\alpha) < \omega \cdot (F'(\alpha) + 1) = f(\alpha)$ . This shows that every  $f \in [A]^{\omega_1}$  is of the correct type. Clearly,  $[\omega_1]^{\omega_1} \approx [A]^{\omega_1}$ . Thus one has an injection of  $[\omega_1]^{\omega_1}$  into  $[\omega_1]_*^{\omega_1}$ . The inclusion map is an injection of  $[\omega_1]_*^{\omega_1}$  into  $[\omega_1]^{\omega_1}$ .  $\square$

**Definition 2.3.** Let  $\epsilon \leq \omega_1$ . Write  $\omega_1 \rightarrow_* (\omega_1)_2^\epsilon$  to indicate that for all  $P : [\omega_1]_*^\epsilon \rightarrow 2$ , there is an  $i \in 2$  and a club  $C \subseteq \omega_1$  so that for all  $f \in [\omega_1]_*^\epsilon$ ,  $P(f) = i$ . In this case, one says that  $C$  is homogeneous for  $P$  taking value  $i$ .

**Fact 2.4.** (Martin; [8] Theorem 12.2, [3] Corollary 4.10 and 4.27) For all  $\epsilon \leq \omega_1$ ,  $\omega_1 \rightarrow_* (\omega_1)_2^\epsilon$ .

**Definition 2.5.** For  $\epsilon \leq \omega_1$ , let  $\mu_\epsilon$  denote the collection of  $X \subseteq [\omega_1]_*^\epsilon$  so that there exists a club  $C \subseteq \omega_1$  so that  $[C]_*^\epsilon \subseteq X$ . As a consequence of  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ ,  $\mu_\epsilon$  is a countably complete ultrafilter on  $[\omega_1]_*^\epsilon$  for all  $\epsilon \leq \omega_1$ .

**Definition 2.6.** Let  $\pi : \omega \times \omega \rightarrow \omega$  be a bijection. If  $R \subseteq \omega \times \omega$ , then  $x \in {}^\omega\omega$  codes  $R$  if and only if  $(a, b) \in R \Leftrightarrow x(\pi(a, b)) = 0$ . This gives a coding of binary relations on  $\omega$  by elements of  $\mathbb{R}$ . For each  $x \in \mathbb{R}$ ,  $\text{field}(x)$  is the set of  $n$  so that there exists some  $m$  such that  $x(\pi(m, n)) = 0$  or  $x(\pi(n, m)) = 0$ .

Let  $\text{WO}$  denote the set of reals coding wellorderings on subsets of  $\omega$ . If  $w \in \text{WO}$ , then  $<_w$  refers to the wellordering on  $\text{field}(w)$  coded by  $w$ . For each  $w \in \text{WO}$  and  $\alpha < \text{ot}(w)$ , let  $n_\alpha^w$  be the element of  $\text{field}(w)$  which has rank  $\alpha$  according to  $<_w$ . For each  $\alpha < \omega_1$ , let  $\text{WO}_\alpha = \{w \in \text{WO} : \text{ot}(w) = \alpha\}$ . Similarly, one can define  $\text{WO}_{<\alpha}$ ,  $\text{WO}_{\leq\alpha}$ ,  $\text{WO}_{>\alpha}$ , and  $\text{WO}_{\geq\alpha}$ . Note that  $\text{WO}$  is  $\Pi_1^1$ . For each  $\alpha < \omega_1$ ,  $\text{WO}_{>\alpha}$  and  $\text{WO}_{\geq\alpha}$  are  $\Pi_1^1$ ; and  $\text{WO}_\alpha$ ,  $\text{WO}_{<\alpha}$ , and  $\text{WO}_{\leq\alpha}$  are  $\Delta_1^1$ .

**Fact 2.7.** ([3] Fact 4.3) Suppose  $\tau$  is a Player 2 strategy with the property that for all  $x \in \text{WO}$ ,  $\tau(x) \in \text{WO}$  and  $\text{ot}(\tau(x)) > \text{ot}(x)$ . Let  $C_\tau = \{\eta < \omega_1 : (\forall w)(w \in \text{WO}_{<\eta} \Rightarrow \tau(w) \in \text{WO}_{<\eta})\}$ . Then  $C_\tau$  is a club.

**Definition 2.8.** Let  $\text{clubcode}_{\omega_1}$  denote the collection  $\tau \in \mathbb{R}$  so that  $\tau$  is a Player 2 strategy with the property that for all  $w \in \text{WO}$ ,  $\tau(w) \in \text{WO}$  and  $\text{ot}(\tau(w)) > \text{ot}(w)$ . Note that  $\text{clubcode}_{\omega_1}$  is a  $\Pi_2^1$  set.

**Fact 2.9.** (Solovay, [3] Fact 4.6) Suppose  $C$  is a club. There is a  $\tau \in \text{clubcode}_{\omega_1}$  so that  $C_\tau \subseteq C$ .

**Fact 2.10.** ([3] Fact 4.7) Suppose  $A \subseteq \text{clubcode}_{\omega_1}$  is  $\Sigma_1^1$ . Then one can find a club  $C$  uniformly in  $A$  (as a set; e.g. not depending on any  $\Sigma_1^1$  representation of  $A$ ) so that for all  $\tau \in A$ ,  $C \subseteq C_\tau$ .

**Definition 2.11.** A function  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is continuous if and only if for all  $f \in [\omega_1]_*^{\omega_1}$ , there is some  $\alpha < \omega_1$  so that for all  $g \in [\omega_1]_*^{\omega_1}$ , if  $g \restriction \alpha = f \restriction \alpha$ , then  $\Phi(g) = \Phi(f)$ .

If one gives  $[\omega_1]_*^{\omega_1}$  the topology generated by  $N_s = \{f \in [\omega_1]_*^{\omega_1} : s \subset f\}$  for each  $s \in [\omega_1]_*^{<\omega_1}$ , then  $\Phi$  is continuous in the above sense if and only if it is continuous in the topological sense with the range  $\omega_1$  given the discrete topology.

$\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is continuous almost everywhere if and only if there is a club  $C \subseteq \omega_1$  so that  $\Phi$  is continuous on  $[C]_*^{\omega_1}$ ; that is, for all  $f \in [C]_*^{\omega_1}$ , there exists an  $\alpha$  so that for all  $g \in [C]_*^{\omega_1}$  with  $g \restriction \alpha = f \restriction \alpha$ ,  $\Phi(g) = \Phi(f)$ .

**Definition 2.12.** Let  $\text{BS}$  denote a coding of bounded sequences in  $\omega_1$  by reals defined as follows.  $\text{BS}$  is the collection of  $(x, y) \in \mathbb{R}^2$  so that

- (i)  $x \in \text{WO}$ .
- (ii) For all  $n \in \text{field}(x)$ ,  $y_n \in \text{WO}$ .
- (iii) For all  $m, n \in \text{field}(x)$ ,  $m <_x n$  if and only if  $\text{ot}(y_m) < \text{ot}(y_n)$ .

Note that  $\text{BS}$  is  $\Pi_1^1$ . For each  $(x, y) \in \text{BS}$ , let  $\sigma_{(x, y)} : \text{ot}(x) \rightarrow \omega_1$  be defined by  $\sigma_{(x, y)}(\alpha) = \text{ot}(y_{n_\alpha^x})$ . Observe that for every  $\sigma \in [\omega_1]^{<\omega_1}$ , there is some  $(x, y) \in \text{BS}$  so that  $\sigma_{(x, y)} = \sigma$ .

**Definition 2.13.** Let  $\kappa$  be a regular cardinal and  $\lambda \leq \kappa$  be an ordinal. A good coding system for  ${}^\lambda\kappa$  consists of  $\Gamma$ ,  $\text{decode}$ , and  $\text{GC}_{\beta, \gamma}$  for each  $\beta < \lambda$  and  $\gamma < \kappa$  with the following properties:

- (1)  $\Gamma$  is a (boldface) pointclass closed under  $\exists^\mathbb{R}$ . Let  $\check{\Gamma}$  denote the dual pointclass. Let  $\Delta = \Gamma \cap \check{\Gamma}$ .
- (2)  $\text{decode} : \mathbb{R} \rightarrow \mathcal{P}(\lambda \times \kappa)$ . For all  $f \in {}^\lambda\kappa$ , there is some  $x \in \mathbb{R}$  so that  $\text{decode}(x) = f$ .
- (3) For all  $\beta < \lambda$  and  $\gamma < \kappa$ ,  $\text{GC}_{\beta, \gamma} \subseteq \mathbb{R}$ ,  $\text{GC}_{\beta, \gamma} \in \Delta$ , and  $\text{GC}_{\beta, \gamma}$  has the property that  $x \in \text{GC}_{\beta, \gamma}$  if and only if

$$\text{decode}(x)(\beta, \gamma) \wedge (\forall \gamma' < \kappa)(\text{decode}(x)(\beta, \gamma') \Rightarrow \gamma = \gamma').$$

For each  $\beta < \lambda$ , let  $\text{GC}_\beta = \bigcup_{\gamma < \kappa} \text{GC}_{\beta, \gamma}$ .

- (4) (Boundedness property) Suppose  $A \in \exists^\mathbb{R}\Delta$  and  $A \subseteq \text{GC}_\beta$ , then there exists some  $\delta < \kappa$  so that  $A \subseteq \bigcup_{\gamma < \delta} \text{GC}_{\beta, \gamma}$ .
- (5)  $\Delta$  is closed under less than  $\kappa$  wellordered unions.

Let  $\text{GC} = \bigcap_{\beta < \lambda} \text{GC}_\beta$ . Note that if  $x \in \text{GC}$ , then  $\text{decode}(x)$  is the graph of a function in  ${}^\lambda\kappa$ . If  $x \in \text{GC}$ , then one will use function notation such as  $\text{decode}(x)(\beta) = \gamma$  to indicate  $(\beta, \gamma) \in \text{decode}(x)$ .

**Definition 2.14.** Suppose  $\kappa$  is a regular cardinal and  $\lambda$  is such that  $\omega \cdot \lambda \leq \kappa$ . Suppose  $f \in {}^{\omega \cdot \lambda} \kappa$ . Let  $\text{block} : {}^{\omega \cdot \lambda} \kappa \rightarrow {}^\lambda \kappa$  be defined by  $\text{block}(f)(\alpha) = \sup\{f(\omega \cdot \alpha + k) : k \in \omega\}$ . Thus  $\text{block}(f)$  is the function returning the supremum of each  $\omega$ -block.

Suppose  $f, g \in {}^{\omega \cdot \lambda} \kappa$ . Let  $\text{joint} : {}^{\omega \cdot \lambda} \kappa \times {}^{\omega \cdot \lambda} \kappa \rightarrow {}^\lambda \kappa$  be defined by

$$\text{joint}(f, g)(\alpha) = \sup\{f(\omega \cdot \alpha + k), g(\omega \cdot \alpha + k) : k \in \omega\} = \max\{\text{block}(f)(\alpha), \text{block}(g)(\alpha)\}.$$

**Theorem 2.15.** (Martin, [3] Theorem 3.7) Suppose  $\lambda, \kappa$  are ordinals such that  $\omega \cdot \lambda \leq \kappa$ . Suppose there is a good coding system  $(\Gamma, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \lambda, \gamma < \kappa)$  for  ${}^{\omega \cdot \lambda} \kappa$ . Then  $\kappa \rightarrow_* (\kappa)_2^\lambda$  holds.

**Theorem 2.16.** ([3] Theorem 3.8) (Almost everywhere uniformization on good codes) Let  $\kappa$  be a regular cardinal and  $\lambda \leq \kappa$ . Let  $(\Gamma, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \lambda, \gamma < \kappa)$  be a good coding system for  ${}^{\omega \cdot \lambda} \kappa$ . Let  $R \subseteq [\kappa]_*^\lambda \times \mathbb{R}$  be a relation.

Then there is a club  $C \subseteq \kappa$  and a Lipschitz continuous function  $F : \mathbb{R} \rightarrow \mathbb{R}$  so that for all  $x \in \text{GC}$  with  $\text{decode}(x) \in [C]_*^{\omega \cdot \lambda}$  and  $\text{block}(\text{decode}(x)) \in [C]_*^\lambda \cap \text{dom}(R)$ ,  $R(\text{block}(\text{decode}(x)), F(x))$ .

One application of Theorem 2.16 is that under certain conditions large fragments of functions can be absorbed into inner models containing all reals.

**Theorem 2.17.** ([3] Theorem 3.9) Let  $\kappa$  be a regular cardinal and  $\lambda \leq \kappa$ . Suppose  $(\Gamma, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \lambda, \gamma < \kappa)$  is a good coding system for  ${}^\lambda \kappa$ . Let  $M \models \text{AD}$  be an inner model containing all the reals and within  $M$ ,  $(\Gamma, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \lambda, \gamma < \kappa)$  is a good coding system.

Then for any  $\Phi : [\kappa]^\lambda \rightarrow \kappa$ , there is a club  $D$ , necessarily in  $M$  by the coding lemma, so that  $\Phi \upharpoonright [D]_*^\lambda \in M$ .

**Definition 2.18.** Let  $\kappa$  be a regular cardinal and  $\lambda \leq \kappa$ .  $\kappa$  is  $\lambda$ -reasonable if and only if there is a good coding system for  ${}^\lambda \kappa$ .

**Theorem 2.19.** (Martin, [3] Fact 4.9, Theorem 4.18, and Theorem 4.26) For any  $\lambda \leq \omega_1$ ,  $\omega_1$  is  $\lambda$ -reasonable.

*Remark 2.20.* One can check that for  $\lambda < \omega_1$ , the natural good coding system for  ${}^{\omega \cdot \lambda} \omega_1$  (given by BS as in Definition 2.12) has the property that for any  $f \in [\omega_1]_*^\lambda$ , the collection of  $x \in \text{GC}$  with  $\text{block}(\text{decode}(x)) = f$  is  $\Delta_1^1$ . See [3] Fact 4.12.

### 3. CLUB UNIFORMIZATION

**Definition 3.1.** Let  $\text{club}_{\omega_1}$  denote the collection of club subsets of  $\omega_1$ . Let  $R \subseteq [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$ . If  $\sigma \in [\omega_1]_*^{<\omega_1}$ , then let  $R_\sigma = \{C \in \text{club}_{\omega_1} : R(\sigma, C)\}$ . Let  $\text{dom}(R) = \{\sigma \in [\omega_1]_*^{<\omega_1} : R_\sigma \neq \emptyset\}$ . A function  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  is a uniformization for  $R$  if and only if for all  $\sigma \in \text{dom}(R)$ ,  $R(\sigma, F(\sigma))$ .  $R$  is  $\subseteq$ -downward closed if and only if for all  $\sigma \in [\omega_1]_*^{<\omega_1}$  and for all  $C \subseteq D$  with  $C, D \in \text{club}_{\omega_1}$ ,  $R(\sigma, D)$  implies  $R(\sigma, C)$ .

$[\omega_1]_*^{<\omega_1}$ -club uniformization is the statement that every  $R \subseteq [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed has a uniformization. Almost everywhere  $[\omega_1]_*^{<\omega_1}$ -club uniformization is the statement that for every  $R \subseteq [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed, there is a club  $C \subseteq \omega_1$  so that the relation  $R \cap ([C]_*^{<\omega_1} \times \text{club}_{\omega_1})$  has a uniformization.

The primary purpose of this section is to establish the almost everywhere  $[\omega_1]_*^{<\omega_1}$ -club uniformization, which will be applied in the next section to establish continuity results. As a warmup, the following is a simple form of club uniformization.

**Definition 3.2.** Let  $\alpha < \omega_1$ . Let  $R \subseteq [\omega_1]_*^\alpha \times \text{club}_{\omega_1}$  be a  $\subseteq$ -downward closed relation. A uniformization for  $R$  is a function  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  so that for all  $\sigma \in \text{dom}(R)$ ,  $R(\sigma, F(\sigma))$ .  $[\omega_1]_*^\alpha$ -club uniformization is the statement that every  $R \subseteq [\omega_1]_*^\alpha \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed has a uniformization. Almost everywhere  $[\omega_1]_*^\alpha$ -club uniformization is the statement that for every  $R \subseteq [\omega_1]_*^\alpha \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed, there is a club  $C \subseteq \omega_1$  so that  $R \cap ([C]_*^\alpha \times \text{club}_{\omega_1})$  has a uniformization.

**Theorem 3.3.** Let  $\alpha < \omega_1$ . Almost everywhere  $[\omega_1]_*^\alpha$ -club uniformization holds.

*Proof.* Let  $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \alpha, \gamma < \omega_1)$  be the natural good coding system for  ${}^{\omega \cdot \alpha} \omega_1$  which satisfies the property mentioned in Remark 2.20 or [3] Fact 4.12. Fix  $R \subseteq [\omega_1]_*^\alpha \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed. Let  $S \subseteq [\omega_1]_*^\alpha \times \text{clubcode}_{\omega_1}$  be defined by  $S(f, z)$  if and only if  $R(f, C_z)$ . By Theorem 2.16, there is

a Lipschitz continuous function  $F : \mathbb{R} \rightarrow \mathbb{R}$  and a club  $C$  so that for all  $x \in \text{GC}$  with  $\text{decode}(x) \in [C]^{\omega \cdot \alpha}$  and  $\text{block}(\text{decode}(x)) \in \text{dom}(S)$ ,  $S(\text{block}(\text{decode}(x)), F(x))$ . Let  $D$  be the set of limit points of  $C$ . Remark 2.20 implies that for all  $f \in \text{dom}(R) \cap [D]_*^\alpha$ , the set  $K_f = \{x \in \mathbb{R} : \text{decode}(x) \in [C]_*^{\omega \cdot \alpha} \wedge \text{block}(\text{decode}(x)) = f\}$  is  $\Delta_1^1$ . Thus  $F[K_f]$  is a  $\Sigma_1^1$  subset of  $\text{clubcode}_{\omega_1}$ . By Fact 2.10, there is a club  $C_f$  obtained uniformly from  $F[K_f]$  (and hence obtained uniformly from  $f$ ) so that  $C_f \subseteq C_z$  for all  $z \in F[K_f]$ . Since for any  $z \in F[K_f]$ ,  $R(f, C_z)$  and  $R$  is  $\subseteq$ -downward closed,  $R(f, C_f)$ . Thus the function mapping  $f$  to  $C_f$  defined by the procedure above is a uniformization for  $R$  on  $\text{dom}(R) \cap [D]_*^\alpha$ .  $\square$

Note that Theorem 3.3 uses only a boundedness principle. It does not use uniformization or any other consequences of scales. This is in contrast to the argument of Theorem 3.10 for the almost everywhere  $[\omega_1]_*^{<\omega_1}$ -club uniformization which seems to require the relevant relations to be Suslin or uniformizable.

**Definition 3.4.** Let  $\alpha < \omega_1$ , let  $s \in {}^{<\omega}\alpha$ . Let  $N_s^\alpha = \{f \in {}^\omega\alpha : s \subset f\}$ .  ${}^\omega\alpha$  is given the topology generated by  $N_s^\alpha$ , and  ${}^\omega\alpha$  is homeomorphic to  ${}^\omega\omega$ . The concepts of meagerness, comeagerness, and nonmeagerness can be defined as usual.

Under AD, a wellordered intersection of comeager subsets of  ${}^\omega\alpha$  is a comeager subset of  ${}^\omega\alpha$ . Note that the set  $\text{surj}_\alpha = \{f \in {}^\omega\alpha : f : \omega \rightarrow \alpha \text{ is a surjection}\}$  is a comeager subset of  ${}^\omega\alpha$ .

The following is the simple generic coding function for  $\omega_1$ .

**Fact 3.5.** *There is a function  $\mathfrak{G} : {}^\omega\omega_1 \rightarrow \text{WO}$  so that for any  $\alpha < \omega_1$ , if  $f \in \text{surj}_\alpha$ , then  $\text{ot}(\mathfrak{G}(f)) = \alpha$ .*

*Proof.* For any  $f \in {}^\omega\omega_1$ , let  $A_f = \{n \in \omega : (\forall m)(f(n) = f(m) \Rightarrow n \leq m)\}$ . Let  $\mathfrak{G}(f)$  code a binary relation with domain  $A_f$  by letting  $m <_{\mathfrak{G}(f)} n$  if and only if  $f(m) < f(n)$ . It is clear that  $\mathfrak{G}(f) \in \text{WO}$  and if  $f \in \text{surj}_\alpha$ , then  $\text{ot}(\mathfrak{G}(f)) = \alpha$ .  $\square$

**Fact 3.6.** *There is a function  $H : [\omega_1]^{<\omega_1} \times \text{WO} \rightarrow \text{BS}$  with the property that for all  $\sigma \in [\omega_1]^{<\omega_1}$  and for all  $w \in \text{WO}$  so that  $\text{ot}(w) = \sup(\sigma) + 2$ ,  $H(\sigma, w)$  codes  $\sigma$ , that is,  $\sigma_{H(\sigma, w)} = \sigma$  (in the notation of Definition 2.12).*

*Proof.* Fix  $\sigma \in [\omega_1]^{<\omega_1}$ . If  $w \in \text{WO}$  and  $\text{ot}(w) \neq \sup(\sigma) + 2$ , then let  $H(\sigma, w)$  be some fixed element of BS as this case is insignificant. Now assume  $\text{ot}(w) = \sup(\sigma) + 2$ . Observe that  $\text{length}(\sigma) < \sup(\sigma) + 2$ . Canonically from  $w$  and  $\sigma$ , one will produce  $(x, y) \in \text{BS}$  as follows: Let  $x \in \text{WO}$  (which is produced canonically from  $w$  and  $\sigma$ ) code a relation on  $\omega$  whose field is  $\{n \in \text{field}(w) : n <_w n_{\text{length}(\sigma)}^w\}$  (here the notation comes from Definition 2.6) and for  $m, n \in \text{field}(x)$ ,  $m <_x n$  if and only if  $m <_w n$ . Then  $\text{ot}(x) = \text{length}(\sigma)$ .

Similarly, produce  $y$  canonically from  $w$  and  $\sigma$  as follows: Fix a  $k \in \omega$ . If  $k \notin \text{field}(x)$ , then let  $y_k = \bar{0}$ , the constant 0 sequence. If  $k \in \text{field}(x)$ , then let  $\alpha < \text{length}(\sigma)$  so that  $k = n_\alpha^w$ . Let  $y_k$  be the unique real coding a binary relation such that  $\text{field}(y_k) = \{n \in \text{field}(w) : n <_w n_{\sigma(\alpha)}^w\}$  and for all  $m, n \in \text{field}(y_k)$ ,  $m <_{y_k} n \Leftrightarrow m <_w n$ . Then  $y_k \in \text{WO}$  and  $\text{ot}(y_k) = \sigma(\alpha)$ . Let  $y \in \mathbb{R}$  be such that for all  $k \in \omega$ , the  $k^{\text{th}}$  section of  $y$  is  $y_k$ . Thus  $(x, y) \in \text{BS}$  and  $\sigma_{(x, y)} = \sigma$ . Let  $H(\sigma, w) = (x, y)$ .  $\square$

**Theorem 3.7.** *Assume ZF and all sets of reals have the Baire property. Let  $R \subseteq [\omega_1]^{<\omega_1} \times \text{club}_{\omega_1}$  be a  $\subseteq$ -downward closed relation. Define  $\tilde{R} \subseteq \mathbb{R} \times \mathbb{R} \times \mathbb{R}$  (which is the coded version of  $R$ ) by*

$$\tilde{R}(x, y, z) \Leftrightarrow (x, y) \in \text{BS} \wedge z \in \text{clubcode}_{\omega_1} \wedge R(\sigma_{(x, y)}, C_z).$$

*Consider  $\tilde{R}$  as a binary relation on  $\text{BS} \times \text{clubcode}_{\omega_1}$ . Suppose there is a  $J : \text{dom}(\tilde{R}) \rightarrow \text{clubcode}_{\omega_1}$  which is a uniformization for  $\tilde{R}$ . Then there is an  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  which is a uniformization for  $R$ .*

*Thus ZF + AD $_{\mathbb{R}}$  proves  $[\omega_1]^{<\omega_1}$ -club uniformization.*

*Proof.* Fix  $\sigma \in \text{dom}(R)$ . The club  $F(\sigma)$  will be defined in the following:

Recall  $\mathfrak{G}$  is the simple generic coding function from Fact 3.5 and  $H$  is the function from Fact 3.6. Let  $A = \text{surj}_{\sup(\sigma)+2}$ . As observed earlier,  $A$  is comeager as a subset of  ${}^\omega(\sup(\sigma) + 2)$ . Let  $O : A \rightarrow \text{BS}$  be defined by  $O(f) = J(H(\sigma, \mathfrak{G}(f)))$ . Note that  $\mathfrak{G}(f) \in \text{WO}_{\sup(\sigma)+2}$  for all  $f \in A$ . Therefore for all  $f \in A$ ,  $H(\sigma, \mathfrak{G}(f)) \in \text{BS}$  and codes  $\sigma$ . Since  $J$  is a uniformization for  $\tilde{R}$ ,  $\tilde{R}(H(\sigma, \mathfrak{G}(f)), O(f))$  holds for all  $f \in A$ . The key observation is that for any  $f \in A$ ,  $O(f) \in \text{clubcode}_{\omega_1}$  and  $R(\sigma, C_{O(f)})$ .

For any club  $C$ , let  $\text{enum}_C : \omega_1 \rightarrow C$  be the increasing enumeration of  $C$ . For each  $\gamma < \omega_1$ , let  $K(\gamma)$  be the least  $\delta < \omega_1$  so that for comeagerly many  $f \in A$  (in the topological space  ${}^\omega(\sup(\sigma) + 2)$ ),  $\text{enum}_{C_{O(f)}}(\gamma) < \delta$ .

Claim 1:  $K$  is a well-defined function.

To see this: Fix  $\gamma$ . For each  $\epsilon < \omega_1$ , let  $T_\epsilon = \{f \in A : \text{enum}_{C_{O(f)}}(\gamma) = \epsilon\}$  and  $T_{\geq \epsilon} = \{f \in A : \text{enum}_{C_{O(f)}}(\gamma) \geq \epsilon\}$ . For the sake of contradiction, suppose  $K(\gamma)$  is not defined. Then for all  $\epsilon < \omega_1$ ,  $T_{\geq \epsilon}$  is nonmeager. Since  $A = T_{\geq 0} = \bigcup_{\epsilon \geq 0} T_\epsilon$  is comeager and wellordered unions of meager sets are meager (by the Baire property for all sets of reals), there exists some  $\epsilon < \omega_1$  so that  $T_\epsilon$  is nonmeager. Let  $\epsilon_0$  be the least  $\epsilon$  so that  $T_\epsilon$  is nonmeager. Suppose  $\beta < \omega_1$  and  $\epsilon_\alpha$  has been defined for all  $\alpha < \beta$ . Let  $\zeta = \sup\{\epsilon_\alpha : \alpha < \beta\}$ . Since by assumption  $T_{\geq \zeta} = \bigcup_{\epsilon \geq \zeta} T_\epsilon$  is nonmeager and wellordered unions of meager sets are meager, there is an  $\epsilon \geq \zeta$  so that  $T_\epsilon$  is nonmeager. Let  $\epsilon_\beta$  be the least  $\epsilon \geq \zeta$  so that  $T_\epsilon$  is nonmeager. This defines a sequence  $\langle \epsilon_\beta : \beta < \omega_1 \rangle$  so that  $\langle T_{\epsilon_\beta} : \beta < \omega_1 \rangle$  is an uncountable collection of disjoint nonmeager sets. Since all sets of reals have the Baire property, this violates the countable chain condition of the topology on  ${}^\omega(\sup(\sigma) + 2)$ . This completes the proof of Claim 1.

Let  $D = \{\eta < \omega_1 : (\forall \nu < \eta)(K(\nu) < \eta)\}$ . Note that since for any club  $C \subseteq \omega_1$ ,  $\text{enum}_C(\gamma) \geq \gamma$ , one can conclude that  $K(\gamma) > \gamma$ . Also if  $\gamma \leq \gamma'$ ,  $K(\gamma) \leq K(\gamma')$ . Let  $\epsilon < \omega_1$ . Let  $\alpha_0 = \epsilon$ . Let  $\alpha_{n+1} = K(\alpha_n)$ . Hence  $\alpha_{n+1} > \alpha_n$ . Let  $\alpha = \sup\{\alpha_n : n \in \omega\}$ . Note that for all  $\nu < \alpha$ ,  $\nu < \alpha_n$  for some  $n$ . Then one has  $K(\nu) \leq K(\alpha_n) = \alpha_{n+1} < \alpha$ . Thus  $\alpha \in D$  and  $\epsilon < \alpha$ . This shows that  $D$  is unbounded.  $D$  is clearly closed.

Claim 2:  $R(\sigma, D)$ .

To see this: Let  $\eta \in D$ . For each  $\beta < \eta$ , let  $F_\beta^\eta = \{f \in A : \text{enum}_{C_{O(f)}}(\beta) < \eta\}$ . Since  $\eta \in D$ , for all  $\beta < \eta$ ,  $K(\beta) < \eta$ . So the set of  $f \in A$  so that  $\text{enum}_{C_{O(f)}}(\beta) < K(\beta) < \eta$  is comeager, i.e.  $F_\beta^\eta$  is comeager. Then  $Y^\eta = \bigcap_{\beta < \eta} F_\beta^\eta$  is a comeager set. For all  $f \in Y^\eta$ , for all  $\beta < \eta$ ,  $\beta \leq \text{enum}_{C_{O(f)}}(\beta) < \eta$ . Since  $C_{O(f)}$  is a club,  $\eta \in C_{O(f)}$ . It has been shown that if  $\eta \in D$ , then  $Y^\eta$  has the property that for all  $f \in Y^\eta$ ,  $\eta \in C_{O(f)}$ . Let  $Y = \bigcap_{\eta \in D} Y^\eta$ . Since wellordered intersection of comeager sets are comeager,  $Y$  is comeager. Pick an  $f \in Y$ . For any  $\eta \in D$ ,  $f \in Y^\eta$ . So  $\eta \in C_{O(f)}$ . Thus  $D \subseteq C_{O(f)}$ . Since  $R(\sigma, C_{O(f)})$  holds and  $R$  is  $\subseteq$ -downward closed,  $R(\sigma, D)$ . This completes the proof of Claim 2.

Note that  $D$  was produced uniformly from  $\sigma$  by the procedure above. So finally, let  $F(\sigma) = D$ . This defines  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$ .  $F$  is a uniformization for  $R$ .

Now assume  $\text{AD}_{\mathbb{R}}$ . This implies  $\text{AD}$  and hence all set of reals has the Baire property. Moreover, the uniformization  $J$  for  $\tilde{R}$  exists since  $\text{AD}_{\mathbb{R}}$  proves uniformization for all relations on  $\mathbb{R} \times \mathbb{R}$ . Club uniformization follows from the first part of the theorem.  $\square$

For  $\alpha < \omega_1$ , let  $\text{BS}_\alpha$  be the subset of  $\text{BS}$  coding elements of  $[\omega_1]^\alpha$ .

**Corollary 3.8.** *Assume ZF and all sets of reals have the Baire property. Let  $\alpha < \omega_1$ . Let  $R \subseteq [\omega_1]_*^\alpha \times \text{club}_{\omega_1}$  be a  $\subseteq$ -closed relation. Define  $\tilde{R} \subseteq \mathbb{R} \times \mathbb{R} \times \mathbb{R}$  by*

$$\tilde{R}(x, y, z) \Leftrightarrow (x, y) \in \text{BS}_\alpha \wedge z \in \text{clubcode}_{\omega_1} \wedge R(\sigma_{(x,y)}, C_z).$$

*Consider  $\tilde{R}$  as a relation on  $\text{BS}_\alpha \times \text{clubcode}_{\omega_1}$ . Suppose there is a  $J : \text{dom}(\tilde{R}) \rightarrow \text{clubcode}_{\omega_1}$  which is a uniformization for  $\tilde{R}$ . Then there is an  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  which is a uniformization for  $R$ .*

*Thus  $\text{AD}_{\mathbb{R}}$  proves  $[\omega_1]^\alpha$ -club uniformization, for all  $\alpha < \omega_1$ .*

**Fact 3.9.** *Assume ZF + AD. Then  $L(\mathbb{R}) \models \text{AD}$  (and even  $\text{AD}^+$ ) and  $L(\mathbb{R})$  does not satisfy  $[\omega_1]^\alpha$ -club uniformization (when  $\omega \leq \alpha < \omega_1$ ) or  $[\omega_1]^{<\omega_1}$ -club uniformization.*

*Proof.* Work in  $L(\mathbb{R})$ . Consider the relation  $S \subseteq \mathbb{R} \times \text{club}_{\omega_1}$  defined by  $S(x, C)$  if and only if for all club  $D \subseteq C$ ,  $D \notin \text{HOD}_x$ .

First, one will show that  $\text{dom}(S) = \mathbb{R}$ . Fix an  $x \in \mathbb{R}$ . Since  $\omega_2$  is measurable in  $V$ , every wellordered sequence of elements of  $\mathcal{P}(\omega_1)$  has length less than  $\omega_2$ . Thus  $(\mathcal{P}(\omega_1))^{\text{HOD}_x}$  has cardinality less than  $\omega_2$  in  $L(\mathbb{R})$ . Let  $\langle C_\alpha : \alpha < \omega_1 \rangle$  be an enumeration of all club subsets of  $\omega_1$  which belong to  $\text{HOD}_x$ . (This enumeration does not belong to  $\text{HOD}_x$ .) One will construct a club  $E \subseteq \omega_1$  as follows. Let  $E_0 = \emptyset$ . If  $\gamma$  is a limit ordinal and  $E_\nu$  has been defined for all  $\nu < \gamma$ , then let  $E_\gamma$  be the closure of  $\bigcup_{\nu < \gamma} E_\nu$ . Now suppose  $\gamma$  is an ordinal so that  $\langle \alpha_\nu : \nu < \gamma \rangle$  and the closed set  $E_\gamma$  has been defined with  $\alpha_\nu \notin E_\gamma$  for any  $\nu < \gamma$ . Let  $\alpha_\gamma$  be least element of  $C_\gamma$  greater than  $\sup E_\gamma$ . Let  $E_{\gamma+1} = E_\gamma \cup \{\alpha_\gamma + 1\}$ . Note that  $\alpha_\gamma \notin E_{\gamma+1}$ .

In the end, one has constructed a sequence  $\langle \alpha_\gamma : \gamma < \omega_1 \rangle$  and a sequence  $\langle E_\gamma : \gamma < \omega_1 \rangle$  so that for all  $\gamma < \omega_1$ ,  $\alpha_\gamma \in C_\gamma$  and for all  $\nu < \gamma < \omega_1$ ,  $\alpha_\nu \notin E_\gamma$ . Let  $E = \bigcup_{\gamma < \omega_1} E_\gamma$ . One can check that  $E$  is club and  $\alpha_\gamma \notin E$  for any  $\gamma < \omega_1$ . Now suppose there was a club  $D$  such that  $D \subseteq E$  and  $D \in \text{HOD}_x$ . Then there



is some  $\gamma < \omega_1$  so that  $D = C_\gamma$ . But  $\alpha_\gamma \notin D$  since  $\alpha_\gamma \notin E$ . But  $\alpha_\gamma \in C_\gamma$ . Hence  $D \neq C_\gamma$  which is a contradiction. This shows that  $E \subseteq \omega_1$  is a club subset with the property that  $E$  has no club subsets that belong to  $\text{HOD}_x$ . That is,  $S(x, E)$ . Since  $x \in \mathbb{R}$  was arbitrary, it has been shown that  $\text{dom}(S) = \mathbb{R}$ .

Observe that  $S$  is  $\subseteq$ -downward closed in the sense that for all  $x \in \mathbb{R}$ ,  $S(x, C)$  and  $D \subseteq C$ , then  $S(x, D)$ .

Suppose there is a function  $F : \mathbb{R} \rightarrow \text{club}_{\omega_1}$  so that  $F$  uniformizes  $S$ . In  $L(\mathbb{R})$ ,  $F$  is  $\text{OD}_z$  for some  $z \in \mathbb{R}$ . Since  $F$  is a uniformization,  $S(z, F(z))$ . Therefore  $F(z)$  is a club subset of  $\omega_1$  which is  $\text{OD}_z$  and thus  $F(z) \in \text{HOD}_z$ . This contradicts the definition of  $S$ .

Considering  $\mathbb{R}$  as  $[\omega]^\omega$ , the set of increasing sequences in  $\omega$ , define  $R \subseteq [\omega_1]^\omega \times \text{club}_{\omega_1}$  by

$$R(x, C) \Leftrightarrow (x \in \mathbb{R} \wedge S(x, C)) \vee (x \notin [\omega]^\omega).$$

$R$  can not be uniformized or else  $S$  could be uniformized. This shows  $[\omega_1]^\omega$ -club uniformization fails. Similar examples give the failure of  $[\omega_1]^\alpha$ -club uniformization for all  $\omega \leq \alpha < \omega_1$  and a failure of  $[\omega_1]^{<\omega_1}$ -club uniformization.  $\square$

Thus almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization is the best one can expect in AD alone. This is verified by the following result.

**Theorem 3.10.** *Almost everywhere  $[\omega_1]^{<\omega_1}$ -club uniformization holds: Let  $R \subseteq [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$  which is  $\subseteq$ -downward closed. Then there is a club  $D \subseteq \omega_1$  so that  $R \cap ([D]_*^{<\omega_1} \times \text{club}_{\omega_1})$  has a uniformization.*

*Proof.* Suppose  $D \subseteq \omega_1$  is a club. Let  $\text{BS}^D$  denote the subset of  $\text{BS}$  which code elements of  $[D]_*^{<\omega_1}$ . If one can find a club  $D \subset \omega_1$  so that  $\tilde{R} \cap (\text{BS}^D \times \text{clubcode}_{\omega_1})$  has a uniformization, then Theorem 3.7 would give the conclusion of this theorem.

Fix a  $\Sigma_2^1$  set  $U \subseteq \mathbb{R}^3$  which is universal for  $\Sigma_2^1$  subsets of  $\mathbb{R}^2$ . Take any  $f \in [\omega_1]^{\omega_1}$ . Let  $T_f \subseteq \text{WO} \times \text{clubcode}_{\omega_1}$  be defined by  $T_f(w, z)$  if and only if

$$f \restriction \text{ot}(w) \in \text{dom}(R) \wedge z \in \text{clubcode}_{\omega_1} \wedge R(f \restriction \text{ot}(w), C_z).$$

That is,  $T_f$  is a relation which attempts to collect reals coding clubs that are associated to all the initial segments of  $f$  according to  $R$ . By the coding lemma applied to the pointclass  $\Sigma_2^1$  and the usual prewellordering on  $\text{WO}$ , there is some  $e$  so that

(1)  $U_e \subseteq T_f$ .

(2) For all  $w \in \text{WO}$ ,  $(T_f)_w \neq \emptyset$  if and only if  $U_{e,w} \neq \emptyset$ .

(Note that  $(T_f)_w = \{c \in \mathbb{R} : T_f(w, c)\}$ . Recall that  $U \subseteq \mathbb{R}^3$  and  $U_{a,b} = \{c \in \mathbb{R} : U(a, b, c)\}$ .) Say that  $e \in \mathbb{R}$  is an  $f$ -selector if and only if (1) and (2) holds for  $e$  and  $f$ .

Fix a good coding system  $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega_1, \gamma < \omega_1)$  for  ${}^{\omega \cdot \omega_1}\omega_1$ . Consider the relation,  $S \subseteq [\omega_1]_*^{\omega_1} \times \mathbb{R}$  defined by  $S(f, e)$  if and only if  $e$  is an  $f$ -selector. By Theorem 2.16, let  $F$  be a Lipschitz function and  $E \subseteq \omega_1$  be a club such that for all  $x \in \text{GC}$  with  $\text{decode}(x) \in [E]^{\omega \cdot \omega_1}$  and  $\text{block}(\text{decode}(x)) \in \text{dom}(S)$ ,  $S(\text{block}(\text{decode}(x)), F(x))$ . By Fact 2.9, let  $z^* \in \text{clubcode}_{\omega_1}$  be such that  $C_{z^*} \subseteq E$ . Let  $D$  be the limit points of  $C_{z^*}$ .

Now consider the relation  $K \subseteq \text{BS} \times \mathbb{R}$  by  $K((x, y), r)$  if and only if the conjunction of the two holds

(1)  $\sigma_{(x, y)} \in [D]_*^{<\omega_1}$ . (That is,  $(x, y) \in \text{BS}^D$ .)

(2)  $r \in \text{GC}$ ,  $\text{decode}(r) \in [C_{z^*}]^{\omega \cdot \omega_1}$ , and  $\sigma_{(x, y)} \subseteq \text{block}(\text{decode}(r))$ .

That is,  $K((x, y), r)$  holds if  $(x, y)$  is a code for a function of length less than  $\omega_1$  of the correct type through  $D$  (which is the set of limit points of  $C_{z^*}$ ) and  $r$  is a code (according to the good coding system) for a full  $\omega_1 = \omega \cdot \omega_1$  length function with the property that  $\sigma_{(x, y)}$  is an initial segment of  $\text{block}(\text{decode}(r))$ .

One can check that  $K$  is projective using  $z^*$  as a parameter. Hence let  $G : \mathbb{R} \rightarrow \mathbb{R}$  be a projective uniformization for this relation. Thus if  $(x, y) \in \text{BS}^D$  is such that  $\sigma_{(x, y)}$  is a bounded function of the correct type, then  $\text{decode}(G(x, y)) \in [C_{z^*}]_*^{\omega \cdot \omega_1}$ , and  $\text{block}(\text{decode}(G(x, y)))$  is an extension of  $\sigma_{(x, y)}$  to a full sequence.

Define  $\tilde{Y} \subseteq \text{BS}^D \times \text{clubcode}_{\omega_1}$  by

$$\tilde{Y}((x, y), v) \Leftrightarrow (x, y) \in \text{BS}^D \wedge v \in U_{F(G(x, y)), x}$$

Note that  $\tilde{Y}$  is projective since  $D$  is the limit points of  $C_{z^*}$ ,  $U$  is  $\Sigma_2^1$ ,  $F$  is a Lipschitz function, and  $G$  is a projective function. Whenever  $(x, y) \in \text{BS}^D$  and  $\sigma_{(x, y)}$  codes a sequence of the correct type of length less than  $\omega_1$  through  $D$ ,  $G(x, y) \in \text{GC}$  is a code for a full function passing through  $C_{z^*}$  so that  $\text{block}(\text{decode}(G(x, y)))$

extends  $\sigma_{(x,y)}$ . Recall that  $x$  is the length of  $\sigma_{(x,y)}$ . Thus  $T_{\text{block}(\text{decode}(G(x,y))),x} = \tilde{R}_{(x,y)}$ . By the property of  $F$ ,  $F(G(x,y))$  is then a  $\text{block}(\text{decode}(G(x,y)))$ -selector. So for all such  $(x,y)$ ,  $U_{F(G(x,y)),x} \subseteq T_{\text{block}(\text{decode}(G(x,y))),x} = \tilde{R}_{(x,y)}$  and  $U_{F(G(x,y)),x} \neq \emptyset$  if and only if  $\tilde{R}_{(x,y)} = T_{\text{block}(\text{decode}(G(x,y))),x} \neq \emptyset$ . Hence  $\tilde{Y} \subseteq \tilde{R} \cap (\text{BS}^D \times \text{clubcode}_{\omega_1})$  and any uniformization for  $\tilde{Y}$  is a uniformization for  $\tilde{R} \cap (\text{BS}^D \times \text{clubcode}_{\omega_1})$ .

However,  $\tilde{Y}$  does have a uniformization since it is projective. Thus  $\tilde{R} \cap (\text{BS}^D \times \text{clubcode}_{\omega_1})$  has a uniformization. By the remarks at the beginning of the proof, this suffices to complete the argument.  $\square$

**Theorem 3.11.** *Assume ZF and all sets of reals have the Baire property. Let  $R \subseteq \omega_1 \times [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$  be a  $\subseteq$ -downward closed relation (on the  $\text{club}_{\omega_1}$ -coordinate). Define  $\tilde{R} \subseteq \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}$  by*

$$\tilde{R}(w, x, y, z) \Leftrightarrow w \in \text{WO} \wedge (x, y) \in \text{BS} \wedge z \in \text{clubcode}_{\omega_1} \wedge R(\text{ot}(w), \sigma_{(x,y)}, C_z).$$

*Consider  $\tilde{R}$  as a relation on  $(\text{WO} \times \text{BS}) \times \text{clubcode}_{\omega_1}$ . Suppose there is a  $J : \text{dom}(\tilde{R}) \rightarrow \text{clubcode}_{\omega_1}$  which is a uniformization for  $\tilde{R}$ . Then there is an  $F : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$  which is a uniformization for  $R$ .*

*Thus under  $\text{AD}_{\mathbb{R}}$ , such relations have a uniformization.*

*Let  $R \subseteq \omega_1 \times [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$  be a  $\subseteq$ -downward closed relation as above. Then there is a club  $D \subseteq \omega_1$  so that  $R \cap (\omega_1 \times [D]_*^{<\omega_1} \times \text{club}_{\omega_1})$  has a uniformization.*

*Proof.* This requires some small modifications in the arguments for Theorem 3.7 and Theorem 3.10.

By an argument similar to Fact 3.6, there is a function  $H : \omega_1 \times [\omega_1]_*^{<\omega_1} \times \text{WO} \rightarrow \text{WO} \times \text{BS}$  with the property that for all  $\alpha < \omega_1$ ,  $\sigma \in [\omega_1]_*^{<\omega_1}$ , and all  $w \in \text{WO}$  so that  $\text{ot}(w) = \max\{\sup(\sigma) + 2, \alpha + 1\}$ , one has that  $H(\alpha, \sigma, w) \in \text{WO} \times \text{BS}$  with the property that  $\text{ot}(\pi_1(H(\alpha, \sigma, w))) = \alpha$  and  $\pi_2(H(\alpha, \sigma, w))$  codes  $\sigma$ , where  $\pi_1, \pi_2 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  are the projection maps onto the first and second coordinate, respectively. Using  $H$ , one can prove the first part by making the necessary modifications to the argument of Theorem 3.7.

For the second part, let  $\rho : \omega_1 \rightarrow \omega_1 \times \omega_1$  be a bijection. Let  $\varpi_1, \varpi_2 : \omega_1 \times \omega_1 \rightarrow \omega_1$  be the projection onto the first and second coordinate, respectively. Define a relation  $T_f \subseteq \text{WO} \times \text{clubcode}_{\omega_1}$  by  $T_f(w, z)$  if and only if

$$(\varpi_1(\rho(\text{ot}(w))), f \upharpoonright \varpi_2(\rho(\text{ot}(w)))) \in \text{dom}(R) \wedge z \in \text{clubcode}_{\omega_1} \wedge R(\varpi_1(\rho(\text{ot}(w))), f \upharpoonright \varpi_2(\rho(\text{ot}(w))), z).$$

With this version of the relation  $T_f$ , one can prove the second statement with a modification of the argument in Theorem 3.10.  $\square$

#### 4. CONTINUITY OF FUNCTIONS $[\omega_1]^{\omega_1} \rightarrow \omega_1$

**Lemma 4.1.** *Suppose  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  has the property that there is a club  $C \subseteq \omega_1$  so that for all  $f \in [C]_*^{\omega_1}$ ,  $\Phi(f) < f(0)$ . Then there is a club  $D \subseteq \omega_1$  and a  $\zeta < \omega_1$  so that for all  $f \in [D]_*^{\omega_1}$ ,  $\Phi(f) = \zeta$ .*

*Proof.* Define a partition  $P : [\omega_1]_*^{\omega_1} \rightarrow 2$  by  $P(\alpha \hat{f}) = 0$  if and only if  $\Phi(f) < \alpha$ . By the strong partition property  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , there is a club  $E \subseteq \omega_1$  which is homogeneous for  $P$ . Let  $\tilde{E} = \{\alpha \in E : \text{enum}_E(\alpha) = \alpha\}$  where  $\text{enum}_E : \omega_1 \rightarrow E$  is the increasing enumeration of  $E$ .  $\tilde{E} \subseteq E$  is also a club subset of  $\omega_1$ . Let  $f \in [\tilde{E} \cap C]_*^{\omega_1}$ . Then  $\Phi(f) < f(0)$  by the assumption on  $C$ . Since  $f$  is a function of the correct type and  $f(0) \in \tilde{E}$ , one can find an  $\alpha \in E$  with  $\Phi(f) < \alpha < f(0)$ . Then  $\alpha \hat{f} \in [E]_*^{\omega_1}$  and  $P(\alpha \hat{f}) = 0$ . Since  $E$  is homogeneous for  $P$ , one must have that  $E$  is homogeneous for  $P$  taking value 0. Let  $E_0 = E \setminus (\min E + 1)$ . For all  $f \in [E_0]_*^{\omega_1}$ , one has that  $\Phi(f) < \min E$  since  $(\min E) \hat{f} \in [E]_*^{\omega_1}$  and  $P((\min E) \hat{f}) = 0$ . By the countable completeness of the strong partition measure on  $\omega_1$ , there is a club  $D \subseteq E_0$  and a  $\zeta < \min E$  so that for all  $f \in [D]_*^{\omega_1}$ ,  $\Phi(f) = \zeta$ .  $\square$

**Example 4.2.** The existence of a function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$  which is not continuous  $\mu_{\omega_1}$ -almost everywhere intuitively amounts to asking whether there is a way to define a map that truly uses all information about  $f$  and not merely an initial segment of  $f$ , for  $\mu_{\omega_1}$ -almost all  $f \in [\omega_1]_*^{\omega_1}$ .

One function that at first glance may appear to use the whole function  $f \in [\omega_1]^{\omega_1}$  is  $\Phi(f) = \omega_1^{L[f]}$ . However, almost everywhere  $\Phi$  uses no information about  $f$ . It is  $\mu_{\omega_1}$ -almost everywhere a constant function.

To see this: Let  $f \in [\omega_1]_*^{\omega_1}$ . For each  $\alpha < \omega_1$ , let  $f_\alpha \in [\omega_1]_*^{\omega_1}$  be defined by  $f_\alpha(\beta) = f(\alpha + \beta)$ . Note that for all  $\alpha < \beta < \omega_1$ ,  $f_\beta \in L[f_\alpha]$ . So  $\omega_1^{L[f_\beta]} \leq \omega_1^{L[f_\alpha]}$ . The sequence  $\langle \omega_1^{L[f_\alpha]} : \alpha < \omega_1 \rangle$  is a nonincreasing sequence of ordinals. It must be eventually constant else one would have an infinite decreasing sequence of ordinals. Let  $\epsilon_f$  be the eventual constant value of this sequence.

Let  $Q : [\omega_1]_*^{\omega_1} \rightarrow 2$  be defined by  $Q(f) = 0 \Leftrightarrow \omega_1^{L[f]} = \epsilon_f$ . By the strong partition property  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , there is some  $D \subseteq \omega_1$  club which is homogeneous for  $Q$ . Let  $f \in [D]_*^{\omega_1}$ . Let  $\alpha$  be minimal so that  $\omega_1^{L[f_\alpha]} = \epsilon_f$ . Note that  $\omega_1^{L[f_\alpha]} = \epsilon_{f_\alpha}$  and  $f_\alpha \in [D]_*^{\omega_1}$ . Thus  $Q(f_\alpha) = 0$ . Thus  $D$  is homogeneous for  $Q$  taking value 0. It has been shown that for all  $f \in [D]_*^{\omega_1}$ ,  $\omega_1^{L[f]} = \omega_1^{L[f_\alpha]}$  for all  $\alpha < \omega_1$ .

Consider  $P : [D]_*^{\omega_1} \rightarrow 2$  defined by  $P(f) = 0 \Leftrightarrow \Phi(f) = \omega_1^{L[f]} < f(0)$ . By  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , let  $C \subseteq D$  be a club such that  $C$  is homogeneous for  $P$ . Take any  $f \in [C]_*^{\omega_1}$ . Note that for all  $\alpha < \omega_1$ ,  $\Phi(f_\alpha) = \omega_1^{L[f_\alpha]} = \omega_1^{L[f]} = \Phi(f)$  since  $f \in [D]_*^{\omega_1}$ . Pick  $\delta$  so that  $f(\delta) > \omega_1^{L[f]}$ . Then  $\Phi(f_\delta) = \Phi(f) < f(\delta) = f_\delta(0)$ . Since  $f_\delta \in [C]_*^{\omega_1}$ , one has that  $C$  is homogeneous for  $P$  taking value 0.

So for all  $f \in [C]_*^{\omega_1}$ ,  $\Phi(f) < f(0)$ . By Lemma 4.1,  $\Phi$  is  $\mu_{\omega_1}$ -almost everywhere a constant function.

With Trang, it has been shown that the constant almost everywhere value of  $\Phi$  is quite large. It is in particular not  $\omega_1^L$ .

Claim 1: There is some  $\epsilon < \omega_1$  and some club  $C_0 \subseteq \omega_1$  so that for all  $f \in [C_0]_*^{\omega_1}$ , the canonical  $L[f]$  wellordering of  $\mathbb{R}^{L[f]}$  has ordertype  $\epsilon$ .

Suppose not. For all  $\alpha < \omega_1$ , there is a club  $C \subseteq \omega_1$  so that the canonical wellordering of  $\mathbb{R}^{L[f]}$  has length greater than  $\alpha$ . By the countable additivity of the strong partition measure, for each  $\alpha < \omega_1$ , there is a real  $r_\alpha$  so that for almost all  $f \in [\omega_1]_*^{\omega_1}$ ,  $r_\alpha$  is the  $\alpha^{\text{th}}$  real in the canonical wellordering. Suppose  $\alpha \neq \beta$ . There is a club  $C_\alpha$  and  $C_\beta$  so that for all  $f \in [C_\alpha]_*^{\omega_1}$  and  $g \in [C_\beta]_*^{\omega_1}$ , the  $\alpha^{\text{th}}$  real of  $L[f]$  is  $r_\alpha$  and the  $\beta^{\text{th}}$  real of  $L[g]$  is  $r_\beta$ . If  $f \in [C_\alpha \cap C_\beta]_*^{\omega_1}$ , then  $r_\alpha$  and  $r_\beta$  are the  $\alpha^{\text{th}}$  and  $\beta^{\text{th}}$  real of  $L[f]$ . Hence  $r_\alpha \neq r_\beta$ . This shows that  $\langle r_\alpha : \alpha < \omega_1 \rangle$  is an  $\omega_1$  sequence of distinct reals. This is impossible under AD. Claim 1 has been shown.

By a countable additivity argument, one can show that there is a club  $C_1$  so that for all  $f, g \in [C_1]_*^{\omega_1}$ ,  $\mathbb{R}^{L[f]} = \mathbb{R}^{L[g]}$ . Let  $\mathbb{R}^*$  denote this common set of reals. Suppose  $x \in \mathbb{R}^*$ . AD implies that  $x^\sharp$  exists. Let  $C_x$  be the club set of Silver indiscernible for  $L[x]$ . Let  $f \in [C_1 \cap C_x]_*^{\omega_1}$ . Note that  $f \restriction \omega \in L[f]$  and  $x \in L[f]$ . Thus  $x^\sharp$  can be defined within  $L[f]$  as the collection of formulas true in  $L[x]$  using  $\{f(n) : n \in \omega\}$  as indiscernibles. So  $x^\sharp \in \mathbb{R}^*$ . It has been shown that  $\mathbb{R}^*$  is closed under sharps. This implies that for  $f \in [C_1]_*^{\omega_1}$ ,  $\omega_1^{L[f]} > \omega_1^L$ .

This example motivates the further study of the stable theory of the partition measures  $\mu_\epsilon$  for  $\epsilon \leq \omega_1$ : Note that for each sentence  $\varphi$  in the language of set theory, since  $\mu_\epsilon$  is an ultrafilter, one has exactly one of the following holds: (i) for  $\mu_\epsilon$ -almost all  $f$ ,  $L[f] \models \varphi$  or (ii) for  $\mu_\epsilon$ -almost all  $f$ ,  $L[f] \models \neg\varphi$ .

In work with Trang, the authors study which natural statements belong to the stable theory of the partition measure  $\mu_\epsilon$ . For example, for all  $\epsilon \leq \omega_1$ , for  $\mu_\epsilon$ -almost all  $f$ ,  $L[f] \models \text{GCH}$ . The most difficult case is  $\epsilon = \omega_1$  where the  $[\omega_1]_*^{\omega_1}$ -almost everywhere club uniformization plays as essential role as a construction principle. This result will appear elsewhere.

As a warmup to showing every function  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is continuous, we will show that elements of  $\prod_{[\omega_1]_*^{\omega_1}} \omega_1 / \mu$  which have representatives that are continuous form an initial segment of the ultraproduct.

**Fact 4.3.** *Suppose  $\Psi, \Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$ . Suppose  $\Phi$  is continuous  $\mu_{\omega_1}$ -almost everywhere and  $\Psi <_{\mu_{\omega_1}} \Phi$ , which means  $\{f \in [\omega_1]_*^{\omega_1} : \Psi(f) < \Phi(f)\} \in \mu_{\omega_1}$ . Then  $\Psi$  is continuous  $\mu_{\omega_1}$ -almost everywhere.*

*Proof.* Let  $C_0 \subseteq \omega_1$  be a club so that  $\Phi$  is continuous on  $[C_0]_*^{\omega_1}$  and  $\Psi(f) < \Phi(f)$  for all  $f \in [C_0]_*^{\omega_1}$ . Let  $K \subseteq [C_0]_*^{\omega_1}$  be the collection of  $\sigma$  so that for all  $f, g \in [C_0]_*^{\omega_1}$  with  $f \restriction |\sigma| = g \restriction |\sigma| = \sigma$ ,  $\Phi(f) = \Phi(g)$ . Since  $\Phi$  is continuous on  $[C_0]_*^{\omega_1}$ ,  $K$  is dense in  $[C_0]_*^{\omega_1}$  in the sense that for all  $f \in [C_0]_*^{\omega_1}$ , there exists an  $\alpha < \omega_1$  so that  $f \restriction \alpha \in K$ . For each  $\sigma \in K$ , let  $d_\sigma = \Phi(f)$  for any  $f \in [C_0]_*^{\omega_1}$  such that  $f \restriction |\sigma| = \sigma$ , which is well defined by the definition of  $K$ .

For each  $\sigma \in K$ , define  $\Gamma_\sigma : [C_0 \setminus \sup(\sigma) + 1]_*^{\omega_1} \rightarrow \omega_1$  by  $\Gamma_\sigma(g) = \Psi(\sigma^\wedge g)$ . Thus for all  $g \in [C_0 \setminus \sup(\sigma) + 1]_*^{\omega_1}$ ,  $\Gamma_\sigma(g) < d_\sigma$ . By the countable additivity of  $\mu_{\omega_1}$ , for  $\mu_{\omega_1}$ -almost all  $g$ ,  $\Gamma_\sigma(g)$  takes a constant value denoted  $c_\sigma$ . Define  $\Psi' : [C_0]_*^{\omega_1} \rightarrow \omega_1$  as follows: For each  $f \in [C_0]_*^{\omega_1}$ , find the least  $\alpha$  so that  $f \restriction \alpha \in K$  (which exists by the density of  $K$ ), and let  $\Psi'(f) = c_{f \restriction \alpha}$ . Note that  $\Psi'$  is continuous.

The claim is that  $\Psi =_\mu \Psi'$ :

Define  $P : [C_0]_*^{\omega_1} \rightarrow 2$  by  $P(f) = 0$  if and only if  $\Psi(f) = \Psi'(f)$ . By the strong partition relation  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , let  $C_1 \subseteq C_0$  be a club on which  $P$  is homogeneous. Let  $f \in [C_1]_*^{\omega_1}$ . Find the least  $\alpha$  so that  $f \restriction \alpha \in K$ . There is some club  $D$  so that  $\Gamma_{f \restriction \alpha}$  takes constant value  $c_{f \restriction \alpha}$  on  $[D]_*^{\omega_1}$ . Let  $D' = (C_1 \setminus \sup(\sigma) + 1) \cap D$ . Let  $g \in [D']_*^{\omega_1}$ . Then  $\Psi(f \restriction \alpha^\wedge g) = \Gamma_{f \restriction \alpha}(g) = c_{f \restriction \alpha} = \Psi'(f \restriction \alpha^\wedge g)$ . So

$P(f \upharpoonright \alpha \hat{g}) = 0$  and  $f \upharpoonright \alpha \hat{g} \in [C_1]_*^{\omega_1}$ . Hence  $C_1$  is homogeneous for  $P$  taking value 0. This implies  $\Psi = \Psi'$  on  $[C_1]_*^{\omega_1}$ , so  $\Psi$  is continuous on  $[C_1]_*^{\omega_1}$  since  $\Psi'$  is continuous.  $\square$

The following is a useful notation:

**Definition 4.4.** Define  $\text{drop} : [\omega_1]^{\omega_1} \times \omega_1 \rightarrow [\omega_1]^{\omega_1}$  by  $\text{drop}(f, \delta)(\alpha) = f(\delta + \alpha)$ . Thus  $\text{drop}(f, \delta)$  is merely  $f$  with its  $\delta^{\text{th}}$ -initial segment,  $f \upharpoonright \delta$ , removed.

Let  $A \subseteq \omega_1$  be an unbounded subsets of  $\omega_1$ . Let  $\text{next}_A : \omega_1 \rightarrow A$  be defined by  $\text{next}_A(\alpha)$  is the smallest element of  $A$  strictly larger than  $\alpha$ . Let  $\text{next}_A^\omega : \omega_1 \rightarrow A$  be defined by  $\text{next}_A^\omega(\alpha)$  is the  $\omega^{\text{th}}$ -element of  $A$  larger than  $\alpha$ .

**Theorem 4.5.** *Every function  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is continuous almost everywhere.*

*Proof.* Let  $P : [\omega_1]_*^{\omega_1} \rightarrow 2$  be defined by  $P(f) = 0$  if and only if there exists  $\alpha < \omega_1$  so that for all club  $C \subseteq \omega_1$ , there exists  $g \in [C]_*^{\omega_1}$  so that  $f \upharpoonright \alpha \hat{g} \in [\omega_1]_*^{\omega_1}$  (i.e. is strictly increasing) and  $\Phi(f \upharpoonright \alpha \hat{g}) < g(0)$ . (The idea is that if  $P(f) = 0$ , then the  $\alpha$  above gives the initial segment of  $f$  which serves as a continuity point. One would like to use Lemma 4.1 to obtain the value of  $\Phi$  at this continuity point; however, the quantifiers in the definition of  $P$  are the opposite of the quantifiers of Lemma 4.1.)

By  $\omega_1 \rightarrow_* (\omega_1)_*^{\omega_1}$ , let  $D \subseteq \omega_1$  be homogeneous for  $P$ .

Claim 1:  $D$  is homogeneous for  $P$  taking value 0.

To prove this: Suppose that it is homogeneous for  $P$  taking value 1. This means that for all  $f \in [D]_*^{\omega_1}$ , for all  $\alpha < \omega_1$ , there exists a club  $C \subseteq [\omega_1]^{\omega_1}$  so that for all  $g \in [C]_*^{\omega_1}$  such that  $f \upharpoonright \alpha \hat{g} \in [\omega_1]_*^{\omega_1}$ ,  $\Phi(f \upharpoonright \alpha \hat{g}) \geq g(0)$ .

Define  $R \subseteq [D]_*^{<\omega_1} \times \text{club}_{\omega_1}$  by  $R(\sigma, C)$  if and only if for all  $g \in [C]_*^{\omega_1}$  with  $\sigma \hat{g} \in [\omega_1]_*^{\omega_1}$ ,  $\Phi(\sigma \hat{g}) \geq g(0)$ .  $R$  is  $\subseteq$ -downward closed. Note that  $\text{dom}(R) = [D]_*^{<\omega_1}$  since  $D$  is homogeneous for  $P$  taking value 1. By Theorem 3.10, there is a club  $E \subseteq D$  so that  $R \cap ([E]_*^{<\omega_1} \times \text{club}_{\omega_1})$  has a uniformization. Let  $\Lambda : [E]_*^{<\omega_1} \rightarrow \text{club}_{\omega_1}$  be such a uniformization function for  $R$  on  $[E]_*^{<\omega_1}$ .

First one will construct an  $h \in [E]_*^{\omega_1}$  by recursion as follows: Let  $F_0 = E \cap \Lambda(\emptyset)$ . Let  $h(0) = \text{next}_{F_0}^\omega(0)$ . Suppose  $h \upharpoonright \alpha$  and clubs  $F_\beta$  for all  $\beta < \alpha$  have been defined. Let  $F_\alpha = \Lambda(h \upharpoonright \alpha) \cap \bigcap_{\beta < \alpha} F_\beta$ . Let  $h(\alpha) = \text{next}_{F_\alpha}^\omega(\sup(h \upharpoonright \alpha))$ . This completes the construction of  $h \in [E]_*^{\omega_1}$  and sequence of clubs  $\langle F_\beta : \beta < \omega_1 \rangle$ .

Since one has the sequence  $\langle F_\beta : \beta < \omega_1 \rangle$ , one also can define the sequence of functions  $\langle \text{next}_{F_\beta} : \beta < \omega_1 \rangle$ . Define  $H : \omega_1 \times \omega \rightarrow \omega_1$  by recursions as follows:  $H(0, 0) = \text{next}_{F_0}(0)$ .  $H(0, n+1) = \text{next}_{F_0}(H(0, n))$ . If for some  $\alpha$ ,  $H(\beta, n)$  has been defined for all  $\beta < \alpha$  and  $n \in \omega$ , then let  $\mu = \sup\{H(\beta, n) : \beta < \alpha \wedge n \in \omega\}$ . Let  $H(\alpha, 0) = \text{next}_{F_\alpha}(\mu)$ . Let  $H(\alpha, n+1) = \text{next}_{F_\alpha}(H(\alpha, n))$ . Now  $H$  witnesses that  $h$  has uniform cofinality  $\omega$ . By the construction, it is clear that  $h$  is increasing and discontinuous everywhere. Thus  $h \in [E]_*^{\omega_1}$ , i.e. is increasing and has correct type.

Now pick any  $\alpha < \omega_1$ . Since  $\text{drop}(h, \alpha) \in [F_\alpha]_*^{\omega_1} \subseteq [\Lambda(h \upharpoonright \alpha)]_*^{\omega_1}$ , the definition of  $\Lambda$  being a uniformization for  $R$  implies that  $\Phi(h) = \Phi(h \upharpoonright \alpha \hat{\text{drop}(h, \alpha)}) \geq \text{drop}(h, \alpha)(0) = h(\alpha)$ . Since  $\alpha < \omega_1$  was arbitrary, this shows that for all  $\alpha < \omega_1$ ,  $\Phi(h) \geq h(\alpha)$ . Since  $h \in [E]_*^{\omega_1}$  is a strictly increasing function,  $\Phi(h) \geq \omega_1$ . This is impossible since  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  is a function which takes values among the countable ordinals. This establishes Claim 1.

Thus  $D$  is homogeneous for  $P$  taking value 0. Let  $K \subseteq [D]_*^{<\omega_1}$  be the collection of  $\sigma$  such that for all club  $C \subseteq \omega_1$ , there is some  $g \in [C]_*^{\omega_1}$  with  $\sigma \hat{g} \in [\omega_1]_*^{\omega_1}$  and  $\Phi(\sigma \hat{g}) < g(0)$ . Note that  $K$  is dense in  $[D]_*^{\omega_1}$  since  $D$  is homogeneous for  $P$  taking value 0. (Elements of  $K$  will be the continuity points of  $\Phi$ . However, the quantifiers in the definition of  $K$  are the opposite of what is needed to use Lemma 4.1. The next partition attempts to resolve this.)

Fix  $\sigma \in K$ . Let  $Q_\sigma : [D \setminus \sup \sigma + \omega]_*^{\omega_1} \rightarrow 2$  be defined by  $Q_\sigma(g) = 0$  if and only if  $\Phi(\sigma \hat{g}) < g(0)$ . By  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , there is some  $E \subseteq D$  club which is homogeneous for  $Q_\sigma$ . By the property of  $\sigma \in K$ , there is some  $g \in [E]_*^{\omega_1}$  so that  $\sigma \hat{g} \in [\omega_1]_*^{\omega_1}$  and  $\Phi(\sigma \hat{g}) < g(0)$ . Thus one has  $Q_\sigma(g) = 0$ . This shows that  $E$  is homogeneous for  $Q_\sigma$  taking value 0. (Now one can apply Lemma 4.1 to find the value of  $\Phi$  associated to  $\sigma$ .) Define  $V_\sigma : [E \setminus \sup \sigma + \omega]_*^{\omega_1} \rightarrow \omega_1$  by  $V_\sigma(g) = \Phi(\sigma \hat{g})$ . For all  $g \in [E \setminus \sup \sigma + \omega]_*^{\omega_1}$ ,  $V_\sigma(g) < g(0)$ . By Lemma 4.1, there is an  $E' \subseteq E$  club so that  $V_\sigma$  is constant on  $[E']_*^{\omega_1}$  taking value  $c_\sigma$ . Note that  $c_\sigma$  does not depend on the choice of  $E$  or  $E'$  in the sense that for any club  $E''$  so that  $V_\sigma$  is constant on  $[E'']_*^{\omega_1}$ , the constant value must be  $c_\sigma$ .

Define  $\Psi : [D]_*^{\omega_1} \rightarrow \omega_1$  as follows: For  $f \in [D]_*^{\omega_1}$ , find the least  $\alpha$  so that  $f \restriction \alpha \in K$  and let  $\Psi(f) = c_{f \restriction \alpha}$ . Such an  $\alpha$  exists by the density of  $K$ .  $\Psi$  is continuous on  $[D]_*^{\omega_1}$ . This is because for any  $f \in [D]_*^{\omega_1}$ , let  $\alpha$  be the least ordinal so that  $f \restriction \alpha \in K$ . For any  $g \in [D]_*^{\omega_1}$  with  $g \restriction \alpha = f \restriction \alpha$ , one has that  $\alpha$  is also the least ordinal so that  $g \restriction \alpha \in K$ . Thus  $\Psi(g) = c_{g \restriction \alpha} = c_{f \restriction \alpha} = \Psi(f)$ .

Claim 2 : For  $\mu_{\omega_1}$ -almost all  $f$ ,  $\Phi(f) = \Psi(f)$ .

To see this: Define  $Y : [D]_*^{\omega_1} \rightarrow 2$  by  $Y(f) = 0$  if and only if  $\Phi(f) = \Psi(f)$ . By  $\omega_1 \rightarrow_* (\omega_1)_*^{\omega_1}$ , there is some club  $F \subseteq D$  which is homogeneous for  $Y$ . Let  $f \in [F]_*^{\omega_1}$ . Let  $\alpha$  be least so that  $f \restriction \alpha \in K$ . Let  $\sigma = f \restriction \alpha$ . There is some  $F' \subseteq F$  club so that  $V_\sigma$  takes constant value  $c_\sigma$  on  $[F']_*^{\omega_1}$ . Let  $g \in [F']_*^{\omega_1}$  be such that  $\sup(\sigma) < g(0)$ . Let  $f' = \sigma \hat{\ } g$ . Then  $\Phi(f') = \Phi(\sigma \hat{\ } g) = V_\sigma(g) = c_\sigma$ . As noted above, the least  $\alpha$  so that  $f' \restriction \alpha \in K$  is the same as the least  $\alpha$  so that  $f \restriction \alpha \in K$ . So  $c_\sigma = \Psi(f')$ . Thus  $Y(f') = 0$ . Since  $f' \in [F]_*^{\omega_1}$ ,  $F$  must be homogeneous for  $Y$  taking value 0.

It has been shown that for all  $f \in [F]_*^{\omega_1}$ ,  $\Phi(f) = \Psi(f)$ . Since  $\Psi$  is a continuous function,  $\Phi$  is  $\mu_{\omega_1}$ -almost equal to a continuous function.  $\square$

Zapletal asked the first author whether every partition of  $[\omega_1]^{\omega_1}$  into  $\omega_1$  many pieces must have at least one piece of cardinality  $[\omega_1]^{\omega_1}$ . The following gives a positive answer.

**Theorem 4.6.** *Suppose  $\langle X_\alpha : \alpha < \omega_1 \rangle$  is a sequence of subsets of  $[\omega_1]^{\omega_1}$  so that  $\bigcup_{\alpha < \omega_1} X_\alpha = [\omega_1]^{\omega_1}$ . Then there is an  $\alpha < \omega_1$  so that  $X_\alpha \approx [\omega_1]^{\omega_1}$ .*

*Proof.* Define  $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$  by letting  $\Phi(f)$  be the least  $\alpha$  such that  $f \in X_\alpha$ . By Theorem 4.5, there is some club  $C \subseteq \omega_1$  so that  $\Phi$  is continuous on  $[C]_*^{\omega_1}$ . Pick any  $f \in [C]_*^{\omega_1}$ . Let  $\delta = \Phi(f)$ . By continuity, there is some  $\alpha$  so that for all  $g \in [C]_*^{\omega_1}$  with  $g \restriction \alpha = f \restriction \alpha$ ,  $\Phi(g) = \Phi(f) = \delta$ . Using Fact 2.2, let  $\Delta : [\omega_1]^{\omega_1} \rightarrow [C \setminus f(\alpha)]_*^{\omega_1}$  be a bijection. Define  $\Gamma : [\omega_1]^{\omega_1} \rightarrow X_\delta$  by  $\Gamma(g) = (f \restriction \alpha) \hat{\ } \Delta(g)$ . Then  $\Gamma$  is an injection. Thus  $|X_\delta| = |[\omega_1]^{\omega_1}|$ .  $\square$

**Theorem 4.7.**  $|[\omega_1]^{<\omega_1}| < |[\omega_1]^{\omega_1}|$ .

*Proof.* For  $\alpha \leq \beta < \omega_1$ , let  $X_{\alpha,\beta} = [\beta]^\alpha$  and note that  $|X_{\alpha,\beta}| \leq |\mathbb{R}|$ . Observe that  $[\omega_1]^{<\omega_1} = \bigcup_{\alpha \leq \beta < \omega_1} X_{\alpha,\beta}$ . By using the Gödel pairing function, one can recognize this union as an  $\omega_1$ -length union of subsets of  $[\omega_1]^{<\omega_1}$  with cardinality less than or equal to  $\mathbb{R}$  (but non-uniformly). Therefore  $|[\omega_1]^{<\omega_1}| = |[\omega_1]^{\omega_1}|$  is impossible since it would violate Theorem 4.6.  $\square$

## 5. CONTINUITY OF FUNCTIONS $[\omega_1]^{\omega_1} \rightarrow \omega_1 \omega_1$

Recall that  $\omega_1 \omega_1$  is the collection of all functions  $f : \omega_1 \rightarrow \omega_1$ .

**Definition 5.1.** A function  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1 \omega_1$  is continuous if and only if for all  $f \in [\omega_1]^{\omega_1}$ , for all  $\epsilon < \omega_1$ , there exists a  $\delta < \omega_1$  so that for all  $g \in [\omega_1]^{\omega_1}$ , if  $f \restriction \delta = g \restriction \delta$ , then  $\Phi(f) \restriction \epsilon = \Phi(g) \restriction \epsilon$ .

If one gives  $[\omega_1]^{\omega_1}$  and  $\omega_1 \omega_1$  the topology indicated in Definition 2.11, then  $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1 \omega_1$  is continuous if and only if  $\Phi$  is continuous in the topological sense.

$\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1 \omega_1$  is continuous almost everywhere if and only if there club  $C \subseteq \omega_1$  so that  $\Phi$  is continuous on  $[C]_*^{\omega_1}$ .

**Lemma 5.2.** *There is no club  $D \subseteq \omega_1$  and no function  $\Lambda : [D]_*^{\omega_1} \rightarrow \omega_1$  with the property that for all  $f \in [D]_*^{\omega_1}$ , for all  $\alpha < \omega_1$ , there exists a club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$ , if  $(f \restriction \alpha) \hat{\ } g \in [\omega_1]_*^{\omega_1}$ , then  $\Lambda((f \restriction \alpha) \hat{\ } g) \geq g(0)$ .*

*Proof.* The proof of Claim 1 in Theorem 4.5 is precisely this lemma. As there, one can prove this by using the almost everywhere  $[\omega_1]_*^{<\omega_1}$ -club uniformization (Theorem 3.10). However, having already established the continuity property in Theorem 4.5, this lemma can be derived easily as follows:

Suppose such a club  $D \subseteq \omega_1$  and function  $\Lambda$  exist. By Theorem 4.5, there is a  $D_0 \subseteq D$  so that  $\Lambda \restriction [D_0]_*^{\omega_1}$  is continuous. Take any  $f \in [D_0]_*^{\omega_1}$ . Let  $\zeta = \Lambda(f)$ . By continuity, there is an  $\alpha < \omega_1$  so that for all  $h \in [D_0]_*^{\omega_1}$ , if  $f \restriction \alpha = h \restriction \alpha$ , then  $\Lambda(h) = \Lambda(f) = \zeta$ . By the hypothesis applied to this  $f$  and  $\alpha$ , there exists some club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$ , if  $(f \restriction \alpha) \hat{\ } g \in [\omega_1]_*^{\omega_1}$ , then  $\Lambda((f \restriction \alpha) \hat{\ } g) \geq g(0)$ . Pick  $g \in [C \cap D_0]_*^{\omega_1}$  such that  $(f \restriction \alpha) \hat{\ } g \in [\omega_1]_*^{\omega_1}$  and  $g(0) > \zeta$ . Then  $\Lambda((f \restriction \alpha) \hat{\ } g) \geq g(0) > \zeta$  by choice of  $C$ . However  $(f \restriction \alpha) \hat{\ } g \in [D_0]_*^{\omega_1}$  and  $((f \restriction \alpha) \hat{\ } g) \restriction \alpha = f \restriction \alpha$ . Since  $f \restriction \alpha$  is a point of continuity for  $\Lambda$  on  $[D_0]_*^{\omega_1}$  for taking value  $\zeta$ , one must have  $\Lambda((f \restriction \alpha) \hat{\ } g) = \zeta$ . Contradiction.  $\square$

**Theorem 5.3.** (With Trang) Every function  $\Phi : [\omega_1]^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  is continuous almost everywhere.

*Proof.* Define a partition  $P_0 : [\omega_1]^{\omega_1} \rightarrow 2$  by  $P_0(f) = 0$  if and only if for all  $\beta < \omega_1$ , for all  $\gamma < \omega_1$ , there exists an  $\alpha > \gamma$  so that for all club  $C \subseteq \omega_1$ , there exists a  $g \in [C]_*^{\omega_1}$  so that  $(f \restriction \alpha)^\wedge g \in [\omega_1]_*^{\omega_1}$  and  $\Phi((f \restriction \alpha)^\wedge g)(\beta) < g(0)$ . (Here  $P_0(f) = 0$  will intuitively mean that for each  $\beta < \omega_1$  and  $\gamma < \omega_1$ , there is an  $\alpha > \gamma$  such that  $f \restriction \alpha$  is a continuity point for the  $\beta^{\text{th}}$  coordinate of  $\Phi$ .)

By  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , there is a club  $D_0 \subseteq \omega_1$  which is homogeneous for  $P_0$ .

Claim 1:  $D_0$  is homogeneous for  $P_0$  taking value 0.

To see this, suppose  $D_0$  is homogeneous for  $P_0$  taking value 1. Negating the definition of  $P_0$ , one see that  $P_0(f) = 1$  if and only if there exists  $\beta < \omega_1$ , there exists  $\gamma < \omega_1$  so that for all  $\alpha > \gamma$ , there exists a club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$ , if  $(f \restriction \alpha)^\wedge g \in [\omega_1]_*^{\omega_1}$ , then  $\Phi((f \restriction \alpha)^\wedge g)(\beta) \geq g(0)$ .

Let  $\Psi_0 : [D_0]_*^{\omega_1} \rightarrow \omega_1$  be defined by letting  $\Psi_0(f)$  be the least  $\beta$  witnessing the first existential quantifier in the definition of  $P_0(f) = 1$ . By Theorem 4.5, there is a club  $D'_0 \subseteq D_0$  so that  $\Psi_0 \restriction [D'_0]_*^{\omega_1}$  is continuous.

Define  $\Psi_1 : [D'_0]_*^{\omega_1} \rightarrow \omega_1$  by  $\Psi_1(f)$  is the least  $\gamma$  witnessing the second existential quantifier in the definition of  $P_0(f) = 1$  for  $\beta = \Psi_0(f)$ . Again by Theorem 4.5, there is a club  $D_1 \subseteq D'_0$  so that  $\Psi_1 \restriction [D_1]_*^{\omega_1}$  is continuous. (Note that  $\Psi_0 \restriction [D_1]_*^{\omega_1}$  is also continuous.)

Now take an  $h \in [D_1]_*^{\omega_1}$ . Let  $\hat{\beta} = \Psi_0(h)$  and  $\hat{\gamma} = \Psi_1(h)$ . By the continuity of  $\Psi_0 \restriction [D_1]_*^{\omega_1}$  and  $\Psi_1 \restriction [D_1]_*^{\omega_1}$ , there is  $\zeta < \omega_1$  so that for all  $g \in [D_1]_*^{\omega_1}$ , if  $g \restriction \zeta = h \restriction \zeta$ , then  $\Psi_0(g) = \Psi_0(h) = \hat{\beta}$  and  $\Psi_1(g) = \Psi_1(h) = \hat{\gamma}$ . Let  $\xi = \max\{\zeta, \hat{\gamma}\}$ . Let  $\sigma = h \restriction \xi$ . Note that for all  $g \in [D_1]_*^{\omega_1}$  so that  $\sigma^\wedge g \in [D_1]_*^{\omega_1}$ , one has that  $\Psi_0(\sigma^\wedge g) = \hat{\beta}$  and  $\Psi_1(\sigma^\wedge g) = \hat{\gamma}$ .

Define  $\Lambda : [D_1 \setminus (\sup \sigma + 1)]_*^{\omega_1} \rightarrow \omega_1$  by  $\Lambda(f) = \Phi(\sigma^\wedge f)(\hat{\beta})$ . Observe that  $\Lambda$  has the property that for all  $\alpha$  and  $f \in [D_1 \setminus (\sup \sigma + 1)]_*^{\omega_1}$ , there is a club  $C \subseteq \omega_1$  so that for all  $g \in [C]_*^{\omega_1}$ , if  $(f \restriction \alpha)^\wedge g \in [\omega_1]_*^{\omega_1}$ , then  $\Lambda((f \restriction \alpha)^\wedge g) \geq g(0)$ . Such a function can not exist by Lemma 5.2. Claim 1 has been shown so  $D_0$  is homogeneous for  $P_0$  taking value 0.

For each  $\beta < \omega_1$ , let  $K_\beta$  be the collection of  $\sigma \in [D_0]_*^{<\omega_1}$  so that for all clubs  $C \subseteq \omega_1$ , there exists a  $g \in [C]_*^{\omega_1}$  so that  $\sigma^\wedge g \in [\omega_1]_*^{\omega_1}$  and  $\Phi(\sigma^\wedge g)(\beta) < g(0)$ . Note that for all  $\beta < \omega_1$ ,  $K_\beta$  is dense in  $[D_0]_*^{\omega_1}$ , which means that for all  $f \in [D_0]_*^{\omega_1}$ , for all  $\gamma < \omega_1$ , there exists an  $\alpha > \gamma$  with  $f \restriction \alpha \in K_\beta$ . To see this: for any  $f \in [D_0]_*^{\omega_1}$  and  $\gamma < \omega_1$ ,  $P_0(f) = 0$  implies there exists some  $\alpha > \gamma$  so that for all club  $C \subseteq \omega_1$ , there exists a  $g \in [C]_*^{\omega_1}$  with  $f \restriction \alpha^\wedge g \in [\omega_1]_*^{\omega_1}$  and  $\Phi((f \restriction \alpha)^\wedge g)(\beta) < g(0)$ . This  $\alpha$  would suffice. ( $K_\beta$  will be the collection of continuity point for the  $\beta^{\text{th}}$ -coordinate of  $\Phi$ . The next partition will use Lemma 4.1 to find the value associated to this continuity point.)

For each  $\beta < \omega_1$  and  $\sigma \in K_\beta$ , define the partition  $Q_\sigma^\beta : [\omega_1 \setminus (\sup \sigma + 1)]_*^{\omega_1} \rightarrow 2$  by  $Q_\sigma^\beta(g) = 0$  if and only if  $\Phi(\sigma^\wedge g)(\beta) < g(0)$ . By  $\omega_1 \rightarrow (\omega_1)_*^{\omega_1}$ , there is a club  $E \subseteq \omega_1$  which is homogeneous for  $Q_\sigma^\beta$ . By definition of  $\sigma \in K_\beta$ , there is a  $g \in [E]_*^{\omega_1}$  so that  $\sigma^\wedge g \in [\omega_1]_*^{\omega_1}$  and  $\Phi(\sigma^\wedge g) < g(0)$ . Thus  $E$  is homogeneous for  $Q_\sigma^\beta$  taking value 0. Define  $\Phi_\sigma^\beta : [E \setminus (\sup \sigma + 1)]_*^{\omega_1} \rightarrow \omega_1$  by  $\Phi_\sigma^\beta(g) = \Phi(\sigma^\wedge g)(\beta)$ . Since  $E$  is homogeneous for  $Q_\sigma^\beta$  taking value 0, one has that for all  $g \in [E \setminus (\sup \sigma + 1)]_*^{\omega_1}$ ,  $\Phi_\sigma^\beta(g) < g(0)$ . By Lemma 4.1, there is a club  $F \subseteq E$  and a  $c_\sigma^\beta < \omega_1$  so that for all  $g \in [F]_*^{\omega_1}$ ,  $\Phi_\sigma^\beta(g) = c_\sigma^\beta$ . (Note that  $c_\sigma^\beta$  does not depend on the choice of clubs  $E$  or  $F$ .)

For each  $f \in [D_0]_*^{\omega_1}$ , define a strictly increasing sequence  $\langle \alpha_f^\beta : \beta < \omega_1 \rangle$  by recursion as follows: Let  $\alpha_f^0$  be the least  $\alpha$  so that  $f \restriction \alpha \in K_0$ , which exists by the density of  $K_0$  in  $[D_0]_*^{\omega_1}$ . Suppose  $\beta < \omega_1$  and for all  $\gamma < \beta$ ,  $\alpha_f^\gamma$  has been defined with the property that  $f \restriction \alpha_f^\gamma \in K_\gamma$ . Let  $\xi = \sup\{\alpha_f^\gamma : \gamma < \beta\}$ . Let  $\alpha_f^\beta$  be the least  $\alpha > \xi$  so that  $f \restriction \alpha \in K_\beta$ , which exists by the density of  $K_\beta$  in  $[D_0]_*^{\omega_1}$ . Note that the map  $f \mapsto \langle \alpha_f^\beta : \beta < \omega_1 \rangle$  is continuous in the sense that for any  $f \in [D_0]_*^{\omega_1}$ , any  $\gamma < \omega_1$ , and for all  $g \in [D_0]_*^{\omega_1}$ , if  $g \restriction \alpha_f^\gamma = f \restriction \alpha_f^\gamma$ , then  $\alpha_f^\beta = \alpha_g^\beta$  for all  $\beta \leq \gamma$ . (Intuitively, the sequence  $\langle \alpha_f^\beta : \beta < \omega_1 \rangle$  collects the length of the various continuity points of  $f$  for the various coordinates of  $\Phi(f)$ .)

Define  $\Gamma : [D_0]_*^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  by  $\Gamma(f)(\beta) = c_{f \restriction \alpha_f^\beta}^\beta$ . Pick any  $f \in [D_0]_*^{\omega_1}$  and  $\gamma < \omega_1$ . As observed above, for all  $g$  so that  $g \restriction \alpha_f^\gamma = f \restriction \alpha_f^\gamma$ , one has that  $\langle \alpha_g^\beta : \beta \leq \gamma \rangle = \langle \alpha_f^\beta : \beta \leq \gamma \rangle$ . Hence  $\Gamma(f) \restriction \gamma + 1 = \Gamma(g) \restriction \gamma + 1$  for all  $g$  so that  $g \restriction \alpha_f^\gamma = f \restriction \alpha_f^\gamma$ . This shows that  $\Gamma : [D_0]_*^{\omega_1} \rightarrow {}^{\omega_1}\omega_1$  is continuous.

Claim 2 : There is a club  $D_2 \subseteq \omega_1$  so that  $\Phi \restriction [D_2]_*^{\omega_1} = \Gamma \restriction [D_2]_*^{\omega_1}$ .

To see Claim 2: Define a partition  $P_1 : [D_0]_*^{\omega_1} \rightarrow 2$  by  $P_1(f) = 0$  if and only if  $\Gamma(f) = \Phi(f)$ . By  $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ , there exists a club  $D_2 \subseteq D_0$  which is homogeneous for  $P_1$ .

Define a relation  $R \subseteq \omega_1 \times [D_2]_{*}^{\omega_1} \times \text{club}_{\omega_1}$  by  $R(\beta, \sigma, C)$  if and only if  $\sigma \in K_\beta$  and  $\Phi_\sigma^\beta$  is constant on  $[C]_{*}^{\omega_1}$  taking value  $c_\sigma^\beta$ . Note that the domain of  $R$  is  $Y = \{(\beta, \sigma) : \sigma \in K_\beta\}$ .  $R$  is  $\subseteq$ -closed in the  $\text{club}_{\omega_1}$ -coordinate. By Theorem 3.11, there is a club  $D_3 \subseteq D_2$  and a uniformization function  $\Sigma : Z \rightarrow \text{club}_{\omega_1}$  so that for all  $(\beta, \sigma) \in Z$ ,  $R(\beta, \sigma, \Sigma(\beta, \sigma))$ , where  $Z = \{(\beta, \sigma) : \beta \in \omega_1 \wedge \sigma \in K_\beta \cap [D_3]_{*}^{<\omega_1}\}$ .

If  $C \subseteq \omega_1$  is a club, then let  $p_C \in [C]_{*}^{\omega_1}$  be defined by  $p_C(\alpha) = \text{enum}_C(\omega \cdot (\alpha + 1))$ .  $p_C$  can be regarded as the canonical correct type function passing through the club  $C$ .

A function  $h \in [D_3]_{*}^{\omega_1}$ , an increasing sequence of ordinals  $\langle \gamma_\delta : \delta < \omega_1 \rangle$ , and a sequence of clubs  $\langle F_\delta : \delta < \omega_1 \rangle$  will be constructed by recursion. (The function  $h$  will be defined by specifying longer initial segments of  $h$  at each stage.)

Let  $g_0 = p_{D_3}$ . Note that  $g_0 \in [D_3]_{*}^{\omega_1}$ . Let  $\gamma_0 = \alpha_{g_0}^0$ . Define  $h \upharpoonright \gamma_0 = g_0 \upharpoonright \gamma_0$ . Note that  $h \upharpoonright \gamma_0 \in K_0$  and therefore  $(0, h \upharpoonright \gamma_0) \in Z$ . Let  $F_0 = \Sigma(0, h \upharpoonright \gamma_0) \cap D_3$ . Note that for any  $h' \supseteq h \upharpoonright \gamma_0$ , one has that  $\alpha_{h'}^0 = \gamma_0$ .

Suppose  $\gamma_\beta$ ,  $h \upharpoonright \gamma_\beta$ , and  $F_\beta$  have been defined for all  $\beta < \delta$ . Suppose it has also been shown that for all  $\beta < \delta$ , for all  $h' \in [D_3]_{*}^{\omega_1}$  such that  $h' \supseteq h \upharpoonright \gamma_\beta$ , one has that  $\alpha_{h'}^\beta = \gamma_\beta$ . Let  $\xi = \sup\{\gamma_\beta : \beta < \delta\}$ . Let  $G_\delta = (\bigcap_{\beta < \delta} F_\beta) \setminus (\sup(h \upharpoonright \xi) + 1)$ . Let  $g_\delta = h \upharpoonright \xi \wedge p_{G_\delta}$ . Note that  $g_\delta \in [D_3]_{*}^{\omega_1}$ . Let  $\gamma_\delta = \alpha_{g_\delta}^\delta$ . Let  $h \upharpoonright \gamma_\delta = g_\delta \upharpoonright \gamma_\delta$ . Since  $h \upharpoonright \gamma_\beta \subseteq g_\delta$  for all  $\beta < \delta$ , one has that  $\gamma_\delta > \gamma_\beta$  for all  $\beta < \delta$ . Note that for all  $h' \supseteq h \upharpoonright \gamma_\delta$ ,  $\alpha_{h'}^\delta = \gamma_\delta$ . Also  $h \upharpoonright \gamma_\delta \in K_\delta$  and therefore  $(\delta, h \upharpoonright \gamma_\delta) \in Z$ . Let  $F_\delta = \Sigma(\delta, h \upharpoonright \gamma_\delta) \cap \bigcap_{\beta < \delta} F_\beta$ .

This completes the construction. Note that  $h \in [D_3]_{*}^{\omega_1}$ . By construction,  $\langle \gamma_\delta : \delta < \omega_1 \rangle = \langle \alpha_h^\delta : \delta < \omega_1 \rangle$ . Fix any  $\delta < \omega_1$ . Due to the construction,  $\text{drop}(h, \alpha_h^\delta) \in [F_\delta]_{*}^{\omega_1} \subseteq [\Sigma(\delta, h \upharpoonright \alpha_h^\delta)]_{*}^{\omega_1}$ . Since  $\Sigma$  is a uniformization for  $R$ , one has that  $\Phi(h)(\delta) = \Phi(h \upharpoonright \alpha_h^\delta \wedge \text{drop}(h, \alpha_h^\delta))(\delta) = \Phi_{h \upharpoonright \alpha_h^\delta}^\delta(\text{drop}(h, \alpha_h^\delta)) = c_{h \upharpoonright \alpha_h^\delta}^\delta = \Gamma(h)(\delta)$ . Since  $\delta$  was arbitrary,  $\Phi(h) = \Gamma(h)$ . Thus  $P_1(h) = 0$ . Since  $D_3$  was homogeneous for  $P_1$  and  $h \in [D_3]_{*}^{\omega_1}$ ,  $D_3$  is homogeneous for  $P_1$  taking value 0. Thus  $\Phi \upharpoonright [D_3]_{*}^{\omega_1} = \Gamma \upharpoonright [D_3]_{*}^{\omega_1}$ .  $\Phi$  is continuous on  $[D_3]_{*}^{\omega_1}$  since  $\Gamma$  is continuous on  $[D_3]_{*}^{\omega_1}$ . This completes the proof.  $\square$

## 6. FAILURE OF CONTINUITY PROPERTY AT $\omega_2$

A natural question would be whether the continuity phenomenon occurs at  $\omega_2$ . That is, for every function  $\Phi : [\omega_2]_{*}^{\omega_2} \rightarrow \omega_2$ , is there is a club  $C \subseteq \omega_2$  so that  $\Phi \upharpoonright [C]_{*}^{\omega_2}$  is continuous?

**Fact 6.1.** (Martin) For all  $\alpha < \omega_2$ ,  $\omega_2 \rightarrow_* (\omega_2)_2^\alpha$ . That is,  $\omega_2$  is a weak partition cardinal.

(Martin and Paris) The partition relation  $\omega_2 \rightarrow_* (\omega_2)_2^{\omega_2}$  fails.

The strong partition property for  $\omega_1$  played an essential role in establishing the continuity property for functions  $\Phi : [\omega_1]_{*}^{\omega_1} \rightarrow \omega_1$ . The failure of the strong partition property at  $\omega_2$  seems to suggest that one should use a counterexample to the strong partition property as a counterexample to the continuity property. However, it is not clear if the fact that a function  $P : [\omega_2]_{*}^{\omega_2} \rightarrow 2$  has no club homogeneous set alone can imply the failure of the continuity property. For this, one needs to analyze explicit counterexamples to the strong partition property at  $\omega_2$ .

The proof of the Martin-Paris theorem roughly shows that if the partition relation  $\omega_2 \rightarrow (\omega_2)_2^{\omega_2}$  holds, then  $\omega_3 \rightarrow (\omega_3)_2^\alpha$  holds for each  $\alpha < \omega_1$ . The partition relation  $\omega_3 \rightarrow (\omega_3)_2^2$  already implies that  $\omega_3$  is regular. However, AD proves that  $\text{cof}(\omega_3) = \omega_2$ . See [8] Section 13.

The second author produced an explicit example of the failure of the strong partition property at  $\omega_2$ . Its proof gives some additional properties that will show this function also fails to have the continuity property. The proof of the following theorem requires an analysis of the ultrapower  $\prod_{\omega_1} \omega_1 / \mu$ , where  $\mu$  is the club measure on  $\omega_1$ , the Kunen tree, and Kunen functions for functions of the form  $f : \omega_1 \rightarrow \omega_1$ .

Since  $\prod_{\omega_1} \omega_1 / \mu = \omega_2$ , every function  $h : \omega_1 \rightarrow \mathcal{P}(\omega_1)$  represents a subset of  $\omega_2$ . If  $A \subseteq \omega_2$ , then denote  $A \in \prod_{\omega_1} \mathcal{P}(\omega_1) / \mu$  if and only if there is a function  $h : \omega_1 \rightarrow \mathcal{P}(\omega_1)$  so that  $A = [h]_\mu$ .

One can also show that  $\prod_{\omega_1} \mathcal{P}(\omega_1) / \mu$  is necessarily missing a subset of  $\omega_2$ . For instance,  $\{\alpha \in \omega_2 : \text{cof}(\alpha) = \omega_1\}$  does not belong to  $\prod_{\omega_1} \mathcal{P}(\omega_1) / \mu$ . These results are essential ingredients of the proof of the follow theorem. The background material and proof of the results mentioned above and the following theorem are given in [3] Section 6.

**Theorem 6.2.** (Jackson) Let  $\mu$  denote the club measure on  $\omega_1$ . Let  $P : [\omega_2]_{*}^{\omega_2} \rightarrow 2$  be defined by  $P(f) = 0$  if and only if  $\text{rang}(f) \in \prod_{\omega_1} \mathcal{P}(\omega_1) / \mu$ . Then there is no club  $C \subseteq \omega_2$  so that  $P$  is constant on  $[C]_{*}^{\omega_2}$ .

As a corollary of the proof of this result, one also has

**Corollary 6.3.** *Let  $P$  denote the function from Theorem 6.2. Let  $\sigma \in [\omega_2]_*^{<\omega_2}$ . Define  $P_\sigma : [\omega_2 \setminus (\sup(\sigma) + \omega)]_*^{\omega_2} \rightarrow 2$  by  $P_\sigma(f) = 0$  if and only if  $P(\sigma \hat{\ } f) = 0$ . Then there is no club  $C \subseteq \omega_2$  so that  $P_\sigma$  is constant on  $[C]_*^{\omega_2}$ .*

With these results, one can show that  $P$  is not continuous on  $[C]_*^{\omega_2}$  for any club  $C \subseteq \omega_2$ .

**Theorem 6.4.** *Let  $P : [\omega_2]_*^{\omega_2} \rightarrow 2$  be the function from Theorem 6.2. Then there is no club  $C \subseteq \omega_2$  so that  $P \upharpoonright [C]_*^{\omega_2}$  is continuous.*

*Proof.* Suppose there was a club  $C$  so that  $P \upharpoonright [C]_*^{\omega_2}$  is continuous. Take any  $f \in [C]_*^{\omega_2}$ . Without loss of generality, say that  $P(f) = 0$ . By continuity, there is some  $\zeta < \omega_2$  so that for all  $g \in [C]_*^{\omega_2}$  with  $g \upharpoonright \zeta = f \upharpoonright \zeta$ ,  $P(f) = P(g)$ . Let  $\sigma = f \upharpoonright \zeta$ . This would mean that  $C \setminus (\sup(\sigma) + \omega)$  would be a club homogeneous for  $P_\sigma$ . This contradicts Corollary 6.3.  $\square$

## REFERENCES

1. Andrés Eduardo Caicedo and Richard Ketchersid, *A trichotomy theorem in natural models of  $\text{AD}^+$* , Set theory and its applications, Contemp. Math., vol. 533, Amer. Math. Soc., Providence, RI, 2011, pp. 227–258. MR 2777751
2. William Chan, *Ordinal definability and combinatorics of equivalence relations*, J. Math. Log. **19** (2019), no. 2, 1950009, 24. MR 4014889
3. William Chan, *An introduction to combinatorics of determinacy*, Trends in Set Theory, vol. 752, 2020, pp. 21–75.
4. William Chan and Stephen Jackson, *Cardinality of Wellordered Disjoint Unions of Quotients of Smooth Equivalence Relations*, arXiv e-prints (2019), arXiv:1903.03902.
5. William Chan, Stephen Jackson, and Nam Trang, *More definable combinatorics around the first and second uncountable cardinal*, In preparation.
6. William Chan and Connor Meehan, *Definable Combinatorics of Some Borel Equivalence Relations*, ArXiv e-prints (2017).
7. Gunter Fuchs, *A characterization of generalized Příkrý sequences*, Arch. Math. Logic **44** (2005), no. 8, 935–971. MR 2193185
8. Alexander S. Kechris, *AD and projective ordinals*, Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77), Lecture Notes in Math., vol. 689, Springer, Berlin, 1978, pp. 91–132. MR 526915
9. Alexander S. Kechris and W. Hugh Woodin, *Generic codes for uncountable ordinals, partition properties, and elementary embeddings*, Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I, Lect. Notes Log., vol. 31, Assoc. Symbol. Logic, Chicago, IL, 2008, pp. 379–397. MR 2463619
10. Itay Neeman, *Inner models and ultrafilters in  $L(\mathbb{R})$* , Bull. Symbolic Logic **13** (2007), no. 1, 31–53. MR 2300902
11. Philip Welch, *Closed Unbounded classes and the Häftig Quantifier Model*, arXiv e-prints (2019), arXiv:1903.02663.

DEPARTMENT OF MATHEMATICS, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PA 15213

Email address: wchan3@andrew.cmu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203

Email address: Stephen.Jackson@unt.edu