

CARDINALITY OF THE SET OF BOUNDED SUBSETS OF A CARDINAL

WILLIAM CHAN

ABSTRACT. If α and β are cardinals, then the cardinal exponentiation ${}^\alpha\beta$ is the set of all functions from α into β . Among the very few behaviors that can be inferred from the definition of cardinal exponentiation and basic set construction principles, one can show that if $\alpha \leq \gamma$ and $\beta \leq \delta$ are cardinals, then $|{}^\alpha\beta| \leq |{}^\gamma\delta|$. The ABCD hypothesis is the aesthetically simplest behavior for cardinal exponentiation which asserts that this is the only relationship between infinite cardinal exponentiations: For any cardinals $\omega \leq \alpha \leq \beta$ and $\omega \leq \gamma \leq \delta$, $|{}^\alpha\beta| \leq |{}^\gamma\delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$.

The axiom of determinacy AD asserts every two-player infinite game on ω has a winning strategy for one of the two players. AD^+ is Woodin's extension of AD. Let Θ be the supremum of the ordinals onto which $\mathbb{R}$ surjects. Determinacy axioms govern an important initial segment of the universe consisting of the sets onto which $\mathbb{R}$ surjects and Θ is the height of this initial segment.

Assume AD^+ . The ABCD hypothesis holds below Θ : For any cardinals $\omega \leq \alpha \leq \beta < \Theta$ and $\omega \leq \gamma < \delta < \Theta$, $|{}^\alpha\beta| \leq |{}^\gamma\delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$. This provides a complete classification of the relationship between any pair of infinite cardinal exponentiations below Θ .

If κ is a cardinal, then let $\mathcal{P}_B(\kappa) = \{A \in \mathcal{P}(\kappa) : \sup(A) < \kappa\}$ be the set of bounded subsets of κ . The previous result follows from the eponymous result of the paper: Assume AD^+ . For all cardinals $\omega < \kappa < \Theta$ and $\epsilon < \kappa$, $\mathcal{P}_B(\kappa)$ does not inject into ${}^\epsilon\text{ON}$, the class of ϵ -sequences of ordinals.

1. INTRODUCTION

The set of natural numbers is denoted ω . A distinguished characteristic of ω and each finite natural number is the ability to make definitions by recursion and proofs by induction. This property is equivalent to the usual ordering on ω being a wellordering. Using recursion, one defines the usual arithmetic operations on ω including addition, multiplication, and exponentiation. Of particular significance here is the exponentiation operation. If m and n are natural numbers, let ${}^m n$ be the set of all functions from an m -element set into an n -element set. By arithmetic considerations, the usual exponentiation n^m is the size of the set ${}^m n$ and in particular the set ${}^m n$ is finite and wellorderable. The birth of set theory began with Cantor's investigation of exponentiation for infinite sets.

Cantor formalized the notion of size for arbitrary infinite sets. If X and Y are two sets, then X and Y have the same cardinality if and only if there is a bijection from X to Y . An equivalence class of the bijection equivalence relation is called a cardinality. If X is a set, then the cardinality of X , denoted $|X|$, is the bijection equivalence class of X . Cardinalities are compared by the injection relation. $|X| \leq |Y|$ if and only if there is an injection of X into Y . $|X| < |Y|$ if and only if $|X| \leq |Y|$ and $\neg(|Y| \leq |X|)$.

The infinite analog of the natural numbers are sets which possess a form of transfinite induction and recursion or equivalently sets endowed with wellorderings. Ordinals are transitive sets α so that its own membership relation $\in$ defines a wellordering. Ordinals are sets with a natural intrinsic wellordering and every wellordering is order isomorphic to an ordinal. Each natural number does not inject into any smaller natural number, but this is not true of all ordinals. A cardinal is an ordinal which does not inject into any smaller ordinal. Cardinals are natural infinite analog of sizes which possess transfinite recursion and induction. If a set X is wellorderable, then $|X|$ has a unique member which is a cardinal. (Thus, if the axiom of choice AC holds, then every cardinality has a cardinal. The investigation of cardinality under AC is tantamount to the study of cardinals.)

For any two set X and Y , ${}^X Y$ is the set of all functions f from X into Y which can be regarded as the general exponentiation of sets. Cantor proved the first theorem of set theory that X does not surject into ${}^X 2$ and thus $|X| < |{}^X 2|$. The behavior of cardinal exponentiation has received substantial investigation by

set theorists. However, there are very few relationships between infinite cardinal exponentiations which can be inferred solely from the definition of cardinal exponentiation and very basic set construction principles:

- (1) For any cardinal κ , ${}^0\kappa = \{\emptyset\} = 1$.
- (2) For any cardinal $\kappa > 0$, ${}^\kappa 0 = \emptyset = 0$.
- (3) For any cardinal κ , $|{}^\kappa 1| = |1|$.
- (4) For any cardinals $2 \leq \delta \leq \kappa$ with κ infinite, $|{}^\kappa \delta| = |\mathcal{P}(\kappa)| = |{}^\kappa 2| = |{}^\kappa \kappa|$.
- (5) For any cardinals α , β , γ , and δ , if $\alpha \leq \gamma$ and $\beta \leq \delta$, then $|{}^\alpha \beta| \leq |{}^\gamma \delta|$.

Exponentiations of finite natural numbers have an intrinsic arithmetic reason for being finite and hence wellorderable. There seems to be no natural way to infer the wellorderability of infinite cardinal exponentiation from the definition of cardinal exponentiation and basic set theoretic construction principles. Cohen [22] formalized this by showing that the Zermelo-Frankel set theory axiom (ZF) cannot prove $\mathbb{R}$ or ω_2 is wellorderable. Thus the aesthetically simplest behavior for infinite cardinal exponentiation compatible with the necessary properties which can be logically inferred from the definition of cardinal exponentiation is called the ABCD hypothesis which states the above behavior completely classifies infinite cardinal exponentiation.

- (ABCD hypothesis) For all cardinals $\omega \leq \alpha \leq \beta$ and $\omega \leq \gamma \leq \delta$, $|{}^\alpha \beta| \leq |{}^\gamma \delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$.

The ABCD hypothesis does provide a complete classification of infinite cardinal exponentiation in the following sense: given any four infinite cardinals α , β , γ , and δ , the ABCD hypothesis and property (4) can be used to determine the injection relation between ${}^\alpha \beta$ and ${}^\gamma \delta$. (See Remark 6.10.) If $\omega \leq \alpha \leq \beta$ and $\omega \leq \gamma \leq \delta$ are cardinals so that $\alpha < \gamma$ and $\delta < \beta$, then the ABCD hypothesis would imply that $|{}^\alpha \beta|$ and $|{}^\gamma \delta|$ are incomparable cardinalities. Thus the ABCD hypothesis implies the failure of the axiom of choice.

If one assumes the axiom of choice, then for any cardinal κ , ${}^\kappa 2$ is wellorderable and hence is in bijection with a unique cardinal which is frequently denoted 2^κ . The cardinal successor κ^+ is the least cardinal which does not inject into κ . By Cantor's theorem, $\kappa < 2^\kappa$ and thus the smallest possible value for 2^κ is κ^+ . Under the axiom of choice, the aesthetically minimal behavior for infinite cardinal exponentiation is the generalized continuum hypothesis (GCH) which asserts that for all infinite cardinals κ , $2^\kappa = \kappa^+$. However, the argument for the aesthetic minimality of the generalized continuum hypothesis assumes the wellorderability of infinite cardinal exponentiation which seems to be an arbitrary judgement as it cannot be naturally inferred from the definition of cardinal exponentiation and basic set construction principles. In contrast, the failure of the axiom of choice from the ABCD hypothesis is a necessary consequence of the aesthetic minimality considerations. The ABCD hypothesis is a more purely minimal behavior for infinite cardinal exponentiation than the generalized continuum hypothesis from an aesthetic view.

The author does not know if the (global) ABCD hypothesis is consistent. If κ is a cardinal, the ABCD hypothesis below κ is the assertion that for all cardinals $\omega \leq \alpha \leq \beta < \kappa$ and $\omega \leq \gamma \leq \delta < \kappa$, $|{}^\alpha \beta| \leq |{}^\gamma \delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$. The ABCD hypothesis below ω is vacuous. The ABCD hypothesis below ω_1 is trivial since the only infinite exponentiation below ω_1 is ${}^\omega \omega$. The first nontrivial instance is the ABCD hypothesis below ω_2 which amounts to showing $|{}^\omega \omega_1| < |{}^{\omega_1} \omega_1|$. Let Θ be the supremum of the ordinals onto which $\mathbb{R}$ surjects. The descriptive set theoretic world can be intuitively regarded as an initial segment of the universe consisting of transitive sets which are images of $\mathbb{R}$. Θ is the ordinal height of the descriptive set theoretic world. $\mathbb{R}$ always surjects onto ω_1 and thus ω_2 is the smallest possible value for Θ . An important instance is whether the ABCD hypothesis holds in the descriptive set theoretic world or namely below Θ . Theorem 3.4 will show that $|{}^\omega \omega_1| < |{}^{\omega_1} \omega_1|$ holds in the Solovay model ([54]) where $\Theta = \omega_2$. Thus the consistency of ZFC and the existence of an inaccessible cardinal implies the consistency of the ABCD hypothesis below $\Theta = \omega_2$.

The classification of cardinal exponentiation is merely one of many other interesting questions of infinitary combinatorics. Rather than investigating consistency strength of the ABCD hypothesis, this paper is more concerned with showing the ABCD hypothesis is a consequence of a practically useful theory that can serve as a foundational framework for many other combinatorial questions. Intuitively, one would like to show that the ABCD hypothesis holds in an idealized universe which possesses all the regularity properties. Descriptive set theory provides an opinion on the nature of this ultimate regular universe at least up to the initial segment of sets which are images of $\mathbb{R}$ (the descriptive set theoretic world). Classical regularity properties include the perfect set property, the property of Baire, and Lebesgue measurability. A particularly prominent regularity property is the axiom of determinacy, AD, which states that all infinite two player games

of perfect information with moves from ω have a winning strategy for one of the two players. AD^+ is Woodin's extension of the basic axiom of determinacy, AD . It is open whether AD and AD^+ are equivalent. AD^+ has proven to be one of the most important theory in the modern development of set theory (with or without the axiom of choice) and is central to many aspects of the inner model theory program, Woodin's Ultimate-L, forcing axioms, and descriptive set theory.

The inclusion of AD^+ among the laws governing the ultimate regular universe empirically seems to yield some remarkably fascinating structure to the descriptive set theoretic world. Many classically pathological or paradoxical sets can no longer exist including wellorderings of the reals, Lebesgue nonmeasurable sets, Banach-Tarski decompositions, Suslin lines ([9]), and maximal almost disjoint families ([53], [20]). It seems empirically that AD^+ is very successful at resolving natural combinatorial questions and independence phenomenon among these natural problems are quite rare. The method of forcing is the most powerful technique for establishing independence results. By the results of the Chan-Jackson ([12]) and Ikegami-Trang ([26]), any nontrivial forcing which is an image $\mathbb{R}$ over a model of AD must destroy AD in the forcing extension. Intuitively, the descriptive set theoretic world satisfying AD is quite fragile to the direct application of nontrivial forcings which are themselves in the descriptive set theoretic world. This perhaps explains the empirical rarity of independence phenomenon for natural combinatorial questions under determinacy assumptions. The classification of cardinal exponentiation below Θ seems to belong to the class of natural problems that determinacy should be able to resolve.

The origin of the ABCD hypothesis came from neither philosophical nor aesthetic considerations. Prior to the formalization, many incremental cardinality computations had slowly suggested the possibility. Woodin ([65]) had known $|\omega_1|^\omega < |\omega_1|^{<\omega_1}$ and $\neg(|S_1| \leq |\omega_{\omega_1}|)$ at least under $\text{AD}_{\mathbb{R}}$ and DC . Jackson and the author ([13]) showed that AD , $\text{DC}_{\mathbb{R}}$, and all subsets of $\mathbb{R}$ have ∞ -Borel codes imply S_1 does not inject into ${}^\omega\text{ON}$ and hence $|\omega_1|^{<\omega_1}$ does not inject into ${}^\omega\text{ON}$. The methods of [13] uses heavily ∞ -Borel codes for subsets of $[\omega_1]^{<\omega_1}$ (which does follow from all subsets of $\mathbb{R}$ having ∞ -Borel codes) and that generic filters exist for countable forcings that belong to an inner model satisfying ZFC . To generalize this proof beyond ω_1 , one would need the existence of generic filters over an inner model satisfying ZFC for a forcing which is uncountable in the real world and the existence of ∞ -Borel codes for sets of subsets of uncountable cardinals. This first proof known to the author from [13] will be the most suitable for generalization in this paper but at the time, both problems seemed intractable. Generally generic filters for uncountable forcings cannot exist. Woodin showed AD^+ implies there are subsets of $\mathcal{P}(\omega_1)$ which do not have ∞ -Borel codes (see [15]). The goal for the next several years was to find forcing-free and ∞ -Borel code-free proofs that $|\omega_1|^\omega < |\omega_1|^{<\omega_1}$ which could generalize to cardinals beyond ω_1 . Jackson and the author ([11]) investigated almost everywhere continuity properties for functions of the form $\Phi : [\omega_1]^\omega \rightarrow \omega_1$. Since continuity is a well controlled failure of injectivity, this continuity result was used to show that AD implies $|\omega_1|^{<\omega_1} < |\mathcal{P}(\omega_1)|$. Motivated by this almost everywhere continuity result, Jackson, Trang, and the author ([19]) established an almost everywhere continuity property for functions of the form $\Phi : [\omega_1]^\epsilon \rightarrow \omega_1$ when $\epsilon < \omega_1$. This was used in [19] to give the first known AD proof that $|\omega_1|^\omega < |\omega_1|^{<\omega_1}$. [19] also established an analogous almost everywhere continuity result for functions of the form $\Phi : [\omega_2]^\epsilon \rightarrow \omega_2$ when $\epsilon < \omega_2$. This continuity result was used to show $|\omega_2|^\omega < |\omega_2|^{<\omega_2}$ in [19] which was the first instance of these computations beyond ω_1 . Using the cardinality computation $|\omega_1|^\omega < |\omega_1|^{<\omega_1}$, [18] showed that AD and $\text{DC}_{\mathbb{R}}$ prove $|\omega_1|^{<\omega_1}$ does not inject into ${}^\omega\text{ON}$, the class of ω -sequences of ordinals. However, the proof in [18] used very specific properties of ω_1 that could not be generalized to ω_2 . [21] then established an almost everywhere continuity property for function of the form $\Phi : [\kappa]^\epsilon \rightarrow \text{ON}$ when $\epsilon < \kappa$, $\text{cof}(\epsilon) = \omega$, and κ satisfies the partition relation $\kappa \rightarrow_* (\kappa)_2^{\epsilon, \epsilon}$. Using this result, [18] showed that if κ satisfies the weak partition property $\kappa \rightarrow_* (\kappa)_2^{<\kappa}$, then for all $\epsilon < \kappa$, $[\kappa]^{<\kappa}$ does not inject into ${}^\epsilon\text{ON}$, the class of ϵ -sequences of ordinals. With this result, these cardinality computations had now been extended to all the familiar weak and strong partition cardinal of determinacy including ω_1 , ω_2 , $\omega_{\omega+1}$, $\omega_{\omega+2}$, the projective ordinals δ_n^1 (for $n \in \omega$), and δ_1^2 . Jackson, Trang, and the author had enough evidence to formulate the seemingly plausible ABCD conjecture which would completely classify cardinal exponentiation below Θ (with the name of the conjecture due to Jackson and Trang):

- (ABCD Conjecture) AD^+ proves that for all cardinals $\omega \leq \alpha \leq \beta < \Theta$ and $\omega \leq \gamma \leq \delta < \Theta$, $|\alpha|^\beta \leq |\gamma|^\delta$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$.

The statement of the ABCD Conjecture is optimal. AD^+ cannot prove the ABCD hypothesis beyond Θ . If AD holds and $V = L(\mathbb{R})$, then for any $\kappa \geq \Theta$, $\kappa^+ = (\kappa^+)^{\text{HOD}}$ (see discussion in the proof of [20] Theorem 6.3) and thus there is even an injection of Θ^+ into $\mathcal{P}(\Theta)$.

The key to solving the ABCD conjecture is to generalize the phenomenon from [21] that $[\kappa]^{<\kappa}$ does not inject into ${}^\epsilon\text{ON}$ when $\epsilon < \kappa$ and κ is a weak partition cardinal. At a singular cardinal, it was not clear if this is the correct formulation. Let $\mathcal{P}_B(\kappa) = \{A \in \mathcal{P}(\kappa) : \sup(A) < \kappa\}$ be the set of bounded subsets of κ . If κ is regular, then $||[\kappa]^{<\kappa}| = |\mathcal{P}_B(\kappa)|$. However, if κ is singular, then $|\mathcal{P}_B(\kappa)| < ||[\kappa]^{<\kappa}|$ (which follows from Theorem 2.23). The following conjecture is the key insight:

- (Bounded Power Set Conjecture) AD^+ proves that for all cardinal κ with $\omega < \kappa < \Theta$ and $\epsilon < \kappa$, $\mathcal{P}_B(\kappa)$ does not inject into ${}^\epsilon\text{ON}$.

The following is the eponymous main theorem of the paper:

- (Theorem 6.7) The Bounded Power Set Conjecture is true.

From the Bounded Power Set Conjecture, one obtains the following:

- (Theorem 6.9) The ABCD Conjecture is true.

The proof of the Bounded Power Set Conjecture will require both the Steel-Woodin inner model theoretic analysis of HOD-type models as a direct system of certain iterable mice ([58], [64]) and Woodin's ∞ -Borel code analysis of HOD-type models. The Bounded Power Set Conjecture is proved first in AD and $V = L(\mathbb{R})$ in Theorem 6.2 where both the inner model theoretic and ∞ -Borel code analysis of the relevant HOD-type model are much simpler. This should provide most of the main ideas and be accessible to those familiar with the inner model theoretic analysis of $\text{HOD}^{L(\mathbb{R})}$ from [60]. The proof of the Bounded Power Set Conjecture under AD^+ will require an analysis of HOD-type models obtained from Suslin-coSuslin reflection, coarse Woodin mice ([62]), their iteration strategies, and directed system of hybrid structures ([48], [49]). The ∞ -Borel code analysis of these HOD-type structures will also be more challenging.

The Nairian models constructed by Blue, Larson, and Sargsyan ([2], [3]) have already been shown to possess on a global scale some of the determinacy features which are used as ingredients of the proof in this paper. There seems to be great promise that the consistency of the (global) ABCD hypothesis could be established by adapting the framework here globally within the Nairian models.

The ABCD conjecture and the Bounded Power Set conjecture merely distinguishes the cardinality of two sets. The existence and nonexistence of injections is the most primitive question of cardinality or infinitary combinatorics. There are more sophisticated combinatorial properties of cardinalities and sets can be endowed with additional mathematical structure. The class of ω -sequences of ordinals corresponds to the ω -exponent cardinal exponentiation and is the simplest nonwellorderable cardinal exponentiation. Many additional combinatorial properties have been established for set of ω -sequences of ordinals under determinacy assumptions.

If κ satisfies $\kappa \rightarrow_* (\kappa)_2^{\omega+\omega}$, then ${}^\omega\kappa$ does not inject into $\mathcal{P}(\delta) \times \text{ON}$ for any $\delta < \kappa$. The conjecture has been that this should be true of all cardinals $\kappa < \Theta$ under AD^+ . For any cardinal κ , let $B(\omega, \kappa)$ be the set of functions $f : \omega \rightarrow \kappa$ which are bounded below κ . Note that $B(\omega, \kappa) \subseteq {}^\omega\kappa$. If $\text{cof}(\kappa) > \omega$, then $B(\omega, \kappa) = {}^\omega\kappa$. Fact 6.11 will show that under AD^+ , if $\kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then $|B(\omega, \kappa)| < |{}^\omega\kappa|$. The methods developed to study the ABCD conjecture will resolve the above conjecture.

- (Theorem 7.19) Assume AD^+ . For all cardinals κ with $2 \leq \kappa < \Theta$ and all $\chi < \kappa$, $B(\omega, \kappa)$ does not inject into $\mathcal{P}(\chi) \times \text{ON}$. In particular, for all cardinals κ with $2 \leq \kappa < \Theta$ and all $\chi < \kappa$, ${}^\omega\kappa$ does not inject into $\mathcal{P}(\chi) \times \text{ON}$.

Jackson, Trang, and the author ([17]) developed a notion of regularity and cofinality in the choiceless framework and calibrated the cofinality of some familiar sets under AD. If X is a set and Y is a class, X has Y -regular cardinality if and only if for any function $\Phi : X \rightarrow Y$, there is a $y \in Y$ so that $|\Phi^{-1}[\{y\}]| = |X|$. [17] shows that regular cardinals, $\mathbb{R}$, and $\mathbb{R}/E_0$ have globally regular cardinality and provided substantial evidence for the conjecture that $\mathcal{P}(\omega_1)$ is also globally regular. Much is also known about the cofinality of ω -sequences through certain cardinals. If $\text{cof}(\kappa) > \omega$, then ${}^\omega\kappa = B(\omega, \kappa)$ does not have $\text{cof}(\kappa)$ -regular cardinality under AD^+ . If $\kappa \rightarrow_* (\kappa)_2^{\omega+\omega}$ holds, then ${}^\omega\kappa = B(\omega, \kappa)$ is δ -regular for all $\delta < \kappa = \text{cof}(\kappa)$. Using the strong partition property $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and Martin's ultrapower representation of ω_{n+1} , [17] can show that ${}^\omega(\omega_{n+1})$ has ω_1 -regular cardinality but not ω_2 -regular cardinality when $1 \leq n < \omega$. (Martin showed

cof(ω_{n+1}) = ω_2 for all $1 \leq n < \omega$.) Upon some reflection, when $\text{cof}(\kappa) = \omega$, there seems to be no witness to ${}^\omega\kappa$ failing any form of ordinal regularity (in analog to ${}^\omega\omega$ being ON-regular and even globally regular). The conjecture seems to be that under AD^+ , if $\kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then ${}^\omega\kappa$ has ON-regular cardinality and if $\omega < \kappa < \Theta$, then $B(\omega, \kappa)$ does not have $\text{cof}(\kappa)$ -regular cardinality but does have δ -regular cardinality for all $\delta < \text{cof}(\kappa)$. For cardinals κ of countable cofinality, this will be fully verified.

- (Theorem 7.29) Assume AD^+ . If $\kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then ${}^\omega\kappa$ has ON-regular cardinality. If $\omega < \kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then $B(\omega, \kappa)$ does not have ω -regular cardinality but does have n -regular cardinality for each $n < \omega$.

Some time after the author had proved Theorem 7.29, Jackson, Trang, and the author ([17]) showed that ${}^\omega(\omega_\omega)$ has ON-regular cardinality under AD and $\text{DC}_\mathbb{R}$ using combinatorial and descriptive set theoretic arguments.

To address the full conjecture concerning ordinal regularity for ω -sequences of ordinals, it remains to consider $B(\omega, \kappa) = {}^\omega\kappa$ when $\text{cof}(\kappa) > \omega$. Steel ([58], [60]) showed that if AD and $V = L(\mathbb{R})$ hold, then for any $\delta < \Theta$, $\text{cof}^{L(\mathbb{R})}((\delta^+)^{\text{HOD}}) = \omega$. In the arguments, one will roughly need ω -cofinal sequences through the successor and double successor in HOD of $\text{cof}(\kappa)$ -many ordinals below κ . The Uniform Cofinality of HOD Successor and Double Successor Hypothesis (see Definition 7.30) roughly states the desired sequence of ω -cofinal maps can be found. The hypothesis seems quite plausibly to be a consequence of AD^+ . When $\text{cof}(\kappa) = \omega$, the hypothesis follows from the appropriate form of countable choice available under AD. The author can verify a substantial instance of the hypothesis for κ with $\text{cof}(\kappa) = \omega_1$ using a simple linear iteration argument and the direct system analysis of $\text{HOD}^{L(\mathbb{R})}$ presented in [60]. In discussion with Schlutzenberg, it seems promising that Schlutzenberg's method of normalization can establish the hypothesis in full generality in $L(\mathbb{R})$. Assuming an appropriate form of the hypothesis, one can fully resolve the extent of ordinal regularity for the exponent ω .

- (Theorem 7.33) Assume AD^+ and the uniform cofinality of HOD successor and double successor hypothesis. Let $\omega \leq \kappa < \Theta$ be a cardinal. Then $B(\omega, \kappa)$ has δ -regular cardinality for all $\delta < \text{cof}(\kappa)$.

The methods developed here should fully resolve many other combinatorial question surrounding the ω -exponent cardinal exponentiation under AD^+ : Assuming $\kappa \rightarrow_* (\kappa)_2^{\omega+\omega}$, Jackson, Trang, and the author ([16]) can show that ${}^\omega\kappa$ is prime (see Definition 7.1). Under AD, [16] also shows that for all $n < \omega$, ${}^\omega(\omega_n)$ is prime and ${}^\omega(\omega_\omega)$ is prime. Jackson, Trang, and the author [19] showed that ${}^\omega\kappa$ is Jónsson if $\kappa \rightarrow_* (\kappa)_2^{\omega \cdot \omega}$. The author ([5]) showed there is a four-element basis for the linear orderings on ${}^\omega\kappa$ under order embeddings if $\kappa \rightarrow_* (\kappa)_2^{\omega+\omega}$. For $1 \leq m < \omega$, let $p(m)$ be the set of prelinear orderings on m (which is also known as the Fubini or ordered Bell number of m). Let $c(1) = 1$ and $c(m) = p(m-1)$ for all $2 \leq m < \omega$. The author [7] showed that $c(m)$ is the optimal coloring number for partitions of m -element subsets of ${}^\omega\kappa$ when $\kappa \rightarrow_* (\kappa)_2^{\omega \cdot \omega}$: For every $1 \leq r < \omega$ and $P : \mathcal{P}^m({}^\omega\kappa) \rightarrow r$, there is an $A \subseteq {}^\omega\kappa$ with $|A| = |\omega\kappa|$ so that $|P[\mathcal{P}^m(A)]| \leq c(m)$ and there is a $Q : \mathcal{P}^m({}^\omega\kappa) \rightarrow c(m)$ so that for all $A \subseteq {}^\omega\kappa$, if $|A| = |\omega\kappa|$, then $|Q[\mathcal{P}^m(A)]| = c(m)$. The extensions of these questions to all cardinals below Θ will be addressed in forthcoming work.


The author would like to thank Steve Jackson and Nam Trang for patiently listening to many presentations of these results and for their helpful comments. The author would also like to thank Takehiko Gappo, Farmer Schlutzenberg, and Benjamin Siskind for many helpful discussions concerning the inner model theoretic tools and results used here.

2. BACKGROUND

Let $\dot{\in}$ be a binary relation symbol. The language of set theory is $\mathcal{L}_{\text{set}} = \{\dot{\in}\}$. ZF is the $\mathcal{L}_{\text{set}}$ -theory of Zermelo-Frankel set theory (without the axiom of choice). ZFC will denote ZF with the axiom of choice, AC. The base theory of this paper will be ZF (and the axiom of choice will never be assumed in the real world). All other set theoretical assumptions beyond ZF will be made explicit. However, the most important tools will involve inner models (of the choiceless real world) satisfying ZFC and their forcing extensions.

Definition 2.1. If X is a set and $A \subseteq {}^\omega X$, consider the game G_A^X in which two players alternate taking turns playing elements $x_0, x_1, x_2, x_3, x_4, \dots$ from X which jointly produces an element $f \in {}^\omega X$ after ω -many moves. Player 1 is said to win this run of G_A^X if and only if $f \in A$. If one of the two players has a winning strategy then the game G_A^X is said to be determined. AD_X is the assertion that for all $A \subseteq {}^\omega X$, G_A^X is

determined.



AD will abbreviate AD_ω which is called the axiom of determinacy. ${}^\omega\omega$ will often be refer to as $\mathbb{R}$. $\text{AD}_\mathbb{R} = \text{AD}_{\omega_\omega}$ is the axiom of real determinacy.

Capturing sets by inner models will be a very important notion throughout this paper. If A is a set and M is an inner model, M captures A if and only if $A \cap M \in A$. Here, one will be most interested in capturing functions; however, capturing merely the graph of a function will not be sufficient. If $\Phi : X \rightarrow Y$ is a function, then let $G_\Phi = \{(x, y) \in X \times Y : \Phi(x) = y\}$ be the graph of Φ . M is closed under Φ if and only if for all $x \in X \cap M$, $\Phi(x) \in M$. M function-captures Φ if and only if M is closed under Φ and M captures G_Φ . If M function-captures Φ , then $\Phi \upharpoonright X \cap M = \{(x, \Phi(x)) : x \in X \cap M\} \in M$. If $\Phi : X \rightarrow \mathcal{P}(\delta)$ where $\delta \in \text{ON}$, then define the strong graph of Φ to be $\text{Gr}_\Phi = \{(x, \alpha) \in X \times \delta : \alpha \in \Phi(x)\}$. If M captures Gr_Φ , then M will function-capture Φ .

The following result is the basic template with the most elementary form of the notion of function-capturing.¹ In this paper, one will frequently consider inner models of ZFC to prove theorems from certain hypothesis. To help distinguish between these inner models and the background universe, the terms “true universe” or “real world” will often refer to the background universe (which typically do not satisfy choice) in which the hypothesis of the theorem being proved holds.

If κ is a cardinal, $\mathcal{P}_B(\kappa) = \{A \in \mathcal{P}(\kappa) : \sup(A) < \kappa\}$.

Theorem 2.2. *Assume κ is a cardinal such that for all $\epsilon < \kappa$, $\neg(|\kappa| \leq |\mathcal{P}(\epsilon)|)$. Then $\neg(|[\kappa]^{\text{cof}(\kappa)}| \leq |\mathcal{P}_B(\kappa)|)$ and therefore for all $\gamma \leq \kappa$ with $\text{cof}(\gamma) = \text{cof}(\kappa)$, $\neg(|[\kappa]^\gamma| \leq |\mathcal{P}_B(\kappa)|)$ and $|\mathcal{P}_B(\kappa)| < |\mathcal{P}(\kappa)|$. If κ is a regular cardinal, then $||\kappa|^{<\kappa}| < |\mathcal{P}(\kappa)|$.*

Proof. Let $\rho : \text{cof}(\kappa) \rightarrow \kappa$ be an increasing cofinal function. Suppose there is an injection $\Phi : [\kappa]^{\text{cof}(\kappa)} \rightarrow \mathcal{P}_B(\kappa)$. Let $\text{Gr}_\Phi = \{(f, \alpha) : \alpha \in \Phi(f)\}$. The relative Gödel constructible universe $L[\rho, \text{Gr}_\Phi]$ is an inner model of ZFC and $L[\rho, \text{Gr}_\Phi] \models \text{cof}(\kappa) = \text{cof}(\kappa)^V$ as witnessed by ρ . For all $\epsilon < \kappa$, $L[\rho, \text{Gr}_\Phi] \models 2^\epsilon < \kappa$. To see this, suppose otherwise that $L[\rho, \text{Gr}_\Phi] \models \kappa \leq 2^\epsilon$. Then there would be an $h \in L[\rho, \text{Gr}_\Phi]$ such that $L[\rho, \text{Gr}_\Phi] \models “h : \kappa \rightarrow \mathcal{P}(\epsilon) \text{ is an injection}”$ and h would still be an injection in the true universe by absoluteness in violation of the hypothesis. Thus $L[\rho, \text{Gr}_\Phi] \models |\mathcal{P}_B(\kappa)| = 2^{<\kappa} = \kappa$. $L[\rho, \text{Gr}_\Phi] \models |[\kappa]^{\text{cof}(\kappa)}| = \kappa^{\text{cof}(\kappa)} > \kappa$ by a basic result of cardinal arithmetic in ZFC ([31] Corollary 5.14). Using comprehension within $L[\rho, \text{Gr}_\Phi]$, let $\tilde{\Phi}$ be defined by $\tilde{\Phi}(f) = \{\alpha \in \kappa : \text{Gr}_\Phi(f, \alpha)\}$ for each $f \in [\kappa]^{\text{cof}(\kappa)}$. Note that $\tilde{\Phi} = \Phi \upharpoonright ([\kappa]^{\text{cof}(\kappa)})^{L[\rho, \text{Gr}_\Phi]}$ and by absoluteness, $L[\rho, \text{Gr}_\Phi] \models \tilde{\Phi} : [\kappa]^{\text{cof}(\kappa)} \rightarrow \mathcal{P}_B(\kappa)$ is an injection. This is a contradiction since $L[\rho, \text{Gr}_\Phi] \models |[\kappa]^{\text{cof}(\kappa)}| > \kappa = |\mathcal{P}_B(\kappa)|$. $\square$

Next, one will describe some important instances in which the hypothesis of Theorem 2.2 holds. A cardinal κ is a measurable cardinal if and only if there is a κ -complete nonprincipal ultrafilter on κ . The next fact shows that the hypothesis of Theorem 2.2 holds at measurable cardinals.

Fact 2.3. *If κ is a measurable cardinal, then κ does not inject into $\mathcal{P}(\epsilon)$ for any $\epsilon < \kappa$.*

Proof. Let μ be a κ -complete nonprincipal ultrafilter on κ . Suppose $\epsilon < \kappa$ and $\Phi : \kappa \rightarrow {}^\epsilon 2$. For each $\gamma < \epsilon$, there is an $i_\gamma \in 2$ so that $A_\gamma = \{\alpha < \kappa : \Phi(\alpha)(\gamma) = i_\gamma\} \in \mu$. By κ -completeness, $A = \bigcap_{\gamma < \epsilon} A_\gamma \in \mu$. Since μ is nonprincipal, there exist $\alpha_0 < \alpha_1$ with $\alpha_0, \alpha_1 \in A$. Then $\Phi(\alpha_0)(\gamma) = i_\gamma = \Phi(\alpha_1)(\gamma)$ for all $\gamma < \epsilon$. Thus $\Phi(\alpha_0) = \Phi(\alpha_1)$ and hence Φ is not an injection. $\square$

Definition 2.4. If κ is an infinite cardinal, then let boldface GCH at κ be the assertion that there is no injection of κ^+ into $\mathcal{P}(\kappa)$. Say that boldface GCH holds below κ if and only if for all cardinals $\omega \leq \delta < \kappa$, boldface GCH at δ holds.

¹In [13] Fact 4.6, Jackson and the author first used ∞ -Borel codes to obtain the necessary function-capturing to prove $||\omega_1|^{<\omega_1}| < |\mathcal{P}(\omega_1)|$. In discussion with the author some years ago, Neeman observed that this earlier proof only requires function-capturing in a single model and he suggested a simplification which requires only a simple cardinality hypothesis. This simplification is presented in [13] Theorem Fact 4.1 which is the precursor Theorem 2.2.

The axiom of countable choice for $\mathbb{R}$, $\text{AC}_\omega^\mathbb{R}$, is the assertion that for all $R \subseteq \omega \times \mathbb{R}$, there is a uniformization function $F : \text{dom}(R) \rightarrow \mathbb{R}$ for R which means that for all $n \in \text{dom}(R)$, $R(n, F(n))$. A remarkably powerful tool under determinacy is the wellordered additivity of the meager ideal which is precisely stated below. It will be very important in Section 3.

Fact 2.5. *Assuming $\text{AC}_\omega^\mathbb{R}$ and all subsets of $\mathbb{R}$ have the property of Baire. Let $\langle A_\alpha : \alpha < \text{ON} \rangle$ be a sequence such that A_α is a meager subset of $\mathbb{R}$ for all $\alpha \in \text{ON}$. Then $\bigcup_{\alpha \in \text{ON}} A_\alpha$ is a meager subset of $\mathbb{R}$.*

Proof. Suppose $\bigcup_{\alpha \in \text{ON}} A_\alpha$ is a nonmeager subset of ${}^\omega\omega$. By the axiom of replacement, there is a $\xi < \text{ON}$ so that $\bigcup_{\alpha < \xi} A_\alpha = \bigcup_{\alpha \in \text{ON}} A_\alpha$. Let δ be the least ordinals ξ so that $\bigcup_{\alpha < \xi} A_\alpha$ is nonmeager. By the property of Baire, there is an $s \in {}^{<\omega}\omega$ so that $\bigcup_{\alpha < \delta} A_\alpha$ is comeager in $N_s = \{f \in {}^\omega\omega : s \subseteq f\}$. Let $\Phi : N_s \rightarrow {}^\omega\omega$ be a homeomorphism and let $B_\alpha = \Phi[A_\alpha]$ for each $\alpha < \delta$. Then for all $\gamma < \delta$, $\bigcup_{\alpha < \gamma} B_\alpha$ is a meager subset of ${}^\omega\omega$ (by the minimality of δ) and $\bigcup_{\alpha < \delta} B_\alpha$ is a comeager subset of ${}^\omega\omega$. Let $\varphi : \bigcup_{\alpha < \delta} B_\alpha \rightarrow \delta$ be defined by $\varphi(x)$ is the least α so that $x \in B_\alpha$. Define $S \subseteq {}^\omega\omega \times {}^\omega\omega$ by $S(x, y)$ if and only if $x, y \in \bigcup_{\alpha < \delta} B_\alpha$ and $\varphi(x) < \varphi(y)$. For all x in the comeager set $\bigcup_{\alpha < \delta} B_\alpha$, $S_x = \{y \in {}^\omega\omega : S(x, y)\} = (\bigcup_{\alpha < \delta} B_\alpha) \setminus (\bigcup_{\alpha \leq \varphi(x)} B_\alpha)$ is comeager since $\bigcup_{\alpha \leq \varphi(x)} B_\alpha$ is meager. By the Kuratowski-Ulam theorem, S is comeager. For each y in the comeager set $\bigcup_{\alpha < \delta} B_\alpha$, $S^y = \{x \in {}^\omega\omega : S(x, y)\} = \bigcup_{\alpha < \varphi(y)} B_\alpha$ is meager. Thus S is meager by the Kuratowski-Ulam theorem. This is a contradiction. $\square$

Boldface GCH at ω is a particularly common and useful hypothesis which can equivalently be restated as there are no uncountable wellorderable subsets of $\mathbb{R}$.

Fact 2.6. *Assume $\text{AC}_\omega^\mathbb{R}$ and all sets of reals have the Baire property and the perfect set property. Then boldface GCH holds at ω .*

Proof. Suppose $A \subseteq \mathbb{R}$ is an uncountable wellorderable set. By the perfect set property, there is an injection $\Phi : \mathbb{R} \rightarrow A$. Since A is wellorderable, $\mathbb{R}$ is wellorderable and thus 2^ω is the unique cardinal in bijection with $\mathbb{R}$. Let $\langle x_\alpha : \alpha < 2^\omega \rangle$ be a wellordering of $\mathbb{R}$. Since singletons are meager, $\mathbb{R} = \bigcup_{\alpha < 2^\omega} \{x_\alpha\}$ is a wellordered union of meager sets. $\mathbb{R}$ is meager by Fact 2.5 which is a contradiction. $\square$

Actually, the perfect set property for subsets of $\mathbb{R}$ already implies the boldface GCH at ω . To see this, as in the proof of Fact 2.6, the failure of boldface GCH at ω and the perfect set property implies $\mathbb{R}$ is wellorderable. A wellordering on $\mathbb{R}$ can be used to create a subset of $\mathbb{R}$ with the perfect set property.

Boldface GCH at ω implies the following cardinality distinctions.

Fact 2.7. *Assume boldface GCH at ω . Then $\neg(|\mathbb{R}| \leq |\omega_1|)$, $\neg(|\omega_1| \leq |\mathbb{R}|)$, $|\mathbb{R}| < |\mathbb{R} \cup \omega_1|$, $|\omega_1| < |\mathbb{R} \cup \omega_1|$, and $|\mathbb{R} \cup \omega_1| < |\mathbb{R} \times \omega_1|$.*

Proof. All except the last are immediate consequences of boldface GCH at ω . Suppose there is an injection $\Phi : \mathbb{R} \times \omega_1 \rightarrow \mathbb{R} \cup \omega_1$. Let $\pi_1 : \mathbb{R} \times \omega_1 \rightarrow \mathbb{R}$ be the projection onto the first coordinate. $\pi_1[\Phi^{-1}[\omega_1]]$ is a wellorderable subset of $\mathbb{R}$ and thus is countable by boldface GCH at ω . Let $r \in \mathbb{R} \setminus \pi_1[\Phi^{-1}[\omega_1]]$. Then $\Phi \upharpoonright \{r\} \times \omega_1 : \{r\} \times \omega_1 \rightarrow \mathbb{R}$ is an injection. This is essentially an injection of ω_1 into $\mathbb{R}$ which is impossible by boldface GCH at ω . $\square$

Let $\text{AC}_\mathbb{R}^\mathbb{R}$ (or Uniformization)² be the assertion that for every relation $R \subseteq \mathbb{R} \times \mathbb{R}$, there is a function $\Lambda : \text{dom}(R) \rightarrow \mathbb{R}$ so that for all $r \in \text{dom}(R)$, $R(r, \Lambda(r))$. $\text{AC}_\mathbb{R}^\mathbb{R}$ implies that $|\mathbb{R}|$, $|\omega_1|$, $|\mathbb{R} \cup \omega_1|$, and $|\mathbb{R} \times \omega_1|$ are the only uncountable cardinalities below $|\mathbb{R} \times \omega_1|$ (see [13] Corollary 7.6). [13] Theorem 7.33 produced many intermediate cardinalities below $|\mathbb{R} \times \omega_1|$ in $L(\mathbb{R}) \models \text{AD}$ where $\text{AC}_\mathbb{R}^\mathbb{R}$ fails.

An important source of measurable cardinals under AD comes from partition properties. (See [6], [8], [39], [37], and [30] for more information concerning partition properties under determinacy.)

² $\text{AD}_{\frac{1}{2}}$ is the statement asserting one of the two players has a winning strategy in games on $\mathbb{R}$ where one of the two players is restricted to playing moves only from ω . Kechris [36] showed that AD and $\text{AC}_\mathbb{R}^\mathbb{R}$ is equivalent to $\text{AD}_{\frac{1}{2}\mathbb{R}}$. $\text{AD}_{\frac{1}{2}\mathbb{R}}$ follows from $\text{AD}_\mathbb{R}$ but is not known to be equivalent. $\text{AD}_\mathbb{R}$ and $\text{AD}_\mathbb{R} + \text{AD}^+$ are also not known to be equivalent. $\text{AD}_\mathbb{R} + \text{DC}$ is strictly stronger than all the aforementioned theories by results of Solovay [55].

Definition 2.8. Let ϵ be an ordinal. A function $f : \epsilon \rightarrow \text{ON}$ is said to be discontinuous everywhere if and only if for all $\alpha < \epsilon$, $\sup(f \upharpoonright \alpha) < f(\alpha)$. f is said to have uniform cofinality ω if and only if there is a function $F : \epsilon \times \omega \rightarrow \text{ON}$ so that for all $\alpha < \epsilon$ and $n \in \omega$, $F(\alpha, n) < F(\alpha, n+1)$ and $f(\alpha) = \sup\{F(\alpha, n) : n \in \omega\}$. f is said to have the correct type if and only if f is both discontinuous everywhere and has uniform cofinality ω . If $X \subseteq \text{ON}$, then let $[X]_*^\epsilon$ be the collection functions $f : \epsilon \rightarrow X$ which have the correct type.

If $\epsilon \leq \kappa$ and $\gamma < \kappa$, let the correct-type partition relation³ $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ be the assertion that for all $P : [\kappa]_*^\epsilon \rightarrow \gamma$, there is a $\beta < \gamma$ and a club $C \subseteq \kappa$ so that for all $f \in [C]_*^\epsilon$, $P(f) = \beta$. If $\epsilon \leq \kappa$, then $\kappa \rightarrow_* (\kappa)_\gamma^{\leq \epsilon}$ is the assertion that for all $\delta < \epsilon$, $\kappa \rightarrow_* (\kappa)_\gamma^\delta$. For $\gamma \leq \kappa$, $\kappa \rightarrow_* (\kappa)_{<\gamma}^\epsilon$ is the assertion that for all $\beta < \gamma$, $\kappa \rightarrow_* (\kappa)_\beta^\epsilon$. It can be shown that for all $\epsilon < \kappa$, $\kappa \rightarrow_* (\kappa)_{2^{\epsilon+\epsilon}}^\epsilon$ implies $\kappa \rightarrow_* (\kappa)_{<\kappa}^\epsilon$.

For $\epsilon \leq \kappa$, let μ_κ^ϵ be the filter on $[\kappa]_*^\epsilon$ defined by $A \in \mu_\kappa^\epsilon$ if and only if there is a club $C \subseteq \kappa$ so that $[C]_*^\epsilon \subseteq A$. μ_κ^ϵ is called the ϵ -exponent partition filter on κ and note that μ_κ^1 is the ω -club filter on κ . If $\kappa \rightarrow_* (\kappa)_2^\epsilon$, then μ_κ^ϵ is an ultrafilter. For any $\delta < \kappa$, $\kappa \rightarrow_* (\kappa)_\delta^\epsilon$ implies μ_κ^ϵ is a δ^+ -complete ultrafilter. Hence $\kappa \rightarrow_* (\kappa)_{<\kappa}^\epsilon$ implies μ_κ^ϵ is κ -complete. $\kappa \rightarrow_* (\kappa)_2^\epsilon$ implies that the ω -club filter μ_κ^1 is a normal measure on κ . κ is said to be a strong partition cardinal if and only if $\kappa \rightarrow_* (\kappa)_2^\kappa$. κ is said to be a very strong partition cardinal if and only if $\kappa \rightarrow_* (\kappa)_{<\kappa}^\kappa$ (or equivalently μ_κ^κ is κ -complete). κ is said to be a weak partition cardinal if and only if $\kappa \rightarrow_* (\kappa)_{2^{\leq \kappa}}^\kappa$.

The exponent two correct type partition relation on an uncountable cardinal already implies the failure of AC by the next result.⁴

Fact 2.9. Let κ be an uncountable cardinal satisfying $\kappa \rightarrow_* (\kappa)_2^\omega$. Then $[\kappa]^\omega$ is not wellorderable.

Proof. Suppose otherwise that $[\kappa]^\omega$ has a wellordering $\prec$. For each $\alpha \in [\kappa]_*^1$ (that is, $\text{cof}(\alpha) = \omega$), let f_α be the $\prec$ -least $f \in [\kappa]^\omega$ so that $\sup(f) = \alpha$. For each $n \in \omega$, let $\Phi_n : [\kappa]_*^1 \rightarrow \kappa$ be defined by $\Phi_n(\alpha) = f_\alpha(n)$. For all $\alpha \in [\kappa]_*^1$, $\Phi_n(\alpha) < \alpha$. Since $\kappa \rightarrow_* (\kappa)_2^\omega$ implies μ_κ^1 is normal, for each $n \in \omega$, there is a λ_n so that $A_n = \{\alpha \in [\kappa]_*^1 : \Phi_n(\alpha) = \lambda_n\} \in \mu_\kappa^1$. Since $\kappa \rightarrow_* (\kappa)_2^\omega$ implies $\kappa \rightarrow_* (\kappa)_{<\kappa}^1$ and hence the κ -completeness of μ_κ^1 , $A = \bigcap_{n \in \omega} A_n \in \mu_\kappa^1$. Let $g : \omega \rightarrow \kappa$ be defined by $g(n) = \lambda_n$. For all $\alpha \in A$, $f_\alpha = g$. Since μ_κ^1 is a nonprincipal measure, let $\alpha_0, \alpha_1 \in A$ with $\alpha_0 \neq \alpha_1$. Then $\alpha_0 = \sup(f_{\alpha_0}) = \sup(g) = \sup(f_{\alpha_1}) = \alpha_1$. Contradiction. $\square$

Fact 2.10. Suppose κ is an uncountable cardinal, $\omega \leq \epsilon \leq \kappa$, $\delta < \kappa$, and $\kappa \rightarrow_* (\kappa)_\delta^\epsilon$ holds. Then there is no injection of ${}^\epsilon \kappa$ into $\mathcal{P}(\delta) \times \text{ON}$.

Proof. Suppose $\Phi : {}^\epsilon \kappa \rightarrow \mathcal{P}(\delta) \times \text{ON}$ is an injection. Let $\pi_1 : \mathcal{P}(\delta) \times \text{ON} \rightarrow \mathcal{P}(\delta)$ be the projection onto the first coordinate. For each $\beta < \delta$, let $P_\beta : [\kappa]_*^\epsilon \rightarrow 2$ be $P_\beta(f) = 1$ if and only if $\beta \in \pi_1(\Phi(f))$. By $\kappa \rightarrow_* (\kappa)_\delta^\epsilon$, there is an $i_\beta \in 2$ and a club $C \subseteq \kappa$ so that for all $f \in [C]_*^\epsilon$, $P_\beta(f) = i_\beta$. Thus $E_\beta = \{f \in [\kappa]_*^\epsilon : P_\beta(f) = i_\beta\} \in \mu_\kappa^\epsilon$. Since $\kappa \rightarrow_* (\kappa)_\delta^\epsilon$ implies that μ_κ^ϵ is δ^+ -complete, $\bigcap_{\beta < \delta} E_\beta \in \mu_\kappa^\epsilon$. There is a club $D \subseteq \kappa$ so that $[D]_*^\epsilon \subseteq \bigcap_{\beta < \delta} E_\beta$. So for all $f \in [D]_*^\epsilon$ and all $\beta < \delta$, $P_\beta(f) = i_\beta$. Let $K = \{\beta \in \delta : i_\beta = 1\}$. Thus for all $f \in [D]_*^\epsilon$, $\pi_1(\Phi(f)) = K$. Hence $\Phi \upharpoonright [D]_*^\epsilon : [D]_*^\epsilon \rightarrow \{K\} \times \text{ON}$. Since $||[\kappa]^\omega| \leq |[D]_*^\epsilon|$ because $\omega \leq \epsilon$, one has a wellordering of $[\kappa]^\omega$. This is impossible by Fact 2.9. $\square$

If $\epsilon \leq \kappa$, Martin defined a good coding system for ${}^\epsilon \kappa$ ([30] Definition 2.33, [6]) which consists of a pointclass and a method of coding elements of ${}^\epsilon \kappa$ by reals with very strict definability constraints relative to the pointclass. Martin showed that if there is a good coding system for ${}^{\omega \cdot \epsilon} \kappa$, then $\kappa \rightarrow_* (\kappa)_2^\epsilon$ holds. Determinacy with the method of good coding systems is the only known set-theoretical setting with strong partition cardinals. If Γ is a pointclass, then let $\tilde{\Gamma}$ be the dual pointclass of Γ consisting of complements of sets of Γ . Let $\delta(\Gamma)$ is the rank of prewellorderings on $\mathbb{R}$ which belong to $\Delta = \Gamma \cap \tilde{\Gamma}$. The projective ordinal δ_n^1 is defined as $\delta(\Sigma_n^1)$. $\delta_1^2 = \delta(\Sigma_1^2)$. If Γ has the prewellordering property and is closed under $\forall^{\mathbb{R}}$, $\wedge$, and $\vee$, then for $\epsilon < \omega_1$, the natural coding of ${}^\epsilon \delta(\Gamma)$ by a real (or rather countably many reals relative to a wellordering of ω of ordertype ϵ) defines a good coding system for ${}^\epsilon \delta(\Gamma)$. Hence for such Γ , $\delta(\Gamma)$ satisfies the countable exponent partition relation. Jackson's description theory ([29], [28], [30]) is used to produce good

³The ordinary partition relation $\kappa \rightarrow (\kappa)_\gamma^\epsilon$ is the assertion that for all $P : [\kappa]^\epsilon \rightarrow \gamma$, there is a $\beta < \gamma$ and an $A \subseteq \kappa$ with $|A| = |\kappa|$ such that for all $f \in [A]^\epsilon$, $P(f) = \beta$.

⁴The ordinary exponent two partition relation at an uncountable cardinal characterizes a weakly compact cardinal which exists under AC assuming the consistency of a weakly compact cardinal which is a relatively mild large cardinal axiom.

coding systems on the odd projective ordinals δ_{2n+1}^1 . Kechris-Kleinberg-Moschovakis-Woodin [37] showed that if Γ is a highly closed pointclass, then $\delta(\Gamma)$ has a good coding system for $\delta^{(\Gamma)}\delta(\Gamma)$ and hence $\delta(\Gamma)$ is a strong partition cardinal. This result will show δ_1^2 is a strong partition cardinal. For each $A \in \mathcal{P}(\mathbb{R})$, let δ_A be the least δ so that $L_\delta(A, \mathbb{R}) \prec_{\Sigma_1} L(A, \mathbb{R})$ in the language with a binary symbol $\dot{\in}$ and unary symbol $\dot{\mathbb{R}}$. [37] also implies that δ_A is a strong partition cardinal for each $A \in \mathcal{P}(\mathbb{R})$. Besides good coding systems, Martin showed that a combinatorial analysis of the ultrapower of a strong partition cardinal by its measures may sometimes yield partition properties. This is used to show that $\omega_2 = \prod_{\omega_1} \omega_1 / \mu_{\omega_1}^1$ is a weak partition cardinal and more generally the even projective ordinals $\delta_{2n+2}^1 = (\delta_{2n+1}^1)^+$ are weak partition cardinals. The following summarizes the most important partition cardinals of determinacy.

Fact 2.11.

- (1) (Martin; [30] Theorem 2.36) Let Γ be a pointclass with the prewellordering property and closed under $\forall^{\mathbb{R}}$, $\wedge$, and $\vee$. Then $\delta(\Gamma) \rightarrow_* (\delta(\Gamma))_2^{<\omega_1}$.
- (2) (Martin; [34] Theorem 12.2, [30] Corollary 4.16, [8] Section 4; [6]) Assume AD. ω_1 is a very strong partition cardinal.
- (3) (Martin-Paris; [8] Theorem 5.19; [11] Theorem 6.2) Assume AD. ω_2 is a weak partition cardinal but is not a strong partition cardinal.
- (4) (Jackson; [28], [29], [30]) Assume AD. For all $n \in \omega$, δ_{2n+1}^1 is a strong partition cardinal. For all $n \in \omega$, δ_{2n+2}^1 is a weak partition cardinal but is not a strong partition cardinal.
- (5) (Kechris-Kleinberg-Moschovakis-Woodin; [37], [6]) Suppose Γ is a pointclass with the prewellordering property and closed under $\forall^{\mathbb{R}}$, $\wedge$, and $\vee$ and $\Delta = \Gamma \cap \check{\Gamma}$ is closed under $\exists^{\mathbb{R}}$ and $\forall^{\mathbb{R}}$. Then $\delta(\Gamma)$ is a very strong partition cardinal.
- (6) (Kechris-Kleinberg-Moschovakis-Woodin; [37], [6], [42] Theorem 5.8) Assume AD. For all $A \in \mathcal{P}(\mathbb{R})$, δ_A is a very strong partition cardinal.
- (7) (Kechris-Kleinberg-Moschovakis-Woodin; [37], [42] Theorem 5.5) Assume AD and $\text{DC}_{\mathbb{R}}$. δ_1^2 is a very strong partition cardinal.

By Fact 2.3, ω_1 does not inject into $\mathcal{P}(\omega)$ and ω_2 does not inject into $\mathcal{P}(\omega_1)$. This gives another proof that boldface GCH holds at ω and shows boldface GCH holds ω_1 . Using Martin's analysis of $\omega_{n+1} = \prod_{[\omega_1]^n} \omega_1 / \mu_{\omega_1}^n$, for each $n \in \omega$, one has that $\text{cof}(\omega_n) = \omega_2$ for all $2 \leq n < \omega$. (Thus ω_3 is the first singular cardinal of determinacy.) Using this ultrapower analysis, Jackson, Trang, and the author ([14]) can show boldface GCH below ω_ω under AD. These arguments generalize using Jackson's description theory ([30], [29], and [28]) to show boldface GCH below $\sup\{\delta_n^1 : n \in \omega\}$. The following is obtained from Theorem 2.2.

Theorem 2.12. Assume AD.

- (1) $|[\omega_1]^{<\omega_1}| < |\mathcal{P}(\omega_1)|$ and $|[\omega_2]^{<\omega_2}| < |\mathcal{P}(\omega_2)|$. For all $3 \leq n < \omega$, $\neg(|[\omega_n]^{\omega_2}| \leq |\mathcal{P}_B(\omega_n)|)$.
- (2) For all $\kappa < \sup\{\delta_n^1 : n \in \omega\}$, $\neg(|[\kappa]^{\text{cof}(\kappa)}| < |\mathcal{P}_B(\kappa)|)$.
- (3) For all $A \in \mathcal{P}(\mathbb{R})$, $|[\delta_A]^{<\delta_A}| < |\mathcal{P}(\delta_A)|$.
- (4) Assume $\text{DC}_{\mathbb{R}}$, $|[\delta_1^2]^{<\delta_1^2}| < |\mathcal{P}(\delta_1^2)|$.

Note that under AD, $\text{cof}(\omega_3) = \omega_2$ and hence Theorem 2.2 and its proof method cannot show that $|[\omega_3]^{\omega_2}| < |\mathcal{P}(\omega_3)|$ or even $|[\omega_3]^{\omega_2}| < |[\omega_3]^{<\omega_3}|$. These cardinality distinctions will be established in Theorem 6.12 as a consequence of the main theorem of the paper.

Since AD implies $\omega_1 \rightarrow_* (\omega_1)_\omega^\omega$, Fact 2.10 implies the following. In the next section, this cardinality computation will be shown to follow from classical regularity properties by a proof which is more suitable for generalization (see Section 7).

Fact 2.13. Assume AD. ω_ω does not inject to $\mathbb{R} \times \text{ON}$. Thus $|\mathbb{R} \times \omega_1| < |[\omega_1]^\omega| = |\omega_\omega|$.

Forcing is a technique developed by Cohen ([22]) to extend a given model of set theory to larger model of set theory in a constructive and minimal manner. Let $\mathbb{P}$ be a nonempty set. Let $\leq_{\mathbb{P}}$ be a partial ordering on $\mathbb{P}$. Assume $\mathbb{P}$ has a $\leq_{\mathbb{P}}$ -largest element denoted by $1_{\mathbb{P}}$. $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}}, 1_{\mathbb{P}})$ is called a forcing. An element $p \in \mathbb{P}$ is often called a condition of $\mathbb{P}$. A $\mathbb{P}$ -filter (in the sense of forcing) is a set $F \subseteq \mathbb{P}$ satisfying the following properties:

- (1) $1_{\mathbb{P}} \in F$.

- (2) For all $p, q \in \mathbb{P}$, if $p \leq_{\mathbb{P}} q$ and $p \in F$, then $q \in F$.
(3) For all $p_0, p_1 \in \mathbb{P}$, there is a $q \in F$ so that $q \leq_{\mathbb{P}} p_0$ and $q \leq_{\mathbb{P}} p_1$.

If $A \subseteq \mathbb{P}$, then let $\langle A \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists q \in A)(q \leq_{\mathbb{P}} p)\}$ which is the $\leq_{\mathbb{P}}$ -upward closure of A . $D \subseteq \mathbb{P}$ is dense if and only if for all $p \in \mathbb{P}$, there is a $q \in D$ such that $q \leq_{\mathbb{P}} p$. $A \subseteq \mathbb{P}$ is a $\mathbb{P}$ -antichain if and only if for all $p_0, p_1 \in A$, if $p_0 \neq p_1$, then p_0 and p_1 are $\mathbb{P}$ -incompatible which means there is no $q \in \mathbb{P}$ with $q \leq_{\mathbb{P}} p_0$ and $q \leq_{\mathbb{P}} p_1$. A $\mathbb{P}$ -antichain A is maximal if and only if A is an antichain and for all $\mathbb{P}$ -antichain B with $B \supseteq A$, $B = A$. Let $\mathcal{D}$ be a collection of dense subsets of $\mathbb{P}$. A $\mathbb{P}$ -filter G is $\mathbb{P}$ -generic over $\mathcal{D}$ if and only if $D \cap G \neq \emptyset$ for all $D \in \mathcal{D}$. Let M be an inner model of ZFC with $\mathbb{P} \in M$. Let $\mathcal{D}_M$ be the collection of all dense subset of $\mathbb{P}$ in M . Let $\mathcal{E}_M$ be the collection of all maximal antichain of $\mathbb{P}$ in M . A filter G is said to be $\mathbb{P}$ -generic over M if and only if G is $\mathbb{P}$ -generic over $\mathcal{D}_M$ or equivalently G intersects every maximal antichain in $\mathcal{E}_M$. If G is $\mathbb{P}$ -generic over M , then $M[G]$ is the $\mathbb{P}$ -generic extension of M by G . $M[G] \models \text{ZFC}$ and is the minimal model of ZFC containing G . See [31] or [41] for more information on the method of forcing, the explicit construction of the generic extension, and details concerning the examples of forcing which will be used below.

An important application of boldface GCH at ω is the existence of generic filters over inner models of the axiom of choice for forcings which are countable in the true universe. The proof of the next result is a well known argument which will be presented in the following manner to motivate Lemma 5.6 later. The use of this fact will be ubiquitous in this paper.

Fact 2.14. *Assume boldface GCH at ω . Let M be an inner model of ZFC and $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}}, 1_{\mathbb{P}}) \in M$ be a forcing which is countable in the true universe. Then there is a $G \subseteq \mathbb{P}$ which is $\mathbb{P}$ -generic over M .*

Proof. Let $\sigma : \mathbb{P} \rightarrow \omega$ be a bijection. Let $\tau : \mathcal{P}(\mathbb{P})^M \rightarrow \mathcal{P}(\omega)$ be defined by $\tau(A) = \sigma[A]$ and observe that τ is an injection. Since $M \models \text{AC}$, there is an injection $\iota : \mathcal{P}(\mathbb{P})^M \rightarrow \gamma$ for some ordinal γ with $\iota \in M$. Then $\iota \circ \tau^{-1} : \tau[\mathcal{P}(\mathbb{P})^M] \rightarrow \gamma$ is an injection. Thus $\tau[\mathcal{P}(\mathbb{P})^M]$ is a wellorderable subset of $\mathcal{P}(\omega)$ and must be countable in the true universe by boldface GCH at ω . Thus $\mathcal{P}(\mathbb{P})^M$ is countable in the true universe. Let $\mathcal{D}$ be the collection of all dense subsets of $\mathbb{P}$ in M . Since $\mathcal{P}(\mathbb{P})^M$ is countable and $\mathcal{D} \subseteq \mathcal{P}(\mathbb{P})^M$, there is a bijection $\nu : \omega \rightarrow \mathcal{D}$. Note the following important observation:

- (1) Every forcing is $< \omega$ -closed and thus $M \models \text{"}\mathbb{P} \text{ is } < \omega\text{-closed"}$.
(2) For every $n \in \omega$, $\nu \restriction n \in M$ since it is finite.

Since $M \models \text{AC}$, fix a wellordering $\sqsubset$ for $\mathbb{P}$ which belongs to M . By the two observations, one can work within M to define a sequence $\langle p_m^n : m < n \rangle$ so that for all $m < n$, p_m^n is the $\sqsubset$ -least $p \in \mathbb{P}$ so that $p \leq_{\mathbb{P}} p_k^n$ for all $k < m$ and $p \in \nu(m)$. By the uniformity obtained by using $\sqsubset$, for all $m < n_0 \leq n_1$, $p_m^{n_0} = p_m^{n_1}$. For each $m \in \omega$, let $p_m = p_m^{m+1}$. Let $G = \langle \{p_n : n \in \omega\} \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists n \in \omega)(p_n \leq_{\mathbb{P}} p)\}$. Since ν enumerates all dense subsets of $\mathbb{P}$ in M and $\{p_n : n \in \omega\} \subseteq G$, G is $\mathbb{P}$ -generic over M . $\square$

Let Θ be the supremum of the ordinals α for which there is a surjection $\pi : \mathbb{R} \rightarrow \alpha$. The Moschovakis coding lemma is a powerful tool of determinacy proved using the recursion theorem. It can be used to show that if AD holds, M is an inner model of AD with $\mathbb{R} \subseteq M$, and $\kappa < \Theta^M$, then $\mathcal{P}(\kappa)^M = \mathcal{P}(\kappa)$. However, one of its frequently used coarse consequences is that if $\kappa < \Theta$ (that is, $\mathbb{R}$ surjects onto κ), then $\mathbb{R}$ surjects onto $\mathcal{P}(\kappa)$. The proof of various forms of the coding lemma and background concerning pointclasses can be found in [45] Section 7D, [30] Theorem 2.12, [37] Lemma 1.5, and [40] Theorem 3.2 and Theorem 3.4.

Fact 2.15. *(Moschovakis coding lemma; [44]) Assume AD. Let Γ be a nonselfdual pointclass closed under $\exists^{\mathbb{R}}$ and $\wedge$. Let $P \subseteq \mathbb{R}$, $\kappa \in \text{ON}$, and $\varphi : P \rightarrow \kappa$ is a surjection. Define $\prec$ on $\mathbb{R}^2$ by $x \prec y$ if and only if $x, y \in P$ and $\varphi(x) \leq \varphi(y)$. Assume $\prec \in \Gamma$. Let $R \subseteq P \times \mathbb{R}$. Then there is an $S \subseteq R$ with $S \in \Gamma$ so that for all $\alpha < \kappa$, if there exists $x \in P$ with $\varphi(x) = \alpha$ and $x \in \text{dom}(R)$, then there exists $y \in P$ with $\varphi(y) = \alpha$ and $y \in \text{dom}(S)$.*

As a consequence, if $\kappa < \Theta$, there is a surjection of $\mathbb{R}$ onto $\mathcal{P}(\kappa)$.

The following illustrates one use of the Moschovakis coding lemma.

Fact 2.16. *Assume AD. For any $\kappa < \Theta$, $\text{AC}_{\omega}^{\mathcal{P}(\kappa)}$ holds: For any $R \subseteq \omega \times \mathcal{P}(\kappa)$, there is a function $\Lambda : \text{dom}(R) \rightarrow \mathcal{P}(\kappa)$ such that for all $n \in \text{dom}(R)$, $R(n, \Lambda(n))$.*

Proof. Since $\kappa < \Theta$, there is a surjection $\pi : \mathbb{R} \rightarrow \mathcal{P}(\kappa)$ by the Moschovakis coding lemma (Fact 2.15). Given $R \subseteq \omega \times \mathcal{P}(\kappa)$, let $S \subseteq \omega \times \mathbb{R}$ be defined by $S(n, x)$ if and only if $R(n, \pi(x))$. Since AD implies

453 $\text{AC}_{\omega}^{\mathbb{R}}$, there is a $\Sigma : \text{dom}(S) \rightarrow \mathbb{R}$ such that for all $n \in \text{dom}(S)$, $S(n, \Sigma(n))$. Since $\text{dom}(R) = \text{dom}(S)$,
 454 for all $n \in \text{dom}(R)$, $R(n, \pi(\Sigma(n)))$ holds. Thus $\Lambda : \text{dom}(R) \rightarrow \mathcal{P}(\kappa)$ defined by $\Lambda = \pi \circ \Sigma$ is the desired
 455 uniformization function. $\square$

456 Under AC , successor cardinals are regular. This is not true under AD since ω_3 is the least singular cardinal
 457 with $\text{cof}(\omega_3) = \omega_2$. However under AD and $\kappa < \Theta$, $\text{AC}_{\omega}^{\mathcal{P}(\kappa)}$ can recover a fragment of the argument to show
 458 $\text{cof}(\kappa^+) > \omega$.

459 **Fact 2.17.** *Assume AD . If $\omega \leq \kappa < \Theta$, then $\text{cof}(\kappa^+) > \omega$.*

460 The following consequence of boldface GCH will be needed later. If $\kappa \in \text{ON}$ and $\alpha \in \text{ON}$, then $\kappa^{+\alpha}$ will
 461 refer to the α^{th} cardinal greater than κ .

462 **Fact 2.18.** *Assume boldface GCH holds at a cardinal κ . Let $M \models \text{ZFC}$ be an inner model. The cardinals of
 463 M below κ^+ is unbounded in κ^+ . Thus for all $\alpha < \text{cof}(\kappa)$, $(\kappa^{+\alpha})^M < \kappa^+$.*

464 *Proof.* Suppose not. Let $\lambda < \kappa^+$ be the largest ordinal $\delta < \kappa^+$ so that $M \models \text{"}\delta \text{ is a cardinal"}$. This
 465 implies $(\lambda^+)^M = \kappa^+$. Since $\kappa \leq \lambda < \kappa^+$, let $\Phi : \lambda \rightarrow \kappa$ be a bijection. Define $\Psi : (\mathcal{P}(\lambda))^M \rightarrow \mathcal{P}(\kappa)$ by
 466 $\Psi(A) = \Phi[A] = \{\Phi(\alpha) : \alpha \in A\}$. Ψ is an injection. By Cantor's theorem in M , $M \models \text{"}\lambda \text{ does not surject onto}$
 467 $\mathcal{P}(\lambda)\text{"}$. Since $M \models \text{AC}$, M has an injection $\Upsilon : (\lambda^+)^M \rightarrow (\mathcal{P}(\lambda))^M$. Since $\kappa^+ = (\lambda^+)^M$, $\Psi \circ \Upsilon : \kappa^+ \rightarrow \mathcal{P}(\kappa)$
 468 is an injection which violates the boldface GCH at κ . $\square$

469 **Fact 2.19.** *Assume AD . Let $\omega \leq \kappa < \Theta$ and boldface GCH holds at κ . Let $M \models \text{ZFC}$ be an inner model.
 470 Then for all $\alpha < \omega_1$, $(\kappa^{+\alpha})^M < \kappa^+$.*

471 *Proof.* The result follows by Fact 2.17 and Fact 2.18. $\square$

472 Becker and Kechris showed that AD implies $\text{cof}(\kappa^+) > \omega_1$ for all $\omega_1 \leq \kappa < \Theta$ using a simple version of
 473 the Kechris-Woodin generic coding function on ω_1 ([38]). Thus Fact 2.19 can be strengthened to show that
 474 $(\kappa^{+\alpha})^M < \kappa^+$ for all $\alpha < \omega_2$ when $\omega_1 \leq \kappa < \Theta$. One can say much more when $M = \text{HOD}^{L(\mathbb{R})}$. Jackson,
 475 Ketchersid, Schlutzenburg, and Woodin (Fact 7.16; [27]) showed that for all $\omega_1 \leq \kappa < \Theta^{L(\mathbb{R})}$, there are κ^+
 476 many cardinals (even measurable cardinals) of $\text{HOD}^{L(\mathbb{R})}$ below κ^+ .

477 **Definition 2.20.** An ∞ -Borel code is a pair $p = (S, \varphi)$ where S is a set of ordinals and φ is a first order
 478 $\mathcal{L}_{\text{set}}$ -formula with two free variables. Let $\mathcal{B}_{\infty}$ denote the set of all ∞ -Borel codes. Let X be a set of sets of
 479 ordinals. Let $\mathfrak{B}_p^X = \{x \in X : L[S, x] \models \varphi(S, x)\}$. A set $A \subseteq X$ is ∞ -Borel if and only if there is an ∞ -Borel
 480 code p so that $A = \mathfrak{B}_p^X$.

481 ∞ -Borel codes provide the most absolute and uniform way to capture a set: Suppose $A \subseteq \mathbb{R}$ and (S, φ) is
 482 an ∞ -Borel code for A . Then any inner model M of ZF with $S \in M$ can capture A by using comprehension
 483 and the observation that $x \in A$ if and only if $L[S, x] \models \varphi(S, x)$.

484 The following is Woodin's theory AD^+ which is a very powerful extension of determinacy.

485 **Definition 2.21.** (Woodin) AD^+ is the conjunction of the following three statements.

- 486 (1) $\text{DC}_{\mathbb{R}}$ (dependent choice for $\mathbb{R}$).
- 487 (2) For all $\kappa < \Theta$, continuous function $\pi : {}^\omega \kappa \rightarrow {}^\omega \omega$, and $A \subseteq {}^\omega \omega$, the game $G_{\pi^{-1}[A]}^{\kappa}$ is determined.
- 488 (3) For all $A \subseteq {}^\omega \omega$, A is ∞ -Borel.

489 Under AD^+ , boldface GCH holds below Θ .

490 **Fact 2.22.** (Steel, Woodin; [61] Theorem 2.16) *Assume AD^+ . For all $\kappa < \Theta$, boldface GCH holds at κ .*

491 Since boldface GCH below κ implies the hypothesis of Theorem 2.2 at κ , the following theorem is obtained
 492 as a corollary.

493 **Theorem 2.23.** *Assume AD^+ . For all cardinals $\kappa < \Theta$, $\neg(|[\kappa]^{\text{cof}(\kappa)}| \leq |\mathcal{P}_B(\kappa)|)$ and therefore for all
 494 $\gamma \leq \kappa$ with $\text{cof}(\gamma) = \text{cof}(\kappa)$, $\neg(|[\kappa]^{\gamma}| \leq |\mathcal{P}_B(\kappa)|)$ and $|\mathcal{P}_B(\kappa)| < |\mathcal{P}(\kappa)|$. If κ is a regular cardinal, then
 495 $|\kappa|^{<\kappa} < |\mathcal{P}(\kappa)|$.*

Theorem 2.12 shows that $|\omega_1|^{<\omega_1} < |\mathcal{P}(\omega_1)| = |\omega_1|^{\omega_1}$. Other proofs of this fact using more substantial amount of determinacy or partition properties can provide more information about the cardinality such as insight into the cofinality of $\mathcal{P}(\omega_1)$. (Cofinality and regularity in the choiceless context is developed in [17].) [11] showed that every function $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ is continuous $\mu_{\omega_1}^{\omega_1}$ -almost everywhere in the sense that there is a club $C \subseteq \omega_1$ so that for all $f \in [C]_*^{\omega_1}$, there is an $\alpha < \omega_1$ so that for all $g \in [C]_*^{\omega_1}$, if $f \restriction \alpha = g \restriction \alpha$, then $\Phi(f) = \Phi(g)$. This implies that $\mathcal{P}(\omega_1)$ has ω_1 -regular cardinality: If $\Phi : \mathcal{P}(\omega_1) \rightarrow \omega_1$, then there is an $\alpha < \omega_1$ so that $|\Phi^{-1}[\{\alpha\}]| = |\mathcal{P}(\omega_1)|$. $\mathcal{P}_B(\omega_1)$ does not have ω_1 -regular cardinality since $\mathcal{P}_B(\omega_1) = \bigcup_{\alpha < \omega_1} \mathcal{P}(\alpha)$ and $|\mathcal{P}(\alpha)| = |\mathcal{P}(\omega)|$ whenever $\alpha < \omega_1$ (and note that $|\mathcal{P}(\omega)| \neq |\mathcal{P}_B(\omega_1)|$ by boldface GCH at ω). This gives another proof that $|\omega_1|^{<\omega_1} = |\mathcal{P}_B(\omega_1)| < |\mathcal{P}(\omega_1)|$.

The next natural cardinality computation to consider would be the relation between $|\omega_1|^\omega = |\omega_1|^\omega$ and $|\mathcal{P}_B(\omega_1)| = |\omega_1|^{<\omega_1}$. Woodin [65] investigated the cardinality below $|\omega_1|^{<\omega_1}$ under $\text{AD}_{\mathbb{R}}$ and DC (which is known to imply AD^+). Woodin also isolated an important subset of $[\omega_1]^{<\omega_1}$ which he called $S_1 = \{f \in [\omega_1]^{<\omega_1} : \omega_1^{L[f]} = \sup(f)\}$. (See Fact 3.10 below for some basic properties of S_1 .)

In [65], Woodin proved two dichotomy results below $[\omega_1]^{<\omega_1}$ under $\text{AD}_{\mathbb{R}}$ and $\text{DC}_{\mathbb{R}}$.

- (1) If $X \subseteq [\omega_1]^\omega$, then either $|X| \leq |\mathbb{R} \times \omega_1|$ or $|X| = |\omega_1|^\omega$.
- (2) If $X \subseteq [\omega_1]^{<\omega_1}$, then $|X| \leq |\omega_1|^\omega$ or $|S_1| \leq |X|$.

The author can prove the first dichotomy for $|\omega_1|^\omega$ under AD and $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$ (or equivalently $\text{AD}_{\frac{1}{2}\mathbb{R}}$) using classical determinacy techniques for ω_1 . The first dichotomy also provides a complete classification of the cardinality below $|\omega_1|^\omega$ under $\text{AD}_{\mathbb{R}}$ and DC (or just $\text{AD}_{\frac{1}{2}\mathbb{R}}$): If $X \subseteq [\omega_1]^\omega$ is uncountable, then $|X|$ is $|\omega_1|$, $|\mathbb{R}|$, $|\mathbb{R} \sqcup \omega_1|$, $|\omega_1 \times \mathbb{R}|$, or $|\omega_1|^\omega$. It was unknown whether the first dichotomy could fail without $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$. As an application of the material developed in Section 4, Theorem 4.23 will produce an intermediate cardinality which is example of this failure.

The second dichotomy of Woodin is only meaningful if $|\omega_1|^\omega < |\omega_1|^{<\omega_1}$ and $|\mathbb{R}| < |S_1|$. Although the proofs of these cardinality distinctions do not seem to be included in [65], Woodin had known these cardinality computations at least under $\text{AD}_{\mathbb{R}}$ and DC. [13] provides a proof that S_1 does not inject into ${}^\omega\text{ON}$, the class of ω -sequences of ordinals, using just AD, $\text{DC}_{\mathbb{R}}$, and all sets of reals have ∞ -Borel code. To motivate the use of function capturing in generic extensions in the main result of the paper, weak versions of the main result will be shown and a survey of the relevant results from [13] will be provided.

Under ZF alone, $\omega_1 < \Theta$ can be proved with a very explicit surjection of $\mathbb{R}$ onto ω_1 . Let $\pi : \omega \times \omega \rightarrow \omega$ be a recursive bijection. If $x \in {}^\omega\omega$, let $\mathcal{R}_x \subseteq \omega \times \omega$ be defined by $R_x(a, b) \Leftrightarrow x(\pi(a, b)) = 1$. Let WO be the set of $w \in \mathbb{R}$ such that $\text{dom}(\mathcal{R}_w) = \omega$ and $(\omega, \mathcal{R}_w)$ is a wellordering. If $w \in \text{WO}$, then let $\text{ot}(w)$ be the ordertype of $(\omega, \mathcal{R}_w)$. $\text{ot} : \text{WO} \rightarrow \omega_1$ is a surjection. If $n \in \omega$ and $w \in \text{WO}$, then let $\text{ot}(w, n)$ be the rank of n in the wellordering $(\omega, \mathcal{R}_w)$. If $\alpha < \text{ot}(w)$, then let $\text{num}(w, \alpha)$ be the unique $n \in \omega$ so that $\text{ot}(w, n) = \alpha$. If $x \in {}^\omega\omega$ and $m \in \omega$, let $x^{[m]} \in {}^\omega\omega$ be defined by $x^{[m]}(n) = x(\pi(m, n))$. Let $u \in \text{BS}$ if and only if $u^{[0]} \in \text{WO}$ and for all $n \in \omega$, $(u^{[1]})^{[n]} \in \text{WO}$. If $u \in \text{BS}$, then let $\text{seq}(u) : \text{ot}(u^{[0]}) \rightarrow \omega_1$ be defined by $\text{seq}(u)(\alpha) = \text{ot}((u^{[1]})^{[\text{num}(u^{[0]}, \alpha)]})$ for each $\alpha < \text{ot}(u^{[0]})$. $\text{seq} : \text{BS} \rightarrow {}^{<\omega_1}\omega_1$ is a surjection. Let $\text{iseq} : \text{BS} \rightarrow \mathcal{P}_B(\omega_1)$ be defined by $\text{iseq}(u) = \text{seq}(u)[\text{ot}(u^{[0]})]$ and note that iseq is a surjection. From the explicit nature of the coding, if $\ell \in {}^{<\omega_1}\omega_1$ and $A \in \mathcal{P}_B(\omega_1)$, then in any inner model M with $\ell \in M$ and $M \models \text{"max}\{\text{dom}(\ell), \text{sup}(\ell)\} \text{ is countable"}$, there is a $u \in \text{BS} \cap M$ so that $\text{seq}(u) = \ell$ and in any inner model M with $A \in M$ and $M \models \text{"sup}(A) \text{ is countable"}$, then there is a $u \in \text{BS} \cap M$ so that $\text{iseq}(u) = A$.

Fact 2.24. *If boldface GCH at ω holds and all subsets of $\mathbb{R}$ have ∞ -Borel codes, then all subsets of $\mathcal{P}_B(\omega_1)$ and all subsets of ${}^{<\omega_1}\omega_1$ have ∞ -Borel codes.*

Proof. Let $X \subseteq \mathcal{P}_B(\omega_1)$. Let $\tilde{X} = \{u \in \text{BS} : \text{iseq}(u) \in X\}$. Since $\tilde{X} \subseteq \mathbb{R}$, the hypothesis implies there is an ∞ -Borel code (S, φ) for $\tilde{X}$. If ξ is an ordinal, then let $\text{Coll}(\omega, \xi)$ be the forcing of finite partial functions from ω into ξ ordered by reverse function inclusion. The claim is that for all $A \in \mathcal{P}_B(\omega_1)$, $A \in X$ if and only if

$$L[S, A] \models 1_{\text{Coll}(\omega, \text{sup}(A))} \Vdash_{\text{Coll}(\omega, \text{sup}(A))} (\exists u \in \text{BS})(\text{iseq}(u) = \check{A} \wedge L[\check{S}, u] \models \varphi(\check{S}, u)).$$

To see this: $(\Rightarrow)$ Suppose $A \in X$. Since $L[S, A] \models \text{AC}$ and $\text{sup}(A)$ is countable in the true world, $\text{Coll}(\omega, \text{sup}(A))$ is countable in the true universe. By Fact 2.14, there is a $G \subseteq \text{Coll}(\omega, \text{sup}(A))$ which is $\text{Coll}(\omega, \text{sup}(A))$ -generic over $L[S, A]$. Since $L[S, A][G] \models \text{sup}(A)$ is countable, there is a $u \in \text{BS} \cap L[S, A][G]$ such that $L[S, A][G] \models \text{iseq}(u) = A$. Since G belongs to the true universe, by absoluteness, $\text{iseq}(u) = A$.

Since $A \in X$, one has $u \in \tilde{X}$. Since (S, φ) is the ∞ -Borel code for $\tilde{X}$, $L[S, u] \models \varphi(S, u)$. By absoluteness, $L[S, A][G] \models \text{"iseq}(u) = A \wedge L[S, u] \models \varphi(S, u)\text{"}$. Thus

$$L[S, A][G] \models (\exists u \in \text{BS})(\text{iseq}(u) = A \wedge L[S, u] \models \varphi(S, u)).$$

By the forcing theorem, there is a $p \in \text{Coll}(\omega, \text{sup}(A))$ so that

$$L[S, A][G] \models p \Vdash_{\text{Coll}(\omega, \text{sup}(A))} (\exists u \in \text{BS})(\text{iseq}(u) = \check{A} \wedge L[S, u] \models \varphi(\check{S}, u)).$$

By the homogeneity of $\text{Coll}(\omega, \text{sup}(A))$,

$$L[S, A][G] \models 1_{\text{Coll}(\omega, \text{sup}(A))} \Vdash_{\text{Coll}(\omega, \text{sup}(A))} (\exists u \in \text{BS})(\text{iseq}(u) = \check{A} \wedge L[S, u] \models \varphi(\check{S}, u)).$$

($\Leftarrow$) Again by Fact 2.14, let $G \subseteq \text{Coll}(\omega, \text{sup}(A))$ be $\text{Coll}(\omega, \text{sup}(A))$ -generic over $L[S, A]$. By the forcing theorem,

$$L[S, A][G] \models (\exists u \in \text{BS})(\text{iseq}(u) = A \wedge L[S, u] \models \varphi(S, u)).$$

Let $u \in \text{BS} \cap L[S, A][G]$ be such that $L[S, A][G] \models \text{iseq}(u) = A \wedge L[S, A] \models \varphi(S, u)$. Since G belongs to the true universe and hence u belongs to the true universe, one has by absoluteness, $\text{iseq}(u) = A$ and $L[S, A] \models \varphi(S, u)$. Since (S, φ) is the ∞ -Borel code for $\tilde{X}$, $u \in \tilde{X}$. By definition of $\tilde{X}$, $\text{iseq}(u) = A \in X$. The completes the proof of the claim.

Let $\phi(S, A)$ be the statement

$$1_{\text{Coll}(\omega, \text{sup}(A))} \Vdash_{\text{Coll}(\omega, \text{sup}(A))} (\exists u \in \text{BS})(\text{iseq}(u) = \check{A} \wedge L[S, u] \models \varphi(\check{S}, u)).$$

The claim implies that (S, ϕ) is an ∞ -Borel code for X . $\square$

The proof of the next result will introduce the forcing and the basic skeleton of the capturing arguments used throughout the paper.

Fact 2.25. *Assume boldface GCH at ω and all subsets of $\mathbb{R}$ have ∞ -Borel codes. Then $|\omega \omega_1| = |[\omega_1]^\omega| < |\mathcal{P}_B(\omega)| = |^{<\omega_1}\omega_1| = |[\omega_1]^{<\omega_1}|$.*

Proof. Suppose $\Phi : \mathcal{P}_B(\omega_1) \rightarrow {}^\omega \omega_1$ is an injection. Let $\text{Gr}_\Phi = \{(A, n, \alpha) \in \mathcal{P}_B(\omega_1) \times \omega \times \omega_1 : \Phi(A)(n) = \alpha\}$ and $\text{Gr}_{\Phi^{-1}} = \{(f, \alpha) \in {}^\omega \omega_1 \times \omega_1 : f \in \Phi[\mathcal{P}_B(\omega_1)] \wedge \alpha \in \Phi^{-1}(f)\}$. By Fact 2.24, let (S_0, φ_0) and (S_1, φ_1) be ∞ -Borel codes for Gr_Φ and $\text{Gr}_{\Phi^{-1}}$, respectively. Let $\text{Coll}(\omega^+, \omega^{++})^{L[S_0, S_1]} = \text{Coll}(\omega_1, \omega_2)^{L[S_0, S_1]}$ be the forcing defined in $L[S_0, S_1]$ consisting of countable partial functions from $\omega_1^{L[S_0, S_1]}$ into $\omega_2^{L[S_0, S_1]}$ ordered by reverse function inclusion. Since this forcing is countable in the real world by boldface GCH at ω and Fact 2.18, Fact 2.14 implies there is a $G \subseteq \text{Coll}(\omega_1, \omega_2)^{L[S_0, S_1]}$ which is generic over $L[S_0, S_1]$. G adds a generic surjection $g : \omega_1^{L[S_0, S_1]} \rightarrow \omega_2^{L[S_0, S_1]}$. Let $A = \{(\alpha, \beta) : g(\alpha) = \beta\}$ be the graph of g which may be regarded as an element of $\mathcal{P}_B(\omega_1)$ (after coding pairs using the Gödel coding function) using the fact that $\omega_2^{L[S_0, S_1]} < \omega_1$ by Fact 2.18. Note that for all $n \in \omega$ and $\alpha \in \omega_1$, $\Phi(A)(n) = \alpha$ if and only if $\text{Gr}_\Phi(A, n, \alpha)$ if and only if $L[S_0, A] \models \varphi_0(S_0, A, n, \alpha)$ if and only if $L[S_0, S_1][G] \models L[S_0, A] \models \varphi_0(S_0, A, n, \alpha)$ by the fact that (S_0, φ_0) is the ∞ -Borel code for Gr_Φ and absoluteness. Thus by applying comprehension in $L[S_0, S_1][G]$, this shows that $\Phi(A) \in {}^\omega \omega_1 \cap L[S_0, S_1][G]$. Since $L[S_0, S_1] \models \text{"Coll}(\omega_1, \omega_2)^{L[S_0, S_1]} \text{ is } < \omega_1\text{-closed"}$, no new ω -sequence through $L[S_0, S_1]$ is added by this forcing. Hence $\Phi(A) \in L[S_0, S_1]$. Note that for all $\alpha \in \omega_1$, $\alpha \in A$ if and only if $\alpha \in \Phi^{-1}(\Phi(A))$ if and only if $\text{Gr}_{\Phi^{-1}}(\Phi(A), \alpha)$ if and only if $L[S_1, \Phi(A)] \models \varphi_1(S_1, \Phi(A), \alpha)$ if and only if $L[S_0, S_1] \models L[S_1, \Phi(A)] \models \varphi_1(S_1, \Phi(A), \alpha)$ since (S_1, φ_1) is the ∞ -Borel code for $\text{Gr}_{\Phi^{-1}}$ and by absoluteness. By comprehension, $A \in L[S_0, S_1]$. Since A is the graph of g , $g \in L[S_0, S_1]$. Since $L[S_0, S_1][G] \models \text{"}g : \omega_1^{L[S_0, S_1]} \rightarrow \omega_2^{L[S_0, S_1]} \text{ is a surjection"}$, $L[S_0, S_1] \models \text{"}g : \omega_1^{L[S_0, S_1]} \rightarrow \omega_2^{L[S_0, S_1]} \text{ is a surjection"}$ by absoluteness. This is a contradiction. $\square$

Inspecting the proof, one found an inner model M which was used to solve two crucial problems:

- (1) (Generic existence problem) There is a generic $G \subseteq \text{Coll}(\omega_1, \omega_2)^M$ which is $\text{Coll}(\omega_1, \omega_2)^M$ -generic over M . (This is provided by Fact 2.14 and boldface GCH at ω .)
- (2) (Capturing problem)
 - $M[G]$ is closed under Φ : for all $A \in \mathcal{P}_B(\omega_1) \cap M[G]$, $\Phi(A) \in M[G]$.
 - M is closed under the partial function Φ^{-1} : For all $f \in \text{dom}(\Phi^{-1}) \cap M$, $\Phi^{-1}(f) \in M$.

Since Gr_Φ and $\text{Gr}_{\Phi^{-1}}$ have ∞ -Borel codes (S_0, φ_0) and (S_1, φ_1) , any inner model of ZF containing S_0 and S_1 can capture Φ and Φ^{-1} in the above sense.

Toward the goal of classifying cardinal exponentiation, one needs to prove a strong instance of Fact 2.25; namely, $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$. Suppose $\Phi : \mathcal{P}_B(\omega_1) \rightarrow {}^\omega\kappa$ where κ is any cardinal below Θ . The challenge is to find an inner model $M \models \text{ZFC}$ which is closed under the partial function $\Phi^{-1} : {}^\omega\kappa \rightarrow \mathcal{P}_B(\kappa)$. $\text{Gr}_{\Phi^{-1}}$ is a subset of ${}^\omega\kappa \times \omega_1$. If $\kappa > \omega_1$, then Fact 2.24 cannot be applied to show that $\text{Gr}_{\Phi^{-1}}$ has an ∞ -Borel code. Instead, [13] used the ∞ -Borel code forcing (see Section 4) and Fact 4.17 to obtain the necessary model closed under Φ^{-1} . However at the time of [13], only Fact 4.17 for the Woodin's classical ∞ -Borel code forcing $\mathbb{B}_n^{\omega;Z}$ which adds reals (for some set Z and $n \in \omega$) was known at least to the authors. The forcing $\text{Coll}(\omega^+, \omega^{++})^M$ (for some suitable inner model M of ZFC) used in the proof of Fact 2.25 does not add any new reals and hence the classical Fact 4.17 for adding reals could not be used. It is for this reason that the more natural forcing $\text{Coll}(\omega^+, \omega^{++})^M$ was not used in [13] but rather the Lévy collapse $\text{Coll}(\omega, < \xi)^M$ where ξ is an inaccessible cardinal of M which is countable in the real world was used since it adds many new reals. One additional benefit of using the Lévy collapse is that it provides a cardinality result even for S_1 .

Fact 2.26. ([13] Theorem 5.7) Assume AD, $\text{DC}_\mathbb{R}$, and all sets of reals have ∞ -Borel codes. S_1 does not inject into ${}^\omega\text{ON}$ and thus $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$.

Several questions arise from these early results.

- (a) Can AD alone prove $[\omega_1]^{<\omega_1}$ does not inject into ${}^\omega\text{ON}$?
- (b) Can AD alone prove S_1 does not inject into ${}^\omega\text{ON}$?
- (c) (Bounded Power Set Conjecture) From any determinacy hypothesis, can it be shown that for all cardinals κ with $\omega < \kappa < \Theta$ and ordinals $\epsilon < \kappa$, $\mathcal{P}_B(\kappa)$ does not inject into ${}^\epsilon\text{ON}$?

Question (a) has been solved using descriptive set theoretic properties and partition properties. These methods also provide partial solutions to Question (c). A few remarks about these combinatorial methods will be mentioned below. Question (b) will be solved in the next section and the techniques will motivate the methods used to solve Question (c). The solution to Question (c) is the main result of this paper.

Using descriptive set theoretic properties of the pointclass Σ_1^1 and Kunen trees ([30]) from the study of measures below ω_ω under AD, the following almost everywhere continuity result with respect to the partition measures $\mu_{\omega_1}^\epsilon$ ($\epsilon < \omega_1$) was established.

Fact 2.27. ([19] Theorem 2.14, [18] Fact 2.5) Assume AD. For every $\epsilon < \omega_1$ and function $\Phi : [\omega_1]_*^\epsilon \rightarrow \omega_1$, there is a club $C \subseteq \omega_1$ and a $\delta < \epsilon$ so that for all $f, g \in [C]_*^\epsilon$, if $f \restriction \delta = g \restriction \delta$ and $\sup(f) = \sup(g)$, then $\Phi(f) = \Phi(g)$.

The continuity property of Fact 2.27 is a very well controlled failure of injectivity. This is used to obtain a proof of Fact 2.25 from AD alone.

Fact 2.28. ([19] Theorem 2.16) Assume AD. $|{}^\omega\omega_1| < |[\omega_1]^{<\omega_1}| = |\mathcal{P}_B(\omega_1)|$.

Building on Fact 2.27 and Martin's ultrapower representation of $\omega_{n+1} = \prod_{[\omega_1]_*^n} \omega_1 / \mu_{\omega_1}^n$ for all $1 \leq n < \omega$, one can show $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega(\omega_\omega)$. Furthermore using Fact 2.27 and extensive descriptive set theoretic properties of ω_1 and the pointclasses of Σ_1^1 and Σ_2^1 , one can show that $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$.

Fact 2.29. ([18] Theorem 2.9 and Theorem 4.4)

- Assume AD. $\neg(|\mathcal{P}_B(\omega_1)| \leq |{}^\omega(\omega_\omega)|)$.
- Assume AD and $\text{DC}_\mathbb{R}$. $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$.

The proof of Fact 2.29 uses many descriptive set theoretic property surrounding ω_1 which are not known to generalize to any higher cardinals. Moreover, $\text{DC}_\mathbb{R}$ is used directly in the proof and indirectly in several descriptive set theoretic property (such as the wellfoundedness of the Wadge degree in pointclass arguments). To remove $\text{DC}_\mathbb{R}$ and answer instances of Question (c) for higher cardinals, a strengthening of Fact 2.27 was established using pure combinatorics.

Fact 2.30. ([21] Theorem 3.7) Assume $\epsilon < \kappa$, $\text{cof}(\epsilon) = \omega$, and $\kappa \rightarrow_* (\kappa)_2^{\epsilon^\epsilon}$ holds. For any function $\Phi : [\kappa]_*^\epsilon \rightarrow \text{ON}$, there is a $\delta < \epsilon$ and a club $C \subseteq \kappa$ such that for all $f, g \in [C]_*^\epsilon$, if $f \restriction \delta = g \restriction \delta$ and $\sup(f) = \sup(g)$, then $\Phi(f) = \Phi(g)$.

This is used to answer Question (a) and Question (c) at weak partition cardinals. The following result gave the most compelling evidence to formulate the Bounded Power Set Conjecture.

Fact 2.31. ([21] Theorem 4.4) *Let κ be a weak partition cardinal (that is, $\kappa \rightarrow_* (\kappa)_2^{<\kappa}$) and $\epsilon < \kappa$. Then $\mathcal{P}_B(\kappa)$ does not inject into ${}^\epsilon\text{ON}$.*

Thus under AD, $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$ and $\mathcal{P}_B(\omega_2)$ does not inject into ${}^{\omega_1}\text{ON}$. Also $\mathcal{P}_B(\omega_3)$ does not inject into ${}^{\omega_1}\omega_3$. Since ω_3 is not a partition cardinal, the first relevant open cardinality computation is whether $\mathcal{P}_B(\omega_3)$ injects into ${}^{\omega_2}\omega_3$ which will be solved here. This is the history of these questions until this paper.

The rest of the paper will prove the main result. An important step is finding an appropriate form of capturing in certain inner models of ZFC and its generic extension without the given function possessing ∞ -Borel codes. In fact, Woodin showed under AD^+ that there is always a subset of $\mathcal{P}(\omega_1)$ which does have an ∞ -Borel code. (See [15].) So Question (c) (the Bounded Power Set Conjecture) cannot be solved using ∞ -Borel codes for functions of the form $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$. Fact 2.26 says that S_1 does not inject into ${}^\omega\text{ON}$ if all subsets of $\mathbb{R}$ have ∞ -Borel codes. Question (b) asks how this can be done without ∞ -Borel codes. The next section will solve Question (b) and the technique will motivate how to resolve the Question (c), the eponymous Bounded Power Set Conjecture.

3. CAPTURING WITHOUT ∞ -BOREL CODE WITHIN THE HEREDITARILY COUNTABLE SETS

This section will provide an answer to question (b) above and show that under AD (and in fact, less than AD), S_1 does not inject into ${}^\omega\text{ON}$. Fact 2.26 used the existence of ∞ -Borel code. A solution must do without ∞ -Borel codes. Capturing will be obtained by using an appropriate HOD-type model. This will motivate some of the techniques of the main theorem of the paper where one will also need to work without the assumption that subsets of the power set of uncountable ordinals have ∞ -Borel codes.

Let $\mathbb{P}$ be a countable forcing. Let $\text{Fin}(\mathbb{P}, 2)$ be the set of all finite partial functions from $\mathbb{P}$ into 2. For each $s \in \text{Fin}(\mathbb{P}, 2)$, let $N_s^\mathbb{P} = \{f \in {}^\mathbb{P}2 : s \subseteq f\}$. Give ${}^\mathbb{P}2$ the topology generated by $\{N_s^\mathbb{P} : s \in \text{Fin}(\mathbb{P}, 2)\}$ as a basis. ${}^\mathbb{P}2$ is homeomorphic to ${}^\omega 2$ with its usual topology and thus ${}^\mathbb{P}2$ is a Polish space. Let $\mathcal{F}_\mathbb{P} \subseteq {}^\mathbb{P}2$ be the set of all $\mathbb{P}$ -filters (or more accurately the characteristic function of $\mathbb{P}$ -filters).

Note that $\mathcal{F}_\mathbb{P}$ is a $\mathbf{\Pi}_2^0$ subset of ${}^\mathbb{P}2$ (using the fact that basic neighborhoods are clopen). Thus $\mathcal{F}_\mathbb{P}$ is a Polish space with the subspace topology inherited from ${}^\mathbb{P}2$. One can then define the usual notions of category for this Polish topology. If $\text{AC}_\omega^\mathbb{R}$ holds and all subsets $\mathcal{F}_\mathbb{P}$ has the property of Baire (defined using its natural Polish topology), then the argument of Fact 2.5 shows that wellordered unions of meager subsets of $\mathcal{F}_\mathbb{P}$ are meager subsets of $\mathcal{F}_\mathbb{P}$. (Under AD, all subsets of all Polish spaces have the Baire property relative to its Polish topology.)

For each $s \in \text{Fin}(\mathbb{P}, 2)$, let $M_s^\mathbb{P} = N_s^\mathbb{P} \cap \mathcal{F}_\mathbb{P}$. It may be the case that some $s \in \text{Fin}(\mathbb{P}, 2)$, $M_s^\mathbb{P} = \emptyset$. Let $\text{sFin}(\mathbb{P}, 2)$ be the collection of $s \in \text{Fin}(\mathbb{P}, 2)$ such that $M_s^\mathbb{P} \neq \emptyset$. Say that $s \in \text{Fin}(\mathbb{P}, 2)$ is $\mathbb{P}$ -suitable if and only if there is a $\tilde{p} \in \mathbb{P}$ so that for all $p \in \text{dom}(s)$ with $s(p) = 1$, $\tilde{p} \leq_\mathbb{P} p$ and for all $p \in \text{dom}(s)$ with $s(p) = 0$, p and $\tilde{p}$ are not compatible. Suppose s is $\mathbb{P}$ -suitable. Let $\tilde{p} \in \mathbb{P}$ be such that for all $p \in \text{dom}(s)$ with $s(p) = 1$, $\tilde{p} \leq_\mathbb{P} p$ and for all $p \in \text{dom}(s)$ with $s(p) = 0$, p and $\tilde{p}$ are not compatible. Let $f : \mathbb{P} \rightarrow 2$ be defined by $f(p) = 1$ if and only if $\tilde{p} \leq_\mathbb{P} p$. Then $f \in \mathcal{F}_\mathbb{P}$ and $s \subseteq f$. This shows that $M_s^\mathbb{P} \neq \emptyset$. Now suppose $s \in \text{Fin}(\mathbb{P}, 2)$ be such that $M_s^\mathbb{P} \neq \emptyset$. Let $f \in M_s^\mathbb{P}$. Since f codes a filter, there is a $\tilde{p}$ in the filter coded by f so that $\tilde{p} \leq_\mathbb{P} p$ for all $p \in \text{dom}(s)$ with $s(p) = 1$ and for all $p \in \text{dom}(s)$ with $s(p) = 0$, p and $\tilde{p}$ are not compatible. This shows that s is $\mathbb{P}$ -suitable. Thus $\{M_s^\mathbb{P} : s \in \text{Fin}_\mathbb{P}\} = \{M_s^\mathbb{P} : s \text{ is } \mathbb{P}\text{-suitable}\}$ is a basis for the topology on $\mathcal{F}_\mathbb{P}$ and $\text{sFin}(\mathbb{P}, 2)$ is the collection of $\mathbb{P}$ -suitable $s \in \text{Fin}(\mathbb{P}, 2)$.

If $D \subseteq \mathbb{P}$ is dense, then let $\mathcal{F}_\mathbb{P}^D = \{f \in \mathcal{F}_\mathbb{P} : (\exists p \in D)(f(p) = 1)\}$. Suppose $f \in \mathcal{F}_\mathbb{P}^D$. Then there is some $p \in D$ so that $f(p) = 1$. Let $s : \{p\} \rightarrow 2$ be defined by $s(p) = 1$. Note $f \in M_s^\mathbb{P} \subseteq \mathcal{F}_\mathbb{P}^D$. This shows $\mathcal{F}_\mathbb{P}^D$ is open. Suppose $s \in \text{sFin}_\mathbb{P}$ or equivalently $\mathbb{P}$ -suitable. There is a $\tilde{p} \in \mathbb{P}$ so that for all $p \in \text{dom}(s)$ with $s(p) = 1$, $\tilde{p} \leq_\mathbb{P} p$ and for all $p \in \text{dom}(s)$ with $s(p) = 0$, $\tilde{p}$ and p are not compatible. Since D is dense, there is a $\bar{p} \in D$ with $\bar{p} \leq_\mathbb{P} \tilde{p}$. Let $s' = s \cup \{(\bar{p}, 1)\}$ and note that s' is still $\mathbb{P}$ -suitable and $M_{s'}^\mathbb{P} \subseteq M_s^\mathbb{P} \cap \mathcal{F}_\mathbb{P}^D$. This shows that $\mathcal{F}_\mathbb{P}^D$ is dense open.

Suppose $\mathcal{D}$ is a countable collection of dense subsets of $\mathbb{P}$. Let $\text{Gen}_\mathbb{P}^\mathcal{D} = \bigcap_{D \in \mathcal{D}} \mathcal{F}_\mathbb{P}^D$ be the collection of $f \in {}^\mathbb{P}2$ which codes $\mathbb{P}$ -generic filter over $\mathcal{D}$. Note that $\text{Gen}_\mathbb{P}^\mathcal{D}$ is comeager since it is a countable intersection of dense open subsets of $\mathcal{F}_\mathbb{P}$.

Recall that if Z is a set, then OD_Z and HOD_Z refer to ordinal definability with parameters from Z . If $Z \in \text{OD}_Z$, then $\text{HOD}_Z \models \text{ZF}$. If $Z \in \text{OD}_Z$ and Z has an OD_Z wellordering, then $\text{HOD}_Z \models \text{ZFC}$.

Lemma 3.1. (Woodin; [40] Theorem 5.42, Claim 2) Assume $\text{AC}_\omega^\mathbb{R}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at ω . Let Z be a set such that there is an OD_Z wellordering of Z . Let $\mathbb{P} \in \text{HOD}_Z$ be a forcing which is countable in the real world. Let $\varsigma_Z^\mathbb{P}$ be the least ordinal γ greater than ω_1^V such that $Z \cup \{\mathbb{P}\} \in V_\gamma$.⁵ Let $\mathcal{D}_Z^\mathbb{P}$ denote the collection of $\mathbb{P}$ -dense subsets of $\mathbb{P}$ in HOD_Z . Let $\mathcal{S}_Z^\mathbb{P}$ be the collection of all tuples $(\varphi, \gamma, \vec{z}, \vec{\xi})$ such that $\gamma \in \text{ON}$ with $\varsigma_Z^\mathbb{P} \leq \gamma$, $\vec{z}$ is a finite tuple of elements from Z , $\vec{\xi}$ is a finite tuple of ordinals less than γ , and φ is a $\mathcal{L}_{\text{set}}$ -formula with $|\vec{z}| + |\vec{\xi}| + 1$ free variables. Uniformly from $\mathbb{P}$ and Z ,⁶ there is a $T_Z^\mathbb{P} \subseteq \text{Gen}_\mathbb{P}^{\mathcal{D}_Z^\mathbb{P}}$ which is comeager in the topology of $\mathcal{F}_\mathbb{P}$ and sequence $\langle P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}, N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} : (\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P} \rangle$ with the following properties:

- (1) For any $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$, $P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} \subseteq \text{sFin}(\mathbb{P}, 2)$ and $N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} \subseteq \text{sFin}(\mathbb{P}, 2)$.
- (2) For any $G \in T_Z^\mathbb{P}$ and $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$, $V_\gamma \models \varphi(G, \vec{z}, \vec{\xi})$ if and only if there is an $s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ so that $G \in M_s^\mathbb{P}$.
- (3) For any $G \in T_Z^\mathbb{P}$ and $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$, $V_\gamma \models \neg \varphi(G, \vec{z}, \vec{\xi})$ if and only if there is an $s \in N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ so that $G \in M_s^\mathbb{P}$.
- (4) For any finite $\vec{z}$ in Z , let $\mathcal{S}_{Z, \vec{z}}^\mathbb{P}$ be the collection of $(\varphi, \gamma, \vec{\xi})$ such that $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$. Then $\mathcal{S}_{Z, \vec{z}}^\mathbb{P} \in \text{HOD}_Z$ and $\langle P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}, N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} : (\varphi, \gamma, \vec{\xi}) \in \mathcal{S}_{Z, \vec{z}}^\mathbb{P} \rangle \in \text{HOD}_Z$.

Proof. Let $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$. Let $P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ be the collection of $s \in \text{sFin}(\mathbb{P}, 2)$ so that $A_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s = \{F \in M_s^\mathbb{P} \cap \text{Gen}_\mathbb{P}^{\mathcal{D}_Z^\mathbb{P}} : V_\gamma \models \varphi(F, \vec{z}, \vec{\xi})\}$ is comeager in $M_s^\mathbb{P}$. Let $A_{\varphi, \gamma, \vec{z}, \vec{\xi}} = \bigcup_{s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}} A_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s$. Let $N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ be the collection of $s \in \text{sFin}(\mathbb{P}, 2)$ so that $B_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s = \{F \in M_s^\mathbb{P} \cap \text{Gen}_\mathbb{P}^{\mathcal{D}_Z^\mathbb{P}} : V_\gamma \models \neg \varphi(F, \vec{z}, \vec{\xi})\}$ is comeager in $M_s^\mathbb{P}$. Let $B_{\varphi, \gamma, \vec{z}, \vec{\xi}} = \bigcup_{s \in N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}} B_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s$. Let $C_{\varphi, \gamma, \vec{z}, \vec{\xi}} = A_{\varphi, \gamma, \vec{z}, \vec{\xi}} \cup B_{\varphi, \gamma, \vec{z}, \vec{\xi}}$. Note that $C_{\varphi, \gamma, \vec{z}, \vec{\xi}} \subseteq \text{Gen}_\mathbb{P}^{\mathcal{D}_Z^\mathbb{P}}$ is a comeager subset of $\mathcal{F}_\mathbb{P}$. Let $T_Z^\mathbb{P} = \bigcap_{(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}} C_{\varphi, \gamma, \vec{z}, \vec{\xi}}$ which is a comeager subset of $\mathcal{F}_\mathbb{P}$ since wellordered intersection of comeager subsets of $\mathcal{F}_\mathbb{P}$ are comeager because all subsets of the Polish space $\mathcal{F}_\mathbb{P}$ has the property of Baire and $\mathcal{S}_Z^\mathbb{P}$ is a wellorderable set (since Z is wellorderable by the hypothesis that there is even an OD_Z wellordering of Z). Note that for all $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$, $P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} \in \text{HOD}_Z$ and $N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} \in \text{HOD}_Z$ and for any finite $\vec{z} \in Z$, $\mathcal{S}_{Z, \vec{z}}^\mathbb{P} \in \text{HOD}_Z$ and $\langle P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}, N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} : (\varphi, \gamma, \vec{\xi}) \in \mathcal{S}_{Z, \vec{z}}^\mathbb{P} \rangle \in \text{HOD}_Z$. Let $G \in T_Z^\mathbb{P}$ and $(\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P}$. Then $G \in T_Z^\mathbb{P} \subseteq C_{\varphi, \gamma, \vec{z}, \vec{\xi}} = A_{\varphi, \gamma, \vec{z}, \vec{\xi}} \cup B_{\varphi, \gamma, \vec{z}, \vec{\xi}}$. Since $A_{\varphi, \gamma, \vec{z}, \vec{\xi}}$ and $B_{\varphi, \gamma, \vec{z}, \vec{\xi}}$ are disjoint, G is in exactly one of the two sets. If $G \in A_{\varphi, \gamma, \vec{z}, \vec{\xi}} = \bigcup_{s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}} A_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s$, then there is some $s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ so that $G \in A_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s \subseteq M_s^\mathbb{P}$ and $V_\gamma \models \varphi(G, \vec{z}, \vec{\xi})$. If $G \in B_{\varphi, \gamma, \vec{z}, \vec{\xi}} = \bigcup_{s \in N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}} B_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s$, then there is some $s \in N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}$ so that $G \in B_{\varphi, \gamma, \vec{z}, \vec{\xi}}^s \subseteq M_s^\mathbb{P}$ and $V_\gamma \models \neg \varphi(G, \vec{z}, \vec{\xi})$. This completes the argument. $\square$

Lemma 3.2. (Woodin; [40] Theorem 5.42, Claim 2) Assume $\text{AC}_\omega^\mathbb{R}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at ω . Let Z be a set such that there is an OD_Z wellordering of Z . Let $\mathbb{P} \in \text{HOD}_Z$ be a forcing which is countable in the real world. Let $T_Z^\mathbb{P} \subseteq \mathcal{F}_\mathbb{P}$ be the comeager set from Lemma 3.1. For all $G \in T_Z^\mathbb{P}$, $\text{HOD}_Z[G] = \text{HOD}_{Z \cup \{G\}}$.

Proof. Let $G \in T_Z^\mathbb{P}$. It is clear that $\text{HOD}_Z[G] \subseteq \text{HOD}_{Z \cup \{G\}}$. Suppose $Y \in \text{HOD}_{Z \cup \{G\}}$. Let U be the transitive closure of $\{Y\}$. Because there is an OD_Z wellordering of Z , $\text{HOD}_{Z \cup \{G\}} \models \text{AC}$. Let $\pi : \delta \rightarrow U$ be a bijection where δ is an ordinal and $\pi \in \text{HOD}_{Z \cup \{G\}}$. Within $\text{HOD}_{Z \cup \{G\}}$, define $R \subseteq \delta \times \delta$ by $R(\alpha, \beta)$ if and only if $\pi(\alpha) \in \pi(\beta)$. There is some ordinal $\gamma > \max\{\delta, \varsigma_Z^\mathbb{P}\}$, finite $\vec{z}$ in Z , and finitely many ordinals $\vec{\xi} < \gamma$,

⁵ V_γ refers to the sets of rank less than γ in the real world.

⁶The comeager set $T_Z^\mathbb{P}$ and the sequence $\langle P_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z}, N_{\varphi, \gamma, \vec{z}, \vec{\xi}}^{\mathbb{P}, Z} : (\varphi, \gamma, \vec{z}, \vec{\xi}) \in \mathcal{S}_Z^\mathbb{P} \rangle$ depend only on $\mathbb{P}$ and Z such that there is an OD_Z wellordering of Z and $\mathbb{P} \in \text{HOD}_Z$ is countable in the real world and does not depend on any witness that $\mathbb{P}$ is countable in the real world.

and a formula with $|\vec{z}| + |\vec{\xi}| + 3$ variables so that for all $\alpha, \beta < \delta$, $R(\alpha, \beta)$ if and only if $V_\gamma \models \varphi(G, \vec{z}, \vec{\xi}, \alpha, \beta)$. For $\alpha, \beta \in \delta$, let $\vec{\xi}_{\alpha, \beta} = \vec{\xi}(\alpha, \beta)$. By Lemma 3.1, the set $\langle P_{\varphi, \gamma, \vec{z}, \vec{\xi}_{\alpha, \beta}}^{\mathbb{P}, Z} : \alpha, \beta \in \delta \rangle \in \text{HOD}_Z$. For all $\alpha, \beta \in \delta$, $R(\alpha, \beta)$ if and only if there is an $s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}_{\alpha, \beta}}^{\mathbb{P}, Z}$ with $G \in M_s^{\mathbb{P}}$ if and only if

$$\text{HOD}_Z[G] \models (\exists s \in P_{\varphi, \gamma, \vec{z}, \vec{\xi}_{\alpha, \beta}}^{\mathbb{P}, Z})(G \in M_s^{\mathbb{P}}).$$

By the axiom of comprehension inside of $\text{HOD}_Z[G]$, $R \in \text{HOD}_Z[G]$. Then $U \in \text{HOD}_Z[G]$ since U is the Mostowski collapse of the wellfounded relation (δ, R) . Finally $Y \in \text{HOD}_Z[G]$ since Y is the unique element of highest rank in U . Since $Y \in \text{HOD}_{Z \cup \{G\}}$ was arbitrary, $\text{HOD}_{Z \cup \{G\}} \subseteq \text{HOD}_Z[G]$. Thus $\text{HOD}_Z[G] = \text{HOD}_{Z \cup \{G\}}$. $\square$

Theorem 3.3. Assume $\text{AC}_\omega^{\mathbb{R}}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at ω . Then there is no injection of $\mathcal{P}_B(\omega_1)$ into ${}^\omega\text{ON}$ and hence $|{}^\omega\omega_1| = |[\omega_1]^\omega| < |\mathcal{P}_B(\omega_1)| = |{}^{<\omega_1}\omega_1| = |[\omega_1]^{<\omega_1}|$.

Proof. Suppose there is an injection $\Phi : \mathcal{P}_B(\omega_1) \rightarrow {}^\omega\text{ON}$. Let $\mathbb{P} = \text{Coll}(\omega_1^+, \omega^{++})^{\text{HOD}_{\{\Phi\}}} = \text{Coll}(\omega_1, \omega_2)^{\text{HOD}_{\{\Phi\}}}$ which is the forcing of countable partial functions from $\omega_1^{\text{HOD}_{\{\Phi\}}}$ into $\omega_2^{\text{HOD}_{\{\Phi\}}}$ as defined within $\text{HOD}_{\{\Phi\}}$. $\mathbb{P}$ is a countable forcing from the view of the real world. By Lemma 3.2, there is $G \subseteq \mathbb{P}$ which is $\mathbb{P}$ -generic over $\text{HOD}_{\{\Phi\}}$ such that $\text{HOD}_{\{\Phi, G\}} = \text{HOD}_{\{\Phi\}}[G]$. $\mathbb{P}$ adds a generic surjection $g : \omega_1^{\text{HOD}_{\{\Phi\}}} \rightarrow \omega_2^{\text{HOD}_{\{\Phi\}}}$. Let $\pi : \omega_1 \times \omega_1 \rightarrow \omega_1$ be the Gödel pair function which belongs to $\text{HOD}_{\{\Phi\}}$. Let $A = \pi[g]$ which is a bounded subset of the true ω_1 and thus $A \in \mathcal{P}_B(\omega_1) \cap \text{HOD}_{\{\Phi\}}[G]$. Since A is ordinal definable from G , $\Phi(A)$ is ordinal definable from Φ and G . Thus $\Phi(A) \in \text{HOD}_{\{\Phi, G\}} = \text{HOD}_{\{\Phi\}}[G]$. So $\Phi(A) \in \text{HOD}_{\{\Phi\}}[G] \cap {}^\omega\text{ON}$. Since $\text{HOD}_{\{\Phi\}} \models \text{“}\mathbb{P} = \text{Coll}(\omega_1, \omega_2) \text{ is } < \omega_1\text{-closed”}$, no new ω -sequences through $\text{HOD}_{\{\Phi\}}$ are added by forcing $\mathbb{P}$ over $\text{HOD}_{\{\Phi\}}$. Thus $\Phi(A) \in \text{HOD}_{\{\Phi\}}$. Since $\Phi(A)$ is ordinal definable from Φ , $A = \Phi^{-1}(\Phi(A))$ is ordinal definable from Φ . Thus $A \in \text{HOD}_{\{\Phi\}}$. Then $g = \pi^{-1}[A]$ is ordinal definable from Φ and so $g \in \text{HOD}_{\{\Phi\}}$. By absoluteness, $\text{HOD}_{\{\Phi\}} \models \text{“}g : \omega_1 \rightarrow \omega_2 \text{ is a surjection”}$. Contradiction. $\square$

The hypothesis of Theorem 3.3 follows from AD but moreover it follows from $\text{AC}_\omega^{\mathbb{R}}$ and all subsets of all Polish spaces have the Baire property and the perfect set property by Fact 2.6. Thus $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$ is a consequence of classical regularity properties.

Theorem 3.4. The consistency of “ZFC and there exists an inaccessible cardinal” implies the consistency of “the Bounded Power Set Conjecture and the ABCD Conjecture below $\Theta = \omega_2$ ”.

Proof. The Solovay model ([54]) constructed from the Lévy collapse of an inaccessible satisfies the hypothesis of Theorem 3.3 and $\Theta = \omega_2$. Theorem 3.3 shows that the Solovay model satisfies the Bounded Powerset Conjecture below ω_2 : $\mathcal{P}_B(\omega_1)$ does not inject into ${}^\omega\text{ON}$. Since $\Theta = \omega_2$, the only nontrivial instance of the ABCD conjecture below Θ is that $|{}^\omega\omega_1| < |{}^{\omega_1}\omega_1|$ which follows from $\mathcal{P}_B(\omega_1)$ not injecting into ${}^\omega\text{ON}$ since $|\mathcal{P}_B(\omega_1)| \leq |{}^{\omega_1}\omega_1|$. $\square$

Theorem 3.5. Assume $\text{AC}_\omega^{\mathbb{R}}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at ω . ${}^\omega\omega_1$ does not inject into $\mathcal{P}(\omega) \times \text{ON}$. Thus $|\mathbb{R} \times \omega_1| < |{}^\omega\omega_1| = |[\omega_1]^\omega|$.

Proof. Suppose otherwise that there is an injection $\Phi : {}^\omega\omega_1 \rightarrow \mathcal{P}(\omega) \times \text{ON}$. Let $\text{Coll}(\omega_1, 2^\omega)^{\text{HOD}_{\{\Phi\}}}$ be the forcing defined in $\text{HOD}_{\{\Phi\}}$ consisting of countable partial functions from $\omega_1^{\text{HOD}_{\{\Phi\}}}$ into the $\mathbb{R}^{\text{HOD}_{\{\Phi\}}}$ ordered by reverse function extension. Let $\dot{\mathbb{N}} \in \text{HOD}_{\{\Phi\}}$ be such that $\text{HOD}_{\{\Phi\}} \models \text{“}\dot{\mathbb{N}} \text{ is the Coll}(\omega_1, \mathbb{R})\text{-name for Namba forcing”}$. (Namba forcing $\mathbb{N}$ consists of $p \subseteq {}^{<\omega}\omega_2$ which are nonempty trees on ω_2 so that each node of p has ω_2 many extensions. Let $\leq_{\mathbb{N}}$ be the subset relation. See [31] Theorem 28.10 for more information on Namba forcing.) Let $\mathbb{P}$ be the two step iterated forcing $\text{Coll}(\omega_1, 2^\omega) * \dot{\mathbb{N}}$. $\mathbb{P}$ is a countable forcing in the real world.⁷ Thus by Theorem 3.2, there is a $\mathbb{P}$ -generic filter over $\text{HOD}_{\{\Phi\}}$ with the property that $\text{HOD}_{\{\Phi\}}[G] = \text{HOD}_{\{\Phi, G\}}$. There are $H \subseteq \text{Coll}(\omega_1, 2^\omega)^{\text{HOD}_{\{\Phi\}}}$ which is $\text{Coll}(\omega_1, 2^\omega)^{\text{HOD}_{\{\Phi\}}}$ -generic over $\text{HOD}_{\{\Phi\}}$ and $K \subseteq \dot{\mathbb{N}}[H]$ which is $\dot{\mathbb{N}}[H]$ -generic over $\text{HOD}_{\{\Phi\}}[H]$ so that $G = H * K$. $\text{Coll}(\omega_1, 2^\omega)^{\text{HOD}_{\{\Phi\}}}$ adds no new reals or even ω -sequences through $\text{HOD}_{\{\Phi\}}$ since $\text{HOD}_{\{\Phi\}} \models \text{“Coll}(\omega_1, 2^\omega) \text{ is } < \omega_1\text{-closed”}$ and

⁷Here, one assumes that some reasonable presentation of the iterated forcing $\text{Coll}(\omega, 2^\omega) * \dot{\mathbb{N}}$ is used to limit the $\text{Coll}(\omega_1, 2^\omega)$ -names for elements of $\dot{\mathbb{N}}$

744 $\text{HOD}_{\{\Phi\}}[H] \models \text{CH}$ (which is the primary purpose of this first forcing). Since $\text{HOD}_{\{\Phi\}}[H] \models \text{CH}$, the $\dot{N}[H]$ -
745 generic filter K adds a generic cofinal function $k : \omega \rightarrow \omega_2^{\text{HOD}_{\{\Phi\}}}$ and adds no new reals by [31] Theorem
746 28.10. Therefore $\mathcal{P}(\omega) \cap \text{HOD}_{\{\Phi\}} = \mathcal{P}(\omega) \cap \text{HOD}_{\{\Phi\}}[G] = \mathcal{P}(\omega) \cap \text{HOD}_{\{\Phi\}}[H * K]$. Since boldface GCH
747 at ω holds, $\omega_2^{\text{HOD}_{\{\Phi\}}}$ is less than the true ω_1 and thus $k \in [\omega_1]^\omega$. Since $\Phi(k) \in \text{OD}_{\{\Phi, K\}} \subseteq \text{OD}_{\{\Phi, G\}}$,
748 $\Phi(k) \in \text{HOD}_{\{\Phi, G\}} = \text{HOD}_{\{\Phi\}}[G]$. Since $\Phi(k) \in \mathcal{P}(\omega) \times \text{ON}$ and $\mathcal{P}(\omega) \cap \text{HOD}_{\{\Phi\}}[G] = \mathcal{P}(\omega) \cap \text{HOD}_{\{\Phi\}}$,
749 $\Phi(k) \in \text{HOD}_{\{\Phi\}}$. Thus $k = \Phi^{-1}(\Phi(k))$ is $\text{OD}_{\{\Phi\}}$ and thus $k \in \text{HOD}_{\{\Phi\}}$. By absoluteness, $\text{HOD}_{\{\Phi\}} \models$
750 “ $k : \omega \rightarrow \omega_2$ is cofinal” which is impossible since successor cardinals are regular in $\text{HOD}_{\{\Phi\}} \models \text{AC}$. $\square$

751 At the moment, the only known proof of the cardinality computation, $|\mathbb{R} \times \omega_1| < |\omega_1|$, under classical
752 regularity property involves HOD-type models and Namba forcing.⁸ However, this proof involving Namba
753 forcing is the only proof of this result suitable for generalization to higher cardinals. Section 7 will use a
754 form of Namba forcing to prove the analog of this result for all cardinals below Θ in AD^+ .

755 Next one will resolve Question (b) from this same hypothesis.

756 **Fact 3.6.** Let $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}}, 1_{\mathbb{P}})$ and $\mathbb{Q} = (\mathbb{Q}, \leq_{\mathbb{Q}}, 1_{\mathbb{Q}})$ be two forcings. Let $\pi : \mathbb{Q} \rightarrow \mathbb{P}$ be a map such that
757 for all $q_0, q_1 \in \mathbb{Q}$, $q_0 \leq_{\mathbb{Q}} q_1$ implies $\pi(q_0) \leq_{\mathbb{P}} \pi(q_1)$ and $\pi(1_{\mathbb{Q}}) = 1_{\mathbb{P}}$. The map $\Lambda_{\pi} : \mathcal{F}_{\mathbb{Q}} \rightarrow \mathcal{F}_{\mathbb{P}}$ defined by
758 $\Lambda_{\pi}(F) = \langle \pi[F] \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists q \in F)(\pi(q) \leq_{\mathbb{P}} p)\}$ is a well defined continuous map (with respect to the
759 topology on $\mathcal{F}_{\mathbb{Q}}$ and the topology on $\mathcal{F}_{\mathbb{P}}$).

760 *Proof.* Suppose $F \in \mathcal{F}_{\mathbb{Q}}$. Then $\langle \pi[F] \rangle_{\mathbb{P}} \in \mathcal{F}_{\mathbb{P}}$. To see this, suppose $p_0, p_1 \in \langle \pi[F] \rangle_{\mathbb{P}}$. Let $q_0, q_1 \in \mathbb{Q}$ be such
761 that $\pi(q_0) \leq_{\mathbb{P}} p_0$ and $\pi(q_1) \leq_{\mathbb{P}} p_1$. Since F is a $\mathbb{Q}$ -filter, there is a $q_2 \in F$ such that $q_2 \leq_{\mathbb{Q}} q_0$ and $q_2 \leq_{\mathbb{Q}} q_1$.
762 Thus $\pi(q_2) \in \langle \pi[F] \rangle_{\mathbb{P}}$, $\pi(q_2) \leq_{\mathbb{P}} \pi(q_0) \leq_{\mathbb{P}} p_0$, and $\pi(q_2) \leq_{\mathbb{P}} \pi(q_1) \leq_{\mathbb{P}} p_1$. $\langle \pi[F] \rangle_{\mathbb{P}}$ is clearly $\leq_{\mathbb{P}}$ -upward closed.
763 Let s be $\mathbb{P}$ -suitable. Suppose $F \in \Lambda_{\pi}^{-1}[M_s^{\mathbb{P}}]$. Since $\Lambda_{\pi}(F) = \langle \pi[F] \rangle_{\mathbb{P}} \in M_s^{\mathbb{P}}$ is a filter, there is a $\tilde{p} \in \Lambda_{\pi}(F)$
764 so that $\tilde{p} \leq_{\mathbb{P}} p$ for all $p \in \text{dom}(s)$ with $s(p) = 1$ and $\tilde{p}$ is incompatible with all $p \in \text{dom}(s)$ with $s(p) = 0$.
765 By definition of $\tilde{p} \in \langle \pi[F] \rangle_{\mathbb{P}}$, there is a $\tilde{q} \in F$ so that $\pi(\tilde{q}) \leq_{\mathbb{P}} \tilde{p}$. Let $t : \{\tilde{q}\} \rightarrow 2$ be define by $t(\tilde{q}) = 1$. It
766 is clear that $F \in M_t^{\mathbb{Q}}$. Now suppose that $H \in M_t^{\mathbb{Q}}$. Since $\pi(\tilde{q}) \leq_{\mathbb{P}} \tilde{p}$, $\tilde{p} \in \langle \pi[H] \rangle_{\mathbb{P}} = \Lambda_{\pi}(H)$. Since $\tilde{p} \leq_{\mathbb{P}} p$
767 for all $p \in \text{dom}(s)$ with $s(p) = 1$ and $\tilde{p}$ is not compatible with any $p \in \text{dom}(s)$ with $s(p) = 0$, one has
768 that $p \in \Lambda_{\pi}(H)$ for all $p \in \text{dom}(s)$ with $s(p) = 1$ and $p \notin \Lambda_{\pi}(H)$ for all $p \in \text{dom}(s)$ with $s(p) = 0$. Thus
769 $\Lambda_{\pi}(H) \in M_s^{\mathbb{Q}}$. It has been shown that $F \in M_t^{\mathbb{Q}} \subseteq \Lambda_{\pi}^{-1}[M_s^{\mathbb{P}}]$. This shows that Λ_{π} is continuous. $\square$

770 Some basic properties of forcing projection will be presented next. (Forcing projection of the ∞ -Borel
771 code forcings will be very important in Section 4.)

772 **Definition 3.7.** Let $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}}, 1_{\mathbb{P}})$ and $\mathbb{Q} = (\mathbb{Q}, \leq_{\mathbb{Q}}, 1_{\mathbb{Q}})$ be two forcings. A function $\pi : \mathbb{Q} \rightarrow \mathbb{P}$ is a forcing
773 projection if and only if the following holds:

- 774 (1) For all $q_0, q_1 \in \mathbb{Q}$, $q_0 \leq_{\mathbb{Q}} q_1$ implies $\pi(q_0) \leq_{\mathbb{P}} \pi(q_1)$ and $\pi(1_{\mathbb{Q}}) = 1_{\mathbb{P}}$.
- 775 (2) For all $q \in \mathbb{Q}$ and $p' \in \mathbb{P}$ such that $p' \leq_{\mathbb{P}} \pi(q)$, there is a $q' \in \mathbb{Q}$ so that $q' \leq_{\mathbb{Q}} q$ and $\pi(q') = p'$.

776 Suppose M is an inner model of ZF so that $\mathbb{P}, \mathbb{Q}, \pi \in M$ and $M \models “\pi : \mathbb{Q} \rightarrow \mathbb{P}$ is a forcing projection”.
777 If $H \subseteq \mathbb{Q}$ be $\mathbb{Q}$ -generic over M , then $\Lambda_{\pi}(H) = \langle \pi[H] \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists q \in H)(\pi(q) \leq_{\mathbb{P}} p)\}$ is a $\mathbb{P}$ -generic
778 filter over M . To see this, suppose $D \in M$ is $\mathbb{P}$ -dense. Let $\tilde{D} = \pi^{-1}[D]$ which belongs to M . Let $q \in \mathbb{Q}$.
779 Since D is dense there is a $p' \in D$ so that $p' \leq_{\mathbb{P}} \pi(q)$. By the properties of a forcing projection, there is a
780 $q' \leq_{\mathbb{Q}} q$ so that $\pi(q') = p'$ and hence $q' \in \tilde{D}$. This shows that $\tilde{D}$ is dense in $\mathbb{Q}$. Since H is $\mathbb{Q}$ -generic over
781 M , $H \cap \tilde{D} \neq \emptyset$. Let $q \in H \cap \tilde{D}$. Then $\pi(q) \in \langle \pi[H] \rangle_{\mathbb{P}} \cap D$.

782 Let $G \subseteq \mathbb{P}$ be $\mathbb{P}$ -generic over M . Let $\mathbb{Q}/_{\pi}G = \{q \in \mathbb{Q} : \pi(q) \in G\}$, $\leq_{\mathbb{Q}/_{\pi}G} = \leq_{\mathbb{Q}} \upharpoonright \mathbb{Q}/_{\pi}G$, and $1_{\mathbb{Q}/_{\pi}G} = 1_{\mathbb{Q}}$.
783 $\mathbb{Q}/_{\pi}G = (\mathbb{Q}/_{\pi}G, \leq_{\mathbb{Q}/_{\pi}G}, 1_{\mathbb{Q}/_{\pi}G})$ is a forcing in $M[G]$ and is called the remainder forcing of $\mathbb{Q}$ by G . Let $\dot{G}$
784 denote the canonical $\mathbb{P}$ -name for its generic filter. Let $\mathbb{Q}/_{\pi}\dot{G}$ be the canonical $\mathbb{P}$ -name such that $1_{\mathbb{P}} \Vdash_{\mathbb{P}} \mathbb{Q}/_{\pi}\dot{G}$
785 $\pi\dot{G} = \{q \in \mathbb{Q} : \check{\pi}(q) \in \dot{G}\}$. Note that if G is $\mathbb{P}$ -generic over M , then $(\mathbb{Q}/_{\pi}\dot{G})[G] = \mathbb{Q}/_{\pi}G$. $\mathbb{Q}$ is forcing
786 equivalent to the two step iterated forcing $\mathbb{P} * (\mathbb{Q}/_{\pi}\dot{G})$. If $H \subseteq \mathbb{Q}$ is $\mathbb{Q}$ -generic over M and $G = \langle \pi[H] \rangle_{\mathbb{P}}$

⁸The important feature of Namba forcing is that it adds a new ω -sequence below the true ω_1 without adding any new reals. Prikry forcing at a measurable cardinal adds a new ω -sequence without adding any bounded subsets below the measurable. Under the hypothesis of Theorem 3.5, there may be no inner models with a measurable. In fact, even in $L(\mathbb{R}) \models \text{AD}$, no ordinals which is countable in the real world can be measurable in $\text{HOD}^{L(\mathbb{R})}$. Steel’s HOD-analysis in $L(\mathbb{R})$ ([60] Lemma 8.25) shows that the real world never has a Prikry generic sequence over $\text{HOD}^{L(\mathbb{R})}$ for any measurable cardinal of $\text{HOD}^{L(\mathbb{R})}$.

787 which is $\mathbb{P}$ -generic over M , then H is $\mathbb{Q}/G$ -generic over $M[G]$, $G * H$ is $\mathbb{P} * (\mathbb{Q}/\pi\dot{G})$ -generic over M , and
 788 $M[H] = M[G][H] = M[G * H]$. (See the introduction of [1] for more information.)

789 **Fact 3.8.** *Suppose $\pi : \mathbb{Q} \rightarrow \mathbb{P}$ is a forcing projection. If E is dense open in the topology on $\mathcal{F}_{\mathbb{P}}$, then*
 790 *$\Lambda_{\pi}^{-1}[E]$ is dense open in the topology of $\mathcal{F}_{\mathbb{Q}}$. Thus for any $C \subseteq \mathcal{F}_{\mathbb{P}}$ which is comeager in the topology of $\mathcal{F}_{\mathbb{P}}$,*
 791 *$\Lambda_{\pi}^{-1}[C] \subseteq \mathcal{F}_{\mathbb{Q}}$ is comeager in the topology of $\mathcal{F}_{\mathbb{Q}}$.*

792 *Proof.* Since Λ_{π} is continuous by Fact 3.6, $\Lambda_{\pi}^{-1}[E]$ is open. Suppose s is $\mathbb{Q}$ -suitable. There is a $\tilde{q}$ so that
 793 $\tilde{q} \leq_{\mathbb{Q}} q$ for all $q \in \text{dom}(s)$ with $s(q) = 1$ and $\tilde{q}$ is not compatible with any $q \in \text{dom}(s)$ with $s(q) = 0$. Let
 794 $u = \{(\pi(\tilde{q}), 1)\}$ which is $\mathbb{P}$ -suitable. Since E is dense open, there is a $\mathbb{P}$ -suitable t so that $M_t^{\mathbb{P}} \subseteq M_u^{\mathbb{P}} \cap E$,
 795 $\pi(\tilde{q}) \in \text{dom}(t)$, and $t(\pi(\tilde{q})) = 1$. Since t is $\mathbb{P}$ -suitable, there is a $p' \in \mathbb{P}$ so that for all $p \in \text{dom}(t)$ with
 796 $t(p) = 1$, $p' \leq_{\mathbb{P}} p$ and p' is not compatible with any $p \in \text{dom}(t)$ with $t(p) = 0$. Since $\pi(\tilde{q}) \in \text{dom}(t)$, one has
 797 $p' \leq_{\mathbb{P}} \pi(\tilde{q})$. By the property of projections, there is a $q' \leq_{\mathbb{Q}} \tilde{q}$ so that $\pi(q') = p'$. Let $v = s \cup \{(q', 1)\}$ which
 798 is $\mathbb{Q}$ -suitable. Note that $\Lambda_{\pi}[M_v^{\mathbb{Q}}] \subseteq M_t^{\mathbb{P}} \subseteq E$. Thus $M_v^{\mathbb{Q}} \subseteq \Lambda_{\pi}^{-1}[E] \cap M_s^{\mathbb{Q}}$. This shows $\Lambda_{\pi}^{-1}[E]$ is dense open.

799 Suppose $C \subseteq \mathcal{F}_{\mathbb{P}}$ is comeager. There is a countable sequence $\langle E_n : n \in \omega \rangle$ of dense open subsets of $\mathcal{F}_{\mathbb{P}}$ such
 800 that $\bigcap_{n \in \omega} E_n \subseteq C$. Then $\bigcap_{n \in \omega} \Lambda_{\pi}^{-1}[E_n] = \pi^{-1}(\bigcap_{n \in \omega} E_n) \subseteq \Lambda_{\pi}^{-1}[C]$. Since $\bigcap_{n \in \omega} \Lambda_{\pi}^{-1}[E_n]$ is a countable
 801 intersection of dense open subsets of $\mathcal{F}_{\mathbb{Q}}$, $\Lambda_{\pi}^{-1}[C]$ is comeager. $\square$

802 **Lemma 3.9.** *Assume $\text{AC}_{\omega}^{\mathbb{R}}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at ω .*
 803 *Let Z be a set such that there is an OD_Z wellordering of Z . Let $\mathbb{Q} \in \text{HOD}_Z$ be a forcing which is countable*
 804 *in the real world. There is a $U_Z^{\mathbb{Q}} \subseteq \text{Gen}_{\mathbb{Q}}^{\mathcal{D}_Z^{\mathbb{Q}}}$ comeager in the topology of $\mathcal{F}_{\mathbb{Q}}$ and obtained uniformly from $\mathbb{Q}$*
 805 *and Z with the property that for all $G \in U_Z^{\mathbb{Q}}$ and forcing projection $\pi : \mathbb{Q} \rightarrow \mathbb{P}$ such that $\pi \in \text{HOD}_Z$ and*
 806 *$\mathbb{P} \in \text{HOD}_Z$ is a countable forcing in the real world, $\text{HOD}_Z[\langle \pi[G] \rangle_{\mathbb{P}}] = \text{HOD}_{Z \cup \{\langle \pi[G] \rangle_{\mathbb{P}}\}}$.*

807 *Proof.* Let $\langle \pi_{\alpha} : \alpha \in \text{ON} \rangle$ be an enumeration of all forcing projections in HOD_Z of $\mathbb{Q}$ into some $\mathbb{P}_{\alpha}$ where $\mathbb{P}_{\alpha}$
 808 is countable in the real world according to the canonical global wellordering of HOD_Z . (The enumeration
 809 does not need to belong to HOD_Z .) Let $T_Z^{\mathbb{P}_{\alpha}}$ be the comeager set from Lemma 3.2 for $\mathbb{P}_{\alpha}$ (which is obtained
 810 uniformly from $\mathbb{P}_{\alpha}$ and Z). For each $\alpha \in \text{ON}$, $\Lambda_{\pi_{\alpha}}^{-1}[T_Z^{\mathbb{P}_{\alpha}}] \subseteq \mathcal{F}_{\mathbb{Q}}$ is comeager by Fact 3.8. Then $U_Z^{\mathbb{Q}} =$
 811 $\text{Gen}_{\mathbb{Q}}^{\mathcal{D}_Z^{\mathbb{Q}}} \cap \bigcap_{\alpha \in \text{ON}} \Lambda_{\pi_{\alpha}}^{-1}[T_Z^{\mathbb{P}_{\alpha}}]$ is a comeager subset of $\text{Gen}_{\mathbb{Q}}^{\mathcal{D}_Z^{\mathbb{Q}}}$ since wellordered intersections of comeager subsets
 812 of $\mathcal{F}_{\mathbb{Q}}$ remain comeager subsets of $\mathcal{F}_{\mathbb{Q}}$ under the given hypothesis. If $\pi : \mathbb{Q} \rightarrow \mathbb{P}$ is such a forcing projection,
 813 then there is an $\alpha \in \text{ON}$ so that $\pi = \pi_{\alpha}$ and $\mathbb{P} = \mathbb{P}_{\alpha}$. Let $G \in U_Z^{\mathbb{Q}}$. Then $\langle \pi[G] \rangle_{\mathbb{P}} = \Lambda_{\pi_{\alpha}}(G) \in T_Z^{\mathbb{P}_{\alpha}} = T_Z^{\mathbb{P}}$.
 814 By the property of $T_Z^{\mathbb{P}}$ coming from Lemma 3.2, $\text{HOD}_Z[\langle \pi[G] \rangle_{\mathbb{P}}] = \text{HOD}_{Z \cup \{\langle \pi[G] \rangle_{\mathbb{P}}\}}$. $\square$

815 **Fact 3.10.** $|\mathbb{R}| \leq |S_1|$. *If boldface GCH at ω holds, then $\neg(|\omega_1| \leq |S_1|)$.*

Proof. For $r \in [\omega]^{\omega}$, let $f_r : \omega_1^{L[r]} \rightarrow \omega_1^{L[r]}$ be defined by

$$f_r(\alpha) = \begin{cases} r(\alpha) & \alpha \in \omega \\ \alpha & \omega \leq \alpha < \omega_1^{L[r]} \end{cases}.$$

816 Note that $f_r \in L[r]$ and that $\omega_1^{L[f_r]} = \omega_1^{L[r]} = \sup(f_r)$. Thus $f_r \in S_1$. The map $\Phi : \mathbb{R} \rightarrow S_1$ defined by
 817 $\Phi(r) = f_r$ is an injection.

818 Now assume boldface GCH at ω . Suppose $\Psi : \omega_1 \rightarrow S_1$ is an injection. The claim is that $\omega_1 =$
 819 $\sup\{\sup(\Psi(\alpha)) : \alpha < \omega_1\}$. If not, then this supremum is some countable ordinal δ . Then one has that
 820 $\Psi : \omega_1 \rightarrow [\delta]^{<\delta}$. Since δ is countable, there is a bijection $\Upsilon : [\delta]^{<\delta} \rightarrow \mathbb{R}$. Then $\Upsilon \circ \Psi : \omega_1 \rightarrow \mathbb{R}$ is an
 821 injection which violates boldface GCH at ω . Let $A = \{(\alpha, \beta, \gamma) \in \omega_1 \times \omega_1 \times \omega_1 : \Psi(\alpha)(\beta) = \gamma\}$. $L[A] \models \text{ZFC}$
 822 and thus by boldface GCH at ω , $\omega_1^{L[A]} \leq (2^{\omega})^{L[A]} < \omega_1$. By the claim, there is some $\alpha < \omega_1$ so that
 823 $\sup(\Psi(\alpha)) > \omega_1^{L[A]}$. Since $\Psi(\alpha) \in L[A]$ and $\Psi(\alpha) \in S_1$, one has that $\omega_1^{L[A]} \geq \omega_1^{L[\Psi(\alpha)]} = \sup(\Psi(\alpha)) > \omega_1^{L[A]}$
 824 which is a contradiction. $\square$

825 **Theorem 3.11.** *Assume $\text{AC}_{\omega}^{\mathbb{R}}$, all subsets of all Polish spaces have the Baire property, and boldface GCH at*
 826 *ω . Then S_1 does not inject into ${}^{\omega}\text{ON}$.*

Proof. Suppose there is an injection $\Phi : S_1 \rightarrow {}^{\omega}\text{ON}$. For $\alpha < \omega_1^{\text{HOD}_{\{\Phi\}}}$, let $\mathbb{K}_{\alpha} = \text{Coll}(\omega, \alpha)$ be the forcing of
 finite partial function p on ω into α . Let $\leq_{\mathbb{K}_{\alpha}}$ be reverse function extension. Let $1_{\mathbb{K}_{\alpha}} = \emptyset$. For $\beta \leq \omega_1^{\text{HOD}_{\{\Phi\}}}$,

let $\mathbb{Q}_\beta = \text{Coll}(\omega, < \beta)$ be the collection of all functions $\mathbf{p}$ with domain β so that for all $\alpha < \beta$, $\mathbf{p}(\alpha) \in \mathbb{K}_\alpha$ and $\text{supp}^\beta(\mathbf{p}) = \{\alpha < \beta : \mathbf{p}(\alpha) \neq 1_{\mathbb{K}_\alpha}\}$ is finite. Let $\mathbf{p} \leq_{\mathbb{Q}_\beta} \mathbf{q}$ if and only if for all $\alpha < \beta$, $\mathbf{p}(\alpha) \leq_{\mathbb{K}_\alpha} \mathbf{q}(\alpha)$. Let $1_{\mathbb{Q}_\beta}$ be defined by $1_{\mathbb{Q}_\beta}(\alpha) = 1_{\mathbb{K}_\alpha}$. $\mathbb{Q}_\beta = (\mathbb{Q}_\beta, \leq_{\mathbb{Q}_\beta}, 1_{\mathbb{Q}_\beta})$ is the Lévy collapse of the ordinals below β to ω . Let $\mathbb{Q} = \mathbb{Q}_{\omega_1^{\text{HOD}\{\Phi\}}} = \text{Coll}(\omega, < \omega_1^{\text{HOD}\{\Phi\}})$.⁹ For $\beta < \omega_1^{\text{HOD}\{\Phi\}}$, let $\pi_\beta : \mathbb{Q} \rightarrow \mathbb{Q}_\beta$ be defined by $\pi_\beta(\mathbf{p}) = \mathbf{p} \upharpoonright \beta$. Suppose $\mathbf{p} \in \mathbb{Q}_\beta$, $\mathbf{q} \in \mathbb{Q}$, and $\mathbf{p} \leq_{\mathbb{Q}_\beta} \mathbf{q} \upharpoonright \beta = \pi_\beta(\mathbf{q})$. Let $\mathbf{q}' \in \mathbb{Q}$ be defined by

$$\mathbf{q}'(\alpha) = \begin{cases} \mathbf{p}(\alpha) & \alpha < \beta \\ \mathbf{p}(\alpha) = \mathbf{q}(\alpha) & \beta \leq \alpha < \omega_1^{\text{HOD}\{\Phi\}} \end{cases}.$$

Then $\mathbf{q}' \leq_{\mathbb{Q}} \mathbf{q}$ and $\pi_\beta(\mathbf{q}') = \mathbf{p}$. This shows that π_β is a forcing projection. By boldface GCH at ω , $\mathbb{Q}_\beta$ and $\mathbb{Q}$ are countable for all $\beta < \omega_1^{\text{HOD}\{\Phi\}}$. By Lemma 3.9, let $G \subseteq \mathbb{Q}$ be $\mathbb{Q}$ -generic over $\text{HOD}_{\{\Phi\}}$ having the property that $\text{HOD}_{\{\Phi\}}[G] = \text{HOD}_{\{\Phi, G\}}$ and for all $\beta < \omega_1^{\text{HOD}\{\Phi\}}$, $\text{HOD}_{\{\Phi\}}[\langle \pi_\beta[G] \rangle_{\mathbb{Q}_\beta}] = \text{HOD}_{\{\Phi, \langle \pi_\beta[G] \rangle_{\mathbb{Q}_\beta}\}}$. The generic G can be coded naturally as a subset of $\omega_1^{\text{HOD}\{\Phi\}}$ and hence as an element $\tilde{g} \in [\omega_1^{\text{HOD}\{\Phi\}}]^{\omega_1^{\text{HOD}\{\Phi\}}}$. Thus $\text{sup}(\tilde{g}) \leq \omega_1^{\text{HOD}\{\Phi\}}$. If done naturally, one will have $L[G] = L[\tilde{g}]$. $\text{HOD}_{\{\Phi\}} \models$ “ $\mathbb{Q}$ has the $\omega_1^{\text{HOD}\{\Phi\}}$ -chain condition” (see [50] Lemma 6.44 using the Δ -system lemma). Thus $\omega_1^{\text{HOD}\{\Phi\}}[G] = \omega_1^{\text{HOD}\{\Phi\}}$. Since $\tilde{g} \in \text{HOD}_{\{\Phi\}}[G]$, $\omega_1^{L[\tilde{g}]} \leq \omega_1^{\text{HOD}_{\{\Phi\}}[G]} = \omega_1^{\text{HOD}\{\Phi\}}$. Since $\tilde{g}$ codes a $\mathbb{K}_\beta$ -generic filter for each $\beta < \omega_1^{\text{HOD}\{\Phi\}}$, $\omega_1^{L[\tilde{g}]} \geq \omega_1^{\text{HOD}\{\Phi\}}$ and also $\text{sup}(\tilde{g}) \geq \omega_1^{\text{HOD}\{\Phi\}}$. This shows that $\text{sup}(\tilde{g}) = \omega_1^{\text{HOD}\{\Phi\}} = \omega_1^{L[\tilde{g}]}$. So $\tilde{g} \in S_1$. Since $\tilde{g}$ is $\text{OD}_{\{G\}}$, $\Phi(\tilde{g}) \in \text{HOD}_{\{\Phi, G\}} = \text{HOD}_{\{\Phi\}}[G]$. $\Phi(\tilde{g}) \in {}^\omega\text{ON}$ and thus there is a $\mathbb{Q}$ -name τ so that $\tau[G] = \Phi(\tilde{g})$. Without loss of generality, $1_{\mathbb{Q}} \Vdash_{\mathbb{Q}}$ “ τ is a function from $\tilde{\omega}$ into ON ”. In $\text{HOD}_{\{\Phi\}}$, for each $n \in \omega$, let A_n be a maximal antichain inside of $\{\mathbf{q} \in \mathbb{Q} : \mathbf{q} \Vdash_{\mathbb{Q}} (\exists \xi \in \text{ON})(\tau(\tilde{n}) = \xi)\}$. Thus $\text{HOD}_{\{\Phi\}} \models$ “ A_n is countable” by the $\omega_1^{\text{HOD}\{\Phi\}}$ -chain condition. Then $\delta_n = \text{sup}\{\max(\text{supp}(\mathbf{q})) : \mathbf{q} \in A_n\} < \omega_1^{\text{HOD}\{\Phi\}}$ since $\omega_1^{\text{HOD}\{\Phi\}}$ is regular in $\text{HOD}_{\{\Phi\}}$. Similarly, $\delta = \text{sup}\{\delta_n : n \in \omega\}$ is less than $\omega_1^{\text{HOD}\{\Phi\}}$. For each $n \in \omega$ and $\mathbf{q} \in A_n$, let α_q^n be the unique α so that $\text{HOD}_{\{\Phi\}} \models \mathbf{q} \Vdash_{\mathbb{Q}} \tau(\tilde{n}) = \alpha$. Let $\sigma = \{((n, \alpha_q^n)^\sim, \mathbf{q} \upharpoonright \delta) : n \in \omega \wedge \mathbf{q} \in A_n\}$. σ is a $\mathbb{Q}_\delta$ -name and $\Phi(\tilde{g}) = \tau[G] = \sigma[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}]$. This shows that $\Phi(\tilde{g}) \in \text{HOD}_{\{\Phi\}}[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}] = \text{HOD}_{\{\Phi, \langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}\}}$. Since $\Phi(\tilde{g})$ is $\text{OD}_{\{\Phi, \langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}\}}$, $\tilde{g} = \Phi^{-1}(\Phi(\tilde{g}))$ is also $\text{OD}_{\{\Phi, \langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}\}}$. Thus $\tilde{g} \in \text{HOD}_{\{\Phi, \langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}\}} = \text{HOD}_{\{\Phi\}}[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}]$. Since $L[\tilde{g}] = L[G]$, one has that $G \in \text{HOD}_{\{\Phi\}}[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}]$ which is impossible since G is $(\mathbb{Q}/\pi_\delta \dot{G})$ -generic over $\text{HOD}_{\{\Phi\}}[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}]$ and $\text{HOD}_{\{\Phi\}}[G] = \text{HOD}_{\{\Phi\}}[\langle \pi_\delta[G] \rangle_{\mathbb{Q}_\delta}][G]$ by the discussion above about remainder forcing applied to the forcing projection π_δ . $\square$

4. ∞ -BOREL CODE FORCING

This section will present ∞ -Borel code forcing (with ordinal definability constraints) on ${}^\omega\kappa$, its directed limit, and the analysis of natural determinacy models as symmetric collapse extension of HOD -type models. These results in various forms are due to Woodin and Ikegami-Trang ([25]). The uniformity of these results will be very important in the main result of the next section.

This section presents detailed proofs of the results in the generality needed in this paper. The results in this section are not very straightforward and simple generalizations of the classical OD ∞ -Borel code forcing on $\mathbb{R}$ in $L(\mathbb{R})$ as presented in [8] which is uniquely simple. There are important differences between Vopěnka forcing and ∞ -Borel code forcing and both are needed in the appropriate situation. The proofs of results for ${}^\omega\kappa$ when $\omega_1 \leq \kappa < \Theta$ require substantial changes from the result for just $\mathbb{R}$ or ${}^\omega\omega$. One needs to be very careful and explicit when relativizing from OD to OD_Z . From the author's experience, argument involving ordinal definability and Vopěnka-style forcings are very susceptible to hidden mistakes which lead to major errors.

To avoid frequent repetition, one makes the following definition.

Definition 4.1. A set Z is a parameter set if and only if $Z \in \text{OD}_Z$ and there is an OD_Z wellordering of Z .

⁹Typically, one would use $\text{Coll}(\omega, < \kappa)$ where κ is an inaccessible cardinal of an inner model of ZFC and κ is countable in the real world. This is the forcing used in [13] Theorem 5.7. Such an inaccessible cardinal exists under AD. However, the hypothesis of Theorem 3.11 cannot prove such an inaccessible exists. For instance, in the minimal Solovay model: Let κ be the least inaccessible cardinal of L . Let $G \subseteq \text{Coll}(\omega, < \kappa)$ be generic over L . Then the Solovay model $L(\mathbb{R}^{L[G]})$ has the property that no countable ordinal is inaccessible in any inner model of ZFC.

After fixing a wellordering $\prec$ of Z which is OD_Z , one can defined a global wellordering of HOD_Z . Thus $\text{HOD}_Z \models \text{ZFC}$. This global wellordering will often be called the canonical wellordering of HOD_Z . Note also that there is a finite $F \subseteq Z$ so that $\prec \in \text{OD}_F$. With $\prec$, all other elements of Z can be defined by referring to their rank in $\prec$. Thus $\text{HOD}_Z = \text{HOD}_F$.

Definition 4.2. Recall that $\mathcal{B}_\infty$ is the class of all ∞ -Borel codes. (That is, $p \in \mathcal{B}_\infty$ if and only if p is a pair (S, φ) where S is a set of ordinals and φ is a $\mathcal{L}_{\text{set}}$ -formula.)

Let κ be an ordinal and Z be a parameter set. Let $n \in \omega$. Let $\mathcal{BC}_n^{\kappa;Z} \subseteq \mathcal{B}_\infty$ be the proper class of ∞ -Borel codes $p = (S, \varphi)$ which belong to HOD_Z (that is, $S \in \text{HOD}_Z$) and $\mathfrak{B}_p^{n(\omega\kappa)} = \{y \in {}^n(\omega\kappa) : L[S, y] \models \varphi(S, y)\} \neq \emptyset$. (If $n = 0$, then ${}^0(\omega\kappa) = \{\emptyset\}$.) For each $n \in \omega$, define a proper class equivalence relation $\sim_n^{\kappa;Z}$ on $\mathcal{BC}_n^{\kappa;Z}$ by $p \sim_n^{\kappa;Z} q$ if and only if $\mathfrak{B}_p^{n(\omega\kappa)} = \mathfrak{B}_q^{n(\omega\kappa)}$. If $p \in \mathcal{BC}_n^{\kappa;Z}$, let $[p]_{\sim_n^{\kappa;Z}}$ be the $\sim_n^{\kappa;Z}$ -equivalence class containing p (which is a proper class). Let $\mathbb{B}_n^{\kappa;Z}$ be the collection of all $p \in \mathcal{BC}_n^{\kappa;Z}$ such that p is the minimal element of $[p]_{\sim_n^{\kappa;Z}}$ according to the canonical wellordering of HOD_Z . (So all elements of $\mathbb{B}_n^{\kappa;Z}$ are ∞ -Borel codes.) For $p, q \in \mathbb{B}_n^{\kappa;Z}$, define $p \leq_{\mathbb{B}_n^{\kappa;Z}} q$ if and only if $\mathfrak{B}_p^{n(\omega\kappa)} \subseteq \mathfrak{B}_q^{n(\omega\kappa)}$. Let $1_{\mathbb{B}_n^{\kappa;Z}}$ be the unique $p \in \mathbb{B}_n^{\kappa;Z}$ such that $\mathfrak{B}_p^{n(\omega\kappa)} = {}^n(\omega\kappa)$. Note that $\mathbb{B}_n^{\kappa;Z}, \leq_{\mathbb{B}_n^{\kappa;Z}}, 1_{\mathbb{B}_n^{\kappa;Z}} \in \text{HOD}_Z$. Let $\mathbb{B}_n^{\kappa;Z}$ also be used to refer to the forcing poset $(\mathbb{B}_n^{\kappa;Z}, \leq_{\mathbb{B}_n^{\kappa;Z}}, 1_{\mathbb{B}_n^{\kappa;Z}})$. $\mathbb{B}_n^{\kappa;Z}$ is called the n -dimensional OD_Z ∞ -Borel code forcing on $\omega\kappa$.

This forcing resembles the classical Vopěnka forcing of non-empty OD_Z subsets of ${}^n(\omega\kappa)$ defined later in Definition 4.8. The ∞ -Borel code forcing will be very important tool in this paper, but the classical Vopěnka forcing will be needed in a few places to establish properties of the ∞ -Borel code forcing. In many important cases in natural models of AD^+ , these two forcings are the same but only after a careful analysis of ∞ -Borel code.

Definition 4.3. Let κ be an ordinal. Let Z be a parameter set. For $0 \leq i < n$, $j \in \omega$, and $\alpha \in \kappa$, let $A_{n,i,j,\alpha} = \{y \in {}^n(\omega\kappa) : y(i)(j) = \alpha\}$ which clearly has an OD_Z ∞ -Borel code. Let $p_{n,i,j,\alpha}^{\kappa;Z} \in \mathbb{B}_n^{\kappa;Z}$ be such that $A_{n,i,j,\alpha} = \mathfrak{B}_{p_{n,i,j,\alpha}^{\kappa;Z}}^{n(\omega\kappa)}$. Let $\dot{\tau}^n = \{((i, j, \alpha), p_{n,i,j,\alpha}^{\kappa;Z}) : i < n \wedge j \in \omega \wedge \alpha \in \kappa\}$.¹⁰ $\dot{\tau}^n$ is the canonical name for adding the generic element of ${}^n(\omega\kappa)$. For $i < n$, let $\dot{\tau}_i^n$ be the canonical name for the element of $\omega\kappa$ which is the i^{th} term of $\dot{\tau}^n$. Note that $\dot{\tau}^n, \dot{\tau}_i^n \in \text{HOD}_Z$.

Definition 4.4. If $\mathcal{J} \subseteq \mathcal{B}_\infty$ is a set of ∞ -Borel codes, κ is an ordinal, and $x \in {}^n(\omega\kappa)$, then define $\mathcal{G}^{\kappa,n}(\mathcal{J}, x) = \{p \in \mathcal{J} : x \in \mathfrak{B}_p^{n(\omega\kappa)}\}$.

Let Z be a parameter set. (Note that $\mathbb{B}_n^{\kappa;Z} \subseteq \mathcal{B}_\infty$.) Let $G_x^{\kappa;Z;n} = \mathcal{G}^{\kappa;n}(\mathbb{B}_n^{\kappa;Z}, x) = \{p \in \mathbb{B}_n^{\kappa;Z} : x \in \mathfrak{B}_p^{n(\omega\kappa)}\}$. $G_x^{\kappa;Z;n}$ is the $\mathbb{B}_n^{\kappa;Z}$ -filter derived from x .

Definition 4.5. Let κ be a cardinal. Let Z be a parameter set. Uniformly let $\hat{\mathbb{B}}^{\kappa;Z}$ be a set of ordinals in HOD_Z which codes $\mathbb{B}_n^{\kappa;Z}$, every $p \in \mathbb{B}_n^{\kappa;Z}$, $\leq_{\mathbb{B}_n^{\kappa;Z}}$, $1_{\mathbb{B}_n^{\kappa;Z}}$, $p_{n,i,j,\alpha}^{\kappa;Z}$, $\dot{\tau}^n$, and $\dot{\tau}_i^n$ for all $j, n \in \omega$ and $i < n$.

This coding can be done in a variety of ways to maintain the following important properties. Uniformly in Z means there is a fixed simple procedure that takes Z and returns $\hat{\mathbb{B}}^{\kappa;Z}$. From $\hat{\mathbb{B}}^{\kappa;Z}$, one should be able to extract all the conditions of $\mathbb{B}_n^{\kappa;Z}$ for all $n \in \omega$. The main property of $\hat{\mathbb{B}}^{\kappa;Z}$ is that in any inner model $M \subseteq \text{HOD}_Z$ with $\hat{\mathbb{B}}^{\kappa;Z} \in M$, one can perform the relevant arguments which involve forcing with $\mathbb{B}_n^{\kappa;Z}$. Moreover, for any such inner model M and $y \in {}^n(\omega\kappa)$, $G_y^{\kappa;Z;n} = \mathcal{G}^{\kappa;n}(\mathbb{B}_n^{\kappa;Z}, y)$ can be defined inside of $M[y]$.

Fact 4.6. Let κ be an ordinal, Z be a parameter set, and $n \in \omega$. Let M be an inner model of HOD_Z with $\hat{\mathbb{B}}^{\kappa;Z} \in M$. Let $y \in {}^n(\omega\kappa)$. Then $G_y^{\kappa;Z;n}$ is $\mathbb{B}_n^{\kappa;Z}$ -generic over M and $\dot{\tau}^n[G_y^{\kappa;Z;n}] = y$. (In particular, this result applies to $M = \text{HOD}_Z$.)

Proof. Let $A \in M$ be a maximal antichain in M . Now work in the real world. Since $A \in M \subseteq \text{HOD}_Z$, let $\langle (S_\alpha, \varphi_\alpha) : \alpha < \delta \rangle$ for some ordinal δ be the enumeration of A according to the canonical global wellordering of HOD_Z . Let $T = \{(\alpha, \beta) : \alpha < \delta \wedge \beta \in S_\alpha\}$ where $\langle \alpha, \beta \rangle$ refers to the Gödel pairing function. Note that $T \in \text{HOD}_Z$ and one can use T to create an OD_Z ∞ -Borel code q_0 such that $\mathfrak{B}_{q_0}^{n(\omega\kappa)} = \bigcup \{\mathfrak{B}_p^{n(\omega\kappa)} : p \in A\}$. For the sake of contradiction, suppose $A \cap G_y^{\kappa;Z;n} = \emptyset$. Then $y \notin \mathfrak{B}_{q_0}^{n(\omega\kappa)}$. From q_0 , one can create an OD_Z ∞ -Borel code q_1 so that $\mathfrak{B}_{q_1}^{n(\omega\kappa)} = \omega\kappa \setminus \mathfrak{B}_{q_0}^{n(\omega\kappa)}$ which is a nonempty set as it contains y . Thus there is a

¹⁰ $\dot{\tau}^n \in \mathbb{B}_n^{\kappa;Z}$ is formally a name for a subset of $n \times \omega \times \kappa$ but will be considered as adding the associated element of ${}^n(\omega\kappa)$.

906 $q \in \mathbb{B}_n^{\kappa;Z}$ such that $\mathfrak{B}_q^{n(\omega\kappa)} = \mathfrak{B}_{q_1}^{n(\omega\kappa)}$. Now return to M . Since $\hat{\mathbb{B}}_n^{\kappa;Z} \in M$, every condition of $\mathbb{B}_n^{\kappa;Z}$ belongs
 907 to M (in particular, $q \in M$) and the forcing ordering $\leq_{\mathbb{B}_n^{\kappa;Z}}$ belongs to M . Then M sees that $A \cup \{q\} \in M$
 908 and is a larger antichain than A which contradicts the maximality of A . The second statement is clear. $\square$

909 One important advantage of $\mathbb{B}_n^{\kappa;Z}$ over the analogous classical Vopěnka forcing is the following fact.

910 **Fact 4.7.** *Let κ be an ordinal, Z be a parameter set, $n \in \omega$, and $y \in {}^n(\omega\kappa)$. Let M be an inner model of*
 911 *HOD_Z with $\hat{\mathbb{B}}_n^{\kappa;Z} \in M$. Then $M[y] = M[G_y^{\kappa;Z;n}]$. (In particular, $\text{HOD}_Z[y] = \text{HOD}[G_y^{\kappa;Z;n}]$.)*

912 *Proof.* Since $M \subseteq M[G_y^{\kappa;Z;n}]$, $\dot{\tau}^n \in M$ (since $\hat{\mathbb{B}}_n^{\kappa;Z} \in M$), and $y = \dot{\tau}^n[G_y^{\kappa;Z;n}] \in M[G_y^{\kappa;Z;n}]$, one has that
 913 $M[y] \subseteq M[G_y^{\kappa;Z;n}]$. Let $p \in \mathbb{B}_n^{\kappa;Z}$ and takes the form (S, φ) where S is a set of ordinals in HOD_Z and
 914 φ is a formula. $p \in G_y^{\kappa;Z;n}$ if and only if $L[S, y] \models \varphi(S, y)$ if and only if $M[y] \models L[S, y] \models \varphi(S, y)$ by the
 915 absoluteness of the satisfaction relation of $L[S, y]$. Thus $G_y^{\kappa;Z;n} \in M[y]$ by comprehension in M . This implies
 916 $M[G_y^{\kappa;Z;n}] \subseteq M[y]$. $\square$

917 The following is the Vopěnka forcing (on ω -sequences of ordinals) which will play a small but necessary
 918 role in this paper.

919 **Definition 4.8.** Let κ be an ordinal, $U \subseteq {}^\omega\kappa$, and Z be a parameter set with $U \in Z$. Let $n \in \omega$. Since Z is
 920 a parameter set, let $\prec$ be an OD_Z wellordering of Z . For each formula, tuple of ordinals, and tuple from Z
 921 (which will be identified with an ordinal by its rank in $\prec$), one can associate a OD_Z set. In this way, there is
 922 a OD_Z definable class surjection $\mathfrak{S}_Z : \text{ON} \rightarrow \text{OD}_Z$. Let $\mathbb{V}_n^{U;Z}$ be the set of ordinals α such that $\mathfrak{S}_Z(\alpha)$ is a
 923 nonempty subset of nU and for all $\alpha' < \alpha$, $\mathfrak{S}_Z(\alpha') \neq \mathfrak{S}_Z(\alpha)$. Note that $\mathbb{V}_n^{U;Z} \in \text{HOD}_Z$. For each $p, q \in \mathbb{V}_n^{U;Z}$,
 924 let $p \leq_{\mathbb{V}_n^{U;Z}} q$ if and only if $\mathfrak{S}_Z(p) \subseteq \mathfrak{S}_Z(q)$. Also note that $\leq_{\mathbb{V}_n^{U;Z}} \in \text{HOD}_Z$. Let $1_{\mathbb{V}_n^{U;Z}}$ be the least ordinal
 925 α so that $\mathfrak{S}_Z(\alpha) = {}^nU$. $\mathbb{V}_n^{U;Z} = (\mathbb{V}_n^{U;Z}, \leq_{\mathbb{V}_n^{U;Z}}, 1_{\mathbb{V}_n^{U;Z}})$ is the n -dimensional OD_Z Vopěnka-forcing on U and
 926 $\mathbb{V}_n^{U;Z} \in \text{HOD}_Z$.

927 If $y \in {}^nU$, let $\tilde{G}_y^{U;Z;n} = \{p \in \mathbb{V}_n^{U;Z} : y \in \mathfrak{S}_Z(p)\}$. This is the $\mathbb{V}_n^{U;Z}$ -filter associated to y .

928 For $0 \leq i < n$, $j \in \omega$, and $\alpha \in \kappa$, let $A_{n,i,j,\alpha} = \{y \in {}^nU : y(i)(j) = \alpha\}$ which is clearly OD_Z . If $A_{n,i,j,\alpha} \neq \emptyset$,
 929 then let $p_{n,i,j,\alpha}^{U;Z} \in \mathbb{V}_n^{U;Z}$ be such that $\mathfrak{S}_Z(p_{n,i,j,\alpha}^{U;Z}) = A_{n,i,j,\alpha}$. Let $\dot{\tau}^n = \{((i, j, \alpha), p_{n,i,j,\alpha}^{U;Z}) : i < n \wedge j \in \omega \wedge \alpha \in$
 930 $\kappa \wedge A_{n,i,j,\alpha} \neq \emptyset\}$. $\dot{\tau}^n$ is the canonical name for adding the generic element of ${}^n(\omega\kappa)$. For $i < n$, let $\dot{\tau}_i^n$ be the
 931 canonical name for the element of ${}^\omega\kappa$ which is the i th term of $\dot{\tau}^n$. Note that $\tau^n, \dot{\tau}_i^n \in \text{HOD}_Z$. (The notation
 932 from the ∞ -Borel code forcing has been reused, but this should not cause confusion in context.)

933 Let $\hat{\mathbb{V}}^{U;Z} \in \text{HOD}_Z$ be a set of ordinals which codes all the conditions of $\mathbb{V}_n^{U;Z}$ for all $n \in \omega$, the set
 934 $\{(n, i, j, \alpha) : j, n \in \omega \wedge i < n \wedge \alpha \in \kappa \wedge A_{n,i,j,\alpha} \neq \emptyset\}$, the map $\langle p_{n,i,j,\alpha}^{U;Z} : j, n \in \omega \wedge i < n \wedge \alpha \in \kappa \wedge A_{n,i,j,\alpha} \neq \emptyset \rangle$.

935 **Fact 4.9.** *Let κ be an ordinal, $U \subseteq {}^\omega\kappa$, Z is a parameter set such that $U \in Z$, and $n \in \omega$. Let M be an inner*
 936 *model of HOD_Z with $\hat{\mathbb{V}}^{U;Z} \in M$. Let $y \in {}^nU$. Then $\tilde{G}_y^{U;Z;n}$ is $\mathbb{V}_n^{U;Z}$ -generic over M and $\dot{\tau}^n[\tilde{G}_y^{U;Z;n}] = y$.*

937 *Proof.* Let $A \in M$ be a maximal antichain in M . Since $M \subseteq \text{HOD}_Z$, $A \in \text{OD}_Z$. Thus $\bigcup\{\mathfrak{S}_Z(p) : p \in A\}$ is an
 938 OD_Z subset of nU . If $\tilde{G}_y^{U;Z;n} \cap A = \emptyset$, then $y \notin \bigcup\{\mathfrak{S}_Z(p) : p \in A\}$. So ${}^nU \setminus \bigcup\{\mathfrak{S}_Z(p) : p \in A\}$ is a nonempty
 939 OD_Z subset of nU since $y \in {}^nU$ and $U \in Z$. Thus there is a $p \in \mathbb{V}_n^{U;Z}$ so that $\mathfrak{S}_Z(p) = {}^nU \setminus \bigcup\{\mathfrak{S}_Z(p) : p \in A\}$.
 940 Then $A \cup \{p\}$ is a $\mathbb{V}_n^{U;Z}$ -antichain which is larger than A . Contradiction. Thus there is some $p \in A$ so that
 941 $y \in \mathfrak{S}_Z(p)$. Thus $\tilde{G}_y^{U;Z;n} \cap A \neq \emptyset$. This shows that $\tilde{G}_y^{U;Z;n}$ is $\mathbb{V}_n^{U;Z}$ -generic over M .

942 Since $\hat{\mathbb{V}}^{U;Z} \in M$, $\dot{\tau}^n \in M$. For all $i < n$, $j \in \omega$, and $\alpha \in \kappa$, $y(i)(j) = \alpha$ if and only if $y \in \mathfrak{S}_Z(p_{n,i,j,\alpha}^{U;Z})$ if
 943 and only if $p_{n,i,j,\alpha}^{U;Z} \in \tilde{G}_y^{U;Z;n}$. Thus $\dot{\tau}^n[\tilde{G}_y^{U;Z;n}] = y$. $\square$

944 As mentioned before Fact 4.7, $\mathbb{V}_n^{U;Z}$ is not as absolute as $\mathbb{B}_n^{\kappa;Z}$. Given $y \in {}^nU$ and $p \in \mathbb{V}_n^{U;Z}$, it is not
 945 clear how $M[y]$ can determine if $p \in \tilde{G}_y^{U;Z;n}$ or equivalently whether $y \in \mathfrak{S}_Z(p)$ since this requires knowing
 946 the truth about y , Z , and ordinals in the real world. So $M[y] \subseteq M[\tilde{G}_y^{U;Z;n}]$ but it is not clear if they are
 947 equal in general.

948 **Definition 4.10.** Let X be a set. Let $\mathcal{P}_{\omega_1}(X) = \{\sigma \in \mathcal{P}(X) : |\sigma| \leq \omega\}$, i.e. the set of all subsets σ of X
 949 such that ω surjects onto σ . Let ν be an ultrafilter on $\mathcal{P}_{\omega_1}(X)$.

950 $\bullet$ ν is a fine ultrafilter on $\mathcal{P}_{\omega_1}(X)$ if and only if for each $x \in X$, $A_x = \{\sigma \in \mathcal{P}_{\omega_1}(X) : x \in \sigma\} \in \nu$.

- ν is a normal ultrafilter on $\mathcal{P}_{\omega_1}(X)$ if and only if for every function $\Phi : \mathcal{P}_{\omega_1}(X) \rightarrow \mathcal{P}_{\omega_1}(X)$ such that $\{\sigma \in \mathcal{P}_{\omega_1}(X) : \emptyset \neq \Phi(\sigma) \subseteq \sigma\} \in \nu$, there is an $x \in X$ so that $\{\sigma \in \mathcal{P}_{\omega_1}(X) : x \in \Phi(\sigma)\} \in \nu$.
- ν is a supercompact measure on $\mathcal{P}_{\omega_1}(X)$ if and only if ν is an ω_1 -complete, fine, and normal ultrafilter on $\mathcal{P}_{\omega_1}(X)$. If there is a supercompact measure on X , then one says that ω_1 is X -supercompact.

If all sets of reals have the Baire property, then there are no nonprincipal ultrafilters on ω which implies ultrafilters on any set must be ω_1 -complete. Thus AD implies all ultrafilters on any set are ω_1 -complete.

Let $\leq_{\text{Turing}}$ denote Turing reduction on $\mathbb{R}$ defined by $x \leq_{\text{Turing}} y$ if and only if x is Turing computable from y . Let $\equiv_{\text{Turing}}$ denote Turing equivalence on $\mathbb{R}$ defined by $x \equiv_{\text{Turing}} y$ if and only if $x \leq_{\text{Turing}} y$ and $y \leq_{\text{Turing}} x$. Let $\mathcal{D}_{\text{Turing}}$ be $\mathbb{R}/\equiv_{\text{Turing}}$ which is the collection of Turing degrees (equivalence classes of $\equiv_{\text{Turing}}$). One can extend $\leq_{\text{Turing}}$ to $\mathcal{D}_{\text{Turing}}$ by $X \leq_{\text{Turing}} Y$ if and only if there exist $x \in X$ and $y \in Y$ so that $x \leq_{\text{Turing}} y$. For $X \in \mathcal{D}_{\text{Turing}}$, let $\mathcal{C}_{\text{Turing}}^X = \{Y \in \mathcal{D}_{\text{Turing}} : X \leq_{\text{Turing}} Y\}$ which is called the Turing cone above X . Define the Martin filter on $\mathcal{D}_{\text{Turing}}$ by $A \in \mu_{\text{Turing}}$ if and only if there is an $X \in \mathcal{D}_{\text{Turing}}$ such that $\mathcal{C}_{\text{Turing}}^X \subseteq A$. Turing determinacy is the assertion that μ_{Turing} is an ultrafilter. Martin showed that under AD, μ_{Turing} is an ω_1 -complete ultrafilter and thus μ_{Turing} is often called the Martin measure in this setting. Peng and Yu ([47]) showed even Turing determinacy alone implies $\text{AC}_{\omega}^{\mathbb{R}}$ and hence μ_{Turing} is ω_1 -complete.

Suppose K is a set which is a surjective image of $\mathbb{R}$. Let $\pi : \mathbb{R} \rightarrow K$ be a surjection. Let $\tilde{\pi} : \mathcal{D}_{\text{Turing}} \rightarrow \mathcal{P}_{\omega_1}(K)$ be defined by $\tilde{\pi}(X) = \{\pi(z) : [z] \equiv_{\text{Turing}} \leq_{\text{Turing}} X\}$. Let $\tilde{\pi}_* \mu_{\text{Turing}}$ be the Rudin-Keisler pushforward of $\tilde{\pi}$ which is an ultrafilter on $\mathcal{P}_{\omega_1}(K)$ defined by $A \in \tilde{\pi}_* \mu_{\text{Turing}}$ if and only if $\tilde{\pi}^{-1}[A] \in \mu_{\text{Turing}}$. Let $x \in K$ and pick an $r \in \mathbb{R}$ such that $\pi(r) = x$. Let $A_x = \{\sigma \in \mathcal{P}_{\omega_1}(K) : x \in \sigma\}$. Let $R = [r] \equiv_{\text{Turing}}$. For all $Y \in \mathcal{C}_R^{\text{Turing}}$, $x \in \tilde{\pi}(Y)$. This shows that $\mathcal{C}_R^{\text{Turing}} \subseteq \tilde{\pi}^{-1}[A_x]$ and hence $A_x \in \tilde{\pi}_* \mu_{\text{Turing}}$. $\tilde{\pi}_* \mu_{\text{Turing}}$ is a fine measure on $\mathcal{P}_{\omega_1}(K)$. It has been shown that under AD, for every set K which is a surjective image of $\mathbb{R}$, there is an ω_1 -complete fine measure on $\mathcal{P}_{\omega_1}(K)$.

If $\prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ is wellfounded, then $\prod_{\mathcal{P}_{\omega_1}(K)} V/\tilde{\pi}_* \mu_{\text{Turing}}$ is wellfounded. To see this: Define $\Psi : \prod_{\mathcal{P}_{\omega_1}(K)} V/\tilde{\pi}_* \mu_{\text{Turing}} \rightarrow \prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ by $\Psi([f]_{\tilde{\pi}_* \mu_{\text{Turing}}}) = [f \circ \tilde{\pi}]_{\mu_{\text{Turing}}}$. Ψ is a $\dot{\epsilon}$ -embedding with each ultrapower structure given the ultrapower interpretation of $\dot{\epsilon}$. Thus it has been shown under AD that assuming $\prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ is wellfounded, then there is an ω_1 -complete fine measure ν on $\mathcal{P}_{\omega_1}(K)$ such that $\prod_{\mathcal{P}_{\omega_1}(K)} V/\nu$ is wellfounded. If DC holds, then $\prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ is wellfounded. Solovay [55] 12.2 showed that the consistency of $\text{AD}_{\mathbb{R}}$ implies the consistency of $\text{AD} + \neg\text{DC}$. Moreover Solovay [55] Lemma 2.3 states that if $\text{cof}(\Theta) = \omega$, then DC fails. Solovay ([42] Theorem 9.2) showed that $\prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ is wellfounded implies $\text{DC}_{\mathbb{R}}$. Woodin ([56] Corollary 6) showed that AD^+ implies $\prod_{\mathcal{D}_{\text{Turing}}} V/\mu_{\text{Turing}}$ is wellfounded. This latter result of Woodin will not be needed here since one will work in the real world satisfying AD and $\text{DC}_{\mathbb{R}}$ and drop into an inner model of determinacy satisfying DC.

To be explicit, if K is a set, then $L(K)$ is Gödel relative hierarchy built over the transitive closure of $\{K\}$. Unlike $L[K]$ which always satisfies AC, $L(K)$ may not satisfy AC. If $\{K_0, \dots, K_{n-1}\}$ is finite, then one will write $L(K_1, \dots, K_n)$ for $L(\{K_1, \dots, K_n\})$.

If $\kappa \in \text{ON}$ and $m \leq n < \omega$, then let $\pi_{n,m}^{\kappa} : {}^n(\omega^{\kappa}) \rightarrow {}^m(\omega^{\kappa})$ be the projection onto the first m coordinates defined by $\pi_{n,m}^{\kappa}(\ell) = \ell \upharpoonright m$ for all $\ell \in {}^n(\omega^{\kappa})$. The following states that projections of ∞ -Borel sets are ∞ -Borel in a uniform manner. In various forms, this result is due to Woodin and Ikegami-Trang ([25] Claim 2 in the proof Theorem 5). A detailed proof in the form needed here will be provided and uniformity will be very important.

Fact 4.11. (Woodin; Ikegami-Trang) Assume boldface GCH at ω . Let κ be an ordinal and assume there is a fine ω_1 -complete measure ν on $\mathcal{P}_{\omega_1}(\omega^{\kappa})$ such that $\prod_{\mathcal{P}_{\omega_1}(\omega^{\kappa})} V/\nu$ is wellfounded. Then there is an $\text{OD}_{\{\nu\}}$ function Υ so that for all $m \leq n < \omega$ and ∞ -Borel code b , $\pi_{n,m}^{\kappa}[\mathfrak{B}_b^{n(\omega^{\kappa})}] = \mathfrak{B}_{\Upsilon(b,n,m)}^{m(\omega^{\kappa})}$.

Proof. Without loss of generality suppose $m = 1$ and $n = 2$. Let b be an ∞ -Borel code of the form $b = (S, \varphi)$. For any $\sigma \in \mathcal{P}_{\omega_1}(\omega^{\kappa})$, let $\mathbb{Q}_{\sigma} = (\mathbb{V}_2^{\sigma; \{b, \sigma\}})^{L(b, \sigma)}$ denote the 2-dimensional $\text{OD}_{\{b, \sigma\}}$ Vopěnka forcing on σ defined in the inner model $L(b, \sigma)$. All sets in $L(b, \sigma)$ are ordinal definable from b , σ , and finitely many members from the countable set σ . After fixing a bijection of ω onto σ from the real world, one can construct externally a global wellordering of $L(b, \sigma)$ in the real world even though internally $L(b, \sigma)$ may fail to satisfy AC. Since σ is countable in the real world, let $B_0 : \sigma \rightarrow \omega$ be a bijection in the real world. Let $B_1 : \mathcal{P}(\sigma)^{L(b, \sigma)} \rightarrow \mathcal{P}(\omega)$ be an injection defined by $B_1(A) = B_0[A]$. Since $L(\sigma, b)$ has a global wellordering

1001 in the real world, $\mathcal{P}(\sigma)^{L(b,\sigma)}$ is wellorderable in the real world. Thus $B_1[\mathcal{P}(\sigma)^{L(b,\sigma)}]$ is a wellorderable subset
1002 of $\mathcal{P}(\omega)$ in the real world. Since boldface GCH at ω holds in the real world, $B_1[\mathcal{P}(\sigma)^{L(b,\sigma)}]$ is countable and
1003 hence $\mathcal{P}(\sigma)^{L(b,\sigma)}$ is countable in the real world. Let $\mathfrak{S}_{\{b,\sigma\}}^{L(b,\sigma)} : \text{ON} \rightarrow \text{OD}_{\{b,\sigma\}}^{L(b,\sigma)}$ be the surjection used to define
1004 $(\mathbb{V}_2^{\sigma;\{b,\sigma\}})^{L(b,\sigma)}$ (as in Definition 4.8). Then since the restriction $\mathfrak{S}_{\{b,\sigma\}}^{L(b,\sigma)} : (\mathbb{V}_2^{\sigma;\{b,\sigma\}})^{L(b,\sigma)} \rightarrow \mathcal{P}(\sigma)^{L(b,\sigma)}$ is
1005 an injection (note that $\mathbb{V}_2^{\sigma;\{b,\sigma\}} \subseteq \text{ON}$ as in Definition 4.8), $\mathbb{Q}_\sigma$ is countable. Let $B_2 : \mathbb{Q}_\sigma \rightarrow \omega$ be a
1006 bijection. Let $B_3 : \mathcal{P}(\mathbb{Q}_\sigma)^{L(b,\sigma)} \rightarrow \mathcal{P}(\omega)$ be defined by $B_3(A) = B_2[A]$ which is an injection. Again by
1007 boldface GCH at ω , $\mathcal{P}(\mathbb{Q}_\sigma)^{L(b,\sigma)}$ is countable in the real world. Let $\mathbb{C}_\sigma = \text{Coll}(\omega, \mathbb{Q}_\sigma)$ which is the forcing of
1008 partial functions from ω into $\mathbb{Q}_\sigma$ ordered by reverse extension. Since $\mathbb{Q}_\sigma$ is countable, $\mathbb{C}_\sigma$ is countable. Let
1009 $\hat{\mathbb{V}}_\sigma^b = (\hat{\mathbb{V}}_\sigma^{\sigma;\{b,\sigma\}})^{L(b,\sigma)}$ which is the code set from Definition 4.8. Note that $\mathbb{Q}_\sigma, \mathbb{C}_\sigma \in L[b, \hat{\mathbb{V}}_\sigma^b]$.
1010 By the fineness of ν , $\{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa) : x \in \sigma\} \in \nu$ for all $x \in \omega^\kappa$. Let $\prec_\sigma$ denote the constructibility
1011 wellordering of $L[b, \hat{\mathbb{V}}_\sigma^b]$. For any $x \in \omega^\kappa$, let $\prec_{\sigma,x}$ be the constructibility wellordering¹¹ on $L[b, \hat{\mathbb{V}}_\sigma^b, x]$. Using
1012 the maps $\sigma \mapsto \prec_\sigma$ and $\sigma \mapsto \prec_{\sigma,x}$ in the usual proof of Loś's theorem, one has that $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b]/\nu$
1013 and $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b, x]/\nu$ for all $x \in \omega^\kappa$ satisfy Loś' theorem. By the hypothesis that $\prod_{\mathcal{P}_{\omega_1}(\omega^\kappa)} V/\nu$
1014 ν is wellfounded, one has that $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b]/\nu$ and $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b, x]/\nu$ for all $x \in \omega^\kappa$ are
1015 wellfounded. For any $v \in V$, let $c_v : \mathcal{P}_{\omega_1}(\omega^\kappa) \rightarrow \{v\}$ be the constant function. Let $j_\nu : V \rightarrow \prod_{\mathcal{P}_{\omega_1}(\omega^\kappa)} V/\nu$
1016 be defined by $j_\nu(v) = [c_v]_\nu$. (Note that $\prod_{\mathcal{P}_{\omega_1}(\omega^\kappa)} V/\nu$ may not satisfy ZF and j_ν may not be elementary
1017 since Loś' theorem may fail for this ultrapower.) Define the following functions on $\mathcal{P}_{\omega_1}(\omega^\kappa)$: $\Phi_\mathbb{Q}(\sigma) = \mathbb{Q}_\sigma$,
1018 $\Phi_\mathbb{C}(\sigma) = \mathbb{C}_\sigma$, $\Phi_{\hat{\mathbb{V}}^b}(\sigma) = \hat{\mathbb{V}}_\sigma^b$. Let $\mathbb{Q}_\infty = [\Phi_\mathbb{Q}]_\nu$, $\mathbb{C}_\infty = [\Phi_\mathbb{C}]_\nu$, and $\hat{\mathbb{V}}_\infty^b = [\Phi_{\hat{\mathbb{V}}^b}]_\nu$. Let $b_\infty = j_\nu(b)$ and
1019 $S_\infty = j_\nu(S)$ (where recall $b = (S, \varphi)$). By Ló's theorem for $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b]/\nu$, b_∞ is an ∞ -Borel code of
1020 the form (S_∞, φ) . Also by Loś' theorem, $\hat{\mathbb{V}}_\infty^b$ is a set of ordinals and for all $x \in \omega^\kappa$, $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b, x]/\nu$
1021 $\nu = L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu(x)]$. $L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu(x)] \models$ “ $j_\nu(x)$ is a function from $j_\nu(\omega)$ into $j_\nu(\kappa)$ ”. Since ν is ω_1 -
1022 complete, $j_\nu(\omega) = \omega$ and thus $j_\nu(x)$ is a function from ω into $j_\nu(\kappa)$. Thus by Loś' theorem, for all $n \in \omega$,
1023 $j_\nu(x)(n) = j_\nu(x)(j_\nu(n)) = j_\nu(x(n))$. Thus $j_\nu(x) = j_\nu \circ x$ and hence $L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu(x)] = L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu \circ x]$.
1024 Let $j_* = j_\nu \upharpoonright \kappa = \{(\alpha, \beta) \in \kappa \times j_\nu[\kappa] : j(\alpha) = \beta\}$. Note that j_* is essentially an $\text{OD}_{\{\nu\}}$ set of ordinals and
1025 $j_\nu(x) = j_\nu \circ x = j_* \circ x$. The maps $b \mapsto b_\infty$ and $b \mapsto \hat{\mathbb{V}}_\infty^b$ are $\text{OD}_{\{\nu\}}$.

Claim: For all $x \in \omega^\kappa$,

$$x \in \pi_{2,1}^\kappa[\mathfrak{B}_b^{2(\omega^\kappa)}] \Leftrightarrow L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_* \circ x] \models 1_{\mathbb{C}_\infty} \Vdash_{\mathbb{C}_\infty} (\exists y) \left(L[\check{S}_\infty, (j_* \circ x), y] \models \varphi(\check{S}_\infty, (j_* \circ x), y) \right).$$

To see this claim: ($\Leftarrow$) Since $j_* \circ x = j_\nu(x)$, one has

$$L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu(x)] \models 1_{\mathbb{C}_\infty} \Vdash_{\mathbb{C}_\infty} (\exists y) \left(L[\check{S}_\infty, j_\nu(x), y] \models \varphi(\check{S}_\infty, j_\nu(x), y) \right).$$

Since $L[b_\infty, \hat{\mathbb{V}}_\infty^b, j_\nu(x)] = \prod_{\sigma \in \mathcal{P}_{\omega_1}(\omega^\kappa)} L[b, \hat{\mathbb{V}}_\sigma^b, x]/\nu$, one has that the following holds for ν -almost all σ by Loś' theorem.

$$L[b, \hat{\mathbb{V}}_\sigma^b, x] \models 1_{\mathbb{C}_\sigma} \Vdash_{\mathbb{C}_\sigma} (\exists y) \left(L[\check{S}, \check{x}, y] \models \varphi(\check{S}, \check{x}, y) \right).$$

Fix such a σ . Since $\mathbb{C}_\sigma$ is countable in the real world and boldface GCH at ω holds, Fact 2.14 implies there is a $g \subseteq \mathbb{C}_\sigma$ (which belongs to the real world) and is $\mathbb{C}_\sigma$ -generic over $L[b, \hat{\mathbb{V}}_\sigma^b, x]$. By the forcing theorem,

$$L[b, \hat{\mathbb{V}}_\sigma^b, x][g] \models (\exists y) \left(L[S, x, y] \models \varphi(S, x, y) \right).$$

There is a $y \in \omega^\kappa \cap L[b, \hat{\mathbb{V}}_\sigma^b, x][g]$ such that

$$L[b, \hat{\mathbb{V}}_\sigma^b, x][g] \models (L[S, x, y] \models \varphi(S, x, y))$$

1026 and note that y belongs to the real world since g belonged to the real world. By absoluteness, $L[S, x, y] \models$
1027 $\varphi(S, x, y)$. Since $b = (S, \varphi)$, this implies that $(x, y) \in \mathfrak{B}_b^{2(\omega^\kappa)}$. Hence $x \in \pi_{2,1}^\kappa[\mathfrak{B}_b^{2(\omega^\kappa)}]$.

($\Rightarrow$) Now suppose $x \in \pi_{2,1}^\kappa[\mathfrak{B}_b^{2(\omega^\kappa)}]$. There is a $z \in \omega^\kappa$ so that $(x, z) \in \mathfrak{B}_b^{2(\omega^\kappa)}$. So $L[S, x, z] \models \varphi(S, x, z)$.
By fineness of ν , there is an $A \in \nu$ so that $x, z \in \sigma$ for all $\sigma \in A$. Fix $\sigma \in A$. Let $H_{\sigma,x,z}$ be the $\mathbb{Q}_\sigma$ -generic

¹¹The constructibility ordering is not independent of the model $L[b, \hat{\mathbb{V}}_\sigma^b, x]$. To be explicit, let $\prec_{\sigma,x}$ be the constructibility ordering using predicate $b, \hat{\mathbb{V}}_\sigma^b$, and x .

associated to (x, z) which is $\mathbb{Q}_\sigma$ -generic over $L[b, \hat{V}_\sigma^b]$ by Fact 4.9 (note that $(x, z) \in {}^2\sigma$ and $L[b, \hat{V}_\sigma^b]$ is an inner model of $\text{HOD}_{\{b, \sigma\}}^{L(\sigma, b)}$). By absoluteness,

$$L[b, \hat{V}_\sigma^b][H_{\sigma, x, z}] \models L[S, x, z] \models \varphi(S, x, z).$$

In $L[b, \mathbb{V}_\sigma^b]$, $\mathbb{Q}_\sigma$ regularly embeds into $\mathbb{C}_\sigma$ by [31] Corollary 26.8. There is some $g \subseteq \mathbb{C}_\sigma$ so that $H_{\sigma, x, z} \in L[b, \hat{V}_\sigma^b][g]$. Again by absoluteness,

$$L[b, \hat{V}_\sigma^b][g] \models L[S, x, z] \models \varphi(S, x, z).$$

Since $x \in L[b, \hat{V}_\sigma^b][g]$, there is an $h \subseteq \mathbb{C}_\sigma$ such that h is $\mathbb{C}_\sigma$ -generic over $L[b, \hat{V}_\sigma^b][x]$ and $L[b, \hat{V}_\sigma^b][g] = L[b, \hat{V}_\sigma^b][x][h]$ by [31] Corollary 26.10. Thus one has

$$L[b, \hat{V}_\sigma^b, x][h] \models L[S, x, z] \models \varphi(S, x, z).$$

Then one has

$$L[b, \hat{V}_\sigma^b, x][h] \models (\exists y) \left(L[S, x, y] \models \varphi(S, x, y) \right).$$

By the forcing theorem, there is a $p \in \mathbb{C}_\sigma$ so that

$$L[b, \hat{V}_\sigma^b, x] \models p \Vdash_{\mathbb{C}_\sigma} (\exists y) \left(L[\check{S}, \check{x}, y] \models \varphi(\check{S}, \check{x}, y) \right).$$

By the homogeneity of $\mathbb{C}_\sigma$, one has

$$L[b, \hat{V}_\sigma^b, x] \models 1_{\mathbb{C}_\sigma} \Vdash_{\mathbb{C}_\sigma} (\exists y) \left(L[\check{S}, \check{x}, y] \models \varphi(\check{S}, \check{x}, y) \right).$$

Since σ was an arbitrary element of $A \in \nu$, one has the following by Loś' theorem.

$$L[b_\infty, \hat{V}_\infty^b, j_\nu(x)] \models 1_{\mathbb{C}_\infty} \Vdash_{\mathbb{C}_\infty} (\exists y) \left(L[\check{S}_\infty, j_\nu(\check{x}), y] \models \varphi(\check{S}_\infty, j_\nu(\check{x}), y) \right).$$

Since $j_\nu(x) = j_* \circ x$,

$$L[b_\infty, \hat{V}_\infty^b, j_* \circ x] \models 1_{\mathbb{C}_\infty} \Vdash_{\mathbb{C}_\infty} (\exists y) \left(L[\check{S}_\infty, (j_* \circ x), y] \models \varphi(\check{S}_\infty, (j_* \circ x), y) \right).$$

1028 This completes the proof of the claim.

Note that $j_* \circ x$ can be obtained from j_* and x . By the claim and absoluteness, $x \in \pi_{2,1}^\kappa[\mathfrak{B}_b^{2(\omega\kappa)}]$ is equivalent to

$$L[b_\infty, \hat{V}_\infty^b, j_*, x] \models L[b_\infty, \hat{V}_\infty^b, j_* \circ x] \models 1_{\mathbb{C}_\infty} \Vdash_{\mathbb{C}_\infty} (\exists y) \left(L[\check{S}_\infty, (j_* \circ x), y] \models \varphi(\check{S}_\infty, (j_* \circ x), y) \right).$$

1029 Let $\langle b_\infty, \hat{V}_\infty^b, j_* \rangle$ be some uniform method of coding three sets of ordinals into one set of ordinals. Let ψ
1030 be a formula such that $\psi(\langle b_\infty, \hat{V}_\infty^b, j_* \rangle, x)$ asserts the inner expression above (behind the first satisfaction
1031 symbol). $(\langle b_\infty, \hat{V}_\infty^b, j_* \rangle, \psi)$ is an ∞ -Borel code for $\pi_{2,1}^\kappa[\mathfrak{B}_b^{2(\omega\kappa)}]$. Define $\Upsilon(b) = (\langle b_\infty, \hat{V}_\infty^b, j_* \rangle, \psi)$. Υ is $\text{OD}_{\{\nu\}}$
1032 since it was observed above that $j_* \in \text{HOD}_{\{\nu\}}$ and the mappings $b \mapsto b_\infty$ and $b \mapsto \hat{V}_\infty^b$ are $\text{OD}_{\{\nu\}}$. $\square$

1033 *Remark 4.12.* The following are some remark about the structure of the proof and important subtle points
1034 of Fact 4.11: In the direction $(\Leftarrow)$, it was important that $\mathbb{C}_\sigma$ was countable in the real world in order to find
1035 $\mathbb{C}_\sigma$ -generic filters over $L[b, \hat{V}_\sigma^b]$. Note that in the real world, the cardinality of $\mathbb{C}_\sigma$ is the same cardinality
1036 as $\mathbb{Q}_\sigma$. In $(\Rightarrow)$, the relevant property of $\mathbb{Q}_\sigma$ is that it needs to make the pair (x, z) -generic over the relevant
1037 inner model of ZFC. There are two forcings mentioned above with this property, the ∞ -Borel code forcing
1038 and the Vopěnka forcing. The Vopěnka forcing and ∞ -Borel code forcing defined in the real world are never
1039 countable. In a first attempt to make this forcing countable, one goes into $L(\sigma, b)$ which has an external
1040 global wellordering. If one was proving the Fact 4.11 for $\mathbb{R}$ or ${}^\omega\omega$, then the corresponding Vopěnka and
1041 ∞ -Borel code forcing defined in $L(b, \sigma)$ are both already countable. (So one could have more simply used
1042 the ∞ -Borel code forcing and even use the more concrete Martin's measure on the Turing degrees for $\mathbb{R}$ or
1043 ${}^\omega\omega$.) When $\kappa > \omega$, the Vopěnka and ∞ -Borel code forcing on ${}^\omega\kappa$ defined in $L(b, \sigma)$ are still uncountable in
1044 the real world. To resolve this, one restricts the forcing to subsets of the countable set σ rather than all
1045 of ${}^\omega\kappa$. However, it is not clear if σ is ∞ -Borel in $L(\sigma, \mathbb{R})$ with an $\text{OD}_{\{b, \sigma\}}^{L(b, \sigma)}$ ∞ -Borel code (or ∞ -Borel at
1046 all). Thus it is unclear how one can create an ∞ -Borel code forcing which only considers subsets of σ . This
1047 problem was resolved by instead using the Vopěnka forcing $\mathbb{Q}_\sigma = (\mathbb{V}_2^{\sigma; \{b, \sigma\}})^{L(b, \sigma)}$.

Definition 4.13. A set Z is a good parameter set if and only if Z is a parameter set and for all $\tilde{z}$ in the transitive closure of $\{Z\}$, there is an $r \in \mathbb{R}$ so that $\tilde{z}$ is $\text{OD}_{Z \cup \{r\}}$.

Recall for any set K , $L(K)$ is the relativized constructibility starting with the transitive closure $\{K\}$. So in $L(Z, \mathbb{R})$, every set is ordinal definable from finitely many parameters from $\mathbb{R}$ and elements of the transitive closure of Z . If Z is a good parameter set, then this can be improved: every set in $L(Z, \mathbb{R})$ is ordinal definable from finitely many reals and finitely many elements of Z . The purpose of the concept of a good parameter set is for Fact 4.14 below.

Fact 4.14. Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a set such that $V = L(Z, \mathbb{R})$ and Z is a good parameter set. Let $\kappa < \Theta$. Then there is an OD_Z function Υ so that for all $m \leq n < \omega$ and ∞ -Borel code p , $\pi_{n,m}^{\kappa}[\mathfrak{B}_p^{n(\omega\kappa)}] = \mathfrak{B}_{\Upsilon(p,n,m)}^{m(\omega\kappa)}$.

Proof. Fix $\kappa < \Theta$. Since $V = L(Z, \mathbb{R})$ and Z is a good parameter set, every set is $\text{OD}_{Z \cup \{r\}}$ for some $r \in \mathbb{R}$ (by the discussion above). For each $r \in \mathbb{R}$, let π_r denote the least $\text{OD}_{Z \cup \{r\}}$ surjective function $\pi : \mathbb{R} \rightarrow \kappa$ according to the canonical wellordering of $\text{OD}_{Z \cup \{r\}}$ if such a surjection exists and otherwise let $\pi_r : \mathbb{R} \rightarrow \kappa$ be the constant 0 function. Let $\varpi(r) = \pi_{r^{[0]}}(r^{[1]})$ (where recall $r^{[n]}$ comes from the coding of ω -many reals by a single real defined before Fact 2.24). ϖ is an OD_Z surjection of $\mathbb{R}$ onto κ . Let $\pi : \mathbb{R} \rightarrow \omega\kappa$ be defined by $\pi(r)(n) = \varpi(r^{[n]})$. π is an OD_Z surjection of $\mathbb{R}$ onto $\omega\kappa$. Let $\tilde{\pi} : \mathcal{D}_{\text{Turing}} \rightarrow \mathcal{P}_{\omega_1}(\omega\kappa)$ be defined by $\tilde{\pi}(X) = \{z : [z] \equiv_{\text{Turing}} X\}$. $\tilde{\pi}$ is OD_Z . As shown above, $\tilde{\pi}_* \mu_{\text{Turing}}$ is an OD_Z fine ω_1 -complete measure on $\mathcal{P}_{\omega_1}(\omega\kappa)$. Since $L(Z, \mathbb{R}) \models \text{DC}_{\mathbb{R}}$, one has that $L(Z, \mathbb{R}) \models \text{DC}$ and hence $\prod_{\mathcal{P}_{\omega_1}(\omega\kappa)} V / \tilde{\pi}_* \mu_{\text{Turing}}$ is wellfounded. Let Υ be the $\text{OD}_{\{\tilde{\pi}_* \mu_{\text{Turing}}\}}$ function given by Fact 4.14 applied to $\tilde{\pi}_* \mu_{\text{Turing}}$. Since $\pi_* \mu_{\text{Turing}} \in \text{OD}_Z$, $\Upsilon \in \text{OD}_Z$. $\square$

Definition 4.15. Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a good parameter set such that $V = L(Z, \mathbb{R})$. Let $\kappa < \Theta$. Let Υ be the OD_Z function obtained from Fact 4.14.

Define $\Pi_{n,m}^{\kappa;Z} : \mathbb{B}_n^{\kappa;Z} \rightarrow \mathbb{B}_m^{\kappa;Z}$ by $\Pi_{n,m}^{\kappa;Z}(p)$ is the least member of the $[\Upsilon(p,n,m)]_{\sim_{\kappa;Z}}$ according to the canonical wellordering of HOD_Z . By Fact 4.14, $\pi_{n,m}^{\kappa}[\mathfrak{B}_p^{n(\omega\kappa)}] = \mathfrak{B}_{\Pi_{n,m}^{\kappa;Z}(p)}^{m(\omega\kappa)}$. If $q_0 \leq_{\mathbb{B}_n^{\kappa;Z}} q_1$, then $\mathfrak{B}_{q_0}^{n(\omega\kappa)} \subseteq \mathfrak{B}_{q_1}^{n(\omega\kappa)}$. Thus $\pi_{n,m}^{\kappa}[\mathfrak{B}_{q_0}^{n(\omega\kappa)}] \subseteq \pi_{n,m}^{\kappa}[\mathfrak{B}_{q_1}^{n(\omega\kappa)}]$ and hence $\Pi_{n,m}^{\kappa;Z}(q_0) \leq_{\mathbb{B}_m^{\kappa;Z}} \Pi_{n,m}^{\kappa;Z}(q_1)$. Let $q \in \mathbb{B}_n^{\kappa;Z}$ and $p \leq_{\mathbb{B}_m^{\kappa;Z}} \Pi_{n,m}^{\kappa;Z}(q)$. Thus $\mathfrak{B}_p^{m(\omega\kappa)} \subseteq \mathfrak{B}_{\Pi_{n,m}^{\kappa;Z}(q)}^{m(\omega\kappa)}$. Suppose $p = (S_0, \varphi_0)$ and $q = (S_1, \varphi_1)$. Let $\psi(A_0, A_1, v)$ be the formula which asserts “ $L[A_0, v \upharpoonright m] \models \varphi_0(A_0, v \upharpoonright m)$ and $L[A_1, v] \models \varphi_1(A_1, v)$ ”. Let $\tilde{r} = (\langle S_0, S_1 \rangle, \psi)$ and note that $\tilde{r}$ is OD_Z . Then $f \in \mathfrak{B}_{\tilde{r}}^{m(\omega\kappa)}$ if and only if $L[S_0, S_1, f] \models \psi(S_0, S_1, f)$ if and only if $L[S_0, f \upharpoonright m] \models \varphi_0(S_0, f \upharpoonright m)$ and $L[S_1, f] \models \varphi_1(S_1, f)$ if and only if $f \upharpoonright m \in \mathfrak{B}_p^{m(\omega\kappa)}$ and $f \in \mathfrak{B}_q^{n(\omega\kappa)}$. Let $r \in \mathbb{B}_n^{\kappa;Z}$ be such that $r \sim_{\mathbb{B}_n^{\kappa;Z}} \tilde{r}$. Then $r \leq_{\mathbb{B}_n^{\kappa;Z}} q$ and $\Pi_{n,m}^{\kappa;Z}(r) = p$. It has been shown that $\Pi_{n,m}^{\kappa;Z} : \mathbb{B}_n^{\kappa;Z} \rightarrow \mathbb{B}_m^{\kappa;Z}$ is a forcing projection. $\langle \mathbb{B}_m^{\kappa;Z}, \Pi_{n,m}^{\kappa;Z} : m \leq n < \omega \rangle$ is a direct system of forcing projection. Let $\mathbb{B}_{\omega}^{\kappa;Z}$ be the collection of all functions $\mathbf{p} \in \prod_{n \in \omega} \mathbb{B}_n^{\kappa;Z}$ such that the following holds:

- $m \leq n < \omega$, $\Pi_{n,m}^{\kappa;Z}(\mathbf{p}(n)) = \mathbf{p}(m)$.
- There is an $N < \omega$ so that for all $N < k < \omega$, $\mathfrak{B}_{\mathbf{p}(k)}^{k(\omega\kappa)} = \mathfrak{B}_{\mathbf{p}(N)}^{n(\omega\kappa)} \times {}^{k-N}(\omega\kappa)$. The least such N will be denoted $\text{dim}(\mathbf{p})$ and is called the dimension of $\mathbf{p}$.

Let $\leq_{\mathbb{B}_{\omega}^{\kappa;Z}}$ be defined by $\mathbf{p} \leq_{\mathbb{B}_{\omega}^{\kappa;Z}} \mathbf{q}$ if and only if for all $n \in \omega$, $\mathbf{p}(n) \leq_{\mathbb{B}_n^{\kappa;Z}} \mathbf{q}(n)$. Let $1_{\mathbb{B}_{\omega}^{\kappa;Z}}$ be defined by $1_{\mathbb{B}_{\omega}^{\kappa;Z}}(n) = 1_{\mathbb{B}_n^{\kappa;Z}}$ for all $n \in \omega$. Note that the directed system $\langle \mathbb{B}_m^{\kappa;Z}, \Pi_{n,m}^{\kappa;Z} : m \leq n < \omega \rangle$ and $\mathbb{B}_{\omega}^{\kappa;Z} = (\mathbb{B}_{\omega}^{\kappa;Z}, \leq_{\mathbb{B}_{\omega}^{\kappa;Z}}, 1_{\mathbb{B}_{\omega}^{\kappa;Z}})$ belong to HOD_Z . This forcing is called the finite support direct limit of the OD_Z ∞ -Borel code forcing on $\omega\kappa$. For each $n \in \omega$, let $\Pi_{\omega,n}^{\kappa;Z} : \mathbb{B}_{\omega}^{\kappa;Z} \rightarrow \mathbb{B}_n^{\kappa;Z}$ be defined by $\Pi_{\omega,n}^{\kappa;Z}(\mathbf{p}) = \mathbf{p}(n)$. $\Pi_{\omega,n}^{\kappa;Z}$ is a forcing projection.

Let $\dot{G}_{\omega}$ be the canonical $\mathbb{B}_{\omega}^{\kappa;Z}$ -name for the generic filter. For each $n \in \omega$, let $\dot{G}_n$ be the $\mathbb{B}_{\omega}^{\kappa;Z}$ -name such that $1_{\mathbb{B}_{\omega}^{\kappa;Z}} \Vdash \dot{G}_n = \langle \Pi_{\omega,n}^{\kappa;Z}[\dot{G}_{\omega}] \rangle_{\Pi_{\omega,n}^{\kappa;Z}}$ (that is, $\dot{G}_n$ is the canonical $\mathbb{B}_{\omega}^{\kappa;Z}$ -name for the generic filter on $\mathbb{B}_n^{\kappa;Z}$ induced by the forcing projection $\Pi_{\omega,n}^{\kappa;Z}$). Let $\dot{\tau}_n^{\omega}$ be the $\mathbb{B}_{\omega}^{\kappa;Z}$ -name such that $1_{\mathbb{B}_{\omega}^{\kappa;Z}} \Vdash \dot{\tau}_n^{\omega} = \dot{\tau}^n[\dot{G}_n]$ where $\dot{\tau}^n$ here refers to the canonical $\mathbb{B}_n^{\kappa;Z}$ -name for the generic element of ${}^n(\omega\kappa)$ naturally understood as a $\mathbb{B}_{\omega}^{\kappa;Z}$ -name. Let $\dot{\tau}^{\omega}$ be the $\mathbb{B}_{\omega}^{\kappa;Z}$ -name such that for all $n \in \omega$, $1_{\mathbb{B}_{\omega}^{\kappa;Z}} \Vdash \dot{\tau}^{\omega} \upharpoonright \check{n} = \dot{\tau}_n^{\omega}$. $\dot{\tau}^{\omega}$ names the generic element of ${}^{\omega}(\omega\kappa)$ added by $\mathbb{B}_{\omega}^{\kappa;Z}$. Let $\omega_{\dot{\tau}}^{\text{sym}}$ be a $\mathbb{B}_{\omega}^{\kappa;Z}$ -name such that $1_{\mathbb{B}_{\omega}^{\kappa;Z}} \Vdash \omega_{\dot{\tau}}^{\text{sym}} = \{\dot{\tau}^{\omega}(n) : n \in \omega\}$.

(If $\kappa = \omega$, one usually writes $\dot{\mathbb{R}}^{\text{sym}}$ rather than $\omega\dot{\kappa}^{\text{sym}}$.) Also let $\dot{\mathbb{R}}^{\kappa;\text{sym}}$ be a $\mathbb{B}_{\omega}^{\kappa;Z}$ -name for the collection of reals inside $\omega\dot{\kappa}^{\text{sym}}$. (Formally $\dot{\mathbb{R}}^{\text{sym}}$ is only defined for $\mathbb{B}_{\omega}^{\omega;Z}$ and names the terms of the generic ω -sequence of $\mathbb{B}_{\omega}^{\omega;Z}$.) Since $\Pi_{\omega,n}^{\kappa;Z}$ is a forcing projection, whenever G is a $\mathbb{B}_{\omega}^{\kappa;Z}$ filter, let $G_n = \dot{G}_n[G] = \langle \Pi_{\omega,n}^{\kappa;Z}[G] \rangle_{\mathbb{B}_n^{\kappa;Z}}$ which is a filter on $\mathbb{B}_n^{\kappa;Z}$. One can form the corresponding remainder forcing, $\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega,n}^{\kappa;Z} G_n$. $\omega\dot{\kappa}^{\text{sym}}$ and $\dot{\mathbb{R}}^{\kappa;\text{sym}}$ can naturally be considered a $(\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega,n}^{\kappa;Z} G_n)$ -name.

Similarly to Definition 4.5, let $\dot{\mathbb{B}}^{\kappa;Z}$ be an OD_Z set of ordinals which codes $\langle \mathbb{B}_n^{\kappa;Z}, \Pi_{n,m}^{\kappa;Z}, \Pi_{\omega,m}^{\kappa;Z} : m \leq n < \omega \rangle$, $\mathbb{B}_{\omega}^{\kappa;Z}$, $\dot{\tau}^{\omega}$, $\dot{\tau}_n^{\omega}$, and $\omega\dot{\kappa}^{\text{sym}}$. Again there are various ways to define $\dot{\mathbb{B}}^{\kappa;Z}$ but it should be uniform in Z . Its main property is that any inner model $M \subseteq \text{HOD}_Z$ with $\dot{\mathbb{B}}^{\kappa;Z} \in M$ can see the direct system of forcing projections.

If $h : \omega \rightarrow (\omega\kappa)^V$ (which may not necessarily belong to the real world V), then let $G_h^{\kappa;Z;\omega} = \{p \in \omega : (\forall n \in \omega)(h \restriction n \in \mathfrak{B}_{p(n)}^{\kappa;Z;\omega})\}$. Note that $\Pi_{\omega,n}^{\kappa;Z}[G_h^{\kappa;Z;\omega}] = G_{h \restriction n}^{\kappa;Z;n}$. Thus $\dot{\tau}_n^{\omega}[G_h^{\kappa;Z;\omega}] = h \restriction n$. Hence $\dot{\tau}^{\omega}[G_h^{\kappa;Z;\omega}] = h$ and $\omega\dot{\kappa}^{\text{sym}}[G_h^{\kappa;Z;\omega}] = h[\omega]$.

Fact 4.16. (Woodin) Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a good parameter set such that $V = L(Z, \mathbb{R})$. Let $\kappa < \Theta$. Let $G \subseteq \text{Coll}(\omega, \omega\kappa)$ be $\text{Coll}(\omega, \omega\kappa)$ -generic over the true universe V . Let $g : \omega \rightarrow \omega\kappa$ be the generic surjection added by G ; namely $g = \bigcup G$. Let M be an inner model of HOD_Z with $\dot{\mathbb{B}}^{\kappa;Z} \in M$. Then $G_g^{\kappa;Z;\omega}$ is $\mathbb{B}_{\omega}^{\kappa;Z}$ -generic over M .

Proof. Let $D \subseteq \mathbb{B}_{\omega}^{\kappa;Z}$ be a dense subset of $\mathbb{B}_{\omega}^{\kappa;Z}$ in M . Although $\text{Coll}(\omega, \omega\kappa)$ is usually defined as the collection of finite partial functions from ω into $\omega\kappa$, here $\text{Coll}(\omega, \omega\kappa)$ will be identified with it dense subforcing of finite partial function s from ω into $\omega\kappa$ such that $\text{dom}(s) \in \omega$ (i.e. a finite sequence in $\omega\kappa$). Let $\tilde{D} \subseteq \text{Coll}(\omega, \omega\kappa)$ be the collection of $t \in \text{Coll}(\omega, \omega\kappa)$ such that there is a $p \in D$ with $\text{dim}(p) \leq |t|$ and $t \in \mathfrak{B}_{p(|t|)}^{|t|(\omega\kappa)}$. First, one will show that $\tilde{D}$ is dense in $\text{Coll}(\omega, \omega\kappa)$. Fix $s \in \text{Coll}(\omega, \omega\kappa)$ and let $n = |s|$. Let $E \subseteq \mathbb{B}_n^{\kappa;Z}$ consists of those $u \in \mathbb{B}_n^{\kappa;Z}$ such that there exists a $q \in D$ with $\Pi_{\omega,n}^{\kappa;Z}(q) = u$. As an intermediate step, one will show that E is dense in $\mathbb{B}_n^{\kappa;Z}$. Let $r \in \mathbb{B}_n^{\kappa;Z}$. Let τ be a function on ω such that $\tau(k) \in \mathbb{B}_k^{\kappa;Z}$ for all $k \in \omega$, $\tau(k) = \Pi_{n,k}^{\kappa;Z}(r)$ for all $k \leq n$, and $\mathfrak{B}_{\tau(k)}^{k(\omega\kappa)} = \mathfrak{B}_r^{n(\omega\kappa)} \times \mathbb{R}^{k-n(\omega\kappa)}$ for all $n < k < \omega$. Thus $\tau \in \mathbb{B}_{\omega}^{\kappa;Z}$ and $\Pi_{\omega,n}^{\kappa;Z}(\tau) = r$. Since D is dense in $\mathbb{B}_{\omega}^{\kappa;Z}$, there is a $q \in D$ with $q \leq_{\mathbb{B}_{\omega}^{\kappa;Z}} \tau$. Then $\Pi_{\omega,n}^{\kappa;Z}(q) \leq_{\mathbb{B}_n^{\kappa;Z}} r$ and $\Pi_{\omega,n}^{\kappa;Z}(q) \in E$ since $q \in D$. Since $r \in \mathbb{B}_n^{\kappa;Z}$ was arbitrary, E has been shown to be dense in $\mathbb{B}_n^{\kappa;Z}$. $G_s^{\kappa;Z;n}$ is $\mathbb{B}_n^{\kappa;Z}$ -generic over M by Fact 4.6. Thus there is some $u \in G_s^{\kappa;Z;n} \cap E$. By definition of $u \in E$, there is a $q \in D$ with $\Pi_{\omega,n}^{\kappa;Z}(q) = u$. Since $u \in G_s^{\kappa;Z;n}$, $s \in \mathfrak{B}_u^{n(\omega\kappa)}$. Let $m = \max\{n, \text{dim}(q)\}$. By definition of $q \in \mathbb{B}_{\omega}^{\kappa;Z}$, $\Pi_{m,n}^{\kappa;Z}(q(m)) = q(n) = \Pi_{\omega,n}^{\kappa;Z}(q) = u$. Thus there is some $t \in \mathfrak{B}_{q(m)}^{m(\omega\kappa)}$ with $\pi_{m,n}^{\kappa;Z}(t) = t \restriction n = s$. Hence $t \leq_{\text{Coll}(\omega, \omega\kappa)} s$ and $t \in \tilde{D}$ since $q \in D$, $\text{dim}(q) \leq m = |t|$, and $t \in \mathfrak{B}_{q(|t|)}^{|t|(\omega\kappa)}$. Since $s \in \text{Coll}(\omega, \omega\kappa)$ was arbitrary, this shows that $\tilde{D}$ is dense in $\text{Coll}(\omega, \omega\kappa)$. Since $G \subseteq \text{Coll}(\omega, \omega\kappa)$ is $\text{Coll}(\omega, \omega\kappa)$ -generic over V , there is some $s \in G \cap \tilde{D}$. By definition of $s \in \tilde{D}$, there is a $p \in D$ with $\text{dim}(p) \leq |s|$ and $s \in \mathfrak{B}_{p(|s|)}^{|s|(\omega\kappa)}$. Since the generic surjection $g : \omega \rightarrow \omega\kappa$ is defined by $g = \bigcup G$, $s \subseteq g$. For all $k \leq |s|$, $g \restriction k = s \restriction k \in \pi_{|s|,k}^{\kappa;Z}[\mathfrak{B}_{p(|s|)}^{|s|(\omega\kappa)}] = \mathfrak{B}_{p(k)}^{k(\omega\kappa)}$ since $\Pi_{|s|,k}^{\kappa;Z}(p(|s|)) = p(k)$ by definition of $\mathbb{B}_{\omega}^{\kappa;Z}$. Since $\text{dim}(p) \leq |s|$, for all $k > |s|$, $g \restriction k \in \mathfrak{B}_{p(|s|)}^{|s|(\omega\kappa)} \times \mathbb{R}^{k-|s|} = \mathfrak{B}_{p(k)}^{k(\omega\kappa)}$. By definition of $G_g^{\kappa;Z;\omega}$, $p \in G_g^{\kappa;Z;\omega}$. Since $p \in D$, $G_g^{\kappa;Z;\omega} \cap D \neq \emptyset$. Since D was an arbitrary dense subset of $\mathbb{B}_{\omega}^{\kappa;Z}$ in M , $G_g^{\kappa;Z;\omega}$ is $\mathbb{B}_{\omega}^{\kappa;Z}$ -generic over M . $\square$

The following fact will be an important tool for proving the necessary capturing of set in the next section (Lemma 5.9).

Fact 4.17. (Woodin) Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a good parameter set such that $V = L(Z, \mathbb{R})$. Let $\kappa < \Theta$. Let M be an inner model of HOD_Z such that $\dot{\mathbb{B}}^{\kappa;Z} \in M$. Let $n \in \omega$, $x \in {}^n(\omega\kappa)$, $Y \in M[x]$, $a \in M[x] \cap L(Y, \omega\kappa)$, and φ is a $\mathcal{L}_{\text{set}}$ -formula. Then

$$L(Y, \omega\kappa) \models \varphi(a) \text{ if and only if } M[x] \Vdash 1_{\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega,n}^{\kappa;Z} G_x^{\kappa;Z;n}} \Vdash \dot{\mathbb{B}}^{\kappa;Z} / \Pi_{\omega,n}^{\kappa;Z} G_x^{\kappa;Z;n} L(\check{Y}, \omega\dot{\kappa}^{\text{sym}}) \models \varphi(\check{a}).$$

1132 (Note that if $\kappa < \Theta^{L(Y, \mathbb{R})}$, then $L(Y, \omega_\kappa) = L(Y, \mathbb{R})$ by the Moschovakis coding lemma, Fact 2.15. Thus in
1133 the above, ω_κ and $\omega_{\check{\kappa}^{\text{sym}}}$ can be replaced with $\mathbb{R}$ and $\mathbb{R}^{\check{\kappa}^{\text{sym}}}$, respectively.)

Proof. ($\Rightarrow$) Suppose $L(Y, \omega_\kappa) \models \varphi(a)$. Let $\mathbf{p} \in \mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}$. By the definition of the quotient forcing, this means that $\mathbf{p}(n) = \Pi_{\omega, n}^{\kappa; Z}(\mathbf{p}) \in G_x^{\kappa; Z; n}$ and thus $x \in \mathfrak{B}_{\mathbf{p}(n)}^{n(\omega_\kappa)}$. Let $m = \max\{n, \dim(\mathbf{p})\}$. Pick $s \in {}^m(\omega_\kappa)$ such that $s \restriction n = x$ and $s \in \mathfrak{B}_{\mathbf{p}(m)}^{m(\omega_\kappa)}$. Let $G \subseteq \text{Coll}(\omega, \omega_\kappa)$ be $\text{Coll}(\omega, \omega_\kappa)$ -generic over the true universe V with $s \in G$. (The generic does not exist in the real world. One will use the usual convenient forcing argument to conclude a fact about the forcing relation via the forcing theorem.) Let $g : \omega \rightarrow \omega_\kappa$ be the generic surjection added by G . $G_g^{\kappa; Z; \omega}$ is $\mathbb{B}_{\omega, n}^{\kappa; Z}$ -generic over M by Fact 4.16. Since $s \in G$, $s \in \mathfrak{B}_{\mathbf{p}(m)}^{m(\omega_\kappa)}$, and $\dim(\mathbf{p}) \leq m$, one has that $g \restriction k \in \mathfrak{B}_{\mathbf{p}(k)}^{k(\omega_\kappa)}$ for all $k \in \omega$. Thus $\mathbf{p} \in G_g^{\kappa; Z; \omega}$. Note $G_x^{\kappa; Z; n} = \langle \Pi_{\omega, n}^{\kappa; Z}[G_g^{\kappa; Z; \omega}] \rangle_{\mathbb{B}_n^{\kappa; Z}}$ and $G_g^{\kappa; Z; \omega}$ is $\mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}$ -generic over $M[G_x^{\kappa; Z; n}]$ by the properties of forcing projections discussed in Definition 3.7. Since $\omega_{\check{\kappa}^{\text{sym}}}[G_g^{\kappa; Z; \omega}] = \omega_\kappa$ and $L(Y, \omega_\kappa) \models \varphi(a)$, one has

$$M[G_x^{\kappa; Z; n}][G_g^{\kappa; Z; \omega}] \models L(\check{Y}, \omega_{\check{\kappa}^{\text{sym}}}[G_g^{\kappa; Z; \omega}]) \models \varphi(\check{a})$$

Since $M[x] = M[G_x^{\kappa; Z; n}]$ by Fact 4.7, $\mathbf{p} \in G_g^{\kappa; Z; \omega}$, and by using the forcing theorem, there is a $\mathbf{q} \leq \mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}$ $\mathbf{p}$ such that

$$M[x] \models \mathbf{q} \Vdash_{\mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}} L(\check{Y}, \omega_{\check{\kappa}^{\text{sym}}}) \models \varphi(\check{a}).$$

1134 Since $\mathbf{p} \in \mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}$ was arbitrary, one has that within $M[x]$, $1_{\mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}}$ forces the inner state-
1135 ment.

1136 ($\Leftarrow$) Considering x as a condition of $\text{Coll}(\omega, \omega_\kappa)$, pick $G \subseteq \text{Coll}(\omega, \omega_\kappa)$ which is $\text{Coll}(\omega, \omega_\kappa)$ -generic over
1137 V with $x \in G$. Let g be the generic surjection associated to G and let $G_g^{\kappa; Z; \omega}$ be the associated $\mathbb{B}_{\omega, n}^{\kappa; Z}$ -
1138 generic filter over M by Fact 4.16. $G_x^{\kappa; Z; n} = \langle \Pi_{\omega, n}^{\kappa; Z}[G_g^{\kappa; Z; \omega}] \rangle_{\mathbb{B}_n^{\kappa; Z}}$ is $\mathbb{B}_n^{\kappa; Z}$ -generic over M by Fact 4.6. Again
1139 $M[x] = M[G_x^{\kappa; Z; n}]$ by Fact 4.7. As before using general properties of forcing projections from Definition 3.7,
1140 $G_g^{\kappa; Z; \omega}$ is $\mathbb{B}_{\omega, n}^{\kappa; Z} / \Pi_{\omega, n}^{\kappa; Z} G_x^{\kappa; Z; n}$ generic over $M[x]$. By the hypothesis and the forcing theorem, $M[x][G_g^{\kappa; Z; \omega}] \models$
1141 $L(Y, \omega_{\check{\kappa}^{\text{sym}}}[G_g^{\kappa; Z; \omega}]) \models \varphi(a)$. Since $\omega_{\check{\kappa}^{\text{sym}}}[G_g^{\kappa; Z; \omega}] = \omega_\kappa$, one has $L(Y, \omega_\kappa) \models \varphi(a)$. $\square$

1142 The significance of Fact 4.17 is that for any inner model M of HOD_Z with $\check{\mathbb{B}}^{\kappa; Z} \in M$, $x \in \omega_\kappa$, and
1143 $Y \in M[x]$, the truth about objects $a \in M[x] \cap L(Y, \omega_\kappa)$ within the larger choiceless universe $L(Y, \omega_\kappa)$ can
1144 be determined within the ZFC model $M[x]$. There are many applications of Fact 4.17.

1145 In applications below, the parameter set Z may contain elements which do not belong to HOD_Z (but they
1146 are of course OD_Z). For example, if boldface GCH at ω holds and there is an uncountable $A \in \mathcal{P}(\mathbb{R})$ such
1147 that $A \in Z$, then $A \in \text{OD}_Z$ but $A \notin \text{HOD}_Z$. Another very relevant example is if Σ is an iteration strategy
1148 for a countable mouse and $\Sigma \in Z$, then $\Sigma \in \text{OD}_Z$ but $\Sigma \notin \text{HOD}_Z$. However, in many cases, there is a set
1149 $Y \in \text{HOD}_Z$ so that $\text{OD}_Z = \text{OD}_Y$. In this situation, Fact 4.17 can be used to more fully understand HOD_Z
1150 as in the following results.

1151 **Definition 4.18.** A set Z is a very good parameter set if and only if Z is a good parameter set and there is
1152 a set Y so that $Y \in \text{HOD}_Z$ and $\text{HOD}_Z = \text{HOD}_Y$. A set Y as above is called a very good parameter-witness
1153 for Z .

1154 Suppose $V = L(Z, \mathbb{R})$ and Z is a very good parameter set with very good parameter-witness Y . Since Z
1155 is a good parameter set, every element of $L(Z, \mathbb{R})$ is an ordinal definable from finitely many elements of Z
1156 and finitely many reals. Since Y is a very good parameter-witness to Z , every element of $L(Z, \mathbb{R})$ is ordinal
1157 definable from finitely many parameters from Y and finitely many reals. Thus Y is also a good parameter
1158 set and $L(Z, \mathbb{R}) \subseteq L(Y, \mathbb{R})$. Since $V = L(Z, \mathbb{R})$ and hence $Y \in L(Z, \mathbb{R})$, one has $L(Z, \mathbb{R}) = L(Y, \mathbb{R})$. Again,
1159 one can always find a finite very good parameter-witness Y . Moreover, since elements of $y \in Y$ belong to
1160 $\text{HOD}_Z \models \text{ZFC}$, after choosing a bijection π of an ordinal δ onto the transitive closure of y , one can code y as
1161 a set of ordinals. Thus the very good parameter-witness can be chosen to be a finite set of sets of ordinals.

1162 Formally, one can always replace a very good parameter Z with its very good parameter-witness Y . In
1163 practice, as in the proof of main result of the paper (Theorem 6.7), one naturally has a good parameter set

1164 Z which contains elements that do not belong to HOD_Z but one must show that Z is a very good parameter
1165 set by finding its very good parameter-witness Y .

1166 **Fact 4.19.** Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set such that $V = L(Z, \mathbb{R})$. Let $\kappa < \Theta$.
1167 If $n \in \omega$ and $x \in {}^n(\omega^\kappa)$, then $\text{HOD}_Z[x] = \text{HOD}_{Z \cup \{x\}}$.

Proof. Let Y be a very good parameter-witness for Z . Since $\text{OD}_Z = \text{OD}_Y$, $\text{HOD}_Z[x] = \text{HOD}_Y[x]$ and
 $\text{HOD}_{Z \cup \{x\}} = \text{HOD}_{Y \cup \{x\}}$. Thus one will work with Y rather than Z . Also note that $Y \in \text{HOD}_Z = \text{HOD}_Y$.
Since $x \in \text{HOD}_{Y \cup \{x\}}$, $\text{HOD}_Y[x] \subseteq \text{HOD}_{Y \cup \{x\}}$. Let $U \in \text{HOD}_{Y \cup \{x\}}$. Let W be the transitive closure of $\{U\}$
and note that $W \in \text{HOD}_{Y \cup \{x\}}$. Since $\text{HOD}_{Y \cup \{x\}} \models \text{AC}$, let $\pi : \delta \rightarrow W$ be a bijection where $\delta \in \text{ON}$ and
 $\pi \in \text{HOD}_{Y \cup \{x\}}$. Define $R \subseteq \delta \times \delta$ by $R(\alpha, \beta)$ if and only if $\pi(\alpha) \in \pi(\beta)$. Note that $R \in \text{HOD}_{Y \cup \{x\}}$. Thus
there is a finite tuple $\vec{y}$ in Y , a finite tuple of ordinals $\vec{\xi}$, and a $\mathcal{L}_{\text{set}}$ -formula φ so that $R(\alpha, \beta)$ if and only if
 $\varphi(\vec{y}, \vec{\xi}, x, \alpha, \beta)$. Since $Y \in \text{HOD}_Z = \text{HOD}_Y$, all the elements of the tuple $\vec{y}$ also belong to $\text{HOD}_Y \subseteq \text{HOD}_Y[x]$.
By Fact 4.17, for all $\alpha, \beta \in \delta$, $\varphi(\vec{y}, \vec{\xi}, x, \alpha, \beta)$ if and only if $L(Y, \mathbb{R}) \models \varphi(\vec{y}, \vec{\xi}, x, \alpha, \beta)$ if and only if

$$\text{HOD}_Y[x] \models 1_{\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega, n}^{\kappa;Z} G_x^{\kappa;Z; n}} L(\check{Y}, \dot{\mathbb{R}}^{\kappa; \text{sym}}) \models \varphi(\check{\vec{y}}, \check{\vec{\xi}}, \check{x}, \check{\alpha}, \check{\beta}).$$

1168 (Note $\vec{\mathbb{B}}^{\kappa;Z} \in \text{HOD}_Y = \text{HOD}_Z$ although one could also have used $\vec{\mathbb{B}}^{\kappa;Y}$.) By comprehension inside $\text{HOD}_Y[x]$,
1169 one has that $R \in \text{HOD}_Y[x]$. $W \in \text{HOD}_Y[x]$ since W is the Mostowski collapse of (δ, R) . Then $U \in \text{HOD}_Y[x]$,
1170 since U is the unique element of highest rank in W . Since $U \in \text{HOD}_{Y \cup \{x\}}$ was arbitrary, this shows that
1171 $\text{HOD}_{Y \cup \{x\}} \subseteq \text{HOD}_Y[x]$. Thus $\text{HOD}_Y[x] = \text{HOD}_{Y \cup \{x\}}$. $\square$

1172 An argument similar to the proof of Fact 4.19 gives the next result.

1173 **Fact 4.20.** Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set with good parameter witness Y such
1174 that $V = L(Z, \mathbb{R})$. Then $\text{HOD}_Z = L[\vec{\mathbb{B}}^{\kappa;Z}, Y]$.

1175 In particular, $\text{HOD}^{L(\mathbb{R})} = L[\vec{\mathbb{B}}^{\kappa; \emptyset}]$ for any $\kappa < \Theta^{L(\mathbb{R})}$ and in particular $\text{HOD}^{L(\mathbb{R})} = L[\vec{\mathbb{B}}^{\omega; \emptyset}]$. By using Fact
1176 4.19, for any $\kappa < \Theta^{L(\mathbb{R})}$, $n \in \omega$, and $x \in {}^n(\omega^\kappa)$, $\text{HOD}_{\{x\}}^{L(\mathbb{R})} = L[\vec{\mathbb{B}}^{\kappa; \emptyset}, x]$.

1177 Fact 4.17 can be used to show all subsets ${}^\omega\kappa$ in $L(Z, \mathbb{R})$ have ∞ -Borel codes in a uniform manner.

1178 **Fact 4.21.** Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set such that $V = L(Z, \mathbb{R})$. Let $\kappa < \Theta$.
1179 All subsets of ${}^\omega\kappa$ have ∞ -Borel codes. Moreover, if Y is a very good parameter-witness for Z and $A \subseteq {}^\omega\kappa$
1180 is $\text{OD}_{Y \cup \{x\}}$ for some $x \in \mathbb{R}$, then A has an ∞ -Borel code which is $\text{OD}_{Y \cup \{x\}}$ (and this ∞ -Borel code can be
1181 obtained uniformly in the sense described in the proof).

Proof. Let $A \subseteq {}^\omega\kappa$. Let Y be a very good parameter-witness for Z . Since $\text{OD}_Z = \text{OD}_Y$ and $V = L(Z, \mathbb{R}) =$
 $L(Y, \mathbb{R})$, there is a $\mathcal{L}_{\text{set}}$ formula φ , finite tuple $\vec{y}$ from Y , finite tuple of ordinals $\vec{\xi}$, and a finite tuple x in $\mathbb{R}$
(say n is the length of the tuple x) so that for all $a \in {}^\omega\kappa$, $a \in A$ if and only if $\varphi(\vec{y}, \vec{\xi}, x, a)$. Fact 4.17 asserts
that $a \in A$ if and only if $L(Y, \mathbb{R}) \models \varphi(\vec{y}, \vec{\xi}, x, a)$ if and only if

$$L[\vec{\mathbb{B}}^{\kappa;Z}, Y, \vec{y}, x, a] \models 1_{\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega, n+1}^{\kappa;Z} G_{x \cdot a}^{\kappa;Z; n+1}} \Vdash_{\mathbb{B}_{\omega}^{\kappa;Z} / \Pi_{\omega, n+1}^{\kappa;Z} G_{x \cdot a}^{\kappa;Z; n+1}} L(\check{Y}, \dot{\mathbb{R}}^{\kappa; \text{sym}}) \models \varphi(\check{\vec{y}}, \check{\vec{\xi}}, \check{x}, \check{a}).$$

1182 The inner expression can be written as a formula $\psi(\vec{\mathbb{B}}^{\kappa;Z}, Y, \vec{y}, \vec{\xi}, x, a)$. One has shown that $a \in A$ if and
1183 only if $L[\vec{\mathbb{B}}^{\kappa;Z}, Y, \vec{y}, \vec{\xi}, x, a] \models \psi(\vec{\mathbb{B}}^{\kappa;Z}, Y, \vec{\xi}, x, a)$. Hence $(\langle \vec{\mathbb{B}}^{\kappa;Z}, Y, x, \vec{\xi} \rangle, \psi)$ is an ∞ -Borel code for A . (Since
1184 $Y \in \text{HOD}_Z = \text{HOD}_Y$, $\langle \vec{\mathbb{B}}^{\kappa;Z}, Y, \vec{y}, x, \vec{\xi} \rangle$ is essentially a set of ordinals and is $\text{OD}_{Y \cup \{x\}}$.) $\square$

1185 **Fact 4.22.** Assume AD, $\text{DC}_{\mathbb{R}}$, and all subsets of $\mathbb{R}$ has ∞ -Borel codes. Let $\kappa < \Theta$. All subsets of ${}^\omega\kappa$ have
1186 ∞ -Borel code.

1187 *Proof.* Let $\kappa < \Theta$ and let $\pi : \mathbb{R} \rightarrow \kappa$ be a surjection. Let $\preceq$ be a prewellordering on $\mathbb{R}$ defined by $x \preceq y$ if
1188 and only if $\pi(x) \leq \pi(y)$. Let $\varpi : \mathbb{R} \rightarrow {}^\omega\kappa$ be a surjection defined by $\varpi(x)(n) = \pi(x^{[n]})$. Let $A \subseteq {}^\omega\kappa$. Define
1189 $\tilde{A} \subseteq \mathbb{R}$ by $\tilde{A} = \{x \in \mathbb{R} : \varpi(x) \in A\}$. By the hypothesis, let S be a set of ordinals and φ_0, φ_1 be two formulas
1190 such that (S, φ_0) is an ∞ -Borel code for $\tilde{A}$ and (S, φ_1) is an ∞ -Borel code for $\preceq$. Then $\tilde{A}, \preceq \in L(S, \mathbb{R})$.
1191 Within $L(S, \mathbb{R})$, $\preceq$ can be used to recover π . $L(S, \mathbb{R})$ can define ϖ . Using $\tilde{A}$, one can show that $A \in L(S, \mathbb{R})$.
1192 In the inner model $L(S, \mathbb{R})$, $\{S\}$ is a very good parameter set. By Fact 4.21, A has an ∞ -Borel code in
1193 $L(S, \mathbb{R})$ and thus A has an ∞ -Borel code in the real world. $\square$

A very important observation from the proof of Fact 4.21 is that there is an explicit procedure that transforms the definition of A (using parameter Y , finitely many elements from Y , finitely many reals, and finitely many ordinals) to an ∞ -Borel code for A . This uniform transformation is used for various combinatorial results under natural models of AD^+ such as in [13] and [10]. This uniformity is especially powerful when combined with the proof of Woodin's perfect set dichotomy ([8] Theorem 8.5, [4]).

Fact 4.17 is also used to show that in models of the form $L(Z, \mathbb{R})$ as above, $\bar{\mathbb{B}}^{\omega; Z}$ can be used to define intermediate cardinalities which provide counterexamples to the cardinality structure below $\mathcal{P}(\omega_1)$ proved under $\text{AD}_{\frac{1}{2}\mathbb{R}}$ (or equivalently $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$). Under $\text{AD}_{\frac{1}{2}\mathbb{R}}$, [13] Corollary 7.6 showed that the only uncountable cardinality below $|\mathbb{R} \times \omega_1|$ are $|\mathbb{R}|$, $|\omega_1|$, $|\mathbb{R} \sqcup \omega_1|$, and $|\mathbb{R} \times \omega_1|$. [13] Theorem 7.33 isolated a very large family of cardinalities between $|\mathbb{R}|$ and $|\mathbb{R} \times \omega_1|$ which do not have an injective copy of ω_1 in $L(\mathbb{R})$. In $L(\mathbb{R})$, let $N_1 = \bigsqcup_{r \in \mathbb{R}} ((\omega_1^{L(\mathbb{R})})^{+})^{L[\bar{\mathbb{B}}^{\omega; 0}, r]}$. [13] Theorem 4.10 shows $|\mathbb{R} \times \omega_1| < |N_1| < |\mathbb{R} \times \omega_2|$, $|N_1| < |\mathcal{P}(\omega_1)|$, $\neg(|N_1| \leq |\omega_1|^{<\omega_1})$, $\neg(|[\omega_1]^\omega| \leq |N_1|)$, and $|\omega_1|^{<\omega_1} < |N_1 \cup [\omega_1]^{<\omega_1}| < |\mathcal{P}(\omega_1)|$. In $L(\mathbb{R})$, let $M_1 = \bigsqcup_{f \in [\omega_1]^{<\omega_1}} ((\omega_1^{L(\mathbb{R})})^{+})^{L[\bar{\mathbb{B}}^{\omega; 0}, f]}$. Then in $L(\mathbb{R})$, $|\omega_1|^{<\omega_1} < |M_1| < |\mathcal{P}(\omega_1)|$ by similar arguments. Woodin [65] under $\text{AD}_{\mathbb{R}}$ and DC and later the author under $\text{AD}_{\frac{1}{2}\mathbb{R}}$ showed that if $|X| \leq |\omega_1|^\omega$, then either $|X| \leq |\mathbb{R} \times \omega_1|$ or $|\omega_1|^\omega \leq |X|$. It was open whether a counterexample existed if $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$ failed. As an application of the Fact 4.17, the next result will produce the required intermediate cardinality and illustrate how $\bar{\mathbb{B}}^{\kappa; Z}$ is used to diagonalized against injections.¹² This will motivate analogous constructions of intermediate cardinalities below $\mathcal{P}_B(\kappa)$ in $L(\mathbb{R})$ using the techniques of the next section. (See Theorem 6.27.)

If Z is a good parameter set and $V = L(Z, \mathbb{R})$ satisfies boldface GCH at ω , then $L(Z, \mathbb{R})$ fails to satisfy $\text{AD}_{\frac{1}{2}\mathbb{R}}$ or equivalently $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$. This follows from the usual example of Kechris and Solovay: Let $R \subseteq \mathbb{R} \times \mathbb{R}$ be defined by $R(x, y)$ if and only if $y \notin \text{OD}_{Z \cup \{x\}}$. Since Z is a parameter set, $\text{HOD}_Z \models \text{ZFC}$ and hence $\mathbb{R} \cap \text{HOD}_Z$ is wellorderable. By boldface GCH at ω , $\mathbb{R} \cap \text{OD}_Z$ is countable. Thus for all $x \in \mathbb{R}$ and $y \notin \text{OD}_{Z \cup \{x\}}$, $R(x, y)$. That is, $\text{dom}(R) = \mathbb{R}$. Suppose $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a uniformization for R . Since Z is a good parameter set, Φ is $\text{OD}_{Z \cup \{x\}}$ for some $x \in \mathbb{R}$. Then $\Phi(x) \in \text{OD}_{Z \cup \{x\}}$ and $R(x, \Phi(x))$ which is a contradiction.

Theorem 4.23. *Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set such that $V = L(Z, \mathbb{R})$. Let Y be a very good parameter-witness for Z which is a finite set of sets of ordinals. Let $\vec{y}$ be a finite enumeration of Y .¹³ Let $E_2^{\vec{y}} = \bigsqcup_{r \in \mathbb{R}} \omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]} = \{(r, f) : r \in \mathbb{R} \wedge f \in \omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]}\}$. Then $|E_2^{\vec{y}}| \leq |\omega \omega_1|$, $\neg(|\omega_1| \leq |E_2^{\vec{y}}|)$, $\neg(|E_2^{\vec{y}}| \leq |\mathbb{R} \times \omega_1|)$, and $|\mathbb{R} \times \omega_1| < |(\mathbb{R} \times \omega_1) \sqcup E_2^{\vec{y}}| < |\omega \omega_1|$.*

Proof. Since $\omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]} < \omega_1$ for all $r \in \mathbb{R}$ by boldface GCH at ω (which holds under just AD as noted in Section 2), $E_2^{\vec{y}} \subseteq \mathbb{R} \times \omega \omega_1$ and thus $|E_2^{\vec{y}}| \leq |\omega \omega_1|$.

Suppose $\Phi_0 : \omega_1 \rightarrow E_2^{\vec{y}}$ be an injection. Let $\pi_1 : \mathbb{R} \times \omega \omega_1 \rightarrow \mathbb{R}$ be the projection onto the first coordinate. $\pi_1[\Phi_0[\omega_1]] \subseteq \mathbb{R}$ is a wellorderable subset of $\mathbb{R}$ and thus must be countable by boldface GCH at ω . Since ω_1 is regular by $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$, there is an $r \in \mathbb{R}$ so that $|(\pi_1 \circ \Phi_0)^{-1}[\{r\}]| = \omega_1$. So $\Phi_0 : (\pi_1 \circ \Phi_0)^{-1}[\{r\}] \rightarrow \{r\} \times \omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]}$ is an injection. Since $\omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]} < \omega_1$, $|\omega_2^{L[\bar{\mathbb{B}}^{\omega_1; \vec{y}}, r]}| = |\mathbb{R}|$. Thus Φ_0 induces an injection of ω_1 into $\mathbb{R}$ which violates boldface GCH at ω . This shows $\neg(|\omega_1| \leq |E_2^{\vec{y}}|)$.

Suppose $\Phi_1 : E_2^{\vec{y}} \rightarrow \mathbb{R} \times \omega_1$ is an injection. Identify $\mathbb{R}$ with $\mathcal{P}(\omega)$. If $(r, \alpha) \in \mathbb{R} \times \omega_1$, then let $P_{(r, \alpha)} \subseteq \omega \times \omega_1$ be defined by $P_{(r, \alpha)} = \{(n, \alpha) : n \in r\}$. If $(s, f) \in \mathbb{R} \times \omega \omega_1$, then let $Q_{(s, f)} \subseteq \omega \times \omega \times \omega_1$ be defined by $Q_{(s, f)} = \{(m, n, \beta) : m \in s \wedge f(n) = \beta\}$. Let $\text{Gr}_{\Phi_1} \subseteq \mathbb{R} \times \omega \omega_1 \times \omega \times \omega_1$ be defined by $\text{Gr}_{\Phi_1} = \{(r, f, m, \alpha) : (r, f) \in E_2^{\vec{y}} \wedge (m, \alpha) \in P_{\Phi_1(r, f)}\}$. Let $\text{Gr}_{\Phi_1^{-1}} \subseteq \mathbb{R} \times \alpha \times \omega \times \omega \times \omega_1$ be defined by $\text{Gr}_{\Phi_1^{-1}} = \{(s, \alpha, m, n, \beta) : (s, \alpha) \in \Phi_1[\mathbb{R} \times \omega_1] \wedge (m, n, \beta) \in Q_{\Phi_1^{-1}(s, \alpha)}\}$. Since $V = L(Z, \mathbb{R}) = L(Y, \mathbb{R}) = L(\vec{y}, \mathbb{R})$, there are formulas φ_0, φ_1 , finite tuple of ordinals $\vec{\xi}_0, \vec{\xi}_1$, and $e_0, e_1 \in \mathbb{R}$ so that for all $(x, m, \alpha) \in (\mathbb{R} \times \omega \omega_1) \times \omega \times \omega_1$ and $(y, m, n, \beta) \in (\mathbb{R} \times \omega_1) \times \omega \times \omega \times \omega_1$, $\text{Gr}_{\Phi_1}(x, m, \alpha) \Leftrightarrow \varphi_0(x, m, \alpha, \vec{y}, e_0, \vec{\xi}_0)$ and

¹²It is worth noting that one advantage of Fact 4.17 is that one can use $\bar{\mathbb{B}}^{\omega_1; Z}$ rather than $\bar{\mathbb{B}}^{\omega; Z}$ which yields a more elegant proof of Theorem 4.23. The result can still be proved using the classical $\bar{\mathbb{B}}^{\omega; Z}$; however, one would need to introduce a coding of ${}^\omega\omega_1$ by reals and the collapse forcing much like in the proof of Fact 2.24.

¹³By the discussion above, if Z is a very good parameters set, there will always be a very good parameter-witness Y which is a finite set of sets of ordinals. It is necessary to use a finite enumeration $\vec{y}$ of Y rather than the finite set Y itself. For example, if r is a nonconstructible reals, then $L[\{r\}] = L$. However, since Y is a finite set of sets of ordinals, Y can be canonically wellordered by the lexicographic ordering on the powerset of some sufficiently large ordinal. Thus each member of Y is OD_Y .

$\text{Gr}_{\Phi_1^{-1}}(y, m, n, \beta) \Leftrightarrow \varphi_1(y, m, n, \beta, \vec{y}, e_1, \vec{\xi}_1)$. Woodin ([40] Corollary 5.10) showed that there is an $e \in \mathbb{R}$ so that $e_0, e_1 \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$ and $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e] \models \text{CH}$. In $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$, let $\mathbb{P}$ be the Namba forcing ([31] Theorem 28.10) consisting of $p \subseteq {}^{<\omega}\omega_2$ which are nonempty trees on ω_2 so that each node of p has ω_2 many extensions. Let $\leq_{\mathbb{P}}$ be the subset relation. $\mathbb{P}$ is countable in the real world and there is a $G \subseteq \mathbb{P}$ which is $\mathbb{P}$ -generic over $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$ and exists in the real world by Fact 2.14. Since $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e] \models \text{CH}$, by [31] Theorem 28.10, $\mathbb{P}$ adds a generic cofinal function $g : \omega \rightarrow \omega_2^{L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]}$ induced by G and $\mathbb{P}$ adds no new reals. By Fact 4.17 and the fact that $e_0 \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$, $(m, \alpha) \in P_{\Phi_1(e, g)}$ if and only if $\text{Gr}_{\Phi_1}((e, g), m, \alpha)$ if and only if $\varphi_0((e, g), m, \alpha, \vec{y}, e_0, \vec{\xi}_0)$ if and only if

$$L[\vec{\mathbb{B}}^{\omega; \vec{y}}, \vec{y}, e][g] \models 1_{\mathbb{B}_{\omega_1; \vec{y}} / \Pi_{\omega_1; 1}^{\omega_1; \vec{y}}} G_g^{\omega_1; \vec{y}, 1} \Vdash_{\mathbb{B}_{\omega_1; \vec{y}} / \Pi_{\omega_1; 1}^{\omega_1; \vec{y}}} L(\vec{y}, \omega_1^{\text{sym}}) \models \varphi_0(\check{e}, \check{g}, \check{m}, \check{\alpha}, \check{\vec{y}}, \check{e}_0, (\check{\xi}_0)).$$

Thus $P_{\Phi_1(e, g)} \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e][g]$ by comprehension and hence $\Phi_1(e, g) \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e][g]$. Since $\Phi_1(e, g) \in \mathbb{R} \times \omega_1$ and $\mathbb{P}$ adds no new reals, $\Phi_1(e, g) \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$. By Fact 4.17, $(m, n, \beta) \in Q_{(e, g)}$ if and only if $(m, n, \beta) \in Q_{\Phi_1^{-1}(\Phi_1(e, g))}$ if and only if $\text{Gr}_{\Phi_1^{-1}}(\Phi_1(e, g), m, n, \beta)$ if and only if $\varphi_1(\Phi_1(e, g), m, n, \beta, \vec{y}, e_1, \vec{\xi}_1)$ if and only if

$$L[\vec{\mathbb{B}}^{\omega; \vec{y}}, \vec{y}, e] \models 1_{\mathbb{B}_{\omega_1; \vec{y}} / \Pi_{\omega_1; 1}^{\omega_1; \vec{y}}} \Vdash_{\mathbb{B}_{\omega_1; \vec{y}} / \Pi_{\omega_1; 1}^{\omega_1; \vec{y}}} L(\vec{y}, \omega_1^{\text{sym}}) \models \varphi_0((\Phi(e, g)), \check{m}, \check{n}, \check{\beta}, \check{\vec{y}}, \check{e}_1, (\check{\xi}_1)).$$

Thus by comprehension, $Q_{(e, g)} \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$ and thus $g \in L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e]$. This is impossible since by absoluteness, $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e] \models "g : \omega \rightarrow \omega_2 \text{ is cofinal}"$. Since $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e] \models \text{AC}$, $L[\vec{\mathbb{B}}^{\omega_1; \vec{y}}, \vec{y}, e] \models \text{cof}(\omega_2) = \omega_2$. Contradiction.

Since $E_2^{\vec{y}}$ does not inject into $\mathbb{R} \times \omega_1$, one has that $|\mathbb{R} \times \omega_1| < |(\mathbb{R} \times \omega_1) \sqcup E_2^{\vec{y}}|$. Since $|\omega_1| = |\omega_1|^\omega = |[\omega_1]_*|^\omega$, suppose there is an injection $\Phi_2 : [\omega_1]_*^\omega \rightarrow (\mathbb{R} \times \omega_1) \sqcup E_2^{\vec{y}}$. Since AD implies $\omega_1 \rightarrow_* (\omega_1)_2^\omega$, $\mu_{\omega_1}^\omega$ is an ultrafilter on $[\omega_1]_*^\omega$. Thus either $\Phi_2^{-1}[\mathbb{R} \times \omega_1] \in \mu_{\omega_1}^\omega$ or $\Phi_2^{-1}[E_2^{\vec{y}}] \in \mu_{\omega_1}^\omega$. Thus $|\Phi_2^{-1}[\mathbb{R} \times \omega_1]| = |\omega_1|$ or $|\Phi_2^{-1}[E_2^{\vec{y}}]| = |\omega_1|$. Thus Φ_2 either induces an injection of ${}^\omega\omega_1$ into $\mathbb{R} \times \omega_1$ or an injection of ${}^\omega\omega_1$ into $E_2^{\vec{y}}$. Thus there is an injection of ${}^\omega\omega_1$ into $\mathbb{R} \times \omega_1$ or an injection of ${}^\omega\omega_1$ into $E_2^{\vec{y}}$. The first is impossible by Fact 2.10 or Fact 3.5. The second is impossible since $|\omega_1| \leq |{}^\omega\omega_1|$ and it was shown above that $\neg(|\omega_1| \leq |E_2^{\vec{y}}|)$. It has been shown that $|(\mathbb{R} \times \omega_1) \sqcup E_2^{\vec{y}}| < |{}^\omega\omega_1|$. $\square$

Woodin showed that natural models of determinacy satisfying AD^+ , $V = L(\mathcal{P}(\mathbb{R}))$, and $\neg\text{AD}_{\mathbb{R}}$ must take the form $L(J, \mathbb{R})$ where J is a set of ordinals. Thus Woodin's dichotomy (the statement that for all $X \subseteq [\omega_1]^\omega$, $|X| \leq |\mathbb{R} \times \omega_1|$ or $|[\omega_1]^\omega| \leq |X|$) must always fails under AD^+ , $V = L(\mathcal{P}(\mathbb{R}))$, and $\neg\text{AD}_{\mathbb{R}}$ (with counterexamples derived from $E_2^{(J)}$ when $V = L(J, \mathbb{R})$).

5. AMENABLE SURJECTIONS, GENERIC FILTER EXISTENCE, AND CAPTURING

This section will prove the basic tools needed for the main result concerning the cardinality of $\mathcal{P}_B(\kappa)$ when $\kappa < \Theta$.

The proof of Fact 2.25 and Theorem 3.3 serve as the basic template for the main argument. In Fact 2.25, the relevant forcing was $\text{Coll}(\omega^+, \omega^{++})^M$ for some inner model M of ZFC. Since boldface GCH at ω was assumed to hold in the real world, Fact 2.14 asserts that there is a $\text{Coll}(\omega^+, \omega^{++})^M$ -generic filter over M which exists in the real world.

For motivation, assume $V = L(\mathbb{R})$ and for the sake of contradiction there is an injection $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ for $\epsilon < \kappa < \Theta$. Φ is ordinal definable from some $x \in \mathbb{R}$, but for simplicity assume $x = \emptyset$ and thus $\Phi \in \text{OD}^{L(\mathbb{R})}$. The analogous forcing will be $\mathbb{P} = \text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}^{L(\mathbb{R})}}$. The first step will be to show a $\mathbb{P}$ -generic filter over $\text{HOD}^{L(\mathbb{R})}$ exists in the real world. Since Steel ([60] Corollary 8.22) showed $\text{HOD}^{L(\mathbb{R})} \models \text{GCH}$, $\text{HOD}^{L(\mathbb{R})} \models |\mathcal{P}(\mathbb{P})| = 2^{\epsilon^{++}} = \epsilon^{+++}$. The collection $\mathcal{D}$ of all dense subsets of $\mathbb{P}$ in $\text{HOD}^{L(\mathbb{R})}$ is internally in bijection with $(\epsilon^{+++})^{\text{HOD}^{L(\mathbb{R})}}$. Since Steel ([60] Theorem 8.26) showed boldface GCH holds below Θ in $\text{HOD}^{L(\mathbb{R})}$, there is a bijection of $(\epsilon^+)^{\text{HOD}^{L(\mathbb{R})}}$ (even of ϵ or $|\epsilon|^{L(\mathbb{R})}$) onto $(\epsilon^{+++})^{\text{HOD}^{L(\mathbb{R})}}$ externally in the real world. Thus there is a surjection $h : (\epsilon^+)^{\text{HOD}^{L(\mathbb{R})}} \rightarrow \mathcal{D}$. It is tempting to use that $\text{HOD}^{L(\mathbb{R})} \models " \mathbb{P} = \text{Coll}(\epsilon^+, \epsilon^{++}) \text{ is } < \epsilon^+ \text{-closed} "$ to extend conditions in an ϵ^+ -length construction to meet the dense set $h(\alpha)$ at stage α . This construction can only continue for $(\epsilon^+)^{\text{HOD}^{L(\mathbb{R})}}$ -stages if $h \restriction \alpha \in \text{HOD}^{L(\mathbb{R})}$ for

each $\alpha < \epsilon^+$ so that the α^{th} -stage of the construction can be internalized into $\text{HOD}^{L(\mathbb{R})}$ to use the fact that $\text{HOD}^{L(\mathbb{R})} \models \text{“}\mathbb{P} \text{ is } < \epsilon^+ \text{-closed”}$. The existence of such a surjection h which is “amenable” relative to $\text{HOD}^{L(\mathbb{R})}$ will be a consequence of one further important consequence ([60] Lemma 8.25) of Steel’s directed system analysis of $\text{HOD}^{L(\mathbb{R})}$: namely, for every $\epsilon < \Theta$, $\text{cof}^{L(\mathbb{R})}((\epsilon^+)^{\text{HOD}^{L(\mathbb{R})}}) = \omega$.

Let $M \models \text{ZF}$ be an inner model. A set $A \subseteq M$ is M -amenable if and only if for all $x \in M$, $A \cap x \in M$. Here, one will be most concerned with functions $f : \epsilon \rightarrow M$ where $\epsilon \in \text{ON}$. A function $f : \epsilon \rightarrow M$ is M -amenable if and only if for all $\alpha < \epsilon$, $f \restriction \alpha = \{(\xi, f(\xi)) : \xi < \alpha\} \in M$.

The next lemma gives a criterion for the existence of certain M -amenable surjections which are uniform in certain objects. A single application of the lemma will only need an appropriate form of countable choice (see Lemma 5.2). However, the uniformity will be important for applying the lemma to obtain uncountably many M -amenable surjections simultaneously which will be needed in Section 7.

Lemma 5.1. *Let $\omega < \kappa < \lambda$ and M is an inner model of ZF such that $M \models \text{“}\kappa \text{ and } \lambda \text{ are cardinals”}$. Suppose ρ_0 , ρ_1 , and ϖ have the following properties.*

- (1) $\rho_0 : \omega \rightarrow \kappa$ is an increasing cofinal function through κ .
- (2) $\rho_1 : \omega \rightarrow \lambda$ is an increasing cofinal function through λ .
- (3) ϖ is a function on ω so that for all $n \in \omega$, $\varpi(n) : \kappa \rightarrow \rho_1(n)$ is an M -amenable surjection of κ onto $\rho_1(n)$.

Then there is a M -amenable surjection of κ onto λ which is produced uniformly from ρ_0 , ρ_1 , and ϖ with the above property.¹⁴

Proof. Since $M \models \text{“}\kappa \text{ is a cardinal”}$, by modifying ρ_0 (uniformly) if necessary, one may assume ρ_0 takes values among the indecomposable limit ordinals. Let $I_0 = \{\alpha : 0 \leq \alpha < \rho_0(0)\}$ and for all $n > 0$, let $I_n = \{\alpha : \rho_0(n-1) \leq \alpha < \rho_0(n)\}$. Note that for all $n \in \omega$, $\text{ot}(I_n) = \rho_0(n)$ by the indecomposability of $\rho_0(n)$. Let $\mathbf{m}_n : I_n \rightarrow \rho_0(n)$ be the Mostowski collapse of I_n . Note that $I_n, \mathbf{m}_n \in M$ for each $n \in \omega$. For each $n \in \omega$, let $b_n : \rho_0(n) \rightarrow \rho_0(n) \times (n+1)$ be the standard bijection of $\rho_0(n)$ onto $\rho_0(n) \times (n+1)$ which belongs to M (or any inner model of ZF). Let $\pi_1 : \kappa \times \omega \rightarrow \kappa$ and $\pi_2 : \kappa \times \omega \rightarrow \omega$ be the projection on the first and second coordinate, respectively. For each $\alpha < \kappa$, let n_α be the unique n so that $\alpha \in I_n$. For each $n \in \omega$, let $f_n : \kappa \rightarrow \rho_1(n)$ be an M -amenable surjection defined by $f_n = \varpi(n)$. Define $\Phi : \kappa \rightarrow \lambda$ by $\Phi(\alpha) = f_{\pi_2(b_{n_\alpha}(\mathbf{m}_{n_\alpha}(\alpha)))}(\pi_1(b_{n_\alpha}(\mathbf{m}_{n_\alpha}(\alpha))))$.

Let $\beta < \lambda$. Let n_0 be the least $n \in \omega$ so that $\beta < \rho_1(n)$. Let n_1 be the least $n \in \omega$ so that $n \geq n_0$ and there is a $\gamma < \rho_0(n)$ so that $f_{n_0}(\gamma) = \beta$. Let γ_0 be the least γ such that $\gamma < \rho_0(n_1)$ and $f_{n_0}(\gamma) = \beta$. Let $\alpha = \mathbf{m}_{n_1}^{-1}(b_{n_1}^{-1}(\gamma_0, n_0))$. Note that $\alpha \in I_{n_1}$ and therefore $n_1 = n_\alpha$. Thus

$$\begin{aligned} \Phi(\alpha) &= f_{\pi_2(b_{n_\alpha}(\mathbf{m}_{n_\alpha}(\alpha)))}(\pi_1(b_{n_\alpha}(\mathbf{m}_{n_\alpha}(\alpha)))) = f_{\pi_2(b_{n_1}(\mathbf{m}_{n_1}(\alpha)))}(\pi_1(b_{n_1}(\mathbf{m}_{n_1}(\alpha)))) \\ &= f_{\pi_2(b_{n_1}(\mathbf{m}_{n_1}(\mathbf{m}_{n_1}^{-1}(b_{n_1}^{-1}(\gamma_0, n_0))))}(\pi_1(b_{n_1}(\mathbf{m}_{n_1}(\mathbf{m}_{n_1}^{-1}(b_{n_1}^{-1}(\gamma_0, n_0)))))) = f_{\pi_2(\gamma_0, n_0)}(\pi_1(\gamma_0, n_0)) = f_{n_0}(\gamma_0) = \beta \end{aligned}$$

Thus Φ is a surjection. Note that the definition of Φ is uniform in ρ_0 , ρ_1 , and ϖ .

Fix $n \in \omega$. Note that $\rho_0 \restriction n+1$, $\langle \mathbf{m}_k : k < n+1 \rangle$, $\langle I_k : k < n+1 \rangle$, and $\langle b_k : k < n+1 \rangle$ are elements of M (since these are finite functions into M). Since for all $k \leq n$, f_k is an M -amenable function, one has that $f_k \restriction \rho_0(n) \in M$ and thus $\langle f_k \restriction \rho_0(n) : k < n+1 \rangle \in M$. Note that $\Phi \restriction \rho_0(n)$ can be defined using $\rho_0 \restriction n+1$, $\langle \mathbf{m}_k : k < n+1 \rangle$, $\langle I_k : k < n+1 \rangle$, $\langle b_k : k < n+1 \rangle$, and $\langle f_k \restriction \rho_0(n) : k < n+1 \rangle$ using the definition of Φ above restricted to $\alpha < \rho_0(n)$. Since $M \models \text{ZF}$, $\Phi \restriction \rho_0(n) \in M$. Since ρ_0 is cofinal through κ , Φ is an M -amenable function. $\square$

Since AD implies $\text{AC}_\omega^\mathbb{R}$, one obtains the following using the Moschovakis coding lemma.

Lemma 5.2. *Assume AD. Let $\omega < \kappa < \lambda < \Theta$ with $\text{cof}(\kappa) = \text{cof}(\lambda) = \omega$ and M be an inner model of ZF such that $M \models \text{“}\kappa \text{ and } \lambda \text{ are cardinals”}$. Suppose for all $\delta < \lambda$, there is an M -amenable surjection of κ onto δ . Then there is an M -amenable surjection of κ onto λ .*

Proof. Since $\text{cof}(\kappa) = \text{cof}(\lambda) = \omega$, let $\rho_0 : \omega \rightarrow \kappa$ and $\rho_1 : \omega \rightarrow \lambda$ be increasing cofinal function through κ and λ , respectively. By the Moschovakis coding lemma (Fact 2.15), let $\varsigma : \mathbb{R} \rightarrow \mathcal{P}(\lambda)$ be a surjection. Define

¹⁴This means that there is a function Υ so that whenever ρ_0 , ρ_1 , and ϖ have the required properties, $\Upsilon(\rho_0, \rho_1, \varpi)$ is an M -amenable surjection from κ onto λ .

1301 $R \subseteq \omega \times \mathbb{R}$ by $R(n, x)$ if and only if $\varsigma(x)$ codes the graph of an M -amenable surjection of κ onto $\rho_1(n)$.
 1302 Observe that $\text{dom}(R) = \omega$ by the hypothesis. By $\text{AC}_\omega^\mathbb{R}$, there is a sequence $\langle x_n : n \in \omega \rangle$ so that $R(n, x_n)$
 1303 holds for all $n \in \omega$. Let $f_n : \kappa \rightarrow \rho_1(n)$ be the M -amenable surjection coded by $\varsigma(x_n)$. Let $\varpi(n) = f_n$. The
 1304 result now following from applying Lemma 5.1 to ρ_0 , ρ_1 , and ϖ . $\square$

1305 **Lemma 5.3.** *Assume AD. Let $\kappa < \Theta$ with $\text{cof}^V(\kappa) = \omega$ and M be an inner model of ZF such that $M \models \text{“}\kappa$
 1306 is a cardinal” . Suppose $\delta < (\kappa^{+\omega_1^V})^M$ and has the property that for all λ with $\kappa \leq \lambda \leq \delta$, if $M \models \text{“}\lambda$ is a
 1307 successor cardinal”, then $\text{cof}^V(\lambda) = \omega$. Then there is an M -amenable surjection of κ onto δ .*

1308 *Proof.* Note that if δ satisfies this hypothesis, then every $\gamma < \delta$ also satisfies this hypothesis. By induction,
 1309 it will be shown that for every $\gamma \leq \delta$, there is an M -amenable surjection of κ onto γ .

1310 If $\gamma < (\kappa^+)^M$, then there is a surjection of κ onto γ which is M -amenable (and even is a member of M).
 1311 Now suppose $(\kappa^+)^M \leq \gamma \leq \delta$ and for all $\beta < \gamma$, there is an M -amenable surjection of κ onto β .

1312 (Case I) γ is not a cardinal of M . There is a $g \in M$ so that $g : |\gamma|^M \rightarrow \gamma$ is a bijection. By the induction
 1313 hypothesis, there is an M -amenable surjection $k : \kappa \rightarrow |\gamma|^M$. Then $g \circ k : \kappa \rightarrow \gamma$ is an M -amenable surjection.

1314 (Case II) γ is a successor cardinal of M . By the hypothesis, $\text{cof}^V(\gamma) = \omega$. Since $\text{cof}^V(\kappa) = \omega$ and the
 1315 induction hypothesis, Lemma 5.2 implies that there is an M -amenable surjection of κ onto γ .

1316 (Case III) γ is a limit cardinal of M . Since $\gamma \leq \delta < (\kappa^{+\omega_1^V})^M$, $\gamma = (\kappa^{+\alpha})^M$ for some limit ordinal $\alpha < \omega_1^V$.
 1317 So $\text{cof}^V(\alpha) = \omega$ and thus $\text{cof}^V(\gamma) = \omega$. Again Lemma 5.2 implies that there is an M -amenable surjection of
 1318 κ onto γ . $\square$

1319 Frequently, one will need an M -amenable surjection of a cardinal of cofinality ω onto the double successor
 1320 of the cardinal. The following lemma isolates a uniform procedure to obtain such M -amenable surjection
 1321 from ω -cofinal functions into the original cardinal, the successor, and the double successor which will be
 1322 useful in the Section 7.

1323 **Lemma 5.4.** *Suppose M is an inner model of ZF and $\kappa \in \text{ON}$ is such that $M \models \text{“}\kappa$ is a cardinal”. Suppose
 1324 $\tau_0 : \omega \rightarrow \kappa$, $\tau_1 : \omega \rightarrow (\kappa^+)^M$, and $\tau_2 : \omega \rightarrow (\kappa^{++})^M$ are increasing cofinal map into κ , $(\kappa^+)^M$, and $(\kappa^{++})^M$,
 1325 respectively. Suppose $\prec$ is a wellordering of $(\mathcal{P}((\kappa^{++})^M))^M$. Then there is an M -amenable surjection of κ
 1326 onto $(\kappa^{++})^M$ which is obtained uniformly from τ_0 , τ_1 , τ_2 , and $\prec$.*

1327 *Proof.* Let $\rho_0^0 = \tau_0$. Let $\rho_1^0 = \tau_1$. Since for all $n \in \omega$, $\tau_1(n) < (\kappa^+)^M$, there is a surjection of κ onto $\tau_1(n)$
 1328 which is a member of M . Let $\varpi^0(n)$ be the $\prec$ -least surjection of κ onto $\tau_1(n) = \rho_1^0(n)$ and note that ϖ^0 is
 1329 produced uniformly in τ_0 , τ_1 , and $\prec$. Lemma 5.1 gives an M -amenable surjection $h_0 : \kappa \rightarrow (\kappa^+)^M$ which
 1330 is produced uniformly in ρ_0^0 , ρ_1^0 , and ϖ^0 . Let $\rho_0^1 = \tau_0$ and $\rho_1^1 = \tau_2$. For all $n \in \omega$, $\tau_2(n) < (\kappa^{++})^M$ and
 1331 hence there is a surjection of $(\kappa^+)^M$ onto $\tau_2(n)$ which belong to M . Let $\sigma_n : (\kappa^+)^M \rightarrow \tau_2(n)$ be the $\prec$ -least
 1332 surjection of $(\kappa^+)^M$ onto $\tau_2(n)$. Let $\varpi^1(n) = \sigma_n \circ h_0$ which is an M -amenable surjection of κ into $\rho_1^1(n)$.
 1333 Lemma 5.1 gives an M -amenable surjection $h_1 : \kappa \rightarrow (\kappa^{++})^M$ which is obtained uniformly in ρ_0^1 , ρ_1^1 , and
 1334 ϖ^1 . Thus h_1 is obtained uniformly in τ_0 , τ_1 , τ_2 , and $\prec$. $\square$

1335 The following lemma uses a closedness condition of a forcing inside of an inner model M of ZFC and the
 1336 existence of an appropriate M -amenable surjection to create generic filters over M which exist in the real
 1337 world. Section 7 will use amenable surjections to create a large set of filters for a type of Namba forcing
 1338 (which is not countably closed) generic over an inner model M by using fusion sequences.

1339 **Lemma 5.5.** *Assume AD. Let M be an inner model of ZFC and $\mathbb{P} \in M$ be a forcing. Let $\mathcal{D}$ denote the
 1340 $\mathbb{P}$ -dense subsets of $\mathbb{P}$ in M . Let $\kappa \leq \delta < \Theta$ be such that $\text{cof}^V(\kappa) = \omega$, $M \models \text{“}\kappa$ is a cardinal, $|\mathcal{D}| \leq \delta$, and
 1341 $\mathbb{P}$ is $< \kappa$ -closed”, and there is an M -amenable surjection of κ onto δ . Then there is a $G \subseteq \mathbb{P}$ so that G is
 1342 $\mathbb{P}$ -generic over M .*

1343 *Proof.* Since $M \models |\mathcal{D}| \leq \delta$, let $f \in M$ be such that $f : \delta \rightarrow \mathcal{D}$ is a surjection. Let $g : \kappa \rightarrow \delta$ be an
 1344 M -amenable surjection given by the hypothesis. Then $h : \kappa \rightarrow \mathcal{D}$ defined by $h = f \circ g$ is an M -amenable
 1345 surjection. Since $M \models \text{ZFC}$, let $\sqsubset \in M$ be such that $M \models \text{“}\sqsubset$ is a wellordering of $\mathbb{P}$ ”. Since for each $\gamma < \kappa$,
 1346 $h \restriction \gamma \in M$ because h is M -amenable and $M \models \text{“}\mathbb{P}$ is $< \kappa$ -closed”, there is a unique sequence $\langle p_\alpha^\gamma : \alpha < \gamma \rangle \in M$
 1347 so that for all $\alpha < \gamma$, p_α^γ is the $\sqsubset$ -least element $p \in \mathbb{P}$ so that for all $\beta < \alpha$, $p \leq_\mathbb{P} p_\beta^\gamma$ and p belongs to the
 1348 dense set $h(\alpha)$. The uniqueness implies the following continuity: for all $\alpha < \gamma_1 \leq \gamma_2 < \kappa$, $p_\alpha^{\gamma_0} = p_\alpha^{\gamma_1}$. For

all $\alpha < \kappa$, let $p_\alpha = p_\alpha^{\alpha+1}$. Let $G = \langle \{p_\alpha : \alpha < \kappa\} \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists \alpha < \kappa) : p_\alpha \leq_{\mathbb{P}} p\}$. Since for all $\alpha < \kappa$, $p_\alpha \in h(\alpha)$ and $h : \kappa \rightarrow \mathcal{D}$ is a surjection, G is a $\mathbb{P}$ -generic filter over M . $\square$

Lemma 5.6. *Assume AD. Let M be an inner model of ZFC. Let ϵ be an infinite ordinal. Suppose $\text{cof}^V((\epsilon^+)^M) = \text{cof}^V((\epsilon^{++})^M) = \text{cof}^V((\epsilon^{+++})^M) = \omega$ and $M \models 2^{\epsilon^+} = \epsilon^{++} \wedge 2^{\epsilon^{++}} = \epsilon^{+++}$. Let $\text{Coll}(\epsilon^+, \epsilon^{++})^M$ be the forcing of partial function $p : \epsilon^+ \rightarrow \epsilon^{++}$ of size less than ϵ^+ ordered by reverse function extension as defined in M . Then there is a $G \subseteq \text{Coll}(\epsilon^+, \epsilon^{++})^M$ which is $\text{Coll}(\epsilon^+, \epsilon^{++})^M$ -generic over M .*

Proof. Since $\text{cof}((\epsilon^+)^M) = \text{cof}((\epsilon^{++})^M) = \text{cof}((\epsilon^{+++})^M) = \omega$, Lemma 5.3 implies that there is an M -amenable surjection $f : (\epsilon^+)^M \rightarrow (\epsilon^{+++})^M$.

Work in M for the moment. For all $\delta < \epsilon^{++}$, $|\epsilon^+ \delta| = 2^{\epsilon^+} = \epsilon^{++}$. $\text{Coll}(\epsilon^+, \epsilon^{++})$ injects into $\epsilon^+ \epsilon^{++}$, the collection of all functions from ϵ^+ into ϵ^{++} . Since ϵ^{++} is regular (as $M \models \text{ZFC}$), $\epsilon^+ \epsilon^{++} = \bigcup_{\delta < \epsilon^{++}} \epsilon^+ \delta$. Then $|\text{Coll}(\epsilon^+, \epsilon^{++})| = \epsilon^{++}$. So $|\mathcal{P}(\text{Coll}(\epsilon^+, \epsilon^{++}))| = 2^{\epsilon^{++}} = \epsilon^{+++}$. Let $\mathcal{D}$ be the collection of dense subsets of $\text{Coll}(\epsilon^+, \epsilon^{++})$. Then $|\mathcal{D}| \leq |\mathcal{P}(\text{Coll}(\epsilon^+, \epsilon^{++}))| = \epsilon^{+++}$.

One has that $\text{cof}^V((\epsilon^+)^M) = \omega$, $M \models "|\mathcal{D}| \leq \epsilon^{+++}"$, and there is an M -amenable surjection of $(\epsilon^+)^M$ onto $(\epsilon^{+++})^M$. The result follows from Lemma 5.5. $\square$

A tree on a set X is a set $T \subseteq {}^{<\omega}X$ which is closed under subsequences. The body of the tree T is $[T] = \{f \in {}^\omega X : (\forall n \in \omega)(f \upharpoonright n \in T)\}$. A set $A \subseteq {}^\omega \omega$ is Suslin if and only if there is a cardinal κ and a tree T on $\omega \times \kappa$ so that $A = \pi_1[[T]] = \{x \in {}^\omega \omega : (\exists f \in {}^\omega \kappa)((x, f) \in [T])\}$. A set $A \subseteq {}^\omega \omega$ is coSuslin if ${}^\omega \omega \setminus A$ is Suslin. If $A = \pi_1[[T]]$ where T is a tree on $\omega \times \kappa$, then one says that A is κ -Suslin. Let $\mathcal{S}_\kappa$ denote the collection of κ -Suslin sets. A cardinal κ is a Suslin cardinal if and only if there is an $A \subseteq {}^\omega \omega$ which is κ -Suslin but not δ -Suslin for any $\delta < \kappa$. Let SN denote the class of all Suslin cardinals. A Suslin cardinal κ is called a limit Suslin cardinal if and only if κ is a limit of Suslin cardinals. A Suslin cardinal which is not a limit Suslin cardinal is called a successor Suslin cardinal. Steel and Woodin showed that SN is a closed subset of $\text{sup}(\text{SN})$ under AD and $\text{DC}_{\mathbb{R}}$. Under AD and $\text{DC}_{\mathbb{R}}$, $\text{sup}(\text{SN})$ is a limit of Suslin cardinals. To see this, suppose for the sake of contradiction that $\lambda = \text{sup}(\text{SN})$ is not a limit of Suslin cardinals. Then λ must be a successor Suslin cardinal. Suppose S_λ is not closed under $\forall^{\mathbb{R}}$. Then $\forall^{\mathbb{R}} S_\lambda$ would have the scale property by the second periodicity theorem of Moschovakis ([45] 6C.3). Thus there is a set $A \in \forall^{\mathbb{R}} S_\lambda \setminus S_\lambda$ which is Suslin. A must be λ' -Suslin for some $\lambda' > \lambda$. Hence λ is not the largest Suslin cardinal which is a contradiction. This shows S_λ must be closed under $\forall^{\mathbb{R}}$. Then λ must be a regular limit Suslin cardinal by [30] Lemma 3.6 which contradicts the assumption that λ is not even a limit of Suslin cardinals. This completes the argument that $\text{sup}(\text{SN})$ is a limit of Suslin cardinals. In fact, $\text{sup}(\text{SN})$ is a limit of measurable cardinals and even measurable successor Suslin cardinals. To see this, let $\kappa < \text{sup}(\text{SN})$. Since $\text{sup}(\text{SN})$ is a limit of Suslin cardinals, let $\langle \kappa_n : n \in \omega \rangle$ be an increasing sequence of Suslin cardinals above κ and let $\kappa_\infty = \text{sup}\{\kappa_n : n \in \omega\}$ which is a limit Suslin cardinal of cofinality ω . Let $\mathcal{S}_{<\kappa} = \bigcup_{\delta < \kappa} \mathcal{S}_\delta$. Let Σ_0 be the pointclass of countable union of sets from $\mathcal{S}_{<\kappa}$. Σ_0 has the scale property with norms in κ (see [30] Theorem 3.28, Case I), closed under countable union, and even closed under $\exists^{\mathbb{R}}$. By the second periodicity theorem of Moschovakis ([45] Theorem 6C.3), $\Pi_1 = \forall^{\mathbb{R}} \Sigma_0$ has the scale property with norm in $\delta(\Pi_1)$. By the Kunen-Martin Theorem, one can show that $\delta(\Pi_1) = \kappa^+$ and κ^+ is the next Suslin cardinal after κ (see [30] Theorem 3.28, Case I). Since Π_1 is closed under $\forall^{\mathbb{R}}$, $\wedge$, $\vee$, and has the prewellordering property (even the scale property), $\kappa^+ = \delta(\Pi_1)$ is measurable and even satisfies $\kappa^+ \rightarrow_* (\kappa^+)_2^\epsilon$ for all $\epsilon < \omega_1$ by [30] Theorem 2.36 (and thus the ω -club filter $\mu_{\kappa^+}^1$ is a normal measure on κ^+). One can even show that $\text{sup}(\text{SN})$ is a supremum of regular limit Suslin cardinals and such cardinals have the strong partition property by [37]. Under AD^+ , if $\text{sup}(\text{SN}) < \Theta$, then $\text{sup}(\text{SN}) \in \text{SN}$, i.e. $\text{sup}(\text{SN})$ is also a Suslin cardinal and SN has a largest element. (See [42] Corollary 11.23 and Remark 11.24.)

Solovay [55] showed that under $\text{AD}_{\mathbb{R}}$, ω_1 is $\mathbb{R}$ -supercompact. If ν is a supercompact measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$ and $\kappa < \Theta$, then an appropriate Rudin-Keisler pushforward of ν will define a supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$. However, AD^+ cannot prove that ω_1 is $\mathbb{R}$ -supercompact since $L(\mathbb{R}) \models "\omega_1 \text{ is not } \mathbb{R}\text{-supercompact}"$ because Solovay [55] showed this hypothesis implies the existence of $\mathbb{R}^\sharp$. Under AD and $\text{DC}_{\mathbb{R}}$, Harrington and Kechris [24] showed that if κ is below a Suslin cardinal, then ω_1 is κ -supercompact. Woodin [66] showed the supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$ is unique if κ is below a Suslin cardinal. (Also see [42] Theorem 13.15.) Uniqueness will be very important as it implies that the supercompact measure ν on $\mathcal{P}_{\omega_1}(\kappa)$ is ordinal definable when κ is below a Suslin cardinal. Under AD^+ , for all $\kappa < \Theta$, Woodin showed that ω_1 is

κ -supercompact and Neeman showed that there is a unique supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$. Except for Theorem 6.27, this result can be avoided by reflecting a counterexample below the largest Suslin cardinal.

Fact 5.7. (Neeman [46] Theorem 2.19; Woodin [61] Lemma 2.19) Assume AD^+ . For any $\kappa < \Theta$, there is a unique supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$.

Fact 5.8. Let X be a set and let ν be a supercompact measure on $\mathcal{P}_{\omega_1}(X)$. Let $j_\nu : V \rightarrow \prod_{\mathcal{P}_{\omega_1}(X)} V/\nu$ be the ultrapower map. Let $A \subseteq X$ and let $\Phi : \mathcal{P}_{\omega_1}(X) \rightarrow \mathcal{P}_{\omega_1}(X)$ be defined by $\Phi(\sigma) = A \cap \sigma$. Then $[\Phi]_\nu = j_\nu[A]$.

Proof. Suppose $\Psi : \mathcal{P}_{\omega_1}(X) \rightarrow V$ and $[\Psi]_\nu \in [\Phi]_\nu$. By definition of the interpretation of the ultrapower membership relation, $B = \{\sigma \in \mathcal{P}_{\omega_1}(X) : \Psi(\sigma) \in \Phi(\sigma)\} \in \nu$. Define $\Upsilon : B \rightarrow \mathcal{P}_{\omega_1}(X)$ by $\Upsilon(\sigma) = \{\Psi(\sigma)\}$. Then for all $\sigma \in B$, $\emptyset \neq \Upsilon(\sigma) \subseteq \Phi(\sigma) \subseteq \sigma$. By normality, there is an $x \in X$ so that $C = \{\sigma \in \mathcal{P}_{\omega_1}(X) : x \in \Upsilon(\sigma)\} \in \nu$. Since for ν -almost all σ , $\Upsilon(\sigma)$ is the singleton $\Psi(\sigma)$, one has that for ν -almost all σ , $\Psi(\sigma) = x$. Thus $[\Psi]_\nu = j_\nu(x)$. Since for ν -almost all $\sigma \in \mathcal{P}_{\omega_1}(X)$, $\Psi(\sigma) \in \Phi(\sigma) = A \cap \sigma$, one has that $x \in A$. This shows that $[\Phi]_\nu \subseteq j_\nu[A]$. For any $x \in A$, $\{\sigma \in \mathcal{P}_{\omega_1}(X) : x \in A \cap \sigma = \Phi(\sigma)\} = \{\sigma \in \mathcal{P}_{\omega_1}(X) : x \in \sigma\} \in \nu$ by fineness. By Łoś' theorem, $j_\nu(x) \in [\Phi]_\nu$. Since $x \in A$ was arbitrary, this shows that $j_\nu[A] \subseteq [\Phi]_\nu$. One can conclude $j_\nu[A] = [\Phi]_\nu$. $\square$

Let $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}}, 1_{\mathbb{P}})$ be a forcing and $F \subseteq \mathbb{P}$ is a $\mathbb{P}$ -filter. F is said to be countably generated if and only if there is a countable $\sigma \subseteq F$ such that $F = \langle \sigma \rangle_{\mathbb{P}} = \{p \in \mathbb{P} : (\exists q \in \sigma)(q \leq_{\mathbb{P}} p)\}$. The next result says in certain circumstances, the truth concerning countably generated filter in the real world can be expressed over a HOD-type model. This result will give the required capturing necessary in the main result.

Lemma 5.9. Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set and $V = L(Z, \mathbb{R})$. Let $\mathbb{P}$ be a forcing belonging to HOD_Z such that there is an ordinal κ with the property that there is a unique supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$ and $\text{HOD}_Z \models |\mathbb{P}| = |\kappa|$. Let F be a countable generated $\mathbb{P}$ -filter. Then $\text{HOD}_Z[F] = \text{HOD}_{Z \cup \{F\}}$.

In particular, under AD and $\text{DC}_{\mathbb{R}}$, the hypothesis on κ holds if κ is below a Suslin cardinal. Under AD^+ , this hypothesis on κ holds for all $\kappa < \Theta$.

Proof. Let Y be a very good parameter-witness for Z . By the hypothesis, let $\tilde{\nu}$ be the unique supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$. (If κ is below a Suslin cardinal, then the results of Harrington-Kechris and Woodin mentioned above imply the existence of such a unique supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$.) Thus $\tilde{\nu} \in \text{OD}$ and hence $\tilde{\nu} \in \text{OD}_Z$. By the hypothesis, let $\varpi : \kappa \rightarrow \mathbb{P}$ be a bijection with $\varpi \in \text{HOD}_Z$. Define a supercompact measure ν on $\mathcal{P}_{\omega_1}(\mathbb{P})$ by $A \in \nu$ if and only if $\{\varpi^{-1}[\sigma] : \sigma \in A\} \in \tilde{\nu}$. Note that $\nu \in \text{OD}_Z$. Since $V = L(Z, \mathbb{R}) = L(Y, \mathbb{R})$ and $\text{DC}_{\mathbb{R}}$ hold, one has that DC holds. Hence $\prod_{\mathcal{P}_{\omega_1}(\mathbb{P})} V/\nu$ is wellfounded. Let $j_\nu : V \rightarrow \prod_{\mathcal{P}_{\omega_1}(\mathbb{P})} V/\nu$ be the ultrapower map. For each $\sigma \in \mathcal{P}_{\omega_1}(\mathbb{P})$, there is uniform procedure that assigns σ to a global wellordering of $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, F \cap \sigma]$. Thus $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{P})} L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, F \cap \sigma]/\nu$ is wellfounded and satisfies Łoś' theorem. Let $\Phi_F : \mathcal{P}_{\omega_1}(\mathbb{P}) \rightarrow \mathcal{P}_{\omega_1}(\mathbb{P})$ be defined by $\Phi_F(\sigma) = \sigma \cap F$. By Fact 5.8, $[\Phi_F]_\nu = j_\nu[F]$. By Łoś' theorem, $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{P})} L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, F \cap \sigma]/\nu = L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), [\Phi_F]_\nu] = L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu[F]]$. For each $\sigma \in \mathcal{P}_{\omega_1}(\kappa)$, let $\text{Coll}(\omega, F \cap \sigma)$ be the forcing of finite partial function from ω into $F \cap \sigma$. Let Φ_{Coll} be defined by $\Phi_{\text{Coll}}(\sigma) = \text{Coll}(\omega, F \cap \sigma)$. By Łoś' theorem, $[\Phi_{\text{Coll}}]_\nu = \text{Coll}(\omega, j_\nu[F])$. For each σ , let $\dot{g}_\sigma$ be the canonical $\text{Coll}(\omega, F \cap \sigma)$ -name for the generic surjection of ω onto $F \cap \sigma$. Let $\Phi_{\dot{g}}$ be defined by $\Phi_{\dot{g}}(\sigma) = \dot{g}_\sigma$. Let $\dot{h} = [\Phi_{\dot{g}}]_\nu$. By Łoś' theorem, $\dot{h}$ is the canonical $\text{Coll}(\omega, j_\nu[F])$ -name for the generic surjection of ω onto $j_\nu[F]$. Thus $\dot{h}$ belongs to any inner model of ZF containing $j_\nu[F]$. By Łoś' theorem applied in $\prod_{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{P})} L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, F \cap \sigma]/\nu$, $j_\nu(\vec{\mathbb{B}}^{\kappa;Z})$ codes a direct system of forcing projection and $j_\nu(\mathbb{B}_n^{\kappa;Z}) \subseteq \mathcal{B}_\infty$ for all $n \in \omega$ (that is, the conditions of the forcing $j_\nu(\mathbb{B}_n^{\kappa;Z})$ are ∞ -Borel codes or equivalently pairs of ordinals and formulas). Recall the operation $\mathcal{G}^{\kappa,1}$ from Definition 4.4.

Claim 1: Let $a \in \text{HOD}_Z$,

$$L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(a), j_\nu[F]] \models 1_{\text{Coll}(\omega, j_\nu[F])} \Vdash$$

$$\mathcal{G}^{j_\nu(\kappa),1}(j_\nu(\mathbb{B}_1^{\kappa;Z}), \dot{h}) \text{ is } j_\nu(\mathbb{B}_1^{\kappa;Z})\text{-generic over } L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(a)]$$

To see Claim 1: By Loś' theorem, it suffices to show that for all $\sigma \in \mathcal{P}_{\omega_1}(\mathbb{P})$,

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma] \models 1_{\text{Coll}(\omega, F \cap \sigma)} \Vdash G^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, \dot{g}_\sigma) \text{ is } \mathbb{B}_1^{\kappa;Z}\text{-generic over } L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a].$$

1442 Fix any $p \in \text{Coll}(\omega, F \cap \sigma)$. Since σ is countable, $\text{Coll}(\omega, F \cap \sigma)$ is a countable forcing in the real world. By Fact
 1443 2.14, there is a $K \subseteq \text{Coll}(\omega, F \cap \sigma)$ with $p \in K$ and is $\text{Coll}(\omega, F \cap \sigma)$ -generic over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a]$. Let $k : \omega \rightarrow$
 1444 $F \cap \sigma$ be the associated generic surjection. Then $k \in {}^1(\omega^\mathbb{P})$ and $\dot{g}_\sigma[K]$. By Fact 4.6, $G_k^{\kappa;Z;1} = \mathcal{G}^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)$
 1445 is $\mathbb{B}_1^{\kappa;Z}$ -generic over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a]$. By absoluteness, $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a][K] \models \mathcal{G}^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)$ is $\mathbb{B}_1^{\kappa;Z}$ -generic
 1446 over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a]$. By the forcing theorem, there is a $q \leq_{\text{Coll}(\omega, F \cap \sigma)} p$ so that $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a] \models q \Vdash$
 1447 $\mathcal{G}^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, \dot{g}_\sigma)$ is $\mathbb{B}_1^{\kappa;Z}$ -generic over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a]$. Since $p \in \text{Coll}(\omega, F \cap \sigma)$ was arbitrary, $1_{\text{Coll}(\omega, F \cap \sigma)}$ forces
 1448 the same statement. This establishes Claim 1.

1449 By Claim 1, in the forcing language, the remainder forcing $j_\nu(\vec{\mathbb{B}}_\omega^{\kappa;Z}) /_{j_\nu(\Pi_{\omega,1}^{\kappa;Z})} \mathcal{G}^{j_\nu(\kappa),1}(j_\nu(\mathbb{B}_1^{\kappa;Z}), \dot{h})$ is well
 1450 defined.

Claim 2: Let $\varphi(v, w)$ be a $\mathcal{L}_{\text{set}}$ -formula and $a \in \text{HOD}_Z$. Let $\tilde{\varphi}$ be the formula defined by $\tilde{\varphi}(x, y, z)$ is “ y is a forcing, x is a function from ω into y , and $\varphi(\langle x[\omega] \rangle_y, z)$ ”. Then

$$\begin{aligned} \varphi(F, a) \Leftrightarrow L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(a), j_\nu[F]] \models 1_{\text{Coll}(\omega, j_\nu[F])} \Vdash L[j_\nu(\vec{\mathbb{B}}^{\kappa;Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(a), \dot{h}] \models \\ 1_{j_\nu(\vec{\mathbb{B}}_\omega^{\kappa;Z}) /_{j_\nu(\Pi_{\omega,1}^{\kappa;Z})} \mathcal{G}^{j_\nu(\kappa),1}(j_\nu(\mathbb{B}_1^{\kappa;Z}), \dot{h})} \Vdash L(j_\nu(Y), j_\nu(\dot{\mathbb{R}}^{\kappa;\text{sym}})) \models \tilde{\varphi}(\dot{h}, j_\nu(\mathbb{P}), j_\nu(a)). \end{aligned}$$

1451 To see Claim 2: Since F is countably generated, there is a countable $\tilde{\sigma} \subseteq F$ so that $\langle \tilde{\sigma} \rangle_{\mathbb{P}} = F$. Since ν is
 1452 fine and countably complete, $A = \{\sigma \in \mathcal{P}_{\omega_1}(\nu) : \tilde{\sigma} \subseteq \sigma\} \in \nu$.

($\Rightarrow$) Suppose $\varphi(F, a)$. Fix $\sigma \in A$. Let $p \in \text{Coll}(\omega, F \cap \sigma)$. Since σ is countable, $\text{Coll}(\omega, F \cap \sigma)$ is a
 countable forcing in the real world. By Fact 2.14, let $K \subseteq \text{Coll}(\omega, F \cap \sigma)$ be $\text{Coll}(\omega, F \cap \sigma)$ -generic over
 $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, F \cap \sigma]$ with $p \in K$. Let $k : \omega \rightarrow F \cap \sigma$ be the associated generic surjection. Note that
 $\langle k[\omega] \rangle_{\mathbb{P}} = \langle F \cap \sigma \rangle_{\mathbb{P}} \supseteq \langle \tilde{\sigma} \rangle_{\mathbb{P}} = F$ since $\tilde{\sigma} \subseteq \sigma$ because $\sigma \in A$. Hence $\langle k[\omega] \rangle_{\mathbb{P}} = F$. Since $L(Z, \mathbb{R}) = L(Y, \mathbb{R})$,
 one has that $L(Y, \mathbb{R}) \models \varphi(\langle k[\omega] \rangle_{\mathbb{P}}, a)$. Thus $L(Y, \mathbb{R}) \models \tilde{\varphi}(k, \mathbb{P}, a)$. Note that $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a] \subseteq \text{HOD}_Z$. Since
 $k \in {}^\omega \mathbb{P}$, Fact 4.17 (applied to $M = L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a]$) gives that

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, k] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}_k^{\kappa;Z;1}} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(k, \mathbb{P}, a).$$

By absoluteness and since $G_k^{\kappa;Z;1} = \mathcal{G}_k^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)$,

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma][K] \models L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, k] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}_k^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(k, \mathbb{P}, a).$$

Since this holds for all $p \in \text{Coll}(\omega, F \cap \sigma)$ and all $\text{Coll}(\omega, F \cap \sigma)$ -generic filter K over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma]$
 with $p \in K$, one has the following by the forcing theorem.

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma] \models 1_{\text{Coll}(\omega, F \cap \sigma)} \Vdash L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, \dot{g}_\sigma] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, \dot{g}_\sigma)} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(\dot{g}_\sigma, \mathbb{P}, a).$$

1453 Since $\sigma \in A \in \nu$ was arbitrary, the right side of the claim follows from Loś' theorem.

($\Leftarrow$) Assume the right side of the claim. By Loś' theorem, there is a $B \in \nu$ such that for all $\sigma \in B$

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma] \models 1_{\text{Coll}(\omega, F \cap \sigma)} \Vdash L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, \dot{g}_\sigma] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, \dot{g}_\sigma)} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(\dot{g}_\sigma, \mathbb{P}, a).$$

Since $A \in \nu$, $A \cap B \in \nu$. Pick any $\sigma \in A \cap B$. Since $\text{Coll}(\omega, F \cap \sigma)$ is countable in the real world, let
 $K \subseteq \text{Coll}(\omega, F \cap \sigma)$ be $\text{Coll}(\omega, F \cap \sigma)$ -generic over $L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma]$ using Fact 2.14. Let k be the generic
 surjection of ω onto $F \cap \sigma$ induced from K and note that $\dot{g}_\sigma[K] = k$. By the forcing theorem, one has the
 following.

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, F \cap \sigma][K] \models L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, k] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}_k^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(k, \mathbb{P}, a).$$

By absoluteness,

$$L[\vec{\mathbb{B}}^{\kappa;Z}, \mathbb{P}, Y, a, k] \models 1_{\vec{\mathbb{B}}_\omega^{\kappa;Z} /_{\Pi_{\omega,n}^{\kappa;Z}} \mathcal{G}_k^{\kappa;1}(\mathbb{B}_1^{\kappa;Z}, k)} \Vdash L(Y, \dot{\mathbb{R}}^{\kappa;\text{sym}}) \models \tilde{\varphi}(k, \mathbb{P}, a).$$

1454 Since $k \in {}^\omega \mathbb{P}$, Fact 4.17 implies $L(Y, \mathbb{R}) \models \tilde{\varphi}(k, \mathbb{P}, a)$. By the definition of $\tilde{\varphi}$, $L(Y, \mathbb{R}) \models \varphi(\langle k[\omega] \rangle_{\mathbb{P}}, a)$.
 1455 Since $\sigma \in A \cap B \subseteq A$ and thus $\tilde{\sigma} \subseteq F \cap \sigma = k[\omega]$, $\langle k[\omega] \rangle_{\mathbb{P}} = F$. So $L(Y, \mathbb{R}) \models \varphi(F, a)$. Finally since
 1456 $V = L(Z, \mathbb{R}) = L(Y, \mathbb{R})$, one has $\varphi(F, a)$. This completes the proof of the Claim 2.

Now to complete the proof of the lemma: It is clear that $\text{HOD}_Z[F] \subseteq \text{HOD}_{Z \cup \{F\}}$. Let $U \in \text{HOD}_{Z \cup \{F\}}$. Let W be the transitive closure of $\{U\}$. Since $\text{HOD}_{Z \cup \{F\}} \models \text{ZFC}$, there is a bijection $\pi : \delta \rightarrow W$ where δ is an ordinal and $\pi \in \text{HOD}_{Z \cup \{F\}}$. Within $\text{HOD}_{Z \cup \{F\}}$, define $R \subseteq \delta \times \delta$ by $R(\alpha, \beta)$ if and only if $\pi(\alpha) \in \pi(\beta)$. By the hypothesis that $\text{OD}_Z = \text{OD}_Y$, one has that $\text{OD}_{Z \cup \{F\}} = \text{OD}_{Y \cup \{F\}}$. There is a finite sequence $\vec{y}$ in Y (and note that all the elements in $\vec{y}$ belong to HOD_Z since $Y \in \text{HOD}_Z = \text{HOD}_Y$), a finite sequence of ordinals $\vec{\xi}$, and a $\mathcal{L}_{\text{set}}$ -formula φ so that $R(\alpha, \beta)$ if and only if $\varphi(F, \vec{y}, \vec{\xi}, \alpha, \beta)$. Note that $j_\nu(\vec{\mathbb{B}}^{\kappa; Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(\vec{y}), j_\nu \upharpoonright \delta \in \text{HOD}_Z$. Since $j_\nu \upharpoonright \mathbb{P} \in \text{HOD}_Z$, $j_\nu[F] \in \text{HOD}_Z[F]$. Also note that $\dot{h}$ belongs to any inner model of ZF containing $j_\nu[F]$. Thus by the above claim and absoluteness, for all $\alpha, \beta \in \delta$, $R(\alpha, \beta)$ if and only if

$$\begin{aligned} \text{HOD}_Z[F] \models L[j_\nu(\vec{\mathbb{B}}^{\kappa; Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(\vec{y}), j_\nu[F]] \models 1_{\text{Coll}(\omega, j_\nu[F])} \Vdash L[j_\nu(\vec{\mathbb{B}}^{\kappa; Z}), j_\nu(\mathbb{P}), j_\nu(Y), j_\nu(\vec{y}), \dot{h}] \models \\ 1_{j_\nu(\vec{\mathbb{B}}^{\kappa; Z}) / j_\nu(\Pi_{\omega, 1}^{\kappa; Z}) \mathcal{G}^{j_\nu(\kappa), 1}(j_\nu(\vec{\mathbb{B}}_1^{\kappa; Z}), \dot{h})} \Vdash L(j_\nu(Y), j_\nu(\dot{\mathbb{R}}^{\kappa; \text{sym}})) \models \tilde{\varphi}(\dot{h}, j_\nu(\mathbb{P}), j_\nu(\vec{y}), j_\nu(\vec{\xi}), j_\nu(\alpha), j_\nu(\beta)). \end{aligned}$$

1457 Since $j_\nu \upharpoonright \delta \in \text{HOD}_Z[F]$, the above formula is obtained uniformly from (α, β) in $\text{HOD}_Z[F]$. By comprehension in $\text{HOD}_Z[F]$, $R \in \text{HOD}_Z[F]$. Thus $W \in \text{HOD}_Z[F]$ since it is the Mostowski collapse of the wellfounded relation (δ, R) . Then $U \in \text{HOD}_Z[F]$ since it is the element of highest rank in W . Since $U \in \text{HOD}_{Z \cup \{F\}}$ was arbitrary, this shows that $\text{HOD}_{Z \cup F} \subseteq \text{HOD}_Z[F]$. This completes the proof of the lemma. $\square$

1461 **Corollary 5.10.** *Assume AD and $\text{DC}_{\mathbb{R}}$. Let Z be a very good parameter set. Let $\delta \leq \kappa < \Theta$ be such that*
 1462 *κ is below a Suslin cardinal, $\text{HOD}_Z \models \text{"}\delta \text{ is a cardinal"}$, and $\text{cof}^V(\delta) = \omega$. Suppose $f : \delta \rightarrow \kappa$ is a function*
 1463 *with the property that for all $\alpha < \delta$, $f \upharpoonright \alpha \in \text{HOD}_Z$. Then $\text{HOD}_Z[f] = \text{HOD}_{Z \cup \{f\}}$.*

1464 *Proof.* Let $\mathbb{P} = \text{Coll}(\delta, \kappa)^{\text{HOD}_Z}$ be the forcing of partial functions from δ into κ of cardinality less than δ .
 1465 Since κ is less than a Suslin cardinal and the supremum of the Suslin cardinals is a limit of measurable
 1466 Suslin cardinals, κ is below a measurable Suslin cardinal λ . Since λ is measurable, there is no injection
 1467 of λ into $\mathcal{P}(\kappa)$ by Fact 2.3. Because $\mathbb{P}$ may be regarded as a wellorderable subset of $\mathcal{P}(\kappa)$, one must have
 1468 $|\mathbb{P}|^{\text{HOD}_Z} < |\lambda|$. By the hypothesis that for all $\alpha < \delta$, $f \upharpoonright \alpha \in \text{HOD}_Z$, one has that $F = \langle \{f \upharpoonright \alpha : \alpha < \delta\} \rangle_{\mathbb{P}}$
 1469 is a $\mathbb{P}$ -filter, $\text{HOD}_Z[F] = \text{HOD}_Z[f]$, and $\text{HOD}_{Z \cup \{F\}} = \text{HOD}_{Z \cup \{f\}}$. By the hypothesis that $\text{cof}^V(\delta) = \omega$, let
 1470 $\rho : \omega \rightarrow \delta$ be an increasing cofinal function. Let $\sigma = \{f \upharpoonright \rho(n) : n \in \omega\}$. Then $\langle \sigma \rangle_{\mathbb{P}} = F$. This shows that F is
 1471 a countable generated $\mathbb{P}$ -filter. Thus by Lemma 5.9, $\text{HOD}_Z[f] = \text{HOD}_Z[F] = \text{HOD}_{Z \cup \{F\}} = \text{HOD}_{Z \cup \{f\}}$. $\square$

1472 6. CARDINALITY OF THE SET OF BOUNDED SUBSETS OF A CARDINAL

1473 This section will prove the eponymous main result of this paper.

1474 Suppose $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon \text{ON}$ where $\epsilon < \kappa$ is an injection. To follow the template of Fact 2.25 to prove the
 1475 main result, it remains to find a HOD-type model M for which Lemma 5.6 can be used to produce generics
 1476 for $\text{Coll}(\epsilon^+, \epsilon^{++})^M$ in the real world and the HOD-type model M codes Φ in some way so that Lemma 5.9
 1477 can be used to capture the function in the $\text{Coll}(\epsilon^+, \epsilon^{++})^M$ -generic extension. The model and its requisite
 1478 properties come from the direct system analysis of HOD-type inner models of certain AD^+ models. Woodin
 1479 and Steel used this analysis of HOD-type models to show boldface GCH below Θ and the existence and
 1480 uniqueness of the supercompact measure on $\mathcal{P}_{\omega_1}(\kappa)$ for $\kappa < \Theta$ under AD^+ which were mentioned earlier.

1481 First, the main result will be proved for $L(\mathbb{R})$ under AD which has the most familiar HOD-analysis.
 1482 Finally, the main result under AD^+ will be given which uses the theory of Γ -Woodin mice, their iteration
 1483 strategy, and hybrid strategy mouse built over these Γ -Woodin mice.

1484 For $L(\mathbb{R}) \models \text{AD}$, the necessary HOD-type model comes from the original analysis of $\text{HOD}^{L(\mathbb{R})}$. δ_1^2 is
 1485 the supremum of the Δ_1^2 prewellorderings on $\mathbb{R}$. By [40] Theorem 2.28, $(\delta_1^2)^{L(\mathbb{R})}$ is also the least Σ_1 -stable
 1486 ordinal of $L(\mathbb{R})$ (that is, the least ordinal δ such that $L_\delta(\mathbb{R}) \prec_1 L(\mathbb{R})$ in the language of set theory with
 1487 constant symbols for reals and the set of reals). By the analysis of scales in $L(\mathbb{R})$ ([43], [57]), $(\delta_1^2)^{L(\mathbb{R})}$ is also
 1488 the largest Suslin cardinal of $L(\mathbb{R})$. The following will be the relevant properties.

1489 **Fact 6.1.** (Steel, Woodin; [58], [64]) *Assume AD. Let $x \in \mathbb{R}$.*

(1) $\text{HOD}_{\{x\}}^{L(\mathbb{R})} \models \text{GCH}$.

(2) For all $\kappa < \Theta^{L(\mathbb{R})}$, $\text{cof}^{L(\mathbb{R})}((\kappa^+)^{\text{HOD}_{\{x\}}^{L(\mathbb{R})}}) = \omega$.

Only the simpler analysis of $\text{HOD}_{\{x\}}^{L(\mathbb{R})} \cap V_{(\delta_1^2)^{L(\mathbb{R})}}$ is necessary for the first statement of Fact 6.1 (see remarks following the proof of [60] Corollary 8.22). This simpler analysis is also sufficient to prove the second statement of Fact 6.1 for $\kappa < (\delta_1^2)^{L(\mathbb{R})}$ (see remarks following [60] Lemma 8.25) which will be enough for the proof of Theorem 6.2. These are proved by showing that a portion of $\text{HOD}_{\{x\}}^{L(\mathbb{R})}$ is a limit of a directed system of mice with certain suitable properties. Moreover, if one assumes $M_1(x)^\sharp$ exists, then the proof can be even further simplified by using an external directed system of mice as in [60]. The following is the main theorem in $L(\mathbb{R})$.

Theorem 6.2. Assume AD. $L(\mathbb{R}) \models$ “For all cardinals κ with $\omega < \kappa < \Theta$ and any $\epsilon < \kappa$, there is no injection of $\mathcal{P}_B(\kappa)$ into ${}^\epsilon\text{ON}$ ”.

Proof. By Kechris [35], if AD holds, then $L(\mathbb{R}) \models \text{DC}_{\mathbb{R}}$. For the rest of the proof, work within $L(\mathbb{R})$ or assume $V = L(\mathbb{R})$. Assume the theorem is false. Since $\kappa \geq \omega_1$, $|\mathbb{R}| \leq |\mathcal{P}_B(\kappa)|$. If $\epsilon < \omega$, then ${}^\epsilon\text{ON}$ has a definable bijection with ON. An injection $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ when $\epsilon < \omega$ would imply there is an injection of $\mathbb{R}$ into ON. This is impossible since AD implies there are no wellorderings of $\mathbb{R}$. Thus any failure of the theorem must take the form $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ where ϵ is infinite. Let ϕ be the $\mathcal{L}_{\text{set}}$ sentence asserting “there exists $\epsilon < \kappa < \Theta$ and injection $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ ”. Let ZF^* denote some sufficiently large finite fragment of ZF. Let φ be the formula in the language $\mathcal{L}_{\text{set}}$ augmented with a symbol for $\mathbb{R}$ asserting “there exist an ordinal α such that $L_\alpha(\mathbb{R}) \models \text{ZF}^*$, AD, and ϕ ”. Note that φ is Σ_1 in this augmented language. Note that $L(\mathbb{R}) \models \phi$. By reflection applied to the hierarchy $\langle L_\alpha(\mathbb{R}) : \alpha \in \text{ON} \rangle$, there is an $\alpha \in \text{ON}$ so that $L_\alpha(\mathbb{R}) \models \text{ZF}^*$, AD, and ϕ . So $L(\mathbb{R}) \models \varphi$. Since $L_{\delta_1^2}(\mathbb{R}) \prec_1 L(\mathbb{R})$, $L_{\delta_1^2}(\mathbb{R}) \models \varphi$. Thus there is an $\alpha < \delta_1^2$, $L_\alpha(\mathbb{R}) \models \phi$. So there are $\epsilon < \kappa < \Theta^{L_\alpha(\mathbb{R})} < \delta_1^2$ and a function $\Phi \in L_\alpha(\mathbb{R})$ such that $L_\alpha(\mathbb{R}) \models$ “ $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ is an injection”. Since $\kappa < \Theta^{L_\alpha(\mathbb{R})}$, let $\psi : \mathbb{R} \rightarrow \kappa$ be a surjection in $L_\alpha(\mathbb{R})$. Let $\preceq \in L_\alpha(\mathbb{R})$ be the prewellordering on $\mathbb{R}$ of length κ associated to ψ defined by $x \preceq y$ if and only if $\psi(x) \leq \psi(y)$. Let Γ be a pointclass in $L_\alpha(\mathbb{R})$ closed under $\exists^{\mathbb{R}}$ and $\wedge$ with $\preceq \in \Gamma$. (For instance, $\Gamma = \Sigma_1^1(\preceq)$.) Let $U \subseteq \mathbb{R}^3$ be universal for Γ and note that $U \in L_\alpha(\mathbb{R})$. Let $\vec{0}$ and $\vec{1}$ be the constant 0 and 1 function, respectively. For each $e \in \mathbb{R}$, let $A_e = \{\alpha \in \kappa : (\exists x)(\psi(x) = \alpha \wedge U(e, x, \vec{1}))\}$. For each $A \in \mathcal{P}(\kappa)$, define $T_A \subseteq \mathbb{R} \times \mathbb{R}$ by $(x, y) \in T_A$ if and only if $\psi(x) \in A \Rightarrow y = \vec{1}$ and $\psi(x) \notin A \Rightarrow y = \vec{0}$. By the Moschovakis coding lemma (Fact 2.15), there is a $S \in \Gamma$ so that $S \subseteq T_A$ and for all $\alpha < \kappa$, there are $x, y \in \mathbb{R}$ such that $\psi(x) = \alpha$ and $S(x, y)$. Since U is universal, there is an $e \in \mathbb{R}$ so that $S = U_e = \{(x, y) : U(e, x, y)\}$. Then $A = A_e$. It has been shown that for all $A \in \mathcal{P}(\kappa)$, there is an $e \in \mathbb{R}$ such that $A = A_e$. Since $\psi, \preceq, U \in L_\alpha(\mathbb{R})$, the absoluteness of the map $e \mapsto A_e$ implies that $\mathcal{P}(\kappa)^{L_\alpha(\mathbb{R})} = \mathcal{P}(\kappa)^{L(\mathbb{R})}$. Thus $\mathcal{P}_B(\kappa)^{L_\alpha(\mathbb{R})} = \mathcal{P}_B(\kappa)^{L(\mathbb{R})}$. By absoluteness, $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\text{ON}$ is an injection in $L(\mathbb{R})$. All sets in $L(\mathbb{R})$ are ordinal definable from a real. Let $x \in \mathbb{R}$ be such that $\Phi \in \text{OD}_x$. $\text{HOD}_{\{x\}} \models \text{GCH}$ and $\text{cof}^{L(\mathbb{R})}((\epsilon^+)^{\text{HOD}_{\{x\}}}) = \text{cof}^{L(\mathbb{R})}((\epsilon^{++})^{\text{HOD}_{\{x\}}}) = \text{cof}^{L(\mathbb{R})}((\epsilon^{+++})^{\text{HOD}_{\{x\}}}) = \omega$ by Fact 6.1. Thus there is a $G \subseteq \text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ (in the real world) which is $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ -generic over $\text{HOD}_{\{x\}}$ by Lemma 5.6. Let $g : (\epsilon^+)^{\text{HOD}_{\{x\}}} \rightarrow (\epsilon^{++})^{\text{HOD}_{\{x\}}}$ be the generic surjection, namely $g = \bigcup G$. Since $\text{cof}^{L(\mathbb{R})}((\epsilon^+)^{\text{HOD}_{\{x\}}}) = \omega$, let $\rho : \omega \rightarrow (\epsilon^+)^{\text{HOD}_{\{x\}}}$ be an increasing cofinal map. Let $\sigma = \{g \upharpoonright \rho(n) : n \in \omega\}$ and note that $\langle \sigma \rangle_{\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}} = G$. Thus G is a countably generated $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ filter. Also $|(\epsilon^{++})^{\text{HOD}_{\{x\}}}|$ is below a Suslin cardinal (for example, δ_1^2) and $\text{HOD}_{\{x\}} \models |\text{Coll}(\epsilon^+, \epsilon^{++})| = \epsilon^{++}$. Thus by Lemma 5.9, $\text{HOD}_{\{x\}}[G] = \text{HOD}_{\{x, G\}}$. Let $\tilde{g} = \{\langle \alpha, g(\alpha) \rangle : \alpha < (\epsilon^+)^{\text{HOD}_{\{x\}}}\}$ (where $\langle \cdot, \cdot \rangle$ refers to the Gödel pairing function). Note that $\tilde{g} \subseteq (\epsilon^{++})^{\text{HOD}_{\{x\}}}$ and thus $\tilde{g} \in \mathcal{P}_B(\kappa)$ since $(\epsilon^{++})^{\text{HOD}_{\{x\}}} < \kappa$ by Fact 2.18. Since Φ is $\text{OD}_{\{x\}}$ and $\tilde{g} \in \text{OD}_{\{x, G\}}$, one has that $\Phi(\tilde{g}) \in \text{OD}_{\{x, G\}}$. Therefore, $\Phi(\tilde{g}) \in \text{HOD}_{\{x, G\}} = \text{HOD}_{\{x\}}[G]$. Since $\Phi(\tilde{g}) \in {}^\epsilon\text{ON} \cap \text{HOD}_{\{x\}}[G]$ and $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ add no new functions from ϵ into $\text{HOD}_{\{x\}}$ since $\text{HOD}_{\{x\}} \models$ “ $\text{Coll}(\epsilon^+, \epsilon^{++})$ is $< \epsilon^+$ -closed”, one has that $\Phi(\tilde{g}) \in \text{HOD}_{\{x\}}$. Since $\Phi(\tilde{g}) \in \text{OD}_x$, $\Phi^{-1} \in \text{OD}_{\{x\}}$, Φ is an injection, and $\tilde{g} = \Phi^{-1}(\Phi(\tilde{g}))$, one has that $\tilde{g} \in \text{OD}_{\{x\}}$ and hence $\tilde{g} \in \text{HOD}_{\{x\}}$. By applying the inverse of the Gödel pairing function, $g \in \text{HOD}_{\{x\}}$ and thus $G \in \text{HOD}_{\{x\}}$. This is impossible since G is a $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ -generic over $\text{HOD}_{\{x\}}$ or alternatively by absoluteness, $\text{HOD}_{\{x\}} \models$ “ $g : \epsilon^+ \rightarrow \epsilon^{++}$ is a surjection”. $\square$

Theorem 6.3. Assume AD. Within $L(\mathbb{R})$, the following holds: Let $\kappa < \Theta = \Theta^{L(\mathbb{R})}$ be a singular cardinal and let $\epsilon < \kappa$ be such that $\text{cof}(\epsilon) = \text{cof}(\kappa)$. Then

- (1) $|\mathcal{P}_B(\kappa)| < |\kappa|^\epsilon \cup \mathcal{P}_B(\kappa)|$.
- (2) $|\kappa|^\epsilon < |\kappa|^\epsilon \cup \mathcal{P}_B(\kappa)|$.
- (3) $|\kappa|^\epsilon \cup \mathcal{P}_B(\kappa) < |\mathcal{P}(\kappa)|$.

Proof. (1) If $\Phi : [\kappa]^\epsilon \cup \mathcal{P}_B(\kappa) \rightarrow \mathcal{P}_B(\kappa)$ is an injection, then $\Phi \upharpoonright [\kappa]^\epsilon : [\kappa]^\epsilon \rightarrow \mathcal{P}_B(\kappa)$ is an injection. This violates Theorem 2.2 which applies since boldface GCH below $\Theta^{L(\mathbb{R})}$ holds by Fact 2.22 and $L(\mathbb{R}) \models \text{AD}^+$ or more explicitly by [60] Theorem 8.26.

(2) If $\Phi : [\kappa]^\epsilon \cup \mathcal{P}_B(\kappa) \rightarrow [\kappa]^\epsilon$ is an injection, then $\Phi \upharpoonright \mathcal{P}_B(\kappa) : \mathcal{P}_B(\kappa) \rightarrow [\kappa]^\epsilon$ is an injection. This violates Theorem 6.2 since $\epsilon < \kappa$.

(3) Assume that statement (3) is false. By a reflection argument as in the beginning of the proof of Theorem 6.2, there is a singular cardinal $\kappa < \delta_1^2$ and an injection $\Phi : \mathcal{P}(\kappa) \rightarrow [\kappa]^\epsilon \cup \mathcal{P}_B(\kappa)$. Since $V = L(\mathbb{R})$, there is an $x \in \mathbb{R}$ so that Φ is $\text{OD}_{\{x\}}$. By Fact 6.1, $\text{HOD}_{\{x\}} \models \text{GCH}$ and $\text{cof}((\epsilon^+)^{\text{HOD}_{\{x\}}}) = \text{cof}((\epsilon^{++})^{\text{HOD}_{\{x\}}}) = \text{cof}((\epsilon^{++})^{\text{HOD}_{\{x\}}}) = \omega$. By Lemma 5.6, let $G \subseteq \text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ (which belongs to $L(\mathbb{R})$) be $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ -generic over $\text{HOD}_{\{x\}}$. Since $\text{cof}^{L(\mathbb{R})}((\epsilon^+)^{\text{HOD}_{\{x\}}}) = \omega$, G is a countably generated $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ filter. Since $\text{HOD}_{\{x\}} \models |\text{Coll}(\epsilon^+, \epsilon^{++})| = |\epsilon^{++}|$ and $(\epsilon^{++})^{\text{HOD}_{\{x\}}}$ is less than a Suslin cardinal, $\text{HOD}_{\{x\}}[G] = \text{HOD}_{\{x, G\}}$ by Lemma 5.9. Let $g : (\epsilon^+)^{\text{HOD}_{\{x\}}} \rightarrow (\epsilon^{++})^{\text{HOD}_{\{x\}}}$ be the generic surjection associated to G . Let $\tilde{g} = \{ \langle \alpha, g(\alpha) \rangle : \alpha < (\epsilon^+)^{\text{HOD}_{\{x\}}} \}$ where $\langle \cdot, \cdot \rangle$ is the Gödel pairing function. So $\tilde{g} \subseteq (\epsilon^{++})^{\text{HOD}_{\{x\}}}$, $\tilde{g} \in \mathcal{P}_B(\kappa)$, and $\tilde{g} \in \text{HOD}_{\{x\}}[G]$. In $\text{HOD}_{\{x\}}[G]$, define $A = \{ \tilde{g} \cup K : K \in \mathcal{P}(\kappa) \wedge \min(K) > (\epsilon^{++})^{\text{HOD}_{\{x\}}} \}$. Note that $\text{HOD}_{\{x\}}[G] \models |A| = |\mathcal{P}(\kappa)|$. Let $\tilde{\Phi} = \Phi \upharpoonright \text{HOD}_{\{x\}}[G]$. Since $\tilde{\Phi} \in \text{OD}_{\{x, G\}}$, one has that $\tilde{\Phi} \in \text{HOD}_{\{x, G\}} = \text{HOD}_{\{x\}}[G]$. $\text{HOD}_{\{x\}}[G] \models “\tilde{\Phi} : \mathcal{P}(\kappa) \rightarrow [\kappa]^\epsilon \cup \mathcal{P}_B(\kappa)”$ by absoluteness. Since $\text{HOD}_{\{x\}}[G] \models \text{AC}$ and boldface GCH holds below κ in $L(\mathbb{R})$, $\text{HOD}_{\{x\}}[G] \models |\mathcal{P}(\lambda)| < \kappa$ for all $\lambda < \kappa$. Thus $\text{HOD}_{\{x\}}[G] \models |\mathcal{P}_B(\kappa)| = |\bigcup_{\lambda < \kappa} \mathcal{P}(\lambda)| = \kappa$. Now one cannot have that $\tilde{\Phi}[A] \subseteq \mathcal{P}_B(\kappa)$ since then $\text{HOD}_{\{x\}}[G] \models \tilde{\Phi} \upharpoonright A : A \rightarrow \mathcal{P}_B(\kappa)$ is an injection”. This is impossible since $\text{HOD}_{\{x\}}[G] \models |A| = |\mathcal{P}(\kappa)| = 2^\kappa > \kappa = |\mathcal{P}_B(\kappa)|$. This implies that $\tilde{\Phi}[A] \setminus \mathcal{P}_B(\kappa) \neq \emptyset$. Let $\ell \in \tilde{\Phi}[A] \setminus \mathcal{P}_B(\kappa)$ and hence $\ell \in [\kappa]^\epsilon$. There is some $K \in \mathcal{P}(\kappa)^{\text{HOD}_{\{x\}}[G]}$ such that $\tilde{g} \cup K \in A$ and $\tilde{\Phi}(\tilde{g} \cup K) = \ell$. Since $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_{\{x\}}}$ does not add any new functions from ϵ into $\text{HOD}_{\{x\}}$ because $\text{HOD}_{\{x\}} \models “\text{Coll}(\epsilon^+, \epsilon^{++})$ is $< \epsilon^+$ closed, $\ell \in \text{HOD}_{\{x\}}$. Then $\tilde{g} \cup K = \Phi^{-1}(\Phi(\tilde{g} \cup K)) = \Phi^{-1}(\ell) = \tilde{\Phi}^{-1}(\ell) \in \text{HOD}_{\{x\}}$. So $\tilde{g} \in \text{HOD}_{\{x\}}$. Then $g \in \text{HOD}_{\{x\}}$ which is impossible since then $\text{HOD}_{\{x\}} \models “g : \epsilon^+ \rightarrow \epsilon^{++}$ is a surjection” by absoluteness. (See the comments after Theorem 6.15 concerning this proof and the conjecture of whether $\mathcal{P}(\kappa)$ has 2-regular cardinality.) $\square$

The first step in the proof of Theorem 6.2 for $L(\mathbb{R})$ used the fact that $(\delta_1^2)^{L(\mathbb{R})}$ is a Σ_1 -stable ordinal to reflect to counterexamples below δ_1^2 which is the largest Suslin cardinal of $L(\mathbb{R})$ by the scale analysis. The analogous reflection property will be given by Woodin’s Σ_1 -reflection into Suslin-coSuslin sets under AD^+ (and this reflection principle is essentially equivalent to AD^+)¹⁵.

Definition 6.4. Define $\mathcal{K}$ to be the collection of all transitive set A such that there are Suslin and coSuslin sets $E \subseteq \mathbb{R}^2$ and $F \subseteq \mathbb{R}^2$ with the following properties.

- (1) E is an equivalence relation on $\mathbb{R}$. F is a binary relation.
- (2) Define $\dot{\in}_F$ on $\mathbb{R}/E$ by $[x]_E \dot{\in}_F [y]_E$ if and only if $F(x, y)$. $(\mathbb{R}/E, \dot{\in}_F)$ is $\mathcal{L}_{\text{set}}$ isomorphic to $(A, \in)$ or in other words, $(A, \in)$ is the Mostowski collapse of $(\mathbb{R}/E, \dot{\in}_F)$.

Let $\mathcal{S} = \bigcup \mathcal{K}$ which will be called the Suslin and coSuslin sets.

Fact 6.5. (Woodin; [56]) Assume AD^+ and $V = L(\mathcal{P}(\mathbb{R}))$. $(\mathcal{S}, \in) \prec_{\Sigma_1} (V, \in)$, that is, $\mathcal{S}$ is a Σ_1 -elementary substructure of the true universe V in the language $\mathcal{L}_{\text{set}}$.

If $X, Y \in \mathcal{P}(\mathbb{R})$, then X is Wadge reducible to Y (denoted $X \leq_W Y$) if and only if there is a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ so that for all $r \in \mathbb{R}$, $r \in X$ if and only if $f(r) \in Y$. Under AD, the Wadge lemma ([42] Theorem 2.4) states that for any $X, Y \in \mathcal{P}(\mathbb{R})$, either $X \leq_W Y$ or $Y \leq_W (\mathbb{R} \setminus X)$. Martin and Monk ([42]

¹⁵The proof of the main theorem never explicitly uses the ordinal determinacy clause of AD^+ or that any set of reals has ∞ -Borels. The only genuine use of AD^+ beyond AD and $\text{DC}_{\mathbb{R}}$ will be at the beginning of the proof by contradiction where the Σ_1 -reflection into Suslin-coSuslin (Fact 6.5) is used to find a Suslin and coSuslin counterexample.

Theorem 2.9) showed that under AD and $\text{DC}_{\mathbb{R}}$, $\leq_W$ is a wellfounded relation. For each $X \in \mathcal{P}(\mathbb{R})$, $\text{rk}_W(X)$ will denote the Wadge rank of X , that is, the rank of X in $\leq_W$. The length of $\leq_W$ is Θ . For each $\alpha \leq \Theta$, let $\mathcal{W}_\alpha = \{B \in \mathcal{P}(\mathbb{R}) : \text{rk}_W(B) < \alpha\}$ and note that $\mathcal{W}_\Theta = \mathcal{P}(\mathbb{R})$.

A (lightface) pointclass Γ is a good pointclass ([62] Definition 3.1) if and only if Γ is ω -parametrized, closed under recursive substitution, $\exists^\omega$, $\forall^\omega$, $\exists^\mathbb{R}$, the category quantifiers¹⁶, and has the scale property.

Next, a very terse sketch will be given to show that for any Suslin and coSuslin set B , there is a lightface analytical hierarchy of good pointclasses containing B . Let κ be the least Suslin cardinal such that B is κ -Suslin. Since B is Suslin and coSuslin, κ is not the largest Suslin cardinal. Let $\tilde{B} \in \mathcal{P}(\mathbb{R})$ be a Suslin set which is not κ -Suslin. Note that $B \leq_W \tilde{B}$ by the Wadge lemma. B is Suslin and coSuslin in $L(\tilde{B}, \mathbb{R})$ (by using the coding lemma to code the trees). In $L(\tilde{B}, \mathbb{R})$, all sets are ordinal definable from $\tilde{B}$ and a real. Let $R \subseteq \mathbb{R} \times \mathbb{R}$ be defined by $R(x, y)$ if and only if $y \notin \text{OD}_{\{\tilde{B}, x\}}$. R does not have a uniformization in $L(\tilde{B}, \mathbb{R})$ by the Kechris and Solovay argument as described earlier. Since $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$ (or Uniformization) fails, $\sup(\text{SN}^{L(\tilde{B}, \mathbb{R})})$, the supremum of the Suslin cardinals of $L(\tilde{B}, \mathbb{R})$, is below $\Theta^{L(\tilde{B}, \mathbb{R})}$. It was shown earlier that $\sup(\text{SN}^{L(\tilde{B}, \mathbb{R})})$ is a limit of Suslin cardinals. Let λ be the supremum of the first ω many Suslin cardinals of $L(\tilde{B}, \mathbb{R})$ above κ and note that $\lambda \leq \sup(\text{SN}^{L(\tilde{B}, \mathbb{R})}) < \Theta^{L(\tilde{B}, \mathbb{R})}$. Since the supremum of a countable set of Suslin cardinals is a Suslin cardinal of $L(\tilde{B}, \mathbb{R})$ if the supremum is below $\Theta^{L(\tilde{B}, \mathbb{R})}$, λ is a Suslin cardinal of cofinality ω . Let $S_{<\lambda}$ be the collection of sets which are α -Suslin for some $\alpha < \lambda$ which forms a pointclass closed under Wadge reduction. Let Λ_0 be the countable union of sets in $S_{<\lambda}$. Λ_0 is closed under Wadge reductions, $\exists^\mathbb{R}$, $\exists^\omega$, has the scale property, but is not closed under $\forall^\omega$. By the second periodicity theorem ([45] Theorem 6C.3), $\Lambda_1 = \forall^\mathbb{R} \Lambda_0$ has the scale property, closed under Wadge reduction, $\forall^\mathbb{R}$, $\exists^\omega$, $\forall^\omega$, but is not closed under $\exists^\mathbb{R}$. By [45] Lemma 6C.1, $\Lambda_2 = \exists^\mathbb{R} \Lambda_1$ has the scale property, is closed under Wadge reduction, $\exists^\mathbb{R}$, $\forall^\omega$, $\exists^\omega$, but is not closed under $\forall^\mathbb{R}$. (See [30] Theorem 3.28 for more details.) Let U be a universal Λ_2 set. Let $\bar{\varphi} = \langle \varphi_n : n \in \omega \rangle$ be a Λ_2 -scale on U . Let $\prec^*$, $\preceq^* \in \Lambda_2$ be the relations witnessing that $\bar{\varphi}$ is a Λ_2 -scale. Pick $x_0, x_1, x_2 \in \mathbb{R}$ such that $U_{x_0} = B$, $U_{x_1} = \prec^*$, and $U_{x_2} = \preceq^*$. Let $\Gamma_{-1} = \text{pos}\Sigma_1^1(U, x_0, x_1, x_2)$. Γ_{-1} is ω -parametrized, has the scale property, closed under $\exists^\mathbb{R}$, $\forall^\omega$, $\exists^\omega$, recursive substitution, contains B , and is not closed under $\forall^\mathbb{R}$. By recursion, define for $n \in \omega$, $\Gamma_n = \exists^\mathbb{R} \forall^\mathbb{R} \Gamma_{n-1}$. For $n \in \omega$, Γ_n is closed under $\exists^\mathbb{R}$, $\forall^\omega$, $\exists^\omega$, recursive substitution, has the scale property by [45] Lemma 6C.1 and the second periodicity theorem, contains B , is not closed under $\forall^\mathbb{R}$, and closed under category quantifiers (which can be seen by the fact that $\Gamma_n = \exists^\mathbb{R} \forall^\mathbb{R} \Gamma_{n-1}$ and that comeagerness can be expressed by the existence of a winning strategy for a certain player in an unfolded Banach-Mazur game).

Let M be a countable transitive model of ZFC and Σ be an (ω, ω_1) -iteration strategy. The pair (M, Σ) Suslin-coSuslin captures a set A at δ via (T, U) if and only if

- (1) $M \models \delta$ is a Woodin cardinal.
- (2) $(T, U) \in M$ are absolutely complementing trees for the forcing $\text{Coll}(\omega, \delta)$.
- (3) For all $x \in \mathbb{R}$, $x \in A$ if and only if there is a countable iteration tree $\mathcal{T}$ on N according to Σ with last model P and iteration map $i : N \rightarrow P$ such that $x \in \pi_1[i(T)]$ (in the real world) where π_1 is the projection onto the first coordinate.

(See [59] Definition 10.1.) The construction of such objects are closely related to Woodin's construction of a coarse Γ -Woodin mouse (see [62] Definition 3.11 or [63] Definition 7.2.3) while controlling the complexity of its iteration strategy which is found in [62] Lemma 3.13.

Fix a Suslin and coSuslin set A . The following sketches the construction of a pair (M, Σ) which Suslin-coSuslin captures A from [62] Theorem 3.13. Let $\Gamma_0, \Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4$ be the first five pointclasses starting an analytical hierarchy of good pointclasses containing A as described above which are not closed under $\forall^\mathbb{R}$. Let T_4 be a tree projecting onto a complete Γ_4 set. Woodin ([40] Theorem 5.40) showed that there is some $z \in \mathbb{R}$ so that $\text{HOD}_{T_4}^{L[T_4, z]} \models \text{"}\omega_2^{L[T, z]}\text{"}$ is a Woodin cardinal". $V_{\omega_2^{L[T_4, z]}}^{\text{HOD}_{T_4}^{L[T_4, z]}}$ is used to create a coarse Γ_4 -Woodin premouse P . Let K_1 denote the universal Γ_1 set. Let $(\bar{T}_1, U_1)$ be trees projecting to the universal Γ_1 set K_1 and its complement which is constructible from T_4 . There is some η and model P such that V_η^P can be used to create a Γ_0 -Woodin premouse. Let w be a wellordering of V_η^P in P and

¹⁶The following is an example of a category quantifier: if $B \subseteq \mathbb{R} \times \mathbb{R}$ belongs to Γ , then $B^* = \{x \in \mathbb{R} : B_x \text{ is comeager}\}$ belongs to Γ .

let $\pi : M \rightarrow L_\alpha(V_\eta^P \cup \{w, T_1, U_1\})$ be elementary for some appropriate α and M countable and transitive. By elementarity, there are $J, \bar{w}, T, U \in M$ such that $\pi(J) = V_\eta^P$, $\pi(\bar{w}) = w$, $\pi(T) = T_1$, $\pi(U) = U_1$ and $M = L_{\bar{\alpha}}(J \cup \{\bar{w}, T, U\})$ for some ordinal $\bar{\alpha}$. Note that M has a definable wellordering of itself using $\bar{w}$ since $L_\alpha(V_\eta^P \cup \{w, T_1, U_1\})$ does using w . M is a Γ_0 -Woodin mouse with strategy defined by choosing the unique π -realizable branch. (M, Σ) will Suslin-coSuslin capture the universal Γ_1 set. (See [62] Lemma 3.13 for more details.) Moreover, the strategy Σ is determined by the C_{Γ_1} -operator¹⁷ in the sense that if b is a cofinal branch π -realizable branch on a countable tree $\mathcal{T}$ using extenders from V_η^M , then b is the unique cofinal branch such that $C_{\Gamma_1}(V_{\delta(\mathcal{T})}^{\mathcal{M}_b^T}) \subseteq \mathcal{M}_b^T$ (in the notation of [62] Lemma 3.13). Also see [63] Lemma 7.2.7. This will be abbreviated by saying “ Σ is the strategy on M guided by C_{Γ_1} ”. Since the strategy Σ picks π -realizable branches, there is embedding of the last model of any iteration tree according to Σ back into $L_\alpha(V_\eta^P \cup \{w, T_1, U_1\})$ which maps the image of (T, U) under the ultrapower map of the iteration tree back into (T_1, U_1) . This shows that (M, Σ) Suslin-CoSuslin captures the universal Γ_1 -set.

As noted in [59] Theorem 10.2 and the proof of [59] Theorem 10.3, (M, Σ) can be constructed to have condensation and the Dodd-Jensen property. There seems to be a few ways to do this which will be briefly described: (M, Σ) is a coarse structure and Schlutzenberg has observed that the restriction of Σ to certain nice trees satisfies a form of comparison. The structure M is a weak coarse premouse of Schlutzenberg presentation ([51] Definition 3.1). When Σ is restricted to a certain nice $\text{OD}_{M \cup \{\Sigma\}}$ family of trees, Schlutzenberg shows that (M, Σ) satisfies a form of comparison ([51] Lemma 5.2) and even a form of comparison necessary for genericity iteration ([51] Section 5.2).¹⁸ Let $Z = M \cup \{\Sigma\}$. Note that $Z \in \text{OD}_Z^{L(Z, \mathbb{R})}$ and there is an $\text{OD}_Z^{L(Z, \mathbb{R})}$ wellordering of Z (since $M = J_{\bar{\alpha}}(J \cup \{\bar{w}, T, U\})$ and $\bar{w}$ is a wellordering of J). As noted in the remarks before [59] Theorem 10.2, comparison arguments can be used to form a directed system of countable iterates of M according to Σ . The universal Γ_1 -set is ordinal definable from Z since $x \in K_1$ if and only if $L(Z, \mathbb{R})$ satisfies “there is a countable iteration tree $\mathcal{T}$ according to Σ with last model P and iteration map $i : M \rightarrow P$ such that $x \in \pi_1[i(T)]$ ”. Since the set A belongs to the ω -parameterized pointclass $\Gamma_0 \subseteq \Gamma_1$, A is OD_Z using the fact that K_1 is OD_Z . As in the remarks before [59] Theorem 10.2, if one let (T_∞, U_∞) be the image of (T, U) in the directed system of countable iterates of M according to Σ (using the comparison of Schlutzenberg [51]), then (T_∞, U_∞) projects onto the universal Γ_1 -set K_1 and its complement. Note that $(T_\infty, U_\infty) \in \text{OD}_Z$ and hence belongs to HOD_Z since they are sets of ordinals. Let $Y = M \cup \{T_\infty, U_\infty\}$. $Y \in \text{HOD}_Z$. It is clear that $\text{OD}_Y \subseteq \text{OD}_Z$. The C_{Γ_1} operator can be defined from the universal Γ_1 -set K_1 and hence can be defined from (T_∞, U_∞) . $\Sigma \in \text{OD}_Y$ since “ Σ is the strategy on M guided by C_{Γ_1} ”. This shows $\text{OD}_Z = \text{OD}_Y$. (Alternatively, Steel ([59] Theorem 10.3) and Sargsyan ([48] Theorem 2.3.8; [49] Theorem 2.25) showed that M_1^Z Suslin-coSuslin captures K_1 . M_1^Z has a unique (ω, ω_1) -iteration strategy which is OD_Z and has fine structure. Thus comparison arguments and condensation work as usual. In the above discussion, one could replace (M, Σ) with a cut-off of M_1^Z and its unique iteration strategy.)

Fix the above (M, Σ) . A Σ -premouse is a hybrid strategy premouse of the form $J_{\alpha}^{\vec{E}, S}(M)$ which codes fragments of Σ in the sense of [48] Definition 2.2.12 and Definition 2.2.13. As in the proof of boldface GCH sketched in [61] Theorem 2.16, $\text{HOD}_Z^{L(Z, \mathbb{R})}$ is a direct limit of a certain system of Σ -mice with suitable properties generalizing the construction in [64] for $L(\mathbb{R})$. Thus $\text{HOD}_Z^{L(Z, \mathbb{R})} \models \text{GCH}$ and for any $\epsilon < \Theta^{L(Z, \mathbb{R})}$, $\text{cof}^V((\epsilon^+)^{\text{HOD}_Z^{L(Z, \mathbb{R})}}) = \text{cof}^{L(Z, \mathbb{R})}((\epsilon^+)^{\text{HOD}_Z^{L(Z, \mathbb{R})}}) = \omega$.

In summary, one has the following by taking $Z = M \cup \{\Sigma\}$ and $Y = M \cup \{T_\infty, U_\infty\}$ from the above discussion.

Lemma 6.6. *Assume AD and $\text{DC}_{\mathbb{R}}$. Let $A \in \mathcal{P}(\mathbb{R})$ be Suslin and coSuslin. Then there are sets Z and Y such that the following holds.*

- (1) *There is an $\text{OD}_Z^{L(Z, \mathbb{R})}$ wellordering of Z .*
- (2) *$Y \in \text{HOD}_Z^{L(Z, \mathbb{R})}$ and $\text{OD}_Z^{L(Z, \mathbb{R})} = \text{OD}_Y^{L(Z, \mathbb{R})} = \text{OD}_Y^{L(Y, \mathbb{R})}$. (Thus Z is a very good parameter set and Y is a very good parameter-witness for Z .)*
- (3) *$A \in \text{OD}_Z^{L(Z, \mathbb{R})}$ and thus $A \in L(Z, \mathbb{R})$.*

¹⁷See [62] Definition 1.2. $C_{\Gamma}(x)$ is the set of $y \in \mathbb{R}$ which is $\Delta(x)$ in some countable ordinals where $\Delta = \Gamma \cap \dot{\Gamma}$. See [32] and [33] for more on C_{Γ} .

¹⁸This comparison for coarse structures is also used in [52] to give a proof of the derived model theorem.

(4) $\text{HOD}_Z^{L(Z, \mathbb{R})} \models \text{GCH}$.

(5) For all $\epsilon < \Theta^{L(Z, \mathbb{R})}$, $\text{cof}^{L(Z, \mathbb{R})}((\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}) = \omega$.

One can now solve the Bounded Power Set conjecture.

Theorem 6.7. Assume AD^+ . For all cardinals κ such that $\omega < \kappa < \Theta$ and all $\epsilon < \kappa$, there is no injection of $\mathcal{P}_B(\kappa)$ into ${}^\epsilon\text{ON}$.

Proof. Suppose the theorem is false. Then there exist an uncountable cardinal $\lambda < \Theta$, $\chi < \lambda$, and an injection $\Psi : \mathcal{P}_B(\lambda) \rightarrow {}^\chi\text{ON}$. As observed in the proof of Theorem 6.2, χ must be infinite. Let $\tau_0 : \mathbb{R} \rightarrow \chi$ be a surjection. Since $\mathbb{R}$ surjects onto $\lambda < \Theta$, the Moschovakis coding lemma (Fact 2.15) implies there is a surjection of $\mathbb{R}$ onto $\mathcal{P}(\lambda)$ and in particular $\mathcal{P}_B(\lambda)$. Pick a surjection $\tau_1 : \mathbb{R} \rightarrow \mathcal{P}_B(\lambda)$. Let $A = \{\Psi(B)(\alpha) : \alpha < \chi \wedge B \in \mathcal{P}_B(\lambda)\}$. Let $\tau_2 : \mathbb{R} \rightarrow A$ be defined by $\tau_2(r) = \Psi(\tau_1(r^{[1]}))(\tau_0(r^{[0]}))$. τ_2 is a surjection and thus $\text{ot}(A) < \Theta$. Let $\eta = \text{ot}(A)$. Let $\tau_3 : A \rightarrow \eta$ be an order preserving bijection. Let $\Upsilon : \mathcal{P}_B(\kappa) \rightarrow {}^\chi\eta$ be defined by $\Upsilon(B)(\alpha) = \tau_3(\Psi(B)(\alpha))$ for all $\alpha < \chi$. Υ is an injection. It has been shown that there exist $\chi < \lambda < \Theta$, $\eta < \Theta$, and injection $\Upsilon : \mathcal{P}_B(\lambda) \rightarrow {}^\chi\eta$. Using the Moschovakis coding lemma, such a function Υ can be coded by a set of reals and thus $\Upsilon \in L(\mathcal{P}(\mathbb{R}))$. Also since AD^+ holds, $L(\mathcal{P}(\mathbb{R})) \models \text{AD}^+$. Let ϕ be the $\mathcal{L}_{\text{set}}$ sentences asserting “there exists $\epsilon < \kappa < \Theta$, $\delta < \Theta$, and an injection $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\delta$ ”. Recall if $\beta \leq \Theta$, then $\mathcal{W}_\beta = \{A \in \mathbb{R} : \text{rk}_W(A) < \beta\}$. Let ZF^* be some sufficiently large finite fragment of ZF . Let ψ be the formula asserting “there exist $\beta, \xi \in \text{ON}$ such that $L_\xi(\mathcal{W}_\beta) \models \text{ZF}^*, \text{AD}$, and ϕ ”. It was shown (as witnessed by the function Υ) that $L(\mathcal{P}(\mathbb{R})) \models \phi$. By using reflection on the hierarchy $\langle L_\xi(\mathcal{P}(\mathbb{R})) : \xi \in \text{ON} \rangle$, there is a $\xi \in \text{ON}$ so that $L_\xi(\mathcal{P}(\mathbb{R})) \models \text{ZF}^*, \text{AD}$, and ϕ . Letting $\beta = \Theta$, $\mathcal{W}_\beta = \mathcal{W}_\Theta = \mathcal{P}(\mathbb{R})$. Thus $L_\xi(\mathcal{W}_\beta) \models \text{ZF}^*, \text{AD}$, and ϕ . This shows that $L(\mathcal{P}(\mathbb{R})) \models \psi$. Since $\mathcal{S} \prec_1 L(\mathcal{P}(\mathbb{R}))$ by Σ_1 -reflection into Suslin-coSuslin sets (Fact 6.5), there exist $\beta, \xi \in \mathcal{S}$ so that $L_\xi(\mathcal{W}_\beta) \models \text{ZF}^*, \text{AD}$, and ϕ . There is a function $\Phi \in L_\xi(\mathcal{W}_\beta)$, $\epsilon < \kappa < \Theta^{L_\xi(\mathcal{W}_\beta)}$ and $\delta < \Theta^{L_\xi(\mathcal{W}_\beta)}$ so that $L_\xi(\mathcal{W}_\beta) \models “\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\delta \text{ is an injection}”$. By an application of the Moschovakis coding lemma (Fact 2.15) in $L(\mathcal{P}(\mathbb{R}))$ using a prewellordering and pointclass in $L_\xi(\mathcal{W}_\beta)$ as in the proof of Theorem 6.2, $\mathcal{P}_B(\kappa) = \mathcal{P}_B(\kappa)^{L_\xi(\mathcal{W}_\beta)}$ and $\Phi : \mathcal{P}_B(\kappa) \rightarrow {}^\epsilon\delta$ is still an injection in the real world by absoluteness. Since $\Phi \in L_\xi(\mathcal{W}_\beta) \subseteq \mathcal{S}$, the transitive closure of $\{\Phi\}$, $\text{tc}(\{\Phi\})$ belong to $\mathcal{K}$ (from Definition 6.4). Thus there are Suslin and coSuslin sets E and F such that $\text{tc}(\{\Phi\})$ is the Mostowski collapse of $(\mathbb{R}/E, \dot{\in}_F)$. Now since $\kappa \in L_\xi(\mathcal{W}_\beta) \subseteq \mathcal{S}$, κ is below a Suslin cardinal. Since the Suslin cardinals are unbounded below its supremum, there is a Suslin cardinal λ such that $\kappa < \lambda$. Thus there is a tree $T \in \mathcal{S}$ so that T is a tree on λ which projects onto the complete S_λ set. Let (E_0, F_0) be Suslin and coSuslin sets so that $\text{tc}(\{T\})$ is the Mostowski collapse of $(\mathbb{R}/E_0, \dot{\in}_{F_0})$. Let $A = \langle E, F, E_0, F_0 \rangle$ be a single set of reals coding these four sets of reals in a simple manner so that A is Suslin and coSuslin. Lemma 6.6 supplies set Z and Y with the properties stated in the lemma for A . Since $A \in L(Z, \mathbb{R})$ and $A \in \text{OD}_Z^{L(Z, \mathbb{R})}$, one has that $E, F, E_0, F_0 \in L(Z, \mathbb{R})$ and also $E, F, E_0, F_0 \in \text{OD}_Z^{L(Z, \mathbb{R})}$. $T \in L(Z, \mathbb{R})$ since T is the element of highest rank in the Mostowski collapse of $(\mathbb{R}/E_0, \dot{\in}_{F_0})$. So the universal S_λ set and the tree T witnessing it is λ -Suslin belong to $L(Z, \mathbb{R})$. So $L(Z, \mathbb{R}) \models “\lambda \text{ is a Suslin cardinal greater than } \kappa”$. Note that $\Phi \in \text{OD}_Z^{L(Z, \mathbb{R})}$ since $E_0, F_0 \in \text{OD}_Z^{L(Z, \mathbb{R})}$ and Φ is the element of highest rank in the Mostowski collapse of $(\mathbb{R}/E, \dot{\in}_F)$. Since $\text{HOD}_Z^{L(Z, \mathbb{R})} \models \text{GCH}$ and $\text{cof}^{L(Z, \mathbb{R})}((\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}) = \text{cof}^{L(Z, \mathbb{R})}((\epsilon^{++}) \text{HOD}_Z^{L(Z, \mathbb{R})}) = \text{cof}^{L(Z, \mathbb{R})}((\epsilon^{+++}) \text{HOD}_Z^{L(Z, \mathbb{R})}) = \omega$, Lemma 5.6 implies there is a $G \subseteq \text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_Z^{L(Z, \mathbb{R})}}$ which belongs to $L(Z, \mathbb{R})$ and is $\text{Coll}(\epsilon^+, \epsilon^{++})$ -generic over $\text{HOD}_Z^{L(Z, \mathbb{R})}$. Let $g : (\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})} \rightarrow (\epsilon^{++}) \text{HOD}_Z^{L(Z, \mathbb{R})}$ be the generic surjection associated to G . Since $\text{cof}^{L(Z, \mathbb{R})}((\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}) = \omega$, let $\rho : \omega \rightarrow (\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}$ be cofinal increasing in $(\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}$ and $\rho \in L(Z, \mathbb{R})$. Let $\sigma = \{g \upharpoonright \rho(n) : n \in \omega\}$. Then $G = \langle \sigma \rangle_{\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_Z^{L(Z, \mathbb{R})}}}$ and thus G is a countably generated $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_Z^{L(Z, \mathbb{R})}}$ -filter. Since $L(Z, \mathbb{R}) = L(Y, \mathbb{R})$ for $Y \in \text{HOD}_Z^{L(Z, \mathbb{R})}$, $\text{HOD}_Z^{L(Z, \mathbb{R})} \models |\text{Coll}(\epsilon^+, \epsilon^{++})| = |\epsilon^{++}|$, and $\epsilon^{++} < \kappa < \lambda$ which is a Suslin cardinal of $L(Z, \mathbb{R})$, Lemma 5.9 implies that $\text{HOD}_Z^{L(Z, \mathbb{R})}[G] = \text{HOD}_{Z \cup \{G\}}^{L(Z, \mathbb{R})}$. Let $\tilde{g} = \{\langle \alpha, g(\alpha) \rangle : \alpha < (\epsilon^+) \text{HOD}_Z^{L(Z, \mathbb{R})}\}$ and note that $\tilde{g} \in \mathcal{P}_B(\kappa)$ (by Fact 2.18) and $\tilde{g} \in \text{OD}_{\{G\}}^{L(Z, \mathbb{R})}$. Since $\Phi \in \text{OD}_Z^{L(Z, \mathbb{R})}$, one has that $\Phi(\tilde{g}) \in \text{OD}_{Z \cup \{G\}}^{L(Z, \mathbb{R})}$. Hence $\Phi(\tilde{g}) \in \text{HOD}_{Z \cup \{G\}}^{L(Z, \mathbb{R})} = \text{HOD}_Z^{L(Z, \mathbb{R})}[G]$. Since $\Phi(\tilde{g}) \in {}^\epsilon\text{ON}$ and $\text{Coll}(\epsilon^+, \epsilon^{++})^{\text{HOD}_Z^{L(Z, \mathbb{R})}}$ adds no new functions from ϵ into $\text{HOD}_Z^{L(Z, \mathbb{R})}$ since $\text{HOD}_Z^{L(Z, \mathbb{R})} \models “\text{Coll}(\epsilon^+, \epsilon^{++}) \text{ is } < \epsilon^+ \text{-closed}”$, one has that $\Phi(\tilde{g}) \in \text{HOD}_Z^{L(Z, \mathbb{R})}$. Then

1725 $\tilde{g} \in \text{OD}_Z^{L(Z, \mathbb{R})}$ since $\tilde{g} = \Phi^{-1}(\Phi(\tilde{g}))$. Thus $\tilde{g} \in \text{HOD}_Z^{L(Z, \mathbb{R})}$ and therefore $g \in \text{HOD}_Z^{L(Z, \mathbb{R})}$ and $G \in \text{HOD}_Z^{L(Z, \mathbb{R})}$.
 1726 This is impossible since $\text{HOD}_Z^{L(Z, \mathbb{R})} \models "g : \epsilon^+ \rightarrow \epsilon^{++} \text{ is a surjection}"$ by absoluteness. $\square$

1727 By adapting the proof of Theorem 6.3 with the proof of Theorem 6.7, one obtains the following result.

1728 **Theorem 6.8.** *Assume AD^+ . Let $\kappa < \Theta$ be a singular cardinal and let $\epsilon < \kappa$ be such that $\text{cof}(\epsilon) = \text{cof}(\kappa)$.
 1729 Then the following holds.*

- 1730 (1) $|\mathcal{P}_B(\kappa)| < |[\kappa]^\epsilon \cup \mathcal{P}_B(\kappa)|$.
- 1731 (2) $|[\kappa]^\epsilon| < |[\kappa]^\epsilon \cup \mathcal{P}_B(\kappa)|$.
- 1732 (3) $|[\kappa]^\epsilon \cup \mathcal{P}_B(\kappa)| < |\mathcal{P}(\kappa)|$.

1733 Finally, one can prove the ABCD conjecture using the Bounded Power Set conjecture.

1734 **Theorem 6.9.** *Assume AD^+ . Let α, β, γ , and δ be cardinals such that $\omega \leq \alpha \leq \beta < \Theta$ and $\omega \leq \gamma \leq \delta < \Theta$.
 1735 $|\alpha\beta| \leq |\gamma\delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$.*

Proof. ($\Leftarrow$) Suppose $\alpha \leq \gamma$ and $\beta \leq \delta$. Let $f : \alpha \rightarrow \beta$ be a function. Define $\Phi(f) : \gamma \rightarrow \delta$ by

$$\Phi(f)(\xi) = \begin{cases} f(\xi) & \xi < \alpha \\ 0 & \alpha \leq \xi \end{cases}$$

1736 This defines $\Phi : \alpha\beta \rightarrow \gamma\delta$. Suppose $f \neq g$. There is a $\nu < \alpha$ so that $f(\nu) \neq g(\nu)$. Thus $\Phi(f)(\nu) = f(\nu) \neq$
 1737 $g(\nu) = \Phi(g)(\nu)$. This shows that Φ is an injection and hence $|\alpha\beta| \leq |\gamma\delta|$.

1738 ($\Rightarrow$) Suppose this is false. There is an injection $\Phi : \alpha\beta \rightarrow \gamma\delta$ and $\neg(\alpha \leq \gamma \wedge \beta \leq \delta)$. Suppose $\delta < \beta$. Then
 1739 $|\beta| \leq |\alpha\beta| \leq |\gamma\delta| \leq |\delta| = |\mathcal{P}(\delta)|$ where the second inequality is witnessed by Φ (and one can use ($\Leftarrow$) from
 1740 above for the third inequality). Since $\delta < \beta$, $\delta^+ \leq \beta$. Hence there is an injection of δ^+ into $\mathcal{P}(\delta)$ which
 1741 violates boldface GCH at δ (Fact 2.22). Suppose $\gamma < \alpha$. $|\mathcal{P}_B(\alpha)| \leq |\mathcal{P}(\alpha)| \leq |\alpha 2| \leq |\alpha\beta| \leq |\gamma\delta|$. Thus there
 1742 is an injection of $\mathcal{P}_B(\alpha)$ into ${}^\gamma\text{ON}$ where $\gamma < \alpha$ in violation of Theorem 6.7. This yields a contradiction. $\square$

1743 *Remark 6.10.* With the ABCD hypothesis below Θ , one can completely classify the cardinality relationship
 1744 between any pair of infinite cardinal exponentiations below Θ under AD^+ : Suppose $\omega \leq \alpha, \beta, \gamma, \delta < \Theta$ are
 1745 four infinite cardinals below Θ . If $\alpha \leq \beta$, then let $\bar{\alpha} = \alpha$ and $\bar{\beta} = \beta$. If $\beta < \alpha$, then let $\bar{\alpha} = \alpha$ and
 1746 $\bar{\beta} = \alpha$. Since $\omega \leq \beta < \alpha$, one has that $|\alpha\beta| = |\alpha 2| = |\mathcal{P}(\alpha)| = |\alpha\alpha| = |\alpha\bar{\beta}|$. Thus one has found cardinals
 1747 $\omega \leq \bar{\alpha} \leq \bar{\beta} < \Theta$ so that $|\alpha\beta| = |\bar{\alpha}\bar{\beta}|$. Similarly, one can find $\omega \leq \bar{\gamma} \leq \bar{\delta} < \Theta$ so that $|\gamma\delta| = |\bar{\gamma}\bar{\delta}|$. Now one can
 1748 apply the ABCD hypothesis below Θ (Theorem 6.7) to $\omega \leq \bar{\alpha} \leq \bar{\beta} < \Theta$ and $\omega \leq \bar{\gamma} \leq \bar{\delta} < \Theta$ to determine
 1749 what is the cardinality relationship between $|\alpha\beta| = |\bar{\alpha}\bar{\beta}|$ and $|\gamma\delta| = |\bar{\gamma}\bar{\delta}|$.

1750 Let $\epsilon \leq \kappa$ be ordinals. Recall that ${}^\epsilon\kappa$ is the set of functions $f : \epsilon \rightarrow \kappa$ and $[\kappa]^\epsilon$ is the set of increasing
 1751 functions $f : \epsilon \rightarrow \kappa$. Let $B(\epsilon, \kappa) = \{f \in {}^\epsilon\kappa : \sup(f) < \kappa\}$, which is the set of functions $f : \epsilon \rightarrow \kappa$ which are
 1752 bounded below κ . Next, $B(\epsilon, \kappa)$, $[\kappa]^\epsilon$, and ${}^\epsilon\kappa$ will be distinguished and classified.

1753 **Fact 6.11.** *Let $\epsilon \leq \kappa$ be limit ordinals.*

- 1754 (1) $|{}^\epsilon\kappa| = |{}^{|\epsilon|}\kappa|$.
- 1755 (2) $|B(\epsilon, \kappa)| = |B(|\epsilon|, \kappa)|$.
- 1756 (3) If $\epsilon \leq |\kappa| < \kappa$, then $|B(\epsilon, \kappa)| = |{}^{|\epsilon|}\kappa|$.
- 1757 (4) If $\text{cof}(\epsilon) \neq \text{cof}(\kappa)$, then $[\kappa]^\epsilon = B(\epsilon, \kappa)$.
- 1758 (5) If $\epsilon \leq \text{cof}(\kappa)$, then $[[\kappa]^\epsilon] = |{}^\epsilon\kappa|$.
- 1759 (6) $|\mathcal{P}_B(|\epsilon|)| \leq |[\kappa]^\epsilon| \leq |{}^\epsilon\kappa|$.
- 1760 (7) If $\text{cof}(\epsilon) = \text{cof}(\kappa)$ and κ is a limit of indecomposable ordinals, then $[[\kappa]^\epsilon] = |{}^\epsilon\kappa|$.
- 1761 (8) Assume for all $\delta < \kappa$, $\neg(|\kappa| \leq |\mathcal{P}(\delta)|)$. If $\text{cof}(\epsilon) = \text{cof}(\kappa)$, then $|B(\epsilon, \kappa)| < |[\kappa]^\epsilon|$.
- 1762 (9) Assume AD^+ . Suppose $\delta < \epsilon \leq \kappa$ and $|\delta| < |\epsilon|$. Then any member of $\{B(\epsilon, \kappa), [\kappa]^\epsilon, {}^\epsilon\kappa\}$ does not
 1763 inject into any member of $\{B(\delta, \kappa), [\kappa]^\delta, {}^\delta\kappa\}$.

1764 *Proof.* (1), (2), (3), and (4) are clear.

1765 (5) $[\kappa]^\epsilon \subseteq {}^\epsilon\kappa$. Let $f \in {}^\epsilon\kappa$. $\Phi(f) \in [\kappa]^\epsilon$ will be defined by induction as follows. Suppose $\alpha < \epsilon$ and
 1766 $\Phi(f) \upharpoonright \alpha \in [\kappa]^\alpha$ has been defined. Since $\alpha < \text{cof}(\kappa)$, $\sup(\Phi(f) \upharpoonright \alpha) < \kappa$. Let $\Phi(\alpha) = \sup(\Phi(f) \upharpoonright \alpha) + f(\alpha)$. This
 1767 completes the definition of $\Phi(f) \in [\kappa]^\epsilon$. $\Phi : {}^\epsilon\kappa \rightarrow [\kappa]^\epsilon$ is an injection. Thus $|{}^\epsilon\kappa| = |[\kappa]^\epsilon|$.

(6) Let k be any element of $[\kappa]^{\epsilon-|\epsilon|}$ such that $\min(k) \geq |\epsilon|$ (if $|\epsilon| = \epsilon$, then $k = \emptyset$). Since $|\epsilon|$ is a cardinal, let A and B be two disjoint subsets of $|\epsilon|$ such that $|A| = |B| = |\epsilon|$. Let $E_A : |\epsilon| \rightarrow A$ and $E_B : |\epsilon| \rightarrow B$ be the increasing enumeration of A and B , respectively. If $\ell \in \mathcal{P}_B(|\epsilon|)$, let $f_\ell : \text{ot}(\ell) \rightarrow \ell$ be the increasing enumeration of ℓ . Define $h_\ell \in [\kappa]^\epsilon$ by

$$h_\ell(\alpha) = \begin{cases} E_A(f_\ell(\alpha)) & \alpha < \text{ot}(\ell) \\ E_B(\sup(E_A \circ f_\ell) + (\alpha - \text{ot}(\ell))) & \text{ot}(\ell) \leq \alpha < |\epsilon| \\ k(\alpha - |\epsilon|) & |\epsilon| \leq \alpha < \epsilon \end{cases}$$

Since $\ell \in \mathcal{P}_B(|\epsilon|)$, $\sup(f_\ell) < |\epsilon|$. Thus $\sup(E_A \circ f_\ell) < |\epsilon|$. These two observations can be used to show that h_ℓ is indeed an increasing function. Define $\Psi : \mathcal{P}_B(|\epsilon|) \rightarrow [\kappa]^\epsilon$ by $\Psi(\ell) = h_\ell$. Ψ is an injection. Thus $|\mathcal{P}_B(|\epsilon|)| \leq |[\kappa]^\epsilon| \leq |\epsilon|^\kappa$.

(7) It is clear that $[\kappa]^\epsilon \subseteq {}^\epsilon\kappa$. If κ is regular, then the result follows from (5). Assume $\text{cof}(\kappa) < \kappa$. Let $\delta = \text{cof}(\epsilon) = \text{cof}(\kappa)$. Let $\rho_0 : \delta \rightarrow \epsilon$ and $\rho_1 : \delta \rightarrow \kappa$ be increasing cofinal maps from δ into ϵ and κ , respectively. One may assume that $\delta < \rho_1(0)$. By the hypothesis, one may assume that the image of ρ_1 consists entirely of indecomposable ordinals (which are closed under the Gödel pairing function, $\langle \cdot, \cdot \rangle$). Let $f \in {}^\epsilon\kappa$. One will define $\Gamma(f) \in [\kappa]^\epsilon$ by recursion as follows. For each $\xi < \delta$, let $B_\xi^f = \{\alpha < \rho_0(\xi) : \sup(\rho_1 \upharpoonright \xi) \leq f(\alpha) < \rho_1(\xi)\}$. For $\xi < \delta$, let $\iota_\xi = \text{ot}(B_\xi^f)$. Let $E_\xi^f : \iota_\xi \rightarrow B_\xi^f$ be the increasing enumeration of B_ξ^f . Suppose $\xi < \delta$ and for all $\gamma < \xi$, an increasing function g_γ^f has been defined so that g_γ^f has domain less than δ and $\sup(g_\gamma^f) < \rho_1(\gamma + 1)$ and for all $\gamma_0 < \gamma_1 < \xi$, $g_{\gamma_0}^f \subseteq g_{\gamma_1}^f$. Let $h = \bigcup_{\gamma < \xi} g_\gamma^f$ and note that $\text{dom}(h) = \sup\{\text{dom}(g_\gamma^f) : \gamma < \xi\} = \sup\{\rho_1(\gamma + 1) : \gamma < \xi\} \leq \rho_1(\xi)$. Define g_ξ^f with domain $\text{dom}(h) + \iota_\xi^f$ by recursion as follows. If $\alpha < \text{dom}(h)$, then $g_\xi^f(\alpha) = h(\alpha)$. Suppose $\beta < \iota_\xi^f$ and $g_\xi^f \upharpoonright \text{dom}(h) + \beta$ has been defined for some $\beta < \iota_\xi^f$. Then let $g_\xi^f(\text{dom}(h) + \beta) = \sup(g_\xi^f \upharpoonright \text{dom}(h) + \beta) + \langle E_\xi^f(\beta), f(E_\xi^f(\beta)) \rangle$. This completes the construction of g_ξ^f . Since $E_\xi^f(\beta) \in B_\xi^f$, $f(E_\xi^f(\beta)) < \rho_1(\xi)$ for all $\beta < \iota_\xi^f$. Since ρ_1 takes image among ordinals which are closed under the Gödel pairing function, $\langle E_\xi^f(\beta), f(E_\xi^f(\beta)) \rangle < \rho_1(\xi)$. Hence $\sup(g_\xi^f) \leq \sup(h) + \rho_1(\xi) \cdot \delta < \rho_1(\xi + 1)$ since $\rho_1(\xi + 1)$ is indecomposable. This completes the construction of $\langle g_\xi^f : \xi < \delta \rangle$. Let $\Gamma(f) = \bigcup_{\xi < \delta} g_\xi^f$. Note that $\Gamma(f) \in [\kappa]^\epsilon$ and $\Gamma : {}^\epsilon\kappa \rightarrow [\kappa]^\epsilon$ is an injection. Thus $|\epsilon|^\kappa \leq |[\kappa]^\epsilon|$.

(8) There is no injection of $[\kappa]^\epsilon$ into $B(\epsilon, \kappa)$ since $|B(\epsilon, \kappa)| \leq |\mathcal{P}_B(\kappa)|$ and there is no injection of $[\kappa]^\epsilon$ into $\mathcal{P}_B(\kappa)$ by Theorem 2.2 when $\text{cof}(\epsilon) = \text{cof}(\kappa)$.

(9) This follows from (6) and Theorem 6.7. $\square$

One can now show that $|[\omega_3]^{\omega_2}| < |[\omega_3]^{<\omega_3}| \leq |\mathcal{P}(\omega_3)|$ under AD^+ .

Theorem 6.12. Assume AD^+ . If $\kappa < \Theta$ is a cardinal and $\epsilon < \kappa$, then $|\epsilon|^\kappa < |{}^{<\kappa}\kappa|$.

Proof. Since $\mathcal{P}_B(\kappa)$ injects into ${}^{<\kappa}\kappa$, ${}^{<\kappa}\kappa$ cannot inject into ${}^\epsilon\kappa$ by Theorem 6.7. $\square$

[17] defined a notion of regularity and described the cofinality of some familiar sets in the choiceless determinacy context.

Definition 6.13. ([17]) Let X be a set and Y be a set or class. X has Y -regular cardinality if and only if for all $\Phi : X \rightarrow Y$, there is a $y \in Y$ so that $|\Phi^{-1}[\{y\}]| = |X|$.

A set X is said to have locally regular cardinality if and only if for all sets Y with $|Y| < |X|$, X has Y -regular cardinality.

A set X is said to have globally regular cardinality if and only if for all sets Y with $\neg(|X| \leq |Y|)$, X has Y -regular cardinality.

Under AD , $\mathbb{R} \sqcup \omega_1$ does not have 2-regular cardinality. Under AD^+ , [17] shows that $\mathbb{R}$ and $\mathbb{R}/E_0$ have globally regular cardinality. Section 7 will show that under AD^+ , if $\omega \leq \kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then ${}^\omega\kappa$ has ON-regular cardinality, $B(\omega, \kappa)$ has n -regular cardinality for all $n \in \omega$, and if $\omega < \kappa$, then $B(\omega, \kappa)$ does not have ω -regular cardinality.

Some information is known about the cofinality of the power set of a strong partition cardinal.

Fact 6.14. ([6]) If $\kappa \rightarrow_* (\kappa)_2^\kappa$ (κ is a strong partition cardinal), then $\mathcal{P}(\kappa)$ is ON-regular: for all $\delta \in \text{ON}$, $\mathcal{P}(\kappa)$ has δ -regular cardinality.

([17]) If $\kappa \rightarrow_* (\kappa)_{<\kappa}^\kappa$ (κ is a very strong partition cardinal), then $\mathcal{P}(\kappa)$ has $<^\kappa \text{ON}$ -regular cardinality: for all $\delta \in \text{ON}$, $\mathcal{P}(\kappa)$ has $<^\kappa \delta$ -regular cardinality.

The main theorem of this section gives the following new ordinal regularity result. It implies that $<^{\omega_\omega}(\omega_{\omega+1})$ does not have ω -regular cardinality which was not previously known.

Theorem 6.15. Assume AD^+ . Let $\delta \leq \kappa < \Theta$ be such that δ is a limit ordinal. Then $<^\delta \kappa$ does not have $\text{cof}(\delta)$ -regular cardinality.

Proof. Let $\rho : \text{cof}(\delta) \rightarrow \delta$ be an increasing cofinal function. Define $\Phi : <^\delta \kappa \rightarrow \text{cof}(\delta)$ by $\Phi(f)$ is the least α so that $\text{dom}(f) < \rho(\alpha)$. For each $\alpha < \text{cof}(\delta)$, $\Phi^{-1}[\{\alpha\}] \subseteq <^{\rho(\alpha)} \kappa$. Thus $|\Phi^{-1}[\{\alpha\}]| \leq |<^{\rho(\alpha)} \kappa| < |<^\delta \kappa|$ since $\mathcal{P}_B(\delta) \subseteq <^\delta \kappa$ and $\mathcal{P}_B(\delta)$ cannot inject into $<^{\rho(\alpha)} \kappa$ or even ${}^{\rho(\alpha)}\text{ON}$ by Theorem 6.7. $\square$

[17] shows that under AD^+ , $\mathcal{P}(\omega_1)$ is regular relative to essentially every set which does not already contain an injective copy of $\mathcal{P}(\omega_1)$ for which one currently have a practical understanding. This gives empirical evidence for the conjecture that $\mathcal{P}(\omega_1)$ has globally regular cardinality under AD^+ .

If $\mathcal{P}(\kappa)$ has 2-regular cardinality, then statement (3) in Theorem 6.3 and Theorem 6.8 would follow immediately from the fact that $\mathcal{P}(\kappa)$ does not inject into $\mathcal{P}_B(\kappa)$ and does not inject into $[\kappa]^\epsilon$ when $\text{cof}(\epsilon) = \text{cof}(\kappa) < \kappa$. However, it is not even known if $\mathcal{P}(\omega_2)$ or $\mathcal{P}(\omega_3)$ has 2-regular cardinality. For this reason, the proofs of statement (3) in Theorem 6.3 and Theorem 6.8 require the more complicated argument resembling the proofs of Theorem 6.2 and Theorem 6.7. Since ω_3 is the first singular cardinal of determinacy, the first new instance of Theorem 6.8 (3) is that AD^+ implies $|[\omega_3]^{\omega_2} \cup \mathcal{P}_B(\omega_3)| < |\mathcal{P}(\omega_3)|$.

At a singular cardinal κ , there are two natural combinatorial sets whose cardinality have not yet been distinguished. They are $|<^\kappa \kappa| = |[\kappa]^{<\kappa}|$ and $|\mathcal{P}(\kappa)|$. The first open case occurs at the first singular cardinal of determinacy. Is $|[\omega_3]^{<\omega_3}| < |\mathcal{P}(\omega_3)|$? Another closely related question is whether $\mathcal{P}(\omega_2)$ injects into $<^{\omega_2} \text{ON}$. The main result of the paper can be used to resolve these questions if one assumes a regularity condition on $\mathcal{P}(\kappa)$.

Fact 6.16. Assume AD^+ . Let $\kappa < \Theta$ and assume $\mathcal{P}(\kappa)$ has κ -regular cardinality. Then $|[\kappa]^{<\kappa}| < |\mathcal{P}(\kappa)|$ and $\mathcal{P}(\kappa)$ does not inject into $<^\kappa \text{ON}$.

Proof. The second statement implies the first. Suppose $\Phi : \mathcal{P}(\kappa) \rightarrow <^\kappa \text{ON}$ is an injection. Let $\Gamma : \mathcal{P}(\kappa) \rightarrow \kappa$ be defined by $\Gamma(f) = |\Phi(f)| = \text{dom}(\Phi(f))$. By the hypothesis that $\mathcal{P}(\kappa)$ is κ -regular, there is an $\epsilon < \kappa$ so that $|\Gamma^{-1}[\{\epsilon\}]| = |\mathcal{P}(\kappa)|$. Let $\Upsilon : \mathcal{P}(\kappa) \rightarrow \Gamma^{-1}[\{\epsilon\}]$ be an injection. Then $\Phi \circ \Upsilon : \mathcal{P}(\kappa) \rightarrow \epsilon \text{ON}$ is an injection. Then $\Phi \circ \Upsilon \upharpoonright \mathcal{P}_B(\kappa) : \mathcal{P}_B(\kappa) \rightarrow \epsilon \text{ON}$ is an injection which violates Theorem 6.7. $\square$

However, Fact 6.16 is applicable for the power set of a strong partition cardinal.

Theorem 6.17. Assume AD^+ . Let $\kappa < \Theta$ and assume $\kappa \rightarrow_* (\kappa)_2^\kappa$. Then $|[\kappa]^{<\kappa}| < |\mathcal{P}(\kappa)|$ and $\mathcal{P}(\kappa)$ does not inject $<^\kappa \text{ON}$.

Proof. Fact 6.14 showed that $\kappa \rightarrow_* (\kappa)_2^\kappa$ implies that $\mathcal{P}(\kappa)$ has ON -regular cardinality. The result follows from Fact 6.16. $\square$

If κ is a strong partition cardinal, then the first statement of Theorem 6.17 can be proved purely combinatorially.

Theorem 6.18. Assume κ satisfies $\kappa \rightarrow_* (\kappa)_2^\kappa$. Then $|[\kappa]^{<\kappa}| < |\mathcal{P}(\kappa)|$.

Proof. Suppose $\Phi : \mathcal{P}(\kappa) \rightarrow [\kappa]^{<\kappa}$ is an injection. Since $\kappa \rightarrow_* (\kappa)_2^\kappa$ implies κ is regular, for all $f \in \mathcal{P}(\kappa)$, $\sup(\Phi(f)) < \kappa$. Let $\Gamma : \mathcal{P}(\kappa) \rightarrow \kappa$ be defined by $\Gamma(f) = \sup(f) + 1$. $\kappa \rightarrow_* (\kappa)_2^\kappa$ implies $\mathcal{P}(\kappa)$ has ON -regular cardinality by Fact 6.14. Thus there is a $\delta < \kappa$ so that $|\Gamma^{-1}[\{\delta\}]| = |\mathcal{P}(\kappa)|$. Let $\Upsilon : \mathcal{P}(\kappa) \rightarrow \Gamma^{-1}[\{\delta\}]$ be an injection. Then $\Phi \circ \Upsilon : \mathcal{P}(\kappa) \rightarrow [\delta]^{<\delta}$ is an injection. This is impossible since $\kappa \rightarrow_* (\kappa)_2^\kappa$ implies that κ is measurable and hence there no injection of κ into $\mathcal{P}(\delta)$ by Fact 2.3. $\square$

If κ is a very strong partition cardinal, then the second statement of Theorem 6.17 can be proved purely combinatorially. (It is open whether every strong partition cardinal is a very strong partition cardinal even under AD or AD^+ .)

Theorem 6.19. Assume κ satisfies $\kappa \rightarrow_* (\kappa)_{<\kappa}^\kappa$. Then $\mathcal{P}(\kappa)$ does not inject into $<^\kappa \text{ON}$.

1853 *Proof.* Fact 6.14 shows that $\kappa \rightarrow_* (\kappa)_{<\kappa}^\kappa$ implies that $\mathcal{P}(\kappa)$ has $<^\kappa$ ON-regular cardinality. So no function
 1854 $\Phi : \mathcal{P}(\kappa) \rightarrow <^\kappa$ ON can be injective. $\square$

1855 Next, one will create a large family of intermediate cardinalities below $\mathcal{P}_B(\kappa)$ when $\kappa < \Theta$ is a cardinal
 1856 in $L(\mathbb{R})$. The following will review some basic facts about J -constructibility reduction when J is a set of
 1857 ordinals.

1858 **Definition 6.20.** Let $J \subseteq \text{ON}$ be a set of ordinals. For $x, y \in \mathbb{R}$, let $x \leq_J y$ if and only if $x \in L[J, y]$ which is
 1859 the J -constructibility degrees. Define $x \equiv_J y$ if and only if $x \leq_J y$ and $y \leq_J x$ which is the J -constructibility
 1860 equivalence relation. Let $\mathcal{D}_J = \mathbb{R} / \equiv_J$ be the set of J -degrees or $\equiv_J$ -equivalence classes. For $X, Y \in \mathcal{D}_J$, let
 1861 $X \leq_J Y$ if and only if for any $x \in X$ and $y \in Y$, $x \leq_J y$. For each $X \in \mathcal{D}_J$, $\mathcal{C}_J(X) = \{Y \in \mathcal{D}_J : X \leq_J Y\}$
 1862 which is the J -constructibility cone above X . Let μ_J be a filter on $\mathcal{D}_J$ defined by $A \in \mu_J$ if and only if there
 1863 is an $X \in \mathcal{D}_J$ so that $\mathcal{C}_J(X) \subseteq A$. A perfect tree p on 2 is J -pointed if and only if for all $x \in [p]$, $p \leq_J x$.
 1864 (Recall that a perfect tree p on 2 is a Turing pointed perfect tree if and only if for all $x \in [p]$, $p \leq_{\text{Turing}} x$,
 1865 where $\leq_{\text{Turing}}$ refers to Turing reduction. Every Turing pointed perfect tree is an J -pointed perfect tree.)
 1866 If p is a J -pointed tree, then $\{[x]_{\equiv_J} : x \in [p]\} = \mathcal{C}_J([p]_{\equiv_J})$. A function f with domain $\mathbb{R}$ is J -invariant if
 1867 and only if for all $x, y \in \mathbb{R}$, if $x \equiv_J y$, then $f(x) = f(y)$. A J -invariant function $f : \mathbb{R} \rightarrow \text{ON}$ is everywhere
 1868 J -increasing if and only if for all $x, y \in \mathbb{R}$, $x \leq_J y$ implies $f(x) \leq f(y)$. A J -invariant function $f : \mathbb{R} \rightarrow \text{ON}$
 1869 is μ_J -almost everywhere increasing if and only if there is an $a \in \mathbb{R}$ so that for all $a \leq_J x \leq_J y$, $f(x) \leq f(y)$.
 1870 If $F, G : \mathcal{D}_J \rightarrow \text{ON}$, then define $F \sim_{\mu_J} G$ if and only if $\{X \in \mathcal{D}_J : F(X) = G(X)\} \in \mu_J$ and $F <_{\mu_J} G$
 1871 if and only if $\{X \in \mathcal{D}_J : F(X) < G(X)\} \in \mu_J$. Let $\prod_{\mathcal{D}_J} \text{ON} / \mu_J$ be the ultrapower of ON by μ_J which are the
 1872 $\sim_{\mu_J}$ -equivalence classes of function $F : \mathcal{D}_J \rightarrow \text{ON}$ ordered by the relation induced by $<_{\mu_J}$. If $f, g : \mathbb{R} \rightarrow \text{ON}$
 1873 are J -invariant functions, then define $f \sim_{\mu_J} g$ if and only if $\{[x]_{\equiv_J} : f(x) = g(x)\} \in \mu_J$ and $f \leq_{\mu_J} g$ if and
 1874 only if $\{[x]_{\equiv_J} : f(x) < g(x)\} \in \mu_J$.

1875 The following basic facts are proved using the same argument as for the corresponding facts for Turing
 1876 reducibility.

1877 **Fact 6.21.** (Martin) Assume AD.

- 1878 (1) For any $B \subseteq \mathbb{R}$, there is a J -pointed perfect tree (even a Turing pointed perfect tree) p so that $[p] \subseteq B$
 1879 or $[p] \subseteq \mathbb{R} \setminus B$.
- 1880 (2) μ_J is a countably complete ultrafilter on $\mathcal{D}_J$.
- 1881 (3) For all $A \subseteq \mathbb{R}$ such that there is a $x \in \mathbb{R}$ with $\{y \in \mathbb{R} : x \leq_J y\} \subseteq A$, then there is a J -pointed tree
 1882 such that $[p] \subseteq A$.

1883 **Fact 6.22.** Assume AD and $\text{DC}_{\mathbb{R}}$. For every J -invariant function $f : \mathbb{R} \rightarrow \text{ON}$, there is a J -invariant
 1884 everywhere J -increasing function $g : \mathbb{R} \rightarrow \text{ON}$ so that $f \sim_{\mu_J} g$.

1885 *Proof.* Let $f : \mathbb{R} \rightarrow \text{ON}$ be J -invariant. Let $\bar{f} : \mathcal{D}_J \rightarrow \text{ON}$ be defined by $\bar{f}([x]_{\equiv_J}) = f(x)$ which is
 1886 well defined since f is J -invariant. Let $A = \{X \in \mathcal{D}_J : (\forall Y \geq X)(\bar{f}(X) \leq \bar{f}(Y))\}$ and $B = \{X \in$
 1887 $\mathcal{D}_J : (\exists Y \geq X)(\bar{f}(Y) < \bar{f}(X))\}$. Since $A \cap B = \emptyset$, $A \cup B = \mathcal{D}_J$, and μ_J is an ultrafilter by Fact 6.21,
 1888 exactly one of A or B belongs to μ_J . Suppose $B \in \mu_J$. Let $Z \in \mathcal{D}_J$ be such that $\mathcal{C}_J(Z) \subseteq B$. Pick
 1889 $z \in Z$. Define $R(x, y)$ if and only if $f(y \oplus z) < f(x \oplus z)$. For any $x \in \mathbb{R}$, $[x \oplus z]_{\equiv_J} \in \mathcal{C}_J(Z) \subseteq B$. So
 1890 there is a $Y \geq [x \oplus z]$ so that $\bar{f}(Y) < \bar{f}([x \oplus z]_{\equiv_J})$. Pick any $y \in Y$ and note that $y \equiv_J y \oplus z$. Then
 1891 $f(y \oplus z) = \bar{f}([y \oplus z]_{\equiv_J}) = \bar{f}([y]_{\equiv_J}) = \bar{f}(Y) < \bar{f}([x \oplus z]_{\equiv_J}) = f(x \oplus z)$. Thus $R(x, y)$. By $\text{DC}_{\mathbb{R}}$, there
 1892 is a sequence $\langle x_n : n \in \omega \rangle$ so that $R(x_n, x_{n+1})$ for all $n \in \omega$. Then $\langle f(x_n \oplus z) : n \in \omega \rangle$ is a descending
 1893 sequence of ordinals which is a contradiction. Thus it must be the case that $A \in \mu_J$. Let $U \in \mathcal{D}_J$ so
 1894 that $\mathcal{C}_J(U) \subseteq A$. Let $u \in U$. Define $g(x) = \min\{f(y) : u \oplus x \leq_J y\}$. g is clearly J -invariant. Suppose
 1895 $x_0 \leq_J x_1$. Then $\{f(y) : u \oplus x_1 \leq_J y\} \subseteq \{f(y) : u \oplus x_0 \leq_J y\}$. Hence $g(x_0) \leq g(x_1)$. Thus g is everywhere
 1896 J -increasing. Suppose $X \geq U$. Let $x \in X$. Note that $x \oplus u \equiv_J x$ and for all $y \in \mathbb{R}$, if $y \geq x \oplus u$, then
 1897 $y \geq_J x \geq_J u$. Since $[x]_{\mu_J} = X \in A$, one has that $f(y) = \bar{f}([y]_{\equiv_J}) \geq \bar{f}([x]_{\equiv_J}) = f(x)$ for all $y \geq_J x \oplus u$.
 1898 Thus $g(x) = \min\{f(y) : y \geq_J x \oplus u\} = f(x)$. This shows that $f \sim_{\mu_J} g$. $\square$

1899 **Definition 6.23.** Let $\text{gp} : \text{ON} \times \text{ON} \rightarrow \text{ON}$ be the Gödel pairing function. If $\delta, \epsilon \in \text{ON}$ and $A \subseteq \text{ON}$, then
 1900 A is said to code a function from δ into ϵ if and only if $\text{gp}^{-1}[A]$ is the graph of a function $f : \delta \rightarrow \epsilon$. If
 1901 $f : \delta \rightarrow \epsilon$, let $\mathfrak{G}_f = \{\text{gp}(\alpha, f(\alpha)) : \alpha < \delta\}$. Note that if ϵ is indecomposable, then $\mathfrak{G}_f \in \epsilon$.

Let $J \subseteq \text{ON}$, $\epsilon \in \text{ON}$, $r \in \mathbb{R}$, and $\alpha < \omega_1$. Let $G_{\alpha,r}^{J,\epsilon}$ be the set of all $\mathfrak{G}_f$ such that $f : \delta \rightarrow (\epsilon^{+(\alpha+1)})^{L[J,r]}$ is a function and $\delta \leq (\epsilon^{+\alpha})^{L[J,r]}$. Note that $G_{\alpha,r}^{J,\epsilon} \subseteq \mathcal{P}((\epsilon^{+(\alpha+1)})^{L[J,r]}) \subseteq \mathcal{P}_B(\epsilon^+)$ since Fact 2.19 implies that $(\epsilon^{+(\alpha+1)})^{L[J,r]} < \epsilon^+$. Thus $|G_{\alpha,r}^{J,\epsilon}| \leq |\mathcal{P}(\epsilon)|$. Also if $A \subseteq \epsilon$, then the characteristic function of A , $\chi_A : \epsilon \rightarrow 2$ belongs to $G_{\alpha,r}^{J,\epsilon}$. This shows that $|\mathcal{P}(\epsilon)| \leq |G_{\alpha,r}^{J,\epsilon}|$. Hence $|G_{\alpha,r}^{J,\epsilon}| = |\mathcal{P}(\epsilon)|$.

Let $h : \mathbb{R} \rightarrow \omega_1$ be a J -invariant everywhere J -increasing function. Then let $F_h^{J,\epsilon} = \bigsqcup_{r \in \mathbb{R}} G_{h(r),r}^{J,\epsilon} = \{(r, f) : r \in \mathbb{R} \wedge f \in G_{h(r),r}^{J,\epsilon}\}$. Note that $|F_h^{J,\epsilon}| \leq |\mathcal{P}_B(\epsilon^+)|$.

Fact 6.24. Assume AD. Let J be a set of ordinals, $\epsilon \in \text{ON}$ with $\omega \leq \epsilon$, and $h : \mathbb{R} \rightarrow \omega_1$ be a J -invariant everywhere J -increasing function. Assume boldface GCH at ϵ holds. Then $\neg(|\epsilon^+| \leq |F_h^{J,\epsilon}|)$ and thus $|\mathcal{P}(\epsilon)| \leq |F_h^{J,\epsilon}| < |\mathcal{P}_B(\epsilon^+)|$.

Proof. Suppose $\Phi : \epsilon^+ \rightarrow F_h^{J,\epsilon}$ is an injection. Let $\pi_1 : \mathbb{R} \times \mathcal{P}(\epsilon^+) \rightarrow \mathbb{R}$ be the projection onto the first coordinate. Thinking of $F_h^{J,\epsilon}$ as a subset of $\mathbb{R} \times \mathcal{P}(\epsilon^+)$, $\pi_1[\Phi[\epsilon^+]]$ is a wellorderable subset of $\mathbb{R}$. Since AD implies boldface GCH at ω (Fact 2.6), $\pi_1[\Phi[\epsilon^+]]$ is countable. By Fact 2.16, $\text{cof}(\epsilon^+) > \omega$. Thus there is an $s \in \mathbb{R}$ so that $|(\pi_1 \circ \Phi)^{-1}[\{s\}]| = |\epsilon^+|$. Then $\Phi : (\pi_1 \circ \Phi)^{-1}[\{s\}] \rightarrow \{s\} \times G_{h(s),s}^{J,\epsilon}$ is an injection. As noted above, $|G_{h(s),s}^{J,\epsilon}| = |\mathcal{P}(\epsilon)|$. Thus this implies there is an injection of ϵ^+ into $\mathcal{P}(\epsilon)$ which violates boldface GCH at ϵ . The last statement follows from the fact that ϵ^+ injects into $\mathcal{P}_B(\epsilon^+)$. $\square$

When $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$ (or Uniformization) holds, the cardinality of $F_h^{J,\epsilon}$ is simply $|\mathcal{P}(\epsilon)|$.

Fact 6.25. Assume $\text{AD}_{\frac{1}{2}\mathbb{R}}$. Let J be a set of ordinals, $\omega \leq \epsilon \in \text{ON}$, and $h : \mathbb{R} \rightarrow \omega_1$ be a J -invariant everywhere J -increasing function. Then $|F_h^{J,\epsilon}| = |\mathcal{P}(\epsilon)|$.

Proof. As shown above, $|\mathcal{P}(\epsilon)| = |G_{h(r),r}^{J,\epsilon}|$ for any $r \in \mathbb{R}$. Thus $|\mathcal{P}(\epsilon)| \leq |F_h^{J,\epsilon}|$. By Fact 2.19, $(\epsilon^{+(h(r)+1)})^{L[J,r]} < \epsilon^+$ for any $r \in \mathbb{R}$. Thus for each $r \in \mathbb{R}$, there is a bijection $(\epsilon^{+(h(r)+1)})^{L[J,r]}$ onto ϵ . By the Moschovakis coding lemma (Fact 2.15), let $\varpi : \mathbb{R} \rightarrow \mathcal{P}(\epsilon^+)$ be a surjection. Define $R \subseteq \mathbb{R} \times \mathbb{R}$ by $R(r, s)$ if and only if $\varpi(s)$ codes the graph of a bijection from $(\epsilon^{+(h(r)+1)})^{L[J,r]}$ to ϵ . Note that $\text{dom}(R) = \mathbb{R}$. By $\text{AC}_{\mathbb{R}}^{\mathbb{R}}$, let $\hat{\Lambda} : \mathbb{R} \rightarrow \mathbb{R}$ be a such that $R(r, \hat{\Lambda}(r))$ for all $r \in \mathbb{R}$. Let $\Lambda(r) : (\epsilon^{+(h(r)+1)})^{L[J,r]} \rightarrow \epsilon$ be the bijection coded by $\hat{\Lambda}(r)$. Define $\Phi : F_h^{J,\epsilon} \rightarrow \mathbb{R} \times \mathcal{P}(\epsilon)$ by $\Phi(r, A) = (r, \Lambda[A])$. Φ is an injection and therefore, $|F_h^{J,\epsilon}| \leq |\mathbb{R} \times \mathcal{P}(\epsilon)| = |\mathcal{P}(\epsilon)|$. $\square$

Fact 6.26. Let $h_0, h_1 : \mathbb{R} \rightarrow \omega_1$ be two J -invariant everywhere J -increasing functions such that $h_0 \leq_{\mu_J} h_1$. Then $|F_{h_0}^{J,\epsilon}| \leq |F_{h_1}^{J,\epsilon}|$. Thus if $h_0 \sim_{\mu_J} h_1$, then $|F_{h_0}^{J,\epsilon}| = |F_{h_1}^{J,\epsilon}|$.

Proof. Let $B = \{x \in \mathbb{R} : h_0(x) \leq h_1(x)\}$. Since $h_0 \leq_{\mu_J} h_1$, there is a $u \in \mathbb{R}$ so that $\{x \in \mathbb{R} : u \leq_J x\} \subseteq B$. By Fact 6.21, there is a J -pointed perfect tree p so that $[p] \subseteq B$. An $s \in p$ is called a split node of p if and only if $s^{\wedge} 0 \in p$ and $s^{\wedge} 1 \in p$. For $n \in \omega$, s is called an n -split node of p if and only if there are exactly n -many proper initial segment of s which are split nodes of p . Let $\text{split}_n(p)$ be the collection of n -split nodes of p . Let $\Xi_p : {}^{<\omega}2 \rightarrow \bigcup_{n \in \omega} \text{split}_n(p)$ be defined as follows: $\Xi_p(\emptyset)$ is the shortest split node node of p . If $\Xi_p(s)$ has been defined and $i \in 2$, then let $\Xi_p(s^{\wedge} i)$ be the shortest split node of p which extends $\Xi_p(s)^{\wedge} i$. Let $\Upsilon_p : {}^{<\omega}2 \rightarrow [p]$ be defined by $\Upsilon_p(f) = \bigcup_{n \in \omega} \Xi_p(f \upharpoonright n)$. Since p is J -pointed and $\Upsilon_p(f) \in [p]$, $p \leq_J \Upsilon_p(f)$. Since f can be recovered from $\Upsilon_p(f)$ and p by a simple computable manner, $f \leq_J \Upsilon_p(f)$.

Define a function Φ on $F_{h_0}^{J,\epsilon}$ by $\Phi(r, A) = \Phi(\Upsilon_p(r), A)$. Φ is clearly an injective function. Let $(r, A) \in F_{h_0}^{J,\epsilon}$. This means that $A \in G_{h(r),r}^{J,\epsilon}$. Thus $A = \mathfrak{G}_f$ where $f : \delta \rightarrow (\epsilon^{+(h_0(r)+1)})^{L[J,r]}$ and $\delta \leq (\epsilon^{+h_0(r)})^{L[J,r]}$. Since h_0 is everywhere J -increasing, $r \leq_J \Upsilon_p(r)$, and $\Upsilon_p(r) \in [p] \subseteq B$, one has that $h_0(r) \leq h_0(\Upsilon_p(r)) \leq h_1(\Upsilon_p(r))$. Since $r \leq_J \Upsilon_p(r)$, $L[J, r] \subseteq L[J, \Upsilon_p(r)]$. Thus $\delta \leq (\epsilon^{+h_0(r)})^{L[J,r]} \leq (\epsilon^{+h_0(r)})^{L[J, \Upsilon_p(r)]} \leq (\epsilon^{+h_1(r)})^{L[J, \Upsilon_p(r)]}$. Similarly, $(\epsilon^{+(h_0(r)+1)})^{L[J,r]} \leq (\epsilon^{+(h_1(r)+1)})^{L[J, \Upsilon_p(r)]}$. Thus $A = \mathfrak{G}_f \in G_{h_1(\Upsilon_p(r)), \Upsilon_p(r)}^{J,\epsilon}$. So $(\Upsilon_p(r), A) \in F_{h_1}^{J,\epsilon}$. $\Phi : F_{h_0}^{J,\epsilon} \rightarrow F_{h_1}^{J,\epsilon}$ is an injection and hence $|F_{h_0}^{J,\epsilon}| \leq |F_{h_1}^{J,\epsilon}|$. $\square$

In contrast to Fact 6.25, if $V = L(\mathbb{R})$ and J is a set of ordinal such that $L[J] = \text{HOD}$, then $F_h^{J,\epsilon}$ will represent a cardinality intermediate between $\mathcal{P}(\epsilon)$ and $\mathcal{P}_B(\epsilon^+)$. By Fact 4.20, $\text{HOD}^{L(\mathbb{R})} = L[\vec{\mathbb{B}}^{\omega; \emptyset}]$ so $\vec{\mathbb{B}}^{\omega; \emptyset}$ is an example of such a set of ordinals J .

Theorem 6.27. Assume AD and $V = L(\mathbb{R})$. Let J be a set of ordinals so that $\text{HOD} = L[J]$, $\omega \leq \epsilon < \Theta$, and $h_0, h_1 : \mathbb{R} \rightarrow \omega_1$ be two J -invariant everywhere J -increasing functions such that $h_0 <_{\mu_J} h_1$, then $|F_{h_0}^{J,\epsilon}| < |F_{h_1}^{J,\epsilon}|$.

Proof. By Fact 6.26, $|F_{h_0}^{J,\epsilon}| \leq |F_{h_1}^{J,\epsilon}|$. Suppose $\Phi : F_{h_1}^{J,\epsilon} \rightarrow F_{h_0}^{J,\epsilon}$ is an injection. Since $V = L(\mathbb{R})$, there is an $\tilde{x} \in \mathbb{R}$ so that Φ is $\text{OD}_{\{\tilde{x}\}}$. Since $h_0 \leq_{\mu_J} h_1$, there exists an $x \in \mathbb{R}$ so that $h_0(x) < h_1(x)$ and $\tilde{x} \leq_J x$. So $\tilde{x} \in \text{HOD}_{\{\tilde{x}\}} = \text{HOD}[\tilde{x}] = L[J][\tilde{x}] = L[J, \tilde{x}] \subseteq L[J, x] = L[J][x] = \text{HOD}[x] = \text{HOD}_{\{x\}}$ by Fact 4.19, the hypothesis on J , and $\tilde{x} \leq_J x$. Thus $\tilde{x} \in \text{OD}_{\{x\}}$. Hence $h_0(x) < h_x(x)$ and $\Phi \in \text{OD}_{\{x\}}$. Consider the forcing $\text{Coll}(\epsilon^{+h_1(x)}, \epsilon^{+(h_1(x)+1)})^{\text{HOD}_{\{x\}}}$. By Lemma 5.3, Lemma 5.5, and Fact 6.1 much like in the proof of Theorem 6.2, there is a $G \subseteq \text{Coll}(\epsilon^{+h_1(x)}, \epsilon^{+(h_1(x)+1)})^{\text{HOD}_{\{x\}}}$ which is generic over $\text{HOD}_{\{x\}}$. Let $g : (\epsilon^{+h_1(x)})^{\text{HOD}_{\{x\}}} \rightarrow (\epsilon^{+(h_1(x)+1)})^{\text{HOD}_{\{x\}}}$ be the generic surjection. Since $\text{HOD}_{\{x\}} = \text{HOD}[x] = L[J, x]$, $g : (\epsilon^{+h_1(x)})^{L[J, x]} \rightarrow (\epsilon^{+(h_1(x)+1)})^{L[J, x]}$. Thus $\mathfrak{G}_g \in G_{h_1(x), x}^{J, \epsilon}$ (using the notation of Definition 6.23). Hence $(x, \mathfrak{G}_g) \in F_{h_1}^{J, \epsilon}$. Then $\Phi(x, \mathfrak{G}_g)$ is $\text{OD}_{\{x, G\}}$. In $\text{HOD}_{\{x\}}$, this forcing is in bijection with $(\epsilon^{+(h_1(x)+1)})^{\text{HOD}_{\{x\}}}$. By Fact 5.7 of Neeman and Woodin, there is a unique supercompact measure on $\mathcal{P}_{\omega_1}((\epsilon^{+(h_1(x)+1)})^{\text{HOD}_{\{x\}}})$. (If ϵ is below a Suslin cardinal or equivalently below δ_1^2 , such supercompact measure exists by Harrington-Kechris [24] and Woodin [66].) By Lemma 5.9, $\Phi(x, \mathfrak{G}_g) \in \text{HOD}_{\{x, G\}} = \text{HOD}_{\{x\}}[G]$. Since $\Phi : F_{h_1}^{J, \epsilon} \rightarrow F_{h_0}^{J, \epsilon}$, $\Phi(x, \mathfrak{G}_g) \in F_{h_0}^{J, \epsilon} \cap \text{HOD}_{\{x\}}[G]$. Thus there is some $r \in \mathbb{R}$ and some f such that $f : \tilde{\delta} \rightarrow (\epsilon^{+(h_0(r)+1)})^{L[J, r]}$, $\tilde{\delta} \leq (\epsilon^{+h_0(r)})^{L[J, r]}$, and $\Phi(x, \mathfrak{G}_g) = (r, \mathfrak{G}_f)$. Since $(r, \mathfrak{G}_f) = \Phi(x, \mathfrak{G}_g) \in \text{HOD}_{\{x\}}[G]$, one has that $r \in \text{HOD}_{\{x\}}[G]$ and $f \in \text{HOD}_{\{x\}}[G]$ (since f is the function whose graph is $\text{gp}^{-1}[\mathfrak{G}_f]$ where gp is the Gödel pairing function). Since $\text{HOD}_{\{x\}} \models \text{“Coll}(\epsilon^{+h_1(x)}, \epsilon^{+(h_1(x)+1)}) \text{ is } < \epsilon^{+h_1(x)}\text{-closed”}$, no new functions from a $\delta < (\epsilon^{+h_1(x)})^{\text{HOD}_{\{x\}}}$ into the ordinals are added by this forcing. No new reals are added by this forcing. So $r \in \text{HOD}_{\{x\}} = L[J, x]$ and thus $r \leq_J x$. Since h_0 is everywhere J -increasing, $h_0(r) \leq h_0(x) < h_1(x)$. So $\tilde{\delta} \leq (\epsilon^{+h_0(r)})^{L[J, r]} \leq (\epsilon^{+h_0(r)})^{L[J, x]} \leq (\epsilon^{+h_0(x)})^{L[J, x]} < (\epsilon^{+(h_0(x)+1)})^{L[J, x]} \leq (\epsilon^{+h_1(x)})^{L[J, x]} = (\epsilon^{+h_1(x)})^{\text{HOD}_{\{x\}}}$. So f is a function from $\tilde{\delta} < (\epsilon^{+h_1(x)})^{\text{HOD}_{\{x\}}}$ into the ordinals. Again since no new functions from a $\delta < (\epsilon^{+h_1(x)})^{\text{HOD}_{\{x\}}}$ into the ordinals are added by this forcing, $f \in \text{HOD}_{\{x\}}$. So $\Phi(x, \mathfrak{G}_g) = (r, \mathfrak{G}_f) \in \text{HOD}_{\{x\}}$. So $(x, \mathfrak{G}_g) = \Phi^{-1}(\Phi(x, \mathfrak{G}_g))$ is $\text{OD}_{\{x\}}$ and so $(x, \mathfrak{G}_g) \in \text{HOD}_{\{x\}}$. In particular, $\mathfrak{G}_g \in \text{HOD}_{\{x\}}$. Then $g \in \text{HOD}_{\{x\}}$ since g is the function whose graph is $\text{gp}^{-1}(\mathfrak{G}_g)$. By absoluteness, $\text{HOD}_{\{x\}} \models \text{“}g : \epsilon^{+h_1(x)} \rightarrow \epsilon^{+(h_1(x)+1)} \text{ is a surjection”}$. Contradiction. $\square$

If $X \in \mathcal{D}_J$, defined $L[J, X]$ to be $L[J, x]$ for any $x \in X$. Note that this is well defined independent of the choice of $x \in X$. (However, the standard global constructibility ordering of $L[J, x]$ does depend on x .) Since $L(\mathbb{R}) \models \text{DC}$, $\prod_{\mathcal{D}_J} \omega_1 / \mu_J$ is wellfounded.

Fact 6.28. (Woodin) Assume AD. Let J be a set of ordinals. $\prod_{X \in \mathcal{D}_J} \omega_1^{L[J, X]} / \mu_J = \omega_1$.

(Woodin) Assume $V = L(\mathbb{R})$ and J is a set of ordinals such that $\text{HOD} = L[J]$, then $\prod_{X \in \mathcal{D}_J} \omega_2^{L[J, X]} = \Theta$.

Proof. The first statement is [42] Theorem 9.6.

Now suppose $V = L(\mathbb{R})$ and J is a set of ordinals such that $\text{HOD} = L[J]$. Let $A \in \mathcal{P}(\mathbb{R})$. A is $\text{OD}_{\{x\}}$ for some $x \in \mathbb{R}$. For any $y \in \mathbb{R}$ such that $x \leq_J y$, $x \in L[J, y] = L[J][y] = \text{HOD}[y] = \text{HOD}_{\{y\}}$ by Fact 4.19. So for any y such that $x \leq_J y$, A is $\text{OD}_{\{y\}}$. Then for all $y \geq_J x$, $A \cap \text{HOD}_{\{y\}}$ is $\text{OD}_{\{y\}}$ and hence $A \cap \text{HOD}_{\{y\}} \in \text{HOD}_{\{y\}}$. For all $y \geq_J x$, $A \cap L[J, y] \in L[J, y]$ since $L[J, y] = \text{HOD}_{\{y\}}$. This verifies that J is a strongly maximal set of ordinals in the terminology of [42]. The result now follow from [42] Corollary 9.26 and Corollary 9.27. $\square$

If $h : \mathbb{R} \rightarrow \text{ON}$ is a J -invariant function, let $\bar{h} : \mathcal{D}_J \rightarrow \text{ON}$ be defined by $\bar{h}([x]_{\equiv_J}) = h(x)$ and note this is well defined by J -invariance. For each $\alpha \in \prod_{\mathcal{D}_J} \omega_1 / \mu_J$, there is a J -invariant everywhere J -increasing function $h : \mathbb{R} \rightarrow \omega_1$ so that $[\bar{h}]_{\sim_{\mu_J}} = \alpha$ by Fact 6.22. Thus for each $\alpha \in \prod_{\mathcal{D}_J} \omega_1 / \mu_J$, let $\mathfrak{F}_\alpha^{J, \epsilon} = |F_h^{J, \epsilon}|$ for any J -invariant everywhere J -increasing function so that $[\bar{h}]_{\sim_{\mu_J}} = \alpha$. $\mathfrak{F}_\alpha^{J, \epsilon}$ is independent of the choice of such h by Fact 6.26. Assume $V = L(\mathbb{R})$ and let J be a set of ordinals such that $L[J] = \text{HOD}$. By Fact 6.24 and Theorem 6.27, the map Σ on $\prod_{\mathcal{D}_J} \omega_1 / \mu_J$ defined by $\Sigma(\alpha) = \mathfrak{F}_\alpha^{J, \epsilon}$ is an order preserving injection of $\prod_{\mathcal{D}_J} \omega_1 / \mu$ with the ultrapower ordering (which is isomorphic to an ordinal larger than Θ by Fact 6.28) into the cardinalities between $\mathcal{P}(\epsilon)$ and $\mathcal{P}_B(\epsilon^+)$ under the cardinality injection ordering.

The ABCD conjecture merely distinguished the cardinality of different cardinal exponentiations. This section will focus on deeper questions concerning cardinality and combinatorics for ω -sequences of ordinals which correspond to the smallest nonwellorderable cardinal exponentiation. The techniques introduced here seem very flexible for resolving many questions concerning ω -sequences of ordinals including primeness, Jónssonness, and basis for linear orderings which will be addressed in forthcoming work.

First, one will consider the question of whether ${}^\omega\kappa$ injects into $\mathcal{P}(\delta) \times \text{ON}$ when $\delta < \kappa$. Recall that if $\delta < \kappa$, then Fact 2.10 says that $\kappa \rightarrow_* (\kappa)_\delta^\omega$ and more directly the δ^+ -completeness of μ_κ^ω imply that ${}^\omega\kappa$ does not inject into $\mathcal{P}(\delta) \times \text{ON}$. However, ω_3 is a singular cardinal and hence does not possess any partition properties. Another approach to the question of whether ${}^\omega\kappa$ injects into $\mathcal{P}(\delta) \times \text{ON}$ is through a combinatorial property called primeness investigated under determinacy in [16].

Definition 7.1. Let A be a set. A is said to be prime if and only if for all sets B and C , if $|A| \leq |B \times C|$, then $|A| \leq |B|$ or $|A| \leq |C|$ (that is, if A injects into $B \times C$, then A already injects into B or C).

Fact 7.2. (Chan-Jackson-Trang; [16]) Assume $\text{AC}_\omega^\mathbb{R}$ and all subsets of $\mathbb{R}$ have the Baire property. $\mathbb{R}$ and $\mathbb{R}/E_0$ are prime.

Fact 7.3. (Chan-Jackson-Trang; [16]) Assume $\kappa \rightarrow_* (\kappa)_2^{\omega+\omega}$. ${}^\omega\kappa$ is prime.

Fact 7.4. (Chan-Jackson-Trang; [16]) Assume AD. For all $n \in \omega$, ${}^\omega(\omega_n)$ is prime. ${}^\omega(\omega_\omega)$ is prime.

Fact 7.5. Assume boldface GCH holds below κ , ${}^\omega\kappa$ is not wellorderable¹⁹, and ${}^\omega\kappa$ is prime. Then ${}^\omega\kappa$ does not inject into $\mathcal{P}(\delta) \times \text{ON}$ for all $\delta < \kappa$.

Proof. Suppose $\Phi : {}^\omega\kappa \rightarrow \mathcal{P}(\delta) \times \text{ON}$ is an injection. By replacement, there is a $\lambda \in \text{ON}$ so that $\Phi : {}^\omega\kappa \rightarrow \mathcal{P}(\delta) \times \lambda$. By primeness, ${}^\omega\kappa$ injects into $\mathcal{P}(\delta)$ or λ . Since $\delta^+ \leq \kappa$, $|\kappa| \leq |{}^\omega\kappa|$, and boldface GCH holds at δ , ${}^\omega\kappa$ cannot inject into $\mathcal{P}(\delta)$. Since ${}^\omega\kappa$ is not wellorderable, ${}^\omega\kappa$ cannot inject into λ . Contradiction. $\square$

Fact 7.6. Assume AD. For all $1 \leq n < \omega$, ${}^\omega(\omega_n)$ does not inject into $\mathcal{P}(\omega_m) \times \text{ON}$ for all $m < n$. ${}^\omega(\omega_\omega)$ does not inject into $\mathcal{P}(\omega_n) \times \text{ON}$ for all $n < \omega$.

Proof. This follows from Fact 7.4 and Fact 7.5. $\square$

It will be shown next using the methods from the previous sections that for all $\kappa < \Theta$, ${}^\omega\kappa$ does not inject into $\mathcal{P}(\delta) \times \text{ON}$ for all $\delta < \kappa$. The basic template for this argument is Theorem 3.5.

Definition 7.7. An measure sequence $\mathbf{u}$ consists of $\vec{\mu}^\mathbf{u} = \langle \mu_n^\mathbf{u} : n \in \omega \rangle$, $\vec{\kappa}^\mathbf{u} = \langle \kappa_n^\mathbf{u} : n \in \omega \rangle$, and $\lambda^\mathbf{u}$ such that for all $n \in \omega$, $\omega < \kappa_n^\mathbf{u}$, $\vec{\kappa}^\mathbf{u}$ is a strictly increasing sequence, $\mu_n^\mathbf{u}$ is a $\kappa_n^\mathbf{u}$ -complete uniform normal measure on $\kappa_n^\mathbf{u}$, and $\lambda^\mathbf{u} = \sup \vec{\kappa}^\mathbf{u} = \sup \{\kappa_n^\mathbf{u} : n \in \omega\}$.

Definition 7.8. Let $\mathbf{u}$ be a measure sequence. A $\mathbf{u}$ -Namba condition is a function $p \in \prod_{n \in \omega} \kappa_n^\mathbf{u} \cup \mu_n^\mathbf{u}$ with the following properties.

- There is an $m \in \omega$ so that for all $i < m$, $p(i) \in \kappa_i^\mathbf{u}$ and for all $i \geq m$, $p(i) \in \mu_i^\mathbf{u}$. The unique such m called the stem-length of p and is denoted $\text{length}(p)$. (In other words, if $i < \text{length}(p)$, then $p(i)$ is an ordinal below $\kappa_i^\mathbf{u}$. If $\text{length}(p) \leq i$, then $p(i)$ is a subset of $\kappa_i^\mathbf{u}$ which belongs to the measure $\mu_i^\mathbf{u}$.)
- $p \upharpoonright \text{length}(p)$ is called the stem of p and is denoted $\text{stem}(p)$.

Let $\mathbb{N}_\mathbf{u}$ denote the collection of all $\mathbf{u}$ -Namba conditions. Define $\leq_{\mathbb{N}_\mathbf{u}}$ on $\mathbb{N}_\mathbf{u}$ as follows. If $p, q \in \mathbb{N}_\mathbf{u}$, then $p \leq_{\mathbb{N}_\mathbf{u}} q$ if and only if the following holds.

- $\text{length}(p) \geq \text{length}(q)$.
- $\text{stem}(q) = \text{stem}(p) \upharpoonright \text{length}(q)$.
- For all $\text{length}(q) \leq i < \text{length}(p)$, $p(i) \in q(i)$.
- For all $\text{length}(p) \leq i < \omega$, $p(i) \subseteq q(i)$.

¹⁹If $\kappa > \omega$, then boldface GCH below κ implies boldface GCH at ω . This implies that $\mathbb{R}$ is not wellorderable. Since $|\mathbb{R}| \leq |{}^\omega\kappa|$, ${}^\omega\kappa$ is not wellorderable.

2034 Let $1_{\mathbb{N}_u}$ be the function defined by $1_{\mathbb{N}_u}(n) = \kappa_n^u$ for all $n \in \omega$. $\mathbb{N}_u = (\mathbb{N}_u, \leq_{\mathbb{N}_u}, 1_{\mathbb{N}_u})$ is u -Namba forcing.

2035 For $k \in \omega$ and $p, q \in \mathbb{N}_u$, let $p \leq_u^k q$ if and only if $\text{stem}(p) = \text{stem}(q)$ and for all $i < k$, $p(i) = q(i)$. A
 2036 sequence $\langle p_k : k \in \omega \rangle$ such that $p_{k+1} \leq_u^k p_k$ for all $k \in \omega$ is a u -fusion sequence. The fusion of the u -fusion
 2037 sequence $\langle p_k : k \in \omega \rangle$ is p_ω defined by $p_\omega(i) = p_{i+1}(i)$. It is clear that p_ω is a u -Namba condition and
 2038 $\text{stem}(p_\omega) = \text{stem}(p_k)$ for any $k \in \omega$. Note that for all $i \in \omega$, $p_\omega \leq_u^i p_i$.

If $s \in {}^{<\omega}(\lambda^u)$ and $p \in \mathbb{N}_u$, then say that $s \in^u p$ if and only if for all $i < \text{length}(p)$, $s(i) = p(i)$ and for all
 $\text{length}(p) \leq i < |s|$, $s(i) \in p(i)$. Define the body of p , denoted $[p]$, to be the set of $f \in {}^\omega(\lambda^u)$ such that for
 all $n \in \omega$, $f \restriction n \in^u p$. If $p \in \mathbb{N}_u$ and $s \in^u p$, then let $(p)_s \in \mathbb{N}_u$ be defined by

$$(p)_s(i) = \begin{cases} s(i) & i < |s| \\ p(i) & |s| \leq i \end{cases}.$$

2039 Note that if $|s| \leq \text{length}(p)$, then $\text{stem}((p)_s) = s$ and $(p)_s = p$, and if $|s| > \text{length}(p)$, then $\text{stem}((p)_s) = s$.

2040 Let F is a $\mathbb{N}_u$ -filter and $p, q \in G$. There is an $r \in F$ so that $r \leq_{\mathbb{N}_u} p$ and $r \leq_{\mathbb{N}_u} q$. Thus for all
 2041 $i < \min\{\text{length}(p), \text{length}(q)\}$, $p(i) = r(i) = q(i)$. Let $f_F = \bigcup \{\text{stem}(p) : p \in F\}$. So there is an $\alpha \leq \omega$ so
 2042 that $f_F : \alpha \rightarrow \lambda^u$. Let $\dot{f}_G$ be the canonical $\mathbb{N}_u$ -name for the ω -sequence added by $\dot{G}$, which is the name for
 2043 the $\mathbb{N}_u$ -generic filter.

2044 *Remark 7.9.* Namba forcing typically refers to forcings whose conditions are trees with some properties. $\mathbb{N}_u$
 2045 more closely resembles some form of Prikry forcing along ω many measures on an increasing ω -sequence of
 2046 measurable cardinals. $\mathbb{N}_u$ is essentially equivalent to diagonal Prikry forcing. $\mathbb{N}_u$ is an instance of a more
 2047 general Namba forcing developed by Cox and Krueger in [23]. Since here u is an increasing sequence of
 2048 measurable cardinals along with its sequence of measures, one has simplified the presentation by removing
 2049 the trees. This Prikry-style presentation seems notationally more practical for handling higher dimensional
 2050 Namba forcing which will be used in forthcoming work.

2051 In [23], a u -Namba-tree is a $T \subseteq {}^{<\omega}(\lambda^u)$ so that for all $s \in T$, $\text{succ}_T(s) = \{\alpha \in \kappa_{|s|}^u : s \hat{\ } \langle \alpha \rangle \in T\} \in \mu_{|s|}^u$.
 2052 Let $\bar{\mathbb{N}}_u$ be the Cox-Krueger u -Namba forcing consisting of u -Namba trees ordered by the subset relation. If
 2053 $p \in \mathbb{N}_u$ is a u -Namba condition, then $T_p = \{s \in {}^{<\omega}(\lambda^u) : s \in^u p\}$ is a u -Namba tree. The map $\Phi : \mathbb{N}_u \rightarrow \bar{\mathbb{N}}_u$
 2054 defined by $\Phi(p) = T_p$ is a dense embedding of forcing posets. Lemma 7.11 and Fact 7.13 corresponds to [23]
 2055 Lemma 6.4 and [23] Proposition 6.14, respectively.

2056 **Fact 7.10.** *Let $M \models \text{ZF}$ be an inner model and $u \in M$ such that $M \models "u \text{ is a measure sequence}"$. Suppose
 2057 $G \subseteq \mathbb{N}_u$ is $\mathbb{N}_u$ -generic over M . Then $f_G : \omega \rightarrow \lambda^u$ is cofinal through λ^u and $f_G \notin M$.*

2058 *Proof.* For each $n \in \omega$, let $D_n = \{p \in \mathbb{N}_u : \text{length}(p) > n\}$. D_n is dense in $\mathbb{N}_u$. If $G \cap D_n \neq \emptyset$ for all $n \in \omega$,
 2059 then $\text{dom}(f_G) = \text{dom}(\bigcup \{\text{stem}(p) : p \in G\}) = \omega$.

For each $n \in \omega$, let $E_n = \{p \in \mathbb{N}_u : \text{length}(p) > n \wedge (\forall i \geq \text{length}(p))(p(i) \subseteq \kappa_i^u \setminus \kappa_n^u)\}$. Let $q \in \mathbb{N}_u$. Let
 $k = \max\{n, \text{length}(q)\}$. Pick $s \in^u q$ with $|s| = k$. Define $p \in \mathbb{N}_u$ as follows:

$$p(i) = \begin{cases} s(i) & i < k \\ q(i) \setminus \kappa_n^u & k \leq i \end{cases}$$

2060 Note that $p \in E_n$ and $p \leq_{\mathbb{N}_u} q$. Thus E_n is dense. Suppose $p \in G \cap E_n$. For all $i \geq \text{length}(p)$ and
 2061 all $s \in^u p$, $s(i) \geq \kappa_n$. Thus $f_G(i) \geq \kappa_n$ for all $i \geq \text{length}(p)$. Thus if $G \cap E_n \neq \emptyset$ for all $n \in \omega$, then
 2062 $\sup(f_G) = \sup\{\kappa_n^u : n \in \omega\} = \lambda^u$.

2063 For each $h \in {}^\omega(\lambda^u)$, let $F_h = \{p \in \mathbb{N}_u : (\exists n < \text{length}(p))(\text{stem}(p)(n) \neq h(n))\}$. Suppose $q \in \mathbb{N}_u$. Since
 2064 $q(\text{length}(q)) \in \mu_{\text{length}(q)}^u$, there exists $\alpha \in q(\text{length}(q))$ such that $h(\text{length}(q)) \neq \alpha$. Let $p \in \mathbb{N}_u$ be defined
 2065 by $p(i) = q(i)$ for $i < \text{length}(q)$, $p(\text{length}(q)) = \alpha$, and $p(i) = q(i) \setminus (\alpha + 1)$ for $i > \text{length}(q)$. Then $p \leq_{\mathbb{N}_u} q$
 2066 and $q \in F_h$. This shows that F_h is a dense subset of $\mathbb{N}_u$. If $G \cap F_h$, then $f_G \neq h$. Thus if $G \cap F_h$ for all
 2067 $h \in ({}^\omega(\lambda^u))^M$, then $f_G \neq h$ for any $h \in ({}^\omega(\lambda^u))^M$. $\square$

2068 **Lemma 7.11.** *Assume AC. Let u be a measure sequence, $D \subseteq \mathbb{N}_u$ be a dense open subset of $\mathbb{N}_u$, and $p \in \mathbb{N}_u$.
 2069 Then there is a $q \leq_{\mathbb{N}_u} p$ and a $k \in \omega$ so that $\text{length}(p) \leq k < \omega$, $\text{length}(q) = \text{length}(p)$, and for all $s \in^u q$
 2070 with $|s| = k$, $(q)_s \in D$.*

Proof. Say that $p \in \mathbb{N}_u$ is D -good if and only if there is a $q \leq_{\mathbb{N}_u} p$ and $\text{length}(p) \leq k < \omega$ so that $\text{length}(q) = \text{length}(p)$ and for all $s \in^u q$ with $|s| = k$, $(q)_s \in D$. Let $\text{bad}_D(p) = \{\alpha \in \kappa_{\text{length}(p)}^u : (p)_{\text{stem}(p) \smallfrown \langle \alpha \rangle} \text{ is not } D\text{-good}\}$.

Claim: If p is not D -good, then $\text{bad}_D(p) \in \mu_{\text{length}(p)}^u$.

To see the claim: Let $m = \text{length}(p)$. Suppose $\text{bad}_D(p) \notin \mu_m^u$. Then $p(m) \setminus \text{bad}_D(p) \in \mu_m^u$. For each $\alpha \in p(m) \setminus \text{bad}_D(p)$, $(p)_{\text{stem}(p) \smallfrown \langle \alpha \rangle}$ is D -good. Let k_α be the least k with $m+1 \leq k < \omega$ so that there exists some $r \leq_{\mathbb{N}_u} (p)_{\text{stem}(p) \smallfrown \langle \alpha \rangle}$ with the property that $\text{length}(r) = m+1$ and for all $s \in^u r$ with $|s| = k$, $(r)_s \in D$. Since $\omega < \kappa_m$ and μ_m^u is κ_m^u -complete, there is a $\bar{k}$ so that $A = \{\alpha \in p(m) \setminus \text{bad}_D(p) : k_\alpha = \bar{k}\} \in \mu_m^u$. Since AC holds, fix a wellordering $\prec$ of $\mathbb{N}_u$. For each $\alpha \in A$, let r_α be the $\prec$ -least $r \in \mathbb{N}_u$ so that $r \leq_{\mathbb{N}_u} (p)_{\text{stem}(p) \smallfrown \langle \alpha \rangle}$ with $\text{length}(r) = m+1$ and for all $s \in^u r$, if $|s| = \bar{k}$, then $(r)_s \in D$. If $i < m$, let $q(i) = p(i)$. Let $q(m) = A$. For all $m < i$, let $q(i) = \bigcap \{r_\alpha(i) : \alpha \in A\}$. Note that for all $m < i$, since $|A| = \kappa_m^u < \kappa_i^u$ and μ_i^u is κ_i^u -complete, $q(i) \in \mu_i^u$. Thus $q \in \mathbb{N}_u$ and $\text{length}(q) = m$. Let $s \in^u q$ with $|s| = \bar{k}$. Since $q(m) = A$, $s(m) \in A$. $(q)_s \leq_{\mathbb{N}_u} (r_{s(m)})_s$. So $(q)_s \in D$ since D is dense open and $(r_{s(m)})_s \in D$ by the choice of $r_{s(m)}$. This shows that $\bar{k}$ and q witness that p is D -good. Contradiction. This completes the proof of the claim.

The fact is equivalent to the assertion that every $p \in \mathbb{N}_u$ is D -good. Suppose $p \in \mathbb{N}_u$ is not D -good. Let $m = \text{length}(p)$. For $j \leq m$, let $p_j = p$. Suppose $m \leq j$ and p_j has been defined so that for all $s \in^u p_j$ with $|s| = j$, $(p_j)_s$ is not D -good and for all $i < j$, $p_{i+1} \leq_u^i p_i$. (This is true for all $j \leq m$, since $p_j = p$.) Thus for all $s \in^u p_j$ with $|s| = j$, $\text{bad}_D((p_j)_s) \in \mu_j^u$. If $i \neq j$, let $p_{j+1}(i) = p_j(i)$. Let $p_{j+1}(j) = \bigcap \{\text{bad}_D((p_j)_s) : s \in^u p_i \wedge |s| = j\}$. Note that $|s| = j$ and $s \in^u p_j$ imply that $s(i) \in \kappa_i^u$ for each $i < j$. Thus $\{s \in^u p_j : |s| = j\} \leq \kappa_{j-1}^u$. Since μ_j^u is κ_j^u -complete, $p_{j+1}(j) \in \mu_j^u$. This shows that $p_{j+1} \in \mathbb{N}_u$. Since $p_{j+1} \upharpoonright j = p_j \upharpoonright j$, $p_{j+1} \leq_u^j p_j$. Suppose $s \in^u p_{j+1}$ with $|s| = j+1$. $s \upharpoonright j \in^u p_j$ and $s(j) \in p_{j+1}(j) \subseteq \text{bad}_D((p_j)_{s \upharpoonright j})$. Thus $(p_{j+1})_s$ is not D -good. This completes the construction of the u -fusion sequence $\langle p_n : n \in \omega \rangle$. Let p_ω be the fusion of $\langle p_n : n \in \omega \rangle$ defined by $p_\omega(i) = p_{i+1}(i)$. Since D is dense, there is a $q \leq_{\mathbb{N}_u} p_\omega$ with $q \in D$. Let $m^* = \text{length}(q)$ and $s^* = \text{stem}(q)$. Since $p_\omega \leq_u^{m^*} p_{m^*}$, $s^* \in^u p_{m^*}$, and $q \leq_{\mathbb{N}_u} (p_\omega)_{s^*} = (p_{m^*})_{s^*}$, one has that q and $m^* = |s^*| = \text{length}((p_{m^*})_{s^*})$ witness that $(p_{m^*})_{s^*}$ is D -good. However, by construction of p_{m^*} , $(p_{m^*})_s$ is not D -good for any $s \in^u p_{m^*}$ with $|s| = m^*$. Contradiction. This completes the proof. $\square$

Lemma 7.12. Assume AC. Let u be a measure sequence, $p \in \mathbb{N}_u$, and φ be a sentence in the forcing language of $\mathbb{N}_u$. Then exactly one of the following holds.

- (1) There is a $q \leq_{\mathbb{N}_u} p$ with $\text{stem}(p) = \text{stem}(q)$ and $q \Vdash_{\mathbb{N}_u} \varphi$.
- (2) There is a $q \leq_{\mathbb{N}_u} p$ with $\text{stem}(p) = \text{stem}(q)$ and $q \Vdash_{\mathbb{N}_u} \neg \varphi$.

Proof. Let $D = \{p \in \mathbb{N}_u : p \Vdash_{\mathbb{P}} \varphi \vee p \Vdash_{\mathbb{P}} \neg \varphi\}$ which is a dense open subset of $\mathbb{N}_u$. By Lemma 7.11, there is a $k \in \omega$ with $\text{length}(p) \leq k < \omega$ and a $q \leq_{\mathbb{N}_u} p$ with $\text{length}(p) = \text{length}(q)$ so that for all $s \in^u q$ with $|s| = k$, $(q)_s \in D$. Let $\bar{k}$ be the least such k . Fix a $\bar{q} \leq_{\mathbb{N}_u} p$ so that $\text{length}(p) = \text{length}(\bar{q})$ and for all $s \in^u \bar{q}$ with $|s| = \bar{k}$, $(\bar{q})_s \in D$. The claim is that $\bar{k} = \text{length}(p) = \text{length}(\bar{q})$. Suppose otherwise that $\bar{k} > \text{length}(\bar{q})$. Since $\text{length}(\bar{q}) < \bar{k}$, $q(\bar{k}-1) \in \mu_{\bar{k}-1}^u$. For each $s \in^u \bar{q}$ with $|s| = \bar{k}-1$ and $\alpha \in q(\bar{k}-1)$, one has that $(\bar{q})_{s \smallfrown \langle \alpha \rangle} \Vdash_{\mathbb{N}_u} \varphi$ or $(\bar{q})_{s \smallfrown \langle \alpha \rangle} \Vdash_{\mathbb{N}_u} \neg \varphi$ by the property of $\bar{k}$ and $\bar{q}$. Let $A_s^0 = \{\alpha \in q(\bar{k}-1) : (q)_{s \smallfrown \langle \alpha \rangle} \Vdash_{\mathbb{N}_u} \neg \varphi\}$ and $A_s^1 = \{\alpha \in q(\bar{k}-1) : (q)_{s \smallfrown \langle \alpha \rangle} \Vdash_{\mathbb{N}_u} \varphi\}$. Since $A_s^0 \cap A_s^1 = \emptyset$ and $A_s^0 \cup A_s^1 = q(\bar{k}-1) \in \mu_{\bar{k}-1}^u$, there is a unique $i_s \in 2$ so that $A_s^{i_s} \in \mu_{\bar{k}-1}^u$. Define $\underline{q}$ as follows: For $i < \bar{k}-1$, $\underline{q}(i) = \bar{q}(i)$. Let $\underline{q}(\bar{k}-1) = \bigcap \{A_s^{i_s} : s \in^u \bar{q} \wedge |s| = \bar{k}-1\}$. For $\bar{k}-1 < i < \omega$, let $\underline{q}(i) = \bar{q}(i)$. Since $\mu_{\bar{k}-1}^u$ is $\kappa_{\bar{k}-1}^u$ -complete and $|\{s \in^u \bar{q} : |s| = \bar{k}-1\}| < \kappa_{\bar{k}-1}^u$, $\underline{q}(\bar{k}-1) \in \mu_{\bar{k}-1}^u$. Thus $\underline{q} \in \mathbb{N}_u$ and $\text{length}(\underline{q}) = \text{length}(p)$. Let $s \in^u \underline{q}$ with $|s| = \bar{k}-1$. Without loss of generality, suppose $i_s = 1$. One seeks to show that $(q)_s \Vdash_{\mathbb{N}_u} \varphi$. Suppose not. Then one can find an $r \leq_{\mathbb{N}_u} (q)_s$ so that $\text{length}(r) \geq \bar{k}$ and $r \Vdash_{\mathbb{N}_u} \neg \varphi$. However $r(\bar{k}-1) \in \underline{q}(\bar{k}-1) \subseteq A_s^1$ and $r \leq_{\mathbb{N}_u} (\underline{q})_{s \smallfrown \langle r(\bar{k}-1) \rangle}$. So $(\underline{q})_{s \smallfrown \langle r(\bar{k}-1) \rangle} \Vdash_{\mathbb{N}_u} \varphi$. Thus $r \Vdash_{\mathbb{N}_u} \varphi$. Contradiction. This means that $(q)_s \in D$. Since $s \in^u \underline{q}$ with $|s| = \bar{k}-1$ was arbitrary, this violates the minimality of $\bar{k}$. This shows that $\bar{k} = \text{length}(\bar{q})$. The only $s \in^u \bar{q}$ with $|s| = \text{length}(\bar{q})$ is $\text{stem}(\bar{q})$. This means $\bar{q} = (\bar{q})_{\text{stem}(\bar{q})} \in D$. Thus $\bar{q} \Vdash_{\mathbb{N}_u} \varphi$ or $\bar{q} \Vdash_{\mathbb{N}_u} \neg \varphi$. $\bar{q}$ is the desired extension of q .

This shows one of the two cases must occur. Suppose both occurred. Let $q_0, q_1 \in \mathbb{N}_u$ be such that $q_0 \leq_{\mathbb{N}_u} p$, $q_1 \leq_{\mathbb{N}_u} p$, $\text{stem}(p) = \text{stem}(q_0) = \text{stem}(q_1)$, $q_0 \Vdash_{\mathbb{N}_u} \neg \varphi$, and $q_1 \Vdash_{\mathbb{N}_u} \varphi$. Let q_2 be defined by $q_2(i) = p(i)$ for all

2120 $i < \text{length}(p)$ and $q_2(i) = q_0(i) \cap q_1(i)$ for all $\text{length}(p) \leq i < \omega$. Then $q_2 \in \mathbb{N}_u$, $q_2 \leq_{\mathbb{N}_u} q_0$, and $q_2 \leq_{\mathbb{N}_u} q_0$.
 2121 Thus $q_2 \Vdash_{\mathbb{N}_u} \varphi$ and $q_2 \Vdash_{\mathbb{N}_u} \neg\varphi$. Contradiction. $\square$

2122 **Fact 7.13.** Assume AC. Let u be a measure sequence. For any $\delta < \lambda^u$, $1_{\mathbb{N}_u} \Vdash_{\mathbb{N}_u} \mathcal{P}(\check{\delta}) = \check{\mathcal{P}}(\delta)$. (Thus $\mathbb{N}_u$
 2123 does not add any new bounded subsets of λ^u .)

2124 *Proof.* Using AC, let $\prec$ be a wellordering of $\mathbb{N}_u$. Let $\delta < \lambda^u$. Let $\tilde{p} \in \mathbb{N}_u$ and $\dot{A}$ be a $\mathbb{N}_u$ -name such that
 2125 $\tilde{p} \Vdash_{\mathbb{N}_u} \dot{A} \subseteq \check{\delta}$. Let $D_{\tilde{p}, \dot{A}}$ be the set of $p \in \mathbb{N}_u$ such that the disjunction of the following holds.

- 2126 (1) p is incompatible with $\tilde{p}$.
 2127 (2) $p \leq_{\mathbb{N}_u} \tilde{p}$ and there exists a $B \in \mathcal{P}(\delta)$ so that $p \Vdash_{\mathbb{N}_u} \dot{A} = \check{B}$.

One will show that $D_{\tilde{p}, \dot{A}}$ is dense. Let $\bar{p} \in \mathbb{N}_u$ be arbitrary. If $\bar{p}$ is incompatible with $\tilde{p}$, then $\bar{p} \in D_{\tilde{p}, \dot{A}}$. Now
 suppose $\bar{p}$ is compatible with $\tilde{p}$. Let $\hat{p}$ be such that $\hat{p} \leq_{\mathbb{N}_u} \bar{p}$ and $\hat{p} \leq_{\mathbb{N}_u} \tilde{p}$. Let $\bar{n}$ be the least n such that
 $\text{length}(\hat{p}) \leq n$ and $\delta < \kappa_n^u$. Fix $\eta < \delta$. By Lemma 7.12, there is a $q \leq_u \hat{p}$ with $\text{length}(q) = \bar{n}$ and an $i_\eta \in 2$
 so that $q \Vdash_{\mathbb{N}_u} \dot{A}(\check{\eta}) = \check{i}_\eta$ (here $\dot{A}$ is regarded as a name for the characteristic function of a subset of δ). Let
 q_η be the $\prec$ -least $q \in \mathbb{N}_u$ so that $\text{length}(q) = \bar{n}$, $q \leq_{\mathbb{N}_u} \hat{p}$, and $q \Vdash_{\mathbb{N}_u} \dot{A}(\check{\eta}) = \check{i}_\eta$. Let $B = \{\eta < \delta : i_\eta = 1\}$.
 Define $\underline{q}$ by

$$\underline{q}(i) = \begin{cases} \hat{p}(i) & i < \bar{n} \\ \bigcap \{q_\eta(i) : \eta < \delta\} & \bar{n} \leq i \end{cases}.$$

2128 By choice of $\bar{n}$, $\delta < \kappa_{\bar{n}}^u$. For all $i \geq \bar{n}$, $\kappa_i^u \geq \kappa_{\bar{n}}^u$ and μ_i^u is κ_i^u -complete. Therefore, $\underline{q}(i) \in \mu_i^u$ for all $\bar{n} \leq i$.
 2129 This shows that $\underline{q} \in \mathbb{N}_u$, $\text{stem}(\underline{q}) = \bar{n}$, and $\underline{q} \leq_{\mathbb{N}_u} q_\eta$ for each $\eta < \delta$. Since $q_\eta \Vdash_{\mathbb{N}_u} \dot{A}(\check{\eta}) = \check{i}_\eta = \check{B}(\check{\eta})$, one has
 2130 that $\underline{q} \Vdash_{\mathbb{N}_u} \dot{A}(\check{\eta}) = \check{B}(\check{\eta})$ for each $\eta < \delta$. Thus $\underline{q} \leq_{\mathbb{N}_u} \tilde{p}$ and $\underline{q} \Vdash_{\mathbb{N}_u} \dot{A} = \check{B}$. Thus $\underline{q} \leq_{\mathbb{N}_u} \bar{p}$ and $\underline{q} \in D_{\tilde{p}, \dot{A}}$. Since
 2131 $\bar{p}$ was arbitrary, this shows that $D_{\tilde{p}, \dot{A}}$ is dense. Since $\tilde{p}$ and $\dot{A}$ were arbitrary with the above properties, this
 2132 shows that $1_{\mathbb{N}_u} \Vdash_{\mathbb{N}_u} \mathcal{P}(\check{\delta}) = \check{\mathcal{P}}(\delta)$. $\square$

2133 The next lemma will provide a setting in which there is an inner model M of ZFC and a $(\mathbb{N}_u)^M$ -generic filter
 2134 over M in the real world. Moreover, it will be shown that there are many such generic filters corresponding
 2135 to each element of the body of a $\mathbb{N}_u$ condition (belonging to the real world but external to M). For later
 2136 purposes, it will be stated to make the uniformity explicit.

2137 **Lemma 7.14.** Let M be an inner model of ZFC. Suppose the following holds.

- 2138 (1) $u \in M$ and $M \models$ “ u is a measure sequence, $2^{\lambda^u} = (\lambda^u)^+$, and $2^{(\lambda^u)^+} = (\lambda^u)^{++}$ ”.
 2139 (2) $\prec$ is a wellordering of all sets in M of rank less than $\lambda^u + 3$.
 2140 (3) $\tau_1 : \omega \rightarrow ((\lambda^u)^+)^M$ is an increasing cofinal function.
 2141 (4) $\tau_2 : \omega \rightarrow ((\lambda^u)^{++})^M$ is an increasing cofinal function.
 2142 (5) $p \in (\mathbb{N}_u)^M$.

2143 Then there is a $q \in \mathbb{N}_u$ (but $q \notin M$) such that $q \leq_{\mathbb{N}_u} p$, $\text{length}(p) = \text{length}(q)$, and for all $g \in [q]$,
 2144 $G_g = \{p \in (\mathbb{N}_u)^M : g \in [p]\}$ is a $(\mathbb{N}_u)^M$ -generic filter over M . q is produced uniformly from u , $\prec$, τ_1 , τ_2 ,
 2145 and p .

2146 *Proof.* Since $M \models$ “ κ_n^u is measurable (and hence strong limit)” for all $n \in \omega$, $M \models$ “ $\mathcal{P}(\kappa_n^u) < \kappa_{n+1}^u$ ”. Thus
 2147 $M \models$ “ $|\mathbb{N}_u| \leq |\mathcal{P}(\lambda^u)| = 2^{\lambda^u} = (\lambda^u)^+$ ” by the hypothesis. Hence $M \models$ “ $|\mathcal{P}(\mathbb{N}_u)| \leq 2^{(\lambda^u)^+} = (\lambda^u)^{++}$ ” by
 2148 the hypothesis. Let $\tau_0 : \omega \rightarrow \lambda^u$ be defined by $\tau_0(n) = \kappa_n^u$ (which is defined uniformly from u). By Lemma
 2149 5.4, there is a M -amenable surjection $\bar{h} : \lambda^u \rightarrow ((\lambda^u)^{++})^M$ which is produced uniformly from τ_0 , τ_1 , τ_2 ,
 2150 and $\prec$. Let $\mathcal{D}$ be the collection of dense open subsets of $(\mathbb{N}_u)^M$. Since $M \models$ “ $|\mathcal{D}| \leq |\mathcal{P}(\mathbb{N}_u)| \leq (\lambda^u)^{++}$ ”, let
 2151 $\hat{h} : (\lambda^u)^{++} \rightarrow \mathcal{D}$ be the $\prec$ -least bijection in M between $(\lambda^u)^{++}$ and $\mathcal{D}$. Then $h : \lambda^u \rightarrow \mathcal{D}$ defined by $h = \hat{h} \circ \bar{h}$
 2152 is an M -amenable surjection produced uniformly from u , $\prec$, τ_1 , and τ_2 . Fix a $p \in (\mathbb{N}_u)^M$. One will build an
 2153 external (to M) u -fusion sequence $\langle p_n : n \in \omega \rangle$ with the following properties.

- 2154 (1) $p_n \in (\mathbb{N}_u)^M$ (that is, $p_n \in M$) .
 2155 (2) $p_0 = p_1 = p$.
 2156 (3) For all $1 \leq n < \omega$, $s \in^u p_{n+1}$ with $|s| = n$, for all $\eta < \kappa_{n-1}^u$, there is an $m \geq n$ so that for all
 2157 $t \in^u (p_{n+1})_s$ with $|t| = m$, $(p_{n+1})_t \in h(\eta)$.

To construct this $\mathbf{u}$ -fusion sequence: Let $p_0 = p_1 = p$. Now suppose $1 \leq n < \omega$ and p_n has been defined. For each $s \in^{\mathbf{u}} p_n$ with $|s| = n$ and $\eta < \kappa_{n-1}^{\mathbf{u}}$, there is some $q \in (\mathbb{N}_{\mathbf{u}})^M$ and some $n \leq m < \omega$ so that $q \leq_{\mathbb{N}_{\mathbf{u}}} (p_n)_s$, $\text{stem}(q) = \text{stem}((p_n)_s) = s$, and for all $t \in^{\mathbf{u}} q$ with $|t| = m$, $(q)_t \in h(\eta)$ by Lemma 7.11 applied in M . Let $q_s^{n,\eta} \in (\mathbb{N}_{\mathbf{u}})^M$ be the $\prec$ -least such q and let $m_s^{n,\eta}$ be the least such m with the above property for $q_s^{n,\eta}$. This defines $\langle q_s^{n,\eta} : s \in^{\mathbf{u}} p_n \wedge |s| = n \wedge \eta < \kappa_{n-1}^{\mathbf{u}} \rangle$ and this sequence belongs to M because $h \restriction \kappa_{n-1}^{\mathbf{u}} \in M$ since h is M -amenable. Define p_{n+1} by

$$p_{n+1}(i) = \begin{cases} p_{n+1}(i) = p_n(i) & i < n \\ p_n(i) = \bigcap \{q_s^{n,\eta}(i) : s \in^{\mathbf{u}} p_n \wedge |s| = n \wedge \eta < \kappa_{n-1}^{\mathbf{u}}\} & n \leq i \end{cases}.$$

Note that for $n \leq i$, $p_{n+1}(i) \in \mu_i$ since $|\{q_s^{n,\eta} : s \in^{\mathbf{u}} p_n \wedge |s| = n \wedge \eta < \kappa_{n-1}^{\mathbf{u}}\}| = \kappa_{n-1}^{\mathbf{u}}$, $\mu_i^{\mathbf{u}}$ is $\kappa_i^{\mathbf{u}}$ -complete, and $\kappa_{n-1}^{\mathbf{u}} < \kappa_i$. Thus $p_{n+1} \in (\mathbb{N}_{\mathbf{u}})^M$ and $p_{n+1} \leq_{\mathbf{u}}^n p_n$. Suppose $s \in^{\mathbf{u}} p_{n+1}$, $|s| = n$, and $\eta < \kappa_{n-1}^{\mathbf{u}}$. So $s \in^{\mathbf{u}} p_n$. By construction, $(p_{n+1})_s \leq_{\mathbb{N}_{\mathbf{u}}} q_s^{n,\eta}$. Let $t \in^{\mathbf{u}} (p_{n+1})_s$ with $|t| = m_s^{n,\eta}$. Then $t \in^{\mathbf{u}} q_s^{n,\eta}$ and $(q_s^{n,\eta})_t \in h(\eta)$ be the properties of $q_s^{n,\eta}$. Since $h(\eta)$ is dense open and $(p_{n+1})_t \leq_{\mathbb{N}_{\mathbf{u}}} q_t^{n,\eta}$, $(p_{n+1})_t \in h(\eta)$. This shows p_{n+1} has property (3). This completes the construction of the $\mathbf{u}$ -fusion sequence $\langle p_n : n \in \omega \rangle$ with the desired properties above. (However this sequence does not belong to M .) Let p_{ω} be the fusion of $\langle p_n : n \in \omega \rangle$ defined by $p_{\omega}(i) = p_{i+1}(i)$ which belongs to $\mathbb{N}_{\mathbf{u}} \setminus (\mathbb{N}_{\mathbf{u}})^M$ (is a $\mathbf{u}$ -Namba condition but not a member of M). Note that p_{ω} is defined uniformly from $\mathbf{u}$, p , h , and $\prec$ and thus uniformly from $\mathbf{u}$, p , $\prec$, τ_1 , and τ_2 . Let $g \in [p]$. $G_g = \{p \in (\mathbb{N}_{\mathbf{u}})^M : g \in [p]\}$ is a $(\mathbb{N}_{\mathbf{u}})^M$ -filter. Let $D \in M$ be a dense open subset of $(\mathbb{N}_{\mathbf{u}})^M$. Since $h : \lambda^{\mathbf{u}} \rightarrow \mathcal{D}$ is a surjection, there is an $\eta < \lambda^{\mathbf{u}}$ so that $h(\eta) = D$. Let $1 \leq n < \omega$ be such that $\eta < \kappa_{n-1}^{\mathbf{u}}$. Since $g \restriction n \in^{\mathbf{u}} p_{\omega}$, $g \restriction n \in^{\mathbf{u}} p_{n+1}$. By the property of p_{n+1} , there is an $m \geq n$ so that for all $t \in^{\mathbf{u}} (p_{n+1})_{g \restriction n}$ with $|t| = m$, $(p_{n+1})_t \in h(\eta)$. Thus $(p_{n+1})_{g \restriction m} \in h(\eta) = D$. Since $p_{\omega} \leq_{\mathbb{N}_{\mathbf{u}}} p_{n+1}$, $g \restriction k \in (p_{n+1})_{g \restriction m}$ for all $k \in \omega$. Thus $g \in [(p_{n+1})_{g \restriction m}]$. By definition of G_g , $(p_{n+1})_{g \restriction m} \in G_g$. Thus $(p_{n+1})_{g \restriction m} \in G_g \cap D$. It has been shown that G_g is $(\mathbb{N}_{\mathbf{u}})^M$ -generic over M . $\square$

In particular, one has the following less uniform version.

Lemma 7.15. *Assume AD. Let $M \models \text{ZFC}$ be an inner model. Let $\mathbf{u} \in M$ be such that $\lambda^{\mathbf{u}} < \Theta$, $M \models \text{"}\mathbf{u}$ is a measure sequence, $2^{\lambda^{\mathbf{u}}} = (\lambda^{\mathbf{u}})^+$, $2^{(\lambda^{\mathbf{u}})^+} = (\lambda^{\mathbf{u}})^{++}$ ", and $\text{cof}(((\lambda^{\mathbf{u}})^+)^M) = \text{cof}(((\lambda^{\mathbf{u}})^{++})^M) = \omega$. For all $p \in (\mathbb{N}_{\mathbf{u}})^M$, there is a $q \in \mathbb{N}_{\mathbf{u}}$ such that $q \leq_{\mathbb{N}_{\mathbf{u}}} p$, $\text{length}(p) = \text{length}(q)$, and for all $g \in [q]$, G_g is $(\mathbb{N}_{\mathbf{u}})^M$ -generic filter over M .*

The following result shows there is an abundance of measurable cardinals of HOD below the true cardinals greater than ω_1 . (ω_1 is the first measurable cardinal of HOD.)

Fact 7.16. *(Jackson, Ketchersid, Schlutzenberg, Woodin; [27] Lemma 2.7²⁰) Assume AD and $V = L(\mathbb{R})$. If $x \in \mathbb{R}$ and κ is a cardinal such that $\omega_1 < \kappa < \Theta$, then $\text{HOD}_x \models \text{"there are } \kappa\text{-many measurable cardinals below } \kappa\text{"}$.*

Let κ be a cardinal. Recall $B(\omega, \kappa) = \{f \in {}^{\omega}\kappa : \sup(f) < \kappa\}$. If $\text{cof}(\kappa) > \omega$, then $B(\omega, \kappa) = {}^{\omega}\kappa$. If $\text{cof}(\kappa) = \omega$, then $|B(\omega, \kappa)| < |{}^{\omega}\kappa|$ by Fact 6.11.

Theorem 7.17. *Assume AD. $L(\mathbb{R}) \models \text{"For all cardinals } \kappa \text{ with } 2 \leq \kappa < \Theta \text{ and all } \chi < \kappa, B(\omega, \kappa) \text{ does not inject into } \mathcal{P}(\chi) \times \text{ON} \text{"}$. In particular, $L(\mathbb{R}) \models \text{"For all cardinals } \kappa \text{ with } 2 \leq \kappa < \Theta \text{ and all } \chi < \kappa, {}^{\omega}\kappa \text{ does not inject into } \mathcal{P}(\chi) \times \text{ON} \text{"}$.*

Proof. By going into the inner model $L(\mathbb{R})$, one may assume $V = L(\mathbb{R})$. Suppose the result fails. There are $2 \leq \kappa < \Theta$, $\chi < \kappa$, and an injection $\Phi : B(\omega, \kappa) \rightarrow \mathcal{P}(\chi) \times \text{ON}$. Since $V = L(\mathbb{R})$, there is an $x \in \mathbb{R}$ so that Φ is $\text{OD}_{\{x\}}$. First, $2 \leq \kappa \leq \omega$ is not possible. This is because $|B(\omega, \kappa)| = |\mathbb{R}|$ and is hence not wellorderable. However, $\mathcal{P}(\chi) \times \text{ON}$ is wellorderable for all $\chi < \kappa \leq \omega$. Thus Φ cannot be an injection. Next, $\kappa = \omega_1$ cannot be possible. This follows from Fact 2.10 or Theorem 3.5.²¹ Thus $\kappa > \omega_1$. By Fact 7.16, $\text{HOD}_{\{x\}} \models \text{"there are } \kappa\text{-many measurable cardinals below } \kappa\text{"}$. Let $\vec{\kappa} = \langle \kappa_n : n \in \omega \rangle$ be the first

²⁰As a corollary of Fact 7.16, every uncountable cardinal in HOD_x is either measurable or a limit of measurable cardinals of HOD_x . [27] showed that this corollary is sufficient for showing every cardinal is Jónsson in $L(\mathbb{R}) \models \text{AD}$. However, since $B(\omega, \kappa) \neq {}^{\omega}\kappa$, for the main result of this section concerning $B(\omega, \kappa)$, one needs the stronger form due to Schlutzenberg in the cofinality ω case.

²¹The case $\kappa = \omega_1$ needs to be considered separately. No ordinal countable in the real world can be measurable in $\text{HOD}_{\{x\}}$ by [60] Lemma 8.25. Thus there is no countable ordinal λ which is a limit of measurable cardinals in $\text{HOD}_{\{x\}}$. The generalized

2193 ω -many measurable cardinals between χ and κ . Let $\lambda = \sup\{\kappa_n : n \in \omega\}$ and note that $\chi < \kappa_0 < \lambda < \kappa$
2194 since $\kappa > \omega$. Let $\vec{\mu} = \langle \mu_n : n \in \omega \rangle \in \text{HOD}_{\{x\}}$ be such that for each $n \in \omega$, μ_n is a κ_n -complete uniform
2195 normal measure on κ_n . Let $\mathbf{u}$ be the measure sequence so that $\kappa_n^{\mathbf{u}} = \kappa_n$, $\mu_n^{\mathbf{u}} = \mu_n$, and $\lambda^{\mathbf{u}} = \lambda$. Let $\mathbb{N}_{\mathbf{u}}$
2196 denote the collection of all $\mathbf{u}$ -Namba conditions and $(\mathbb{N}_{\mathbf{u}})^{\text{HOD}_{\{x\}}}$ be the $\mathbf{u}$ -Namba conditions that belong to
2197 $\text{HOD}_{\{x\}}$. By Fact 6.1, the inner model $\text{HOD}_{\{x\}}$ satisfies the hypothesis of Lemma 7.14 or Lemma 7.15.
2198 Thus by applying Lemma 7.14 or even the less uniform Lemma 7.15, there is a $p \leq_{\mathbb{N}_{\mathbf{u}}} 1_{\mathbb{N}}$ such that for
2199 any $g \in [p]$, G_g is $(\mathbb{N}_{\mathbf{u}})^{\text{HOD}_{\{x\}}}$ -generic over $\text{HOD}_{\{x\}}$. Fix a $g \in [p]$. Note that $g = f_{G_g}$. Since $\lambda < \Theta$ and
2200 $g \in {}^\omega \lambda$, $\text{HOD}_{\{x\}}[g] = \text{HOD}_{\{x,g\}}$ by Fact 4.19²². Moreover, since $\lambda < \kappa$, $g \in B(\omega, \kappa)$. Since $\Phi \in \text{OD}_{\{x\}}$,
2201 $\Phi(g) \in \text{OD}_{\{x,g\}} \cap (\mathcal{P}(\chi) \times \text{ON})$. Thus $\Phi(g) \in \text{HOD}_{\{x,g\}} \cap (\mathcal{P}(\chi) \times \text{ON}) = \text{HOD}_{\{x\}}[g] \cap (\mathcal{P}(\chi) \times \text{ON}) \subseteq$
2202 $\text{HOD}_{\{x\}}[G_g] \cap (\mathcal{P}(\chi) \times \text{ON})$. Since $(\mathbb{N}_{\vec{\mu}})^{\text{HOD}_{\{x\}}}$ does not add any new bounded subsets of λ by Fact 7.13 and
2203 $\chi < \lambda$, $\text{HOD}_{\{x\}}[G_g] \cap (\mathcal{P}(\chi) \times \text{ON}) = \text{HOD}_{\{x\}} \cap (\mathcal{P}(\chi) \times \text{ON})$. Thus $\Phi(g) \in \text{HOD}_{\{x\}}$. Since $\Phi(g) \in \text{OD}_{\{x\}}$
2204 and $\Phi^{-1} \in \text{OD}_{\{x\}}$, one has that $g = \Phi^{-1}(\Phi(g)) \in \text{OD}_{\{x\}}$. Thus $g \in \text{HOD}_{\{x\}}$. This contradicts Fact 7.10
2205 which asserts that $g = f_{G_g} \notin \text{HOD}_{\{x\}}$. $\square$

2206 **Corollary 7.18.** Assume AD. $L(\mathbb{R}) \models$ “For all cardinals κ with $\omega \leq \kappa < \Theta$, all $\chi < \kappa$, all $\omega \leq \epsilon \leq \kappa$,
2207 $\nu \leq \chi$, and $\zeta \in \text{ON}$, $\neg(|[\kappa]^\epsilon| \leq |[\chi]^\nu \times \zeta|)$ ”.

2208 **Theorem 7.19.** Assume AD^+ . For all cardinals κ with $2 \leq \kappa < \Theta$ and all $\chi < \kappa$, $B(\omega, \kappa)$ does not inject
2209 into $\mathcal{P}(\chi) \times \text{ON}$. In particular, for all cardinals κ with $2 \leq \kappa < \Theta$ and all $\chi < \kappa$, ${}^\omega \kappa$ does not inject into
2210 $\mathcal{P}(\chi) \times \text{ON}$.

2211 *Proof.* The proof of this in relation to the proof of Theorem 7.17 is analogous to the proof of Theorem 6.7
2212 in relation to the proof of Theorem 6.2. An initial Σ_1 -reflection was not used in the proof of Theorem 7.17.
2213 Here Σ_1 -reflection into the Suslin-coSuslin sets (Fact 6.5) is still needed to bring a counterexample into a
2214 nice model of the form $L(Z, \mathbb{R})$ as in the proof of Theorem 6.7 so that $\text{HOD}_Z^{L(Z, \mathbb{R})}$ possesses a direct system
2215 HOD-analysis. With such a HOD-analysis, $\text{HOD}_Z^{L(Z, \mathbb{R})}$ will satisfy the analog of Fact 7.16. The proof then
2216 proceeds much like the proof of Theorem 7.17. $\square$

2217 **Corollary 7.20.** Assume AD^+ . For all cardinals κ with $\omega \leq \kappa < \Theta$, all $\chi < \kappa$, all $\omega \leq \epsilon \leq \kappa$, $\nu \leq \chi$, and
2218 $\zeta \in \text{ON}$, $\neg(|[\kappa]^\epsilon| \leq |[\chi]^\nu \times \zeta|)$.

2219 Lemma 7.14 shows that for suitable inner models M , there is a $p \in \mathbb{N}_{\mathbf{u}}$ (which does not belong to M) so
2220 that every $g \in [p]$ induces a $(\mathbb{N}_{\mathbf{u}})^M$ -generic filter over M . The next result computes the cardinality of $[p]$,
2221 the body of p .

2222 **Fact 7.21.** Let $M \models \text{ZFC}$ be an inner model. Let $\mathbf{u} \in M$ be such that $M \models$ “ $\mathbf{u}$ is a measure sequence”’. Let
2223 $p \in \mathbb{N}_{\mathbf{u}}$ (which may not belong to M). Then $|[p]| = |{}^\omega(\lambda^{\mathbf{u}})|$.

Proof. It is clear that $[p] \subseteq {}^\omega(\lambda^{\mathbf{u}})$ and thus $|[p]| \leq |{}^\omega(\lambda^{\mathbf{u}})|$. For each $f \in {}^\omega(\lambda^{\mathbf{u}})$, define $\iota_f : \omega \rightarrow \omega$ by
induction as follows: Let $\iota_f(0)$ be the least $n \in \omega$ so that $\text{slength}(p) \leq n$ and $f(0) < \kappa_n^{\mathbf{u}}$. Suppose $\iota_f(i)$ has
been defined. Let $\iota_f(i+1)$ be the least n greater than $\iota_f(i)$ so that $f(i+1) < \kappa_n^{\mathbf{u}}$. For each $n \in \omega$ with
 $\text{slength}(p) \leq n < \omega$, let $\text{enum}_{p(n)} : \kappa_n^{\mathbf{u}} \rightarrow p(n)$ be the increasing enumeration of $p(n)$. Define $\Phi(f) \in [p]$ by

$$\Phi(f)(n) = \begin{cases} \text{enum}_{p(n)}(0) & n \notin \iota_f[\omega] \\ \text{enum}_{p(n)}(1 + f(k)) & n \in \iota_f[\omega] \wedge n = \iota_f(k) \end{cases}.$$

2224 $\Phi : {}^\omega(\lambda^{\mathbf{u}}) \rightarrow [p]$ is an injection and thus $|{}^\omega(\lambda^{\mathbf{u}})| \leq |[p]|$. Thus $|{}^\omega(\lambda^{\mathbf{u}})| = |[p]|$. $\square$

2225 **Fact 7.22.** Suppose κ is a cardinal, $1 \leq \epsilon < \kappa$, and κ does not inject into $\mathcal{P}(\delta)$ for any $\delta < \kappa$. Then $B(\epsilon, \kappa)$
2226 does not have $\text{cof}(\kappa)$ -regular cardinality.

Namba forcings considered in this section cannot be used. However, as in the proof of Theorem 3.5, one can use the classical
Namba forcing defined in $\text{HOD}_{\{x\}}$ of trees on $(\omega_2)^{\text{HOD}_{\{x\}}}$. Note the existence of generic filters for this Namba forcing over
 $\text{HOD}_{\{x\}}$ is given by Fact 2.14 since this forcing is countable in the real world.

²²The generic object of $(\mathbb{N}_{\vec{\mu}})^{\text{HOD}_{\{x\}}}$ is an ω -sequence through $\lambda < \Theta$. Although the generic filter existence problem is more
difficult here than for Theorem 6.2, the capturing problem has a much simpler solution using Fact 4.19.

2227 *Proof.* Let $\rho : \text{cof}(\kappa) \rightarrow \kappa$ be an increasing cofinal sequence. By definition of $B(\epsilon, \kappa)$, $\sup(f) < \kappa$ for all
 2228 $f \in B(\epsilon, \kappa)$. Let $\Phi : B(\epsilon, \kappa) \rightarrow \text{cof}(\kappa)$ be defined by $\Phi(f)$ is the least α so that $\sup(f) < \rho(\alpha)$. For
 2229 each $\alpha < \text{cof}(\kappa)$, $|\Phi^{-1}[\{\alpha\}]| < |B(\epsilon, \kappa)|$. Since otherwise, κ would inject into $\mathcal{P}(\max\{\epsilon, \rho(\alpha)\})$ since $|\kappa| \leq$
 2230 $|B(\epsilon, \kappa)| = |\Phi^{-1}[\{\alpha\}]| \leq |\epsilon \rho(\alpha)| \leq |\mathcal{P}(\max\{\epsilon, \rho(\alpha)\})|$ which violates the hypothesis since $\max\{\epsilon, \rho(\alpha)\} < \kappa$.
 2231 Φ witnesses that $B(\epsilon, \kappa)$ does not have $\text{cof}(\kappa)$ -regular cardinality. $\square$

2232 **Corollary 7.23.** *Assume κ is a cardinal, $1 \leq \epsilon < \kappa$, κ does not inject into $\mathcal{P}(\delta)$ for any $\delta < \kappa$, and*
 2233 *$\text{cof}(\kappa) > \epsilon$. Then ${}^\epsilon\kappa$ does not have $\text{cof}(\kappa)$ -regular cardinality. In particular, if $\text{cof}(\kappa) > \omega$, ${}^\omega\kappa$ does not have*
 2234 *$\text{cof}(\kappa)$ -regular cardinality.*

2235 *Proof.* If $\text{cof}(\kappa) > \epsilon$, then ${}^\epsilon\kappa = B(\epsilon, \kappa)$. The result follows from Fact 7.22. $\square$

2236 Under AD^+ , $B(\omega, \kappa)$ does not have ω -regular cardinality if $\kappa < \Theta$ and $\text{cof}(\kappa) = \omega$. However, under AD^+ ,
 2237 it will be shown that if $\kappa < \Theta$ and $\text{cof}(\kappa) = \omega$, then ${}^\omega\kappa$ has a much greater degree of ordinal regularity.

2238 **Lemma 7.24.** *Let $M \models \text{ZFC}$ be an inner model. Let $\mathbf{u} \in M$ be such that $\lambda^{\mathbf{u}} < \Theta$, $M \models \text{“}\mathbf{u} \text{ is a measure}$
 2239 *sequence, $2^{\lambda^{\mathbf{u}}} = (\lambda^{\mathbf{u}})^+$, $2^{(\lambda^{\mathbf{u}})^+} = (\lambda^{\mathbf{u}})^{++}$ ”, and $\text{cof}(((\lambda^{\mathbf{u}})^+)^M) = \text{cof}(((\lambda^{\mathbf{u}})^{++})^M) = \omega$. Let $R \subseteq {}^\omega(\lambda^{\mathbf{u}}) \times \text{ON}$*
 2240 *be such that there is an ∞ -Borel code (S, φ) for R with $S \in M$. Let $p \in (\mathbb{N}_{\mathbf{u}})^M$ be such that $[p] \subseteq \text{dom}(R)$*
 2241 *(here $[p]$ is computed in the real world).**

- 2242 (1) *There is a $\gamma \in \text{ON}$ and a $q \in (\mathbb{N}_{\mathbf{u}})^M$ with $q \leq_{\mathbb{N}_{\mathbf{u}}} p$ so that $M \models q \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\gamma})$.*
 2243 (2) *There is a unique $\gamma \in \text{ON}$ so that there is a $q \in (\mathbb{N}_{\mathbf{u}})^M$ with $q \leq_{\mathbb{N}_{\mathbf{u}}} p$, $\text{length}(q) = \text{length}(p)$, and*
 2244 *$M \models q \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\gamma})$.*

2245 *Proof.* To see statement (1): The first claim is that there exists $q_0 \in (\mathbb{N}_{\mathbf{u}})^M$ with $q_0 \leq_{\mathbb{N}_{\mathbf{u}}} p$ and $M \models q_0 \Vdash_{\mathbb{N}_{\mathbf{u}}}$
 2246 $(\exists \beta \in \text{ON})(L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \beta))$.

2247 To see this claim, suppose not. Then there is an $r \in (\mathbb{N}_{\mathbf{u}})^M$ with $r \leq_{\mathbb{N}_{\mathbf{u}}} p$ so that $r \Vdash_{\mathbb{N}_{\mathbf{u}}} (\forall \beta \in$
 2248 $\text{ON})(L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \beta))$. Applying Lemma 7.14 to the inner model M and the condition $r \in (\mathbb{N}_{\mathbf{u}})^M$
 2249 to obtain a condition $\bar{r} \in \mathbb{N}_{\mathbf{u}}$ (but not in M) so that $\bar{r} \leq_{\mathbb{N}_{\mathbf{u}}} r$ and for all $g \in [\bar{r}]$, G_g is $(\mathbb{N}_{\mathbf{u}})^M$ -generic over
 2250 M .²³ Pick a $g \in [\bar{r}]$. Since $g \in [\bar{r}] \subseteq [r] \subseteq [p] \subseteq \text{dom}(R)$, there is a $\gamma \in \text{ON}$ so that $R(g, \gamma)$. However,
 2251 since $g \in [\bar{r}] \subseteq [r]$ implies $r \in G_g$ and $\check{f}_{\check{G}}[G_g] = g$, one has by the forcing theorem that $M[G_g] \models (\forall \beta \in$
 2252 $\text{ON})L[S, g] \models \neg \varphi(S, g, \gamma)$. Thus $M[G] \models L[S, g] \models \neg \varphi(S, g, \gamma)$. By absoluteness, $L[S, g] \models \neg \varphi(S, g, \gamma)$. Since
 2253 (S, φ) is the ∞ -Borel code for R , $\neg R(g, \gamma)$. Contradiction.

2254 Fix q_0 as in the claim above. There is $q_1 \in (\mathbb{N}_{\mathbf{u}})^M$ and $\gamma \in \text{ON}$ so that $q_1 \leq_{\mathbb{N}_{\mathbf{u}}} q_0$ and $M \models q_1 \Vdash_{\mathbb{N}_{\mathbf{u}}}$
 2255 $L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\gamma})$. This establishes the first statement. Now to prove the second statement. Work in
 2256 M . Let D be the set of $q \in \mathbb{N}_{\mathbf{u}}$ so that one of the following holds:

- 2257 (1) q is incompatible with p .
 2258 (2) There exists a $\beta \in \text{ON}$ such that $q \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta})$.

2259 By the first statement just proved above, D is a dense open subset of $\mathbb{N}_{\mathbf{u}}$. By lemma 7.11, there is a $\hat{p} \leq_{\mathbb{N}_{\mathbf{u}}} p$
 2260 with $\text{length}(p) = \text{length}(\hat{p})$ and a $\text{length}(p) \leq k < \omega$ so that for all $s \in {}^u \hat{p}$ with $|s| = k$, $(\hat{p})_s \in D$. Since
 2261 $\hat{p} \leq_{\mathbb{N}_{\mathbf{u}}} p$, the first condition above cannot hold. Thus for each $s \in {}^u \hat{p}$ with $|s| = k$, there is a $\beta_s \in \text{ON}$ so that
 2262 $(\hat{p})_s \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_s)$. By Lemma 7.12, for each $s \in {}^u \hat{p}$ with $|s| = k$, exactly one of the following
 2263 holds:

- 2264 (a) There is a $q \leq_{\mathbb{N}_{\mathbf{u}}} \hat{p}$ with $\text{length}(q) = \text{length}(\hat{p}) = \text{length}(p)$ and $q \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_s)$.
 2265 (b) There is a $q \leq_{\mathbb{N}_{\mathbf{u}}} \hat{p}$ with $\text{length}(q) = \text{length}(\hat{p}) = \text{length}(p)$ and $q \Vdash_{\mathbb{N}_{\mathbf{u}}} L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_s)$.

2266 The claim is that there is an $s \in {}^u \hat{p}$ with $|s| = k$ so that Case (a) holds for s . Since $M \models \text{AC}$, let $\prec$ be a
 2267 wellordering of $\mathbb{N}_{\mathbf{u}}$ in M . Suppose for all $s \in {}^u \hat{p}$, Case (b) holds. Let q_s be the $\prec$ -least such $q \in \mathbb{N}_{\mathbf{u}}$ which
 2268 witnesses Case (b) for s . Define $\bar{q}$ by $\bar{q}(i)$ for all $i < k$ and $\bar{q}(i) = \bigcap \{q_s(i) : s \in {}^u \hat{p} \wedge |s| = k\}$ if $k \leq i$. Since
 2269 $|\{s \in {}^u \hat{p} : |s| = k\}| < |\kappa_i^{\mathbf{u}}|$, $\mu_i^{\mathbf{u}}$ is $\kappa_i^{\mathbf{u}}$ -complete, and $\kappa_k^{\mathbf{u}} \leq \kappa_i^{\mathbf{u}}$ when $k \leq i$, one has that $\bar{q} \in \mathbb{N}_{\mathbf{u}}$ and $\bar{q} \leq_{\mathbb{N}_{\mathbf{u}}} q_s$ for
 2270 all $s \in {}^u \hat{p}$ with $|s| = k$.

2271 Now return to the real world. By Lemma 7.15, there is a $\tilde{q} \in \mathbb{N}_{\mathbf{u}}$ with $\tilde{q} \leq_{\mathbb{N}_{\mathbf{u}}} \bar{q}$ so that $\text{length}(\tilde{q}) =$
 2272 $\text{length}(\bar{q}) = \text{length}(p)$ and for all $g \in [\tilde{q}]$, G_g is $(\mathbb{N}_{\mathbf{u}})^M$ -generic over M . Pick any $g \in [\tilde{q}]$. For any $s \in {}^u \hat{p}$

²³When applying Lemma 7.14, choose any τ_1 and τ_2 witnessing $\text{cof}(((\lambda^{\mathbf{u}})^+)^M) = \omega$ and $\text{cof}(((\lambda^{\mathbf{u}})^{++})^M) = \omega$. Since $M \models \text{AC}$, let $\prec$ be any internal wellordering of the element of M of rank less than $\lambda^{\mathbf{u}} + 3$. Uniformity is not relevant in this lemma.

with $|s| = k$, $g \in [\tilde{q}] \subseteq [\bar{q}] \subseteq [q_s]$ and thus $q_s \in G_g$. For each $s \in {}^u \hat{p}$ with $|s| = k$, $M \models q_s \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_s)$. In particular, since $g \restriction k \in {}^u \hat{p}$ and $|g \restriction k| = k$, $M \models q_{g \restriction k} \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})$. Also $g \in [\tilde{q}] \subseteq [\bar{q}] \subseteq [\hat{p}]$. In particular, $g \in [(\hat{p})_{g \restriction k}]$ and thus $(\hat{p})_{g \restriction k} \in G_g$. Since $g \restriction k \in {}^u \hat{p}$, $|g \restriction k| = k$, and by the definition of $\beta_{g \restriction k}$, $M \models (\hat{p})_{g \restriction k} \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})$. It has been shown that $M \models "q_{g \restriction k} \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})"$, $M \models "(\hat{p})_{g \restriction k} \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})"$, and $q_{g \restriction k}, (\hat{p})_{g \restriction k} \in G$. Since G is a filter, there is an $r \in (\mathbb{N}_u)^M$ with $r \leq_{\mathbb{N}_u} q_{g \restriction k}$ and $r \leq_{\mathbb{N}_u} (\hat{p})_{g \restriction k}$. Then $r \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})$ and $r \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \neg \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_{g \restriction k})$. Contradiction. This establishes the claim. By the claim, there is an $s \in {}^u \hat{p}$ with $|s| = k$ and a $\underline{q} \leq_{\mathbb{N}_u} \hat{p}$ with $\text{length}(\underline{q}) = \text{length}(p)$ so that $M \models \underline{q} \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\beta}_s)$. This $\underline{q}$ and ordinal β_s verify the second statement. The uniqueness of γ follows from the fact that if $q_0, q_1 \in (\mathbb{N}_u)^M$ with $\text{length}(q_0) = \text{length}(q_1)$, then q_0 and q_1 are $\leq_{\mathbb{N}_u}$ -compatible via the condition q_2 defined by $q_2(i) = q_0(i) = q_1(i)$ when $i < \text{length}(q_0) = \text{length}(q_1)$ and $q_2(i) = q_0(i) \cap q_1(i)$ when $\text{length}(q_0) = \text{length}(q_1) \leq i < \omega$. $\square$

Lemma 7.24 statement (1) is sufficient for the following results concerning regularity; however, it is prudent to have the stronger statement (2) which does not change the stem for other constructions. The proof of statement (2) uses statement (1). The proof of statement (1) has an essential use of the ∞ -Borel code.

Lemma 7.24 asserts the existence of an internal object $q \in M$ with the desired property and thus uniformity is not relevant. Uniformity in the next lemma will be important.

Lemma 7.25. *Let M be an inner model of ZFC. Suppose the following holds.*

- (a) $\mathbf{u} \in M$ and $M \models$ “ $\mathbf{u}$ is a measure sequence, $2^{\lambda^{\mathbf{u}}} = (\lambda^{\mathbf{u}})^+$, and $2^{(\lambda^{\mathbf{u}})^+} = (\lambda^{\mathbf{u}})^{++}$ ”.
- (b) $\prec$ is a wellordering of all sets in M of rank less than $\lambda^{\mathbf{u}} + 3$.
- (c) $\tau_1 : \omega \rightarrow ((\lambda^{\mathbf{u}})^+)^M$ is an increasing cofinal function.
- (d) $\tau_2 : \omega \rightarrow ((\lambda^{\mathbf{u}})^{++})^M$ is an increasing cofinal function.
- (e) $p \in (\mathbb{N}_u)^M$.
- (f) (S, φ) is an ∞ -Borel code with $S \in M$. If $R \subseteq {}^\omega(\lambda^{\mathbf{u}}) \times \text{ON}$ is the ∞ -Borel subset of ${}^\omega(\lambda^{\mathbf{u}}) \times \text{ON}$ defined by (S, φ) , then $[p] \subseteq \text{dom}(R)$.

Then there is a $\gamma \in \text{ON}$ and $q \in \mathbb{N}_u$ (but $q \notin M$) with $q \leq_{\mathbb{N}_u} p$, $\text{length}(q) = \text{length}(p)$, and for all $g \in [q]$, $R(g, \gamma)$. Also q is produced uniformly from $\mathbf{u}$, $\prec$, τ_1 , τ_2 , p , and (S, φ) .

Proof. By Lemma 7.24, there is a unique $\gamma \in \text{ON}$ such that there is a $q' \in (\mathbb{N}_u)^M$ with $q' \leq_{\mathbb{N}_u} p$, $\text{length}(q') = \text{length}(p)$, and $M \models q' \Vdash_{\mathbb{N}_u} L[\check{S}, \check{f}_{\check{G}}] \models \varphi(\check{S}, \check{f}_{\check{G}}, \check{\gamma})$. Let $\bar{q}$ be the $\prec$ -least such q' witnessing this property for γ . Note that γ and $\bar{q}$ are obtained uniformly from $\mathbf{u}$, $\prec$, and (S, φ) . By Lemma 7.14, there is a $q \leq_{\mathbb{N}_u} \bar{q}$ with $q \leq_{\mathbb{N}_u} p$, $\text{length}(q) = \text{length}(\bar{q})$, and for all $g \in [q]$, G_g is $(\mathbb{N}_u)^M$ -generic over M and this q is produced uniformly from $\mathbf{u}$, $\prec$, τ_1 , and τ_2 , and $\bar{q}$. Thus q is produced uniformly from $\mathbf{u}$, $\prec$, τ_1 , τ_2 , p , and (S, φ) . Pick any $g \in [q]$. Since $g \in [\bar{q}] \subseteq [\bar{q}]$, $\bar{q} \in G_g$. By the forcing theorem, $M[G_g] \models L[S, g] \models \varphi(S, g, \gamma)$. By absoluteness, $L[S, g] \models \varphi(S, g, \gamma)$. Since (S, φ) is the ∞ -Borel code for R , $R(g, \gamma)$. Since $g \in [q]$ was arbitrary, the result follows. $\square$

Jackson, Trang, and the author ([17]) can show that ${}^\omega(\omega_\omega)$ has ω_ω -regular cardinality under AD using Martin's ultrapower analysis for ω_ω and that ${}^\omega(\omega_\omega)$ has ON-regular cardinality under AD and $\text{DC}_{\mathbb{R}}$ using Steel's result concerning Suslin bounded prewellordering. The next theorem shows that under AD^+ , for all $\kappa < \Theta$ with $\text{cof}(\kappa) = \omega$, ${}^\omega \kappa$ has ON-regular cardinality.

Theorem 7.26. *Assume AD and $V = L(\mathbb{R})$. Let $\omega \leq \kappa < \Theta$ be a cardinal with $\text{cof}(\kappa) = \omega$. Then ${}^\omega \kappa$ has ON-regular cardinality.*

Proof. First, consider $\kappa = \omega$. $|{}^\omega \omega| = |\mathbb{R}|$. Fact 2.5 states that wellordered unions of meager sets are meager. So for any function $\Phi : \mathbb{R} \rightarrow \text{ON}$, there is some $\beta \in \text{ON}$ so that $\Phi^{-1}[\{\beta\}]$ is nonmeager. Then $|\Phi^{-1}[\{\beta\}]| = |\mathbb{R}|$.

Now suppose $\omega < \kappa < \Theta$ and $\text{cof}(\kappa) = \omega$. Since $\text{AC}_{\omega}^{\mathbb{R}}$ implies ω_1 is regular and $\text{cof}(\kappa) = \omega$, one must have that $\kappa > \omega_1$. Let $\rho : \omega \rightarrow \kappa$ be an increasing cofinal sequence. Suppose $\Phi : {}^\omega \kappa \rightarrow \text{ON}$ is a function. Since $\kappa < \Theta$, there is a surjection $\pi : \mathbb{R} \rightarrow \kappa$. Define $\varsigma : \mathbb{R} \rightarrow {}^\omega \kappa$ by $\varsigma(x)(n) = \pi(x^{[n]})$ (where recall that $\langle x^{[n]} : n \in \omega \rangle$ is the recursive function which codes an ω -sequence in $\mathbb{R}$ as an element of $\mathbb{R}$). ς is a surjection. So $\mathbb{R}$ surjects onto $\Phi[{}^\omega \kappa]$ and hence $\text{ot}(\Phi[{}^\omega \kappa]) < \Theta$. By composing Φ with the Mostowski collapse

of $\Phi[\omega\kappa]$ if necessary, one will assume without loss of generality that $\Phi : \omega\kappa \rightarrow \zeta$ where $\zeta < \Theta$. Since $V = L(\mathbb{R})$, there is an $x \in \mathbb{R}$ so that Φ and ρ are $\text{OD}_{\{x\}}$. Thus $\rho \in \text{HOD}_{\{x\}}$ and $\text{HOD}_{\{x\}} \models \text{cof}(\kappa) = \omega$. Let $R = \{(f, \alpha) \in \omega\kappa \times \zeta : \Phi(f) = \alpha\}$. Note that $\text{dom}(R) = \omega\kappa$ since Φ is a function on $\omega\kappa$. R is $\text{OD}_{\{x\}}$ and essentially a subset of ${}^\omega(\max\{\kappa, \zeta\})$. By Fact 4.21, R has an $\text{OD}_{\{x\}}$ ∞ -Borel code (S, φ) . Since $\text{HOD}_{\{x\}} \models \text{cof}(\kappa) = \omega$, Fact 7.16 gives an increasing sequence $\langle \kappa_n : n \in \omega \rangle$ and $\langle \mu_n : n \in \omega \rangle$ which both belong to $\text{HOD}_{\{x\}}$ so that $\sup\{\kappa_n : n \in \omega\} = \kappa$ and $\text{HOD}_{\{x\}} \models \mu_n$ is a normal measure on κ_n for each $n \in \omega$. Let $\mathbf{u} \in \text{HOD}_{\{x\}}$ be the associated measure sequence and $\dot{f}_G$ be the $(\mathbb{N}_{\mathbf{u}})^{\text{HOD}_{\{x\}}}$ -name for the generic ω -sequence. Since the hypothesis of Lemma 7.25 holds for $\text{HOD}_{\{x\}}$ and $1_{\mathbb{N}_{\mathbf{u}}}$ by Fact 6.1, there is a $\gamma < \zeta$ and $q \in \mathbb{N}_{\mathbf{u}}$ so that for all $g \in [q]$, $R(g, \gamma)$. By definition of R , $\Phi(g) = \gamma$ for all $g \in [q]$. It has been shown that $[q] \subseteq \Phi^{-1}[\{\gamma\}]$. Since $[q] = |\omega\kappa|$ by Fact 7.21, $|\Phi^{-1}[\{\gamma\}]| = |\omega\kappa|$. Since Φ was arbitrary, this shows that $\omega\kappa$ has ON-regular cardinality. $\square$

If $\text{cof}(\kappa) = \omega$ and $\omega < \kappa$, Fact 7.22 shows that $B(\omega, \kappa)$ does not have ω -regular cardinality. Under AD, Jackson, Trang, and the author ([17]) can show that $B(\omega, \omega_\omega)$ has n -regular cardinal for all $n \in \omega$ using the Martin's ultrapower analysis for each ω_m when $m < \omega$. This will be generalized next to show under determinacy assumption that $B(\omega, \kappa)$ has n -regular cardinality for all $n \in \omega$. (Note that under AD, there are sets which do not have 2-regular cardinality such as $\mathbb{R} \sqcup \omega_1$.)

Fact 7.27. *Let $M \models \text{ZFC}$ be an inner model. Let $\delta \in \text{ON}$ and $\langle \mathbf{u}_\alpha : \alpha \in \delta \rangle$ be a sequence so that $M \models \mathbf{u}_\alpha$ is a measure sequence for each $\alpha \in \delta$ and $\langle \lambda^{\mathbf{u}_\alpha} : \alpha \in \delta \rangle$ is a strictly increasing sequence. Let $\kappa = \sup\{\lambda^{\mathbf{u}_\alpha} : \alpha \in \delta\}$. Let $\langle p_\alpha : \alpha \in \delta \rangle$ be such that $p_\alpha \in \mathbb{N}_{\mathbf{u}_\alpha}$ (but not necessarily in M) for each $\alpha \in \delta$. Then $|\bigcup_{\alpha \in \delta} [p_\alpha]| = |B(\omega, \kappa)|$.*

Proof. Since $\bigcup_{\alpha \in \delta} [p_\alpha] \subseteq B(\omega, \kappa)$, it is clear that $|\bigcup_{\alpha \in \delta} [p_\alpha]| \leq |B(\omega, \kappa)|$. If $f \in B(\omega, \kappa)$, let $\alpha_f \in \delta$ be the least $\alpha \in \delta$ so that $\sup(f) < \lambda_{\alpha_f}$. Let ι_f be the least $n \in \omega$ so that $\text{length}(p_{\alpha_f}) \leq n$ and $f(0) < \kappa_n^{\alpha_f}$. Suppose $\iota_f(k)$ has been defined. Let $\iota_f(k+1)$ be the least n greater than $\iota_f(k)$ so that $f(k+1) < \kappa_n^{\mathbf{u}_{\alpha_f}}$. For each $\alpha \in \delta$ and $k \in \omega$ with $\text{length}(p_\alpha) \leq k < \omega$, let $\text{enum}_{p_\alpha(k)} : \kappa_k^{\mathbf{u}_\alpha} \rightarrow p_\alpha(k)$ be the increasing enumeration of $p_\alpha(k)$. Define $\Phi : B(\omega, \kappa) \rightarrow \bigcup_{\alpha \in \delta} [p_\alpha]$ by

$$\Phi(f)(n) = \begin{cases} \text{enum}_{p_{\alpha_f}(n)}(0) & n \notin \iota_f[\omega] \\ \text{enum}_{p_{\alpha_f}(n)}(1 + f(k)) & n \in \iota_f[\omega] \wedge n = \iota_f(k) \end{cases}.$$

Φ is an injection and therefore $|B(\omega, \kappa)| \leq |\bigcup_{\alpha \in \delta} [p_\alpha]|$. Thus $|B(\omega, \kappa)| = |\bigcup_{\alpha \in \delta} [p_\alpha]|$. $\square$

Theorem 7.28. *Assume AD and $V = L(\mathbb{R})$. Let $\omega \leq \kappa < \Theta$ be a cardinal with $\text{cof}(\kappa) = \omega$. For all $n < \omega$, $B(\omega, \kappa)$ has n -regular cardinality. If $\omega < \kappa$, then $B(\omega, \kappa)$ does not have ω -regular cardinality.*

Proof. First, suppose $\kappa = \omega$. Note that $|B(\omega, \omega)| = |\mathbb{R}|$. As noted in Theorem 7.26, $\mathbb{R}$ even has ON-regular cardinality.

Next, suppose that $\omega < \kappa < \Theta$ and $\text{cof}(\kappa) = \omega$. Since $\text{AC}_\omega^\mathbb{R}$ implies $\text{cof}(\omega_1) = \omega_1$, one must have $\omega_1 < \kappa$. Let $\rho : \omega \rightarrow \kappa$ be an increasing cofinal function. Let $j \in \omega$. Let $\Phi : B(\omega, \kappa) \rightarrow j$ be a function. Since $V = L(\mathbb{R})$, there is an $x \in \mathbb{R}$ so that ρ and Φ are $\text{OD}_{\{x\}}$. Thus $\rho \in \text{HOD}_{\{x\}}$ and $\text{HOD}_{\{x\}} \models \text{cof}(\kappa) = \omega$. Let $R = \{(f, i) \in B(\omega, \kappa) \times j : \Phi(f) = i\}$. Note that R may be regarded as a subset of $\omega\kappa$. Also $\text{dom}(R) = B(\omega, \kappa)$ since Φ is a function on $B(\omega, \kappa)$. By Fact 4.21, R has an $\text{OD}_{\{x\}}$ ∞ -Borel code (S, φ) . By Fact 7.16, $\text{HOD}_{\{x\}} \models$ “there are κ -many measurable cardinals below κ ”. For each $m, n \in \omega$, let κ_n^m be the $(\omega \cdot \rho(m) + n)^{\text{th}}$ measurable cardinal of $\text{HOD}_{\{x\}}$ and let μ_n^m be the least normal measure on κ_n^m according to the canonical global wellordering of $\text{HOD}_{\{x\}}$. For each $m \in \omega$, let $\lambda_m = \sup\{\kappa_n^m : n \in \omega\}$. For each $m \in \omega$, let $\mathbf{u}_m$ be the measure sequence so that $\kappa_n^{\mathbf{u}_m} = \kappa_n^m$, $\mu_n^{\mathbf{u}_m} = \mu_n^m$, and $\lambda^{\mathbf{u}_m} = \lambda_m$. Since $\rho \in \text{HOD}_{\{x\}}$, one has that $\mathbf{u}_m \in \text{HOD}_{\{x\}}$ for each $m \in \omega$. Note that $\sup\{\lambda_m : m \in \omega\} = \kappa$. For each $m \in \omega$, let $R_m = R \cap (\omega(\lambda_m) \times j)$. (S, φ) essentially remains an ∞ -Borel code for R_m . The hypothesis of Lemma 7.25 holds for $\text{HOD}_{\{x\}}$ by Fact 6.1. For each $m \in \omega$, by applying Lemma 7.25 to $\text{HOD}_{\{x\}}$, $(\mathbb{N}_{\mathbf{u}_m})^{\text{HOD}_{\{x\}}}$, and R_m with ∞ -Borel code (S, φ) , one has a $q \in \mathbb{N}_{\mathbf{u}_m}$ and $i \in j$ so that for all $g \in [q]$, $R_m(g, i)$ or equivalently $R(g, i)$. Let i_m be the least such $i < j$ so that there exists a q with the above property for m . Let $S \subseteq \omega \times \mathcal{P}(\kappa)$ be defined by $S(m, q)$ if and only if $q \in \mathbb{N}_{\mathbf{u}_m}$ and for all $g \in [q]$, $R_m(q, i_m)$. By the discussion above, $\text{dom}(S) = \omega$.

By the Moschovakis coding lemma and $\text{AC}_\omega^\mathbb{R}$ in the real world,²⁴, there is a sequence $\langle q_m : m \in \omega \rangle$ so that for all $m \in \omega$, $S(m, q_m)$. There is an $\bar{i} \in j$ so that $A = \{m \in \omega : i_m = \bar{i}\}$ is infinite. Let $Z = \bigcup_{m \in A} [q_m]$. Suppose $g \in Z$. There is a unique $m \in A$ so that $g \in [q_m]$. By the property q_m and since $i_m = \bar{i}$, $R(g, \bar{i})$. By definition of R , $\Phi(g) = \bar{i}$. This shows that $Z \subseteq \Phi^{-1}[\{\bar{i}\}]$. Since $|Z| = |\bigcup_{m \in A} [q_m]| = |B(\omega, \kappa)|$ by Fact 7.27 (applied to the sequence $\langle u_m : m \in A \rangle$), one has that $|\Phi^{-1}[\{\bar{i}\}]| = |B(\omega, \kappa)|$. Since Φ was arbitrary, this shows that $B(\omega, \kappa)$ has j -regular cardinality. Since $j \in \omega$ was arbitrary, the theorem follows. $\square$

The following is obtained by making similar modifications as in Theorem 6.7 and Theorem 7.19 to Theorem 7.26 and Theorem 7.28.

Theorem 7.29. Assume AD^+ . Suppose $\omega \leq \kappa < \Theta$ with $\text{cof}(\kappa) = \omega$. ${}^\omega\kappa$ has ON-regular cardinality. $B(\omega, \kappa)$ has n -regular cardinality for each $n \in \omega$. If $\omega < \kappa$, then $B(\omega, \kappa)$ does not have ω -regular cardinality.

The results above have been shown for ${}^\omega\kappa$ and $B(\omega, \kappa)$ when $\kappa < \Theta$ with $\text{cof}(\kappa) = \omega$. When $\text{cof}(\kappa) > \omega$, ${}^\omega\kappa = B(\omega, \kappa)$. Fact 7.22 shows that $B(\omega, \kappa)$ does not have $\text{cof}(\kappa)$ -regular cardinality. Under AD, using $\omega_1 \rightarrow_* (\omega_1)_\omega^\omega$ or the countably completeness of $\mu_{\omega_1}^\omega$, one has that ${}^\omega\omega_1$ has ω -regular cardinality. Under AD, using $\omega_2 \rightarrow_* (\omega_1)_{\omega_1}^\omega$ or the ω_2 -completeness of the $\mu_{\omega_2}^\omega$, one has that ${}^\omega\omega_2$ has ω_1 -regular cardinality. Jackson, Trang, and the author can show that for all $2 \leq n < \omega$, ${}^\omega\omega_n$ has ω_1 -regular cardinality. The conjecture is that for all cardinal κ , $B(\omega, \kappa)$ has δ -regular cardinality for all $\delta < \text{cof}(\kappa)$. The methods developed here for $B(\omega, \kappa)$ when $\text{cof}(\kappa) = \omega$ can be extended to ${}^\omega\kappa$ for all cardinal $\kappa < \Theta$ if one assumes an additional hypothesis. The following is the version of this conjecture for $L(\mathbb{R})$.

Definition 7.30. The Uniform Cofinality of HOD Successor and Double Successor Hypothesis for $L(\mathbb{R})$ is the following assertion:

Assume AD and $V = L(\mathbb{R})$. Let $x \in \mathbb{R}$ and $\omega_1 < \kappa < \Theta$ be a cardinal such that $\text{HOD}_{\{x\}} \models \text{cof}(\kappa) = \text{cof}(\kappa)^{L(\mathbb{R})}$. There are functions $h : \text{cof}(\kappa) \rightarrow \kappa$, $H : \text{cof}(\kappa) \times \omega \rightarrow \kappa$, $S_1 : \text{cof}(\kappa) \times \omega \rightarrow \kappa$, and $S_2 : \text{cof}(\kappa) \times \omega \rightarrow \kappa$ so that for all $\alpha \in \text{cof}(\kappa)$ and $n \in \omega$, the following holds.

- (1) $H(\alpha, n) < H(\alpha, n+1)$, $S_1(\alpha, n) < S_1(\alpha, n+1)$, $S_2(\alpha, n) < S_2(\alpha, n+1)$.
- (2) $\langle H(\alpha, n) : n \in \omega \rangle \in \text{HOD}_{\{x\}}$.
- (3) $\text{HOD}_{\{x\}} \models H(\alpha, n)$ is a measurable cardinal.
- (4) $h(\alpha) = \sup\{H(\alpha, n) : n \in \omega\}$.
- (5) h is cofinal in κ .
- (6) $(h(\alpha)^+)^{\text{HOD}_{\{x\}}} = \sup\{S_1(\alpha, n) : n \in \omega\}$.
- (7) $(h(\alpha)^{++})^{\text{HOD}_{\{x\}}} = \sup\{S_2(\alpha, n) : n \in \omega\}$.

Remark 7.31. Recall that Steel [60] Lemma 8.25 shows that successor cardinals of HOD have cofinality ω in the real world. The above conjecture asserts one can uniformly find a sequence of witnesses to many successor cardinals of HOD having cofinality ω .

The following outline the usefulness of the conjecture. Let $\alpha < \text{cof}(\kappa)$ and $n \in \omega$. Let $\tau_0^\alpha(n) = H(\alpha, n)$. Let μ_n^α be the least normal measure on $H(\alpha, n)$ according to the canonical global wellordering of $\text{HOD}_{\{x\}}$. Let $\tau_1^\alpha(n) = S_1(\alpha, n)$. Let $\tau_2^\alpha(n) = S_2(\alpha, n)$. Let u_α be the measure sequence such that $\kappa^{u_\alpha} = H(\alpha, n)$, $\mu_n^{u_\alpha} = \mu_n^\alpha$, and $\lambda^{u_\alpha} = h(\alpha)$. Since for each α , $\langle H(\alpha, n) : n \in \omega \rangle \in \text{HOD}_{\{x\}}$, one has $u_\alpha \in \text{HOD}_{\{x\}}$ which is important since one needs $\text{HOD}_{\{x\}}$ to be able to define $\mathbb{N}_{u_\alpha}$. In the real world, one has the following two sequences $\langle \tau_i^\alpha : \alpha < \text{cof}(\kappa) \wedge i < 3 \rangle$ and $\langle u_\alpha : \alpha < \text{cof}(\kappa) \rangle$. By the uniformity asserted in Lemma 5.4, one has a sequence $\langle e_\alpha : \alpha < \text{cof}(\kappa) \rangle$ so that $e_\alpha : h(\alpha) \rightarrow (h(\alpha)^{++})^{\text{HOD}_{\{x\}}}$ is a $\text{HOD}_{\{x\}}$ -amenable surjection for each $\alpha < \text{cof}(\kappa)$. Similarly, one can apply Lemma 7.15 and Lemma 7.24 uniformly.

Note also that the conjecture is stated with the domain of h being $\text{cof}(\kappa)$. This will be sufficient for the following application. Moreover, for higher dimensional problem such as primeness and Jónssonness addressed in forthcoming work, the measurability of $\text{cof}(\kappa)$ in $\text{HOD}_{\{x\}}$ will be needed.

The belief is that the conjecture will be proved using the directed system analysis of $\text{HOD}_{\{x\}}^{L(\mathbb{R})}$ using certain iterable mice. Theorem 7.34 will provide some evidence for this conjecture by establishing that it holds for those $\kappa < \Theta$ with $\text{cof}(\kappa) = \omega_1$ assuming externally that $M_\omega^\sharp$ exists. For those set Z which appeared

²⁴The use of the coding lemma and $\text{AC}_\omega^\mathbb{R}$ in the real world is necessary. S does not belong to $\text{HOD}_{\{x\}}$ and hence one cannot make the choice of $\langle q_m : m \in \omega \rangle$ using the wellordering in $\text{HOD}_{\{x\}}$. Moreover, a q_m such that $S(m, q_m)$ cannot belong to $(\mathbb{N}_{u_m})^{\text{HOD}_{\{x\}}}$.

in the discussion before Lemma 6.6, $\text{HOD}_Z^{L(Z, \mathbb{R})}$ possesses a similar directed system analysis. Hence one can also formulate the conjecture for such models of the form $L(Z, \mathbb{R})$ and successor and double successor in $\text{HOD}_Z^{L(Z, \mathbb{R})}$. Granting such a conjecture, one can extend as usual the results from $L(\mathbb{R})$ to AD^+ .

Next, one will verify the extent of regularity for $B(\omega, \kappa)$ in $L(\mathbb{R}) \models \text{AD}$ assuming the uniform cofinality of HOD successor and double successor hypothesis in $L(\mathbb{R})$.

Theorem 7.32. *Assume AD and $V = L(\mathbb{R})$, and the uniform cofinality of HOD successor and double successor hypothesis in $L(\mathbb{R})$. Let $\omega \leq \kappa < \Theta$ be a cardinal. Then $B(\omega, \kappa)$ has δ -regular cardinality for all $\delta < \text{cof}(\kappa)$.*

Proof. When $\text{cof}(\kappa) = \omega$, Theorem 7.29 already showed that $B(\omega, \kappa)$ has n -regular cardinality for all $n \in \omega$.

Suppose $\kappa = \omega_1$. Let $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega$. Since $\omega_1 \rightarrow_* (\omega_1)_\omega^\omega, \mu_{\omega_1}^\omega$ is a countably complete measure. Thus there is a club $C \subseteq \omega_1$ and an $m \in \omega$ so that for all $f \in [C]_*^\omega$, $\Phi(f) = m$. Thus $|\Phi^{-1}[\{m\}]| = |\omega_{\omega_1}|$. ω_{ω_1} has ω -regular cardinality.

Assume $\omega_1 < \kappa$ and $\text{cof}(\kappa) > \omega$. Let $\delta < \text{cof}(\kappa)$ and $\Phi : B(\omega, \kappa) \rightarrow \delta$. Let $\rho : \text{cof}(\kappa) \rightarrow \kappa$ be an increasing cofinal map. Since $V = L(\mathbb{R})$, there is an $x \in \mathbb{R}$ so that Φ and ρ are $\text{OD}_{\{x\}}$. Thus $\rho \in \text{HOD}_{\{x\}}$ and $\text{HOD}_{\{x\}} \models \text{“cof}(\kappa) = \text{cof}(\kappa)^{L(\mathbb{R})}\text{”}$. Let $R \subseteq B(\omega, \kappa) \times \delta$ be defined by $R(f, \beta)$ if and only if $\Phi(f) = \beta$. R is $\text{OD}_{\{x\}}$ and thus there is an ∞ -Borel code $(S, \varphi) \in \text{HOD}_{\{x\}}$ for R by Fact 4.21. By the uniform cofinality of HOD successor and double successor hypothesis, let h, H, S_1 , and S_2 be as in Definition 7.30 for $\text{HOD}_{\{x\}}$ and κ . For each $\alpha < \text{cof}(\kappa)$ and $n \in \omega$, let μ_n^α be the least normal measure on $H(\alpha, n)$ according to the canonical global wellordering of $\text{HOD}_{\{x\}}$. Let $\mu_\alpha \in \text{HOD}_{\{x\}}$ be the measure sequence defined by $\kappa_n^\alpha = H(\alpha, n)$, $\mu_n^\alpha = \mu_n^\alpha$, and $\lambda^\alpha = h(\alpha)$. Since for each $\alpha < \text{cof}(\kappa)$, $\langle H(\alpha, n) : n \in \omega \rangle \in \text{HOD}_{\{x\}}$, $\mathbf{u}_\alpha \in \text{HOD}_{\{x\}}$. For each $\alpha < \text{cof}(\kappa)$, let $\tau_1^\alpha : \omega \rightarrow (h(\alpha)^+)^{\text{HOD}_{\{x\}}}$ be defined by $\tau_1^\alpha(n) = S_1(\alpha, n)$. For each $\alpha < \text{cof}(\kappa)$, let $\tau_2^\alpha : \omega \rightarrow (h(\alpha)^+)^{\text{HOD}_{\{x\}}}$ be defined by $\tau_2^\alpha(n) = S_2(\alpha, n)$. Note that $\langle \tau_i^\alpha : \alpha < \text{cof}(\kappa) \wedge i \in \{1, 2\} \rangle$ is a sequence that exists in the real world $L(\mathbb{R})$ but does not belong to $\text{HOD}_{\{x\}}$. Let $\prec$ be the global wellordering of $\text{HOD}_{\{x\}}$. By applying Lemma 7.25 to $\mathbf{u}_\alpha, \prec, \tau_1^\alpha, \tau_2^\alpha, 1_{\mathbb{N}_{\mathbf{u}_\alpha}}$, and (S, φ) uniformly for each α , one has sequences $\langle \gamma_\alpha : \alpha < \text{cof}(\kappa) \rangle$ and $\langle q_\alpha : \alpha < \text{cof}(\kappa) \rangle$ so that for all $\alpha < \text{cof}(\kappa)$ and $g \in [q_\alpha]$, $R(g, \gamma_\alpha)$ holds. By the definition of R , for all $\alpha < \text{cof}(\kappa)$ and $g \in [q_\alpha]$, $\Phi(g) = \gamma_\alpha$. Since $\langle \gamma_\alpha : \alpha < \text{cof}(\kappa) \rangle$ is a sequence below $\delta < \text{cof}(\kappa)$ and $\text{cof}(\kappa)$ is regular, there is a $\bar{\gamma}$ so that the set $A = \{\alpha < \text{cof}(\kappa) : \gamma_\alpha = \bar{\gamma}\}$ has cardinality $\text{cof}(\kappa)$. Let $Z = \bigcup_{\alpha \in A} [q_\alpha]$. Let $g \in Z$. There is a unique $\alpha \in A$ so that $g \in [q_\alpha]$. By properties of q_α and since $\gamma_\alpha = \bar{\gamma}$, one has that $\Phi(g) = \bar{\gamma}$. Thus $Z \subseteq \Phi^{-1}[\{\bar{\gamma}\}]$. Since $|Z| = |B(\omega, \kappa)|$ by Fact 7.27, one has that $|\Phi^{-1}[\{\bar{\gamma}\}]| = |B(\omega, \kappa)|$. Since $\Phi : B(\omega, \kappa) \rightarrow \delta$ was arbitrary, this shows that $B(\omega, \kappa)$ has δ -regular cardinality. Since $\delta < \text{cof}(\kappa)$ was arbitrary, this completes the proof. $\square$

Theorem 7.33. *Assume AD^+ and uniform cofinality of $\text{HOD}_Z^{L(Z, \mathbb{R})}$ successor and double successor conjecture for $L(Z, \mathbb{R})$ when Z has the property from the discussion before Lemma 6.6. Let $\omega \leq \kappa < \Theta$ be a cardinal. Then $B(\omega, \kappa)$ has δ -regular cardinality for all $\delta < \text{cof}(\kappa)$.*

Next, one will provide a sketch of the argument that if $M_\omega^\sharp$ exists, then the uniform cofinality of HOD successor and double successor hypothesis hold for $L(\mathbb{R})$ below δ_1^2 at cofinality ω_1 .

Theorem 7.34. *Assume $M_\omega^\sharp$ exists. Then $L(\mathbb{R}) \models \text{“if } \kappa < \delta_1^2 \text{ is a cardinal and } \text{cof}(\kappa) = \omega_1, \text{ then the uniform cofinality of HOD successor and double successor hypothesis holds at } \kappa\text{”}$.*

Proof. First one will review the relevant theory and notation of the internal directed system from [60] Section 8. By [60] Lemma 8.3, a countable, properly small, and $\mathfrak{D}^{\mathbb{R}}\Pi_1^1$ -iterable mouse M has a unique iteration strategy for a certain game $G^*(M, \omega_1, \omega_1)$ in $L(\mathbb{R})$ which is uniformly Σ_1 definable in $L(\mathbb{R})$ with M as a parameter. See [60] Definition 8.4 for the definition of correct iteration for countable, properly small, and $\mathfrak{D}^{\mathbb{R}}\Pi_1^1$ -iterable mouse. [60] Lemma 8.5 states that the notion of correct iteration is Σ_1 -definable over $L(\mathbb{R})$. Let $\mathcal{F}$ be the set of countable, properly small, $\mathfrak{D}^{\mathbb{R}}\Pi_1^1$ -iterable, and full mouse. The ordering on $\mathcal{F}$ is $\prec^*$ defined by $M \prec^* N$ if and only if M iterates correctly to a cutpoint of N (see [60] Definition 8.7), and in this case, $\pi_{M, N}$ will refer to the unique iteration map of M to a cutpoint of N . $\mathcal{F}, \prec^*$, and the map $\langle (M, N, \pi_{M, N}) : M, N \in \mathcal{F} \wedge M \prec^* N \rangle$ are ordinal definable in $L(\mathbb{R})$. Let F^+ consists of the correct iterates of M_ω by δ_0 -bounded trees. For $M, N \in F^+$, define $M \prec^+ N$ if and only if M iterates correctly to N by a δ_0 -bounded tree. (See [60] Definition 8.10, 8.11, and 8.12.) $(\mathcal{F}^+, \prec^+)$ is external to $L(\mathbb{R})$ in the sense that it

is not even definable in $L(\mathbb{R})$. However, using $(F^+, \prec^+)$, [60] Lemma 8.14 implies that the internal directed system $\langle M, N, \pi_{M,N} : M, N \in \mathcal{F} \wedge M \prec^* N \rangle$ is countably directed with wellfounded direct limit denoted M_∞ . Steel ([60] Theorem 8.20) showed that $\text{HOD}^{L(\mathbb{R})} \cap V_{\delta_1^2} = M_\infty \cap V_{\delta_1^2}$. (The discussion above can be relativized for $\text{HOD}_{\{x\}}^{L(\mathbb{R})}$ where $x \in \mathbb{R}$.)

Now work in $L(\mathbb{R})$. Let $\omega_1 < \kappa < \delta_1^2$ with $\text{cof}(\kappa) = \omega_1$. Since every cofinal map of ω_1 into κ is $\text{OD}_{\{x\}}$ for some $x \in \mathbb{R}$, there is an $x \in \mathbb{R}$ so that $\text{HOD}_{\{x\}} \models \text{cof}(\kappa) = \omega_1^{L(\mathbb{R})}$. Without loss of generality, suppose $\text{HOD} \models \text{cof}(\kappa) = \omega_1^{L(\mathbb{R})}$. Since $\omega_1 < \kappa < \Theta$, there are κ many measurable cardinal of HOD below κ by Fact 7.16. Let $\rho \in \text{HOD} \cap V_{\delta_1^2}$ be such that $\text{HOD} \cap V_{\delta_1^2} \models \text{“}\rho : \omega_1^{L(\mathbb{R})} \rightarrow \kappa \text{ is increasing cofinal and for all } \alpha < \omega_1^{L(\mathbb{R})}, \rho(\alpha) \text{ is a measurable cardinal”}$. Since $\text{HOD} \cap V_{\delta_1^2} = M_\infty \cap V_{\delta_1^2}$, there is an $N_0 \in \mathcal{F}$ so that the following holds.

- (1) Let $j_{0,\infty} : N_0 \rightarrow M_\infty$ denote the direct system map of N_0 into M_∞ .
- (2) Let $\bar{\delta}$ be the least measurable cardinal of N_0 . It can be shown that $j_{0,\infty}(\bar{\delta}) = \omega_1^{L(\mathbb{R})}$.
- (3) There is a $\bar{\kappa} \in N_0$ so that $j_{0,\infty}(\bar{\kappa}) = \kappa$.
- (4) There is a $\bar{\rho} \in N_0$ so that $j_{0,\infty}(\bar{\rho}) = \rho$. Thus $N_0 \models \text{“}\bar{\rho} : \bar{\delta} \rightarrow \bar{\kappa} \text{ is cofinal increasing and for all } \alpha < \bar{\delta}, \bar{\rho}(\alpha) \text{ is measurable”}$.
- (5) Let $\bar{\mu} \in N_0$ be such that $N_0 \models \text{“}\bar{\mu} \text{ is a measure on } \bar{\delta}\text{”}$ (on the extender sequence of N_0).
- (6) By Fact 7.16, $\text{HOD} \cap V_{\delta_1^2} \models \text{“there are } \kappa \text{ many measurable cardinals below } \kappa$. By the elementarity of $j_{0,\infty}$, $N_0 \models \text{“there are } \bar{\kappa} \text{ many measurable cardinals below } \bar{\kappa}\text{”}$. Let $i_{\bar{\mu}} : N_0 \rightarrow \text{Ult}(N_0, \bar{\mu})$ be the ultrapower map. By elementarity, $\text{Ult}(N_0, \bar{\mu}) \models \text{“there are } i_{\bar{\mu}}(\bar{\kappa})\text{-many measurable cardinals below } i_{\bar{\mu}}(\bar{\kappa})\text{”}$. Since $\text{cof}(\bar{\kappa}) = \bar{\delta}$, $i_{\bar{\mu}}(\bar{\kappa}) > \bar{\kappa}$. Since $\bar{\mu} \in N_0$, let $\iota \in \text{Ult}(N_0, \bar{\mu}) \subseteq N_0$ be such that $\text{Ult}(N_0, \bar{\mu}) \models \text{“}\iota : \omega \rightarrow i_{\bar{\mu}}(\bar{\kappa}) \text{ enumerates the first } \omega\text{-many measurable cardinals above } \bar{\kappa}\text{”}$. Note this property of ι is first order expressible in N_0 since $\bar{\mu} \in N_0$.
- (7) Let $\lambda = \sup(\iota)$ and note that $\lambda < i_{\bar{\mu}}(\kappa)$. Let $\lambda^* = (\lambda^+)^{\text{Ult}(N_0, \bar{\mu})}$ and let $\lambda^{**} = (\lambda^{++})^{\text{Ult}(N_0, \bar{\mu})}$. λ^* and λ^{**} are the ordinals which become the successor and double successor of λ in the ultrapower of N_0 by $\bar{\mu}$. Note that these facts about $\bar{\mu}$, λ , λ^* , and λ^{**} are first order expressible in N_0 .
- (8) Since N_0 is countable in $L(\mathbb{R})$, let $\ell_1 : \omega \rightarrow \lambda^*$ and $\ell_2 : \omega \rightarrow \lambda^{**}$ be increasing cofinal functions through λ^* and λ^{**} , respectively, which belong to $L(\mathbb{R})$.

For each $\alpha < \omega_1$, let $j_{0,\alpha} : N_0 \rightarrow N_\alpha$ be the α^{th} -length linear iteration of N_0 by the image of $\bar{\mu}$. Note that $N_\alpha \in \mathcal{F}$ for all $\alpha < \beta$. Let $j_{\alpha,\infty} : N_\alpha \rightarrow M_\infty$ be the direct limit map of N_α into M_∞ . Note that for all $\alpha < \omega_1$, $j_{\alpha,\infty}$ belongs to $L(\mathbb{R})$ (and this is an important reason to use the internal direct system $\mathcal{F}$ rather than the external direct system $\mathcal{F}^+$). For $\alpha < \beta < \omega$, let $j_{\alpha,\beta} : N_\alpha \rightarrow N_\beta$ denote $(\beta - \alpha)^{\text{th}}$ -length linear iteration of N_α by the image of $j_{0,\alpha}(\bar{\mu})$. Furthermore, since these are linear iterations, one can check that $j_{\alpha,\beta} = \pi_{N_\alpha, N_\beta}$ in the notation above; that is, $j_{\alpha,\beta}$ is the unique map of the correct iteration witnessing $N_\alpha \prec^* N_\beta$ according to the unique iteration strategy for the appropriate game. Thus for all $\alpha < \beta < \gamma$, $j_{\alpha,\gamma} = j_{\beta,\gamma} \circ j_{\alpha,\beta}$ and $j_{\alpha,\infty} = j_{\beta,\infty} \circ j_{\alpha,\beta}$. Define the following objects:

- (a) For $\alpha < \omega_1$ and $n \in \omega$, let $H(\alpha, n) = j_{\alpha+1,\infty}(j_{0,\alpha}(\iota))(n)$.
- (b) For $\alpha < \omega_1$, let $h(\alpha) = \sup\{H(\alpha, n) : n \in \omega\} = \sup(j_{\alpha+1,\infty}(j_{0,\alpha}(\iota))) = j_{\alpha+1,\infty}(\sup(j_{0,\alpha}(\iota))) = j_{\alpha+1,\infty}(j_{0,\alpha}(\lambda))$.
- (c) For $\alpha < \omega_1$ and $n \in \omega$, let $S_1(\alpha, n) = j_{\alpha+1,\infty}(j_{0,\alpha}(\ell_1(n)))$.
- (d) For $\alpha < \omega_1$ and $n \in \omega$, let $S_n(\alpha, n) = j_{\alpha+1,\infty}(j_{0,\alpha}(\ell_2(n)))$.

Note that $j_{0,\alpha}(\iota)$ enumerates the next ω -many measurable above $\sup(j_{0,\alpha}(\bar{\rho})) = j_{0,\alpha}(\bar{\kappa})$ in $\text{Ult}(N_\alpha, j_{0,\alpha}(\bar{\mu})) = N_{\alpha+1}$ by (6) above. Note that $\sup(j_{0,\alpha+1}(\bar{\rho})) = j_{0,\alpha+1}(\bar{\kappa}) > j_{0,\alpha}(\bar{\kappa}) = \sup(j_{0,\alpha}(\bar{\rho}))$. By (4), $N_{\alpha+1}$ sees that $j_{0,\alpha+1}(\bar{\kappa}) = \sup(j_{0,\alpha+1}(\bar{\rho}))$ is a limit of measurable cardinals. So $\sup(j_{0,\alpha}(\iota)) = j_{0,\alpha}(\lambda) < j_{0,\alpha+1}(\bar{\kappa})$ since the supremum of the first ω -many measurable cardinals of $N_{\alpha+1}$ above $j_{0,\alpha}(\bar{\kappa})$ is less than $j_{0,\alpha+1}(\bar{\kappa})$ which is above $j_{0,\alpha}(\bar{\kappa})$ and is a limit of $j_{0,\alpha}(\bar{\delta})$ many measurable cardinals of $N_{\alpha+1}$. Similarly, $j_{0,\alpha}(\lambda^*) < j_{0,\alpha}(\lambda^{**}) < j_{0,\alpha+1}(\bar{\kappa})$ since these are the successor and double successor of $j_{0,\alpha}(\lambda)$ in $N_{\alpha+1}$. Then

$$H(\alpha, n) = j_{\alpha+1,\infty}(j_{0,\alpha}(\iota))(n) < h(\alpha) = j_{\alpha+1,\infty}(\sup(j_{0,\alpha}(\iota))) = j_{\alpha+1,\infty}(j_{0,\alpha}(\lambda))$$

$$< j_{\alpha+1,\infty}(j_{0,\alpha}(\lambda^*)) < j_{\alpha+1,\infty}(j_{0,\alpha}(\lambda^{**})) < j_{\alpha+1,\infty}(j_{0,\alpha+1}(\bar{\kappa})) = \kappa.$$

This shows that that $H : \omega_1 \times \omega_1 \rightarrow \kappa$ and $h : \omega_1 \rightarrow \kappa$. Also $S_1(\alpha, n) = j_{\alpha+1, \infty}(j_{0, \alpha}(\ell_1(n))) < j_{\alpha+1, \infty}(j_{0, \alpha}(\lambda^*)) < \kappa$. Similarly, $S_2(\alpha, n) < \kappa$. This shows that $S_1 : \omega_1 \times \omega \rightarrow \kappa$ and $S_2 : \omega_1 \times \omega \rightarrow \kappa$.

Since ι , ℓ_1 , and ℓ_2 are increasing functions, Definition 7.30 property (1) holds by the elementary $j_{0, \alpha}$ and $j_{\alpha+1, \infty}$. Since $j_{0, \alpha}(\iota)$ enumerates the next ω -many measurable cardinals above $j_{0, \alpha}(\bar{\kappa})$ in $\text{Ult}(N_\alpha, j_{0, \alpha}(\bar{\mu})) = N_{\alpha+1}$, $j_{\alpha+1, \infty}(j_{0, \alpha}(\iota)) \in M_\infty$ enumerates the next ω -many measurable cardinals above $j_{\alpha+1, \infty}(j_{0, \alpha}(\bar{\kappa}))$ in $M_\infty \cap V_{\delta_1^2} = \text{HOD} \cap V_{\delta_1^2}$. Thus $\langle H(\alpha, n) : n \in \omega \rangle = j_{\alpha+1, \infty}(j_{0, \alpha}(\iota))$ belongs to HOD which establishes Definition 7.30 property (2) and it enumerates the next ω -many measurable cardinal in HOD above $j_{\alpha+1, \infty}(j_{0, \alpha}(\bar{\kappa}))$. In particular, for all $n \in \omega$, $H(\alpha, n) = j_{\alpha+1, \infty}(j_{0, \alpha}(\iota))(n)$ is a measurable cardinal in HOD which establishes Definition 7.30 property (3). Definition 7.30 property (4) holds by definition of h . For any $\alpha < \omega_1$, $j_{0, \alpha}(\bar{\delta}) > \alpha$ and $j_{0, \alpha}(\bar{\delta}) = \text{dom}(j_{0, \alpha}(\bar{\rho}))$. By (4) above, $N_\alpha \models j_{0, \alpha}(\bar{\rho})(\alpha)$ is a measurable cardinal. Since $j_{\alpha, \alpha+1} : N_\alpha \rightarrow \text{Ult}(N_\alpha, j_{0, \alpha}(\bar{\mu})) = N_{\alpha+1}$ is just the ultrapower map of $j_{0, \alpha}(\bar{\mu})$ and this ultrapower map fixes all strong limit cardinals of cofinality not equal to $j_{0, \alpha}(\bar{\delta})$, one has that $j_{0, \alpha+1}(\bar{\rho})(\alpha) = j_{\alpha, \alpha+1}(j_{0, \alpha}(\bar{\rho})(\alpha)) = j_{0, \alpha}(\bar{\rho})(\alpha)$. Clearly, $j_{0, \alpha}(\bar{\rho})(\alpha) < \sup(j_{0, \alpha}(\bar{\rho})) = j_{0, \alpha}(\bar{\kappa}) < j_{0, \alpha}(\lambda)$. Thus one has

$$\rho(\alpha) = j_{0, \infty}(\bar{\rho})(\alpha) = j_{\alpha+1, \infty}(j_{0, \alpha+1}(\bar{\rho})(\alpha)) = j_{\alpha+1, \infty}(j_{0, \alpha}(\bar{\rho})(\alpha)) < j_{\alpha+1, \infty}(j_{0, \alpha}(\lambda)) = h(\alpha)$$

Since $\alpha < \omega_1^{L(\mathbb{R})}$ was arbitrary and $\sup(\rho) = \kappa$, one has that $\sup(h) = \kappa$. Thus h is cofinal in κ which establishes Definition 7.30 (5). Note that since $\text{Ult}(N_0, \bar{\mu}) \models \lambda^* = \lambda^+$ by (7), $\text{Ult}(N_0, \bar{\mu}) \models \lambda^*$ is regular. Since N_0 and $\text{Ult}(N_0, \bar{\mu})$ have the same $\bar{\delta}$ -sequences, $N_0 \models \text{cof}(\lambda^*) > \bar{\delta}$. Since $j_{0, \alpha}$ is a linear iteration by $\bar{\mu}$ which is a measure on $\bar{\delta}$ and its image, $j_{0, \alpha}$ is continuous at any ordinal whose cofinality is greater than $\bar{\delta}$. Hence $j_{0, \alpha}$ is continuous at λ^* . Since $\sup(\ell_1) = \lambda^*$, $\sup[j_{0, \alpha} \circ \ell_1] = \sup\{j_{0, \alpha}(\ell_1(n)) : n \in \omega\} = j_{0, \alpha}(\lambda^*)$. By (6), $N_{\alpha+1} \models j_{0, \alpha}(\lambda^*) = j_{0, \alpha}(\lambda)^+$. Thus $N_{\alpha+1} \models "j_{0, \alpha}(\lambda^*) \text{ is regular non-measurable cardinal}"$. The direct limit map $j_{\alpha+1, \infty}$ is continuous at all regular non-measurable cardinals of $N_{\alpha+1}$ (see the remark in the proof of [60] Lemma 8.25). Since $\sup[j_{0, \alpha} \circ \ell_1] = j_{0, \alpha}(\lambda^*)$, $\sup[j_{\alpha+1, \infty} \circ j_{0, \alpha} \circ \ell_1] = j_{\alpha+1, \infty}(j_{0, \alpha}(\lambda^*))$. Thus $\sup\{S_1(\alpha, n) : n \in \omega\} = \sup\{j_{\alpha+1, \infty}(j_{0, \alpha}(\ell_1(n))) : n \in \omega\} = j_{\alpha+1, \infty}(j_{0, \alpha}(\lambda^*))$. Since $N_{\alpha+1} \models j_{0, \alpha}(\lambda^*) = j_{0, \alpha}(\lambda)^+$, one has that $M_\infty \models j_{\alpha+1, \infty}(j_{0, \alpha}(\lambda^*)) = j_{0, \alpha+1}(j_{0, \alpha}(\lambda))^+ = h(\alpha)^+$. Since $M_\infty \cap V_{\delta_1^2} = \text{HOD} \cap V_{\delta_1^2}$, $\text{HOD} \models j_{0, \alpha+1}(j_{0, \alpha}(\lambda^*)) = h(\alpha)^+$. This establishes Definition 7.30 (6). Definition 7.30 (7) is shown in an analogous manner. This completes the proof. $\square$

To remove the restriction that $\kappa < \delta_1^2$ in Theorem 7.34, one can perhaps use the more complex internal directed system of [64] Section 6. To remove the use of the external assumption of $M_\omega^\sharp$, one can possibly use the idea of [64] Section 7. Linear iterations cannot be used once the cofinality of κ is greater than ω_1 . It is anticipated that the uniform cofinality of HOD successor and double successor hypothesis in its full generality should be established through the direct system analysis of HOD . In conversation with Schlutzenburg, it seems that Schlutzenburg's result on normalization [51] which implies HOD is a normal iterate of M_ω may suggest a path forward.

REFERENCES

1. Uri Abraham, *Proper forcing*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 333–394. MR 2768684
2. Douglas Blue, Paul Larson, and Grigor Sargsyan, *The failure of square at all uncountable cardinals is weaker than a woodin limit of woodin cardinals*, 2026.
3. Douglas Blue, Paul B. Larson, and Grigor Sargsyan, *Nairian models*, 2025.
4. Andrés Eduardo Caicedo and Richard Ketchersid, *A trichotomy theorem in natural models of AD^+* , Set theory and its applications, Contemp. Math., vol. 533, Amer. Math. Soc., Providence, RI, 2011, pp. 227–258. MR 2777751
5. William Chan, *Basis for uncountable linear orders*, In preparation.
6. ———, *Definable combinatorics of strong partition cardinals*, In preparation.
7. William Chan, *Optimal partition relations for omega sequences through uncountable partition cardinals*, In preparation.
8. ———, *An introduction to combinatorics of determinacy*, Trends in Set Theory, Contemp. Math., vol. 752, Amer. Math. Soc., Providence, RI, 2020, pp. 21–75. MR 4132099
9. William Chan and Stephen Jackson, *$L(\mathbb{R})$ with determinacy satisfies the Suslin hypothesis*, Adv. Math. **346** (2019), 305–328. MR 3910797
10. William Chan and Stephen Jackson, *Cardinality of wellordered disjoint unions of quotients of smooth equivalence relations*, Ann. Pure Appl. Logic **172** (2021), no. 8, 102988. MR 4256846
11. ———, *Definable combinatorics at the first uncountable cardinal*, Trans. Amer. Math. Soc. **374** (2021), no. 3, 2035–2056. MR 4216731

- 2537 12. ———, *The destruction of the axiom of determinacy by forcings on $\mathbb{R}$ when Θ is regular*, Israel J. Math. **241** (2021),
2538 no. 1, 119–138. MR 4242147
- 2539 13. ———, *Applications of infinity-Borel codes to definability and definable cardinals*, Fund. Math. **265** (2024), no. 3, 215–258.
2540 MR 4771874
- 2541 14. William Chan, Stephen Jackson, and Nam Trang, *Boldface GCH below the first uncountable limit cardinal*, In preparation.
- 2542 15. ———, *∞ -Borel codes in natural models of AD^+* , Submitted.
- 2543 16. William Chan, Stephen Jackson, and Nam Trang, *The primeness property for cardinalities*, In preparation.
- 2544 17. William Chan, Stephen Jackson, and Nam Trang, *A search for the cofinality of the power set of the first uncountable*
2545 *cardinal*, In preparation.
- 2546 18. William Chan, Stephen Jackson, and Nam Trang, *The size of the class of countable sequences of ordinals*, Trans. Amer.
2547 Math. Soc. **375** (2022), no. 3, 1725–1743. MR 4378077
- 2548 19. ———, *More definable combinatorics around the first and second uncountable cardinals*, J. Math. Log. **23** (2023), no. 3,
2549 Paper No. 2250029, 31. MR 4603918
- 2550 20. ———, *Almost disjoint families under determinacy*, Adv. Math. **437** (2024), Paper No. 109410, 34. MR 4669334
- 2551 21. ———, *Almost Everywhere Behavior of Functions According to Partition Measures*, Forum Math. Sigma **12** (2024), Paper
2552 No. e16. MR 4696011
- 2553 22. Paul Cohen, *The independence of the continuum hypothesis*, Proc. Nat. Acad. Sci. U.S.A. **50** (1963), 1143–1148.
2554 MR 0157890
- 2555 23. Sean Cox and John Krueger, *Namba forcing, weak approximation, and guessing*, J. Symb. Log. **83** (2018), no. 4, 1539–1565.
2556 MR 3893287
- 2557 24. Leo A. Harrington and Alexander S. Kechris, *On the determinacy of games on ordinals*, Ann. Math. Logic **20** (1981), no. 2,
2558 109–154. MR 622782
- 2559 25. Daisuke Ikegami and Nam Trang, *On supercompactness of ω_1* , Advances in mathematical logic, Springer Proc. Math. Stat.,
2560 vol. 369, Springer, Singapore, [2021] ©2021, pp. 27–45. MR 4378920
- 2561 26. Daisuke Ikegami and Nam Trang, *Preservation of AD via forcings*, 2023.
- 2562 27. S. Jackson, R. Ketchersid, F. Schlutzenberg, and W. H. Woodin, *Determinacy and Jónsson cardinals in $L(\mathbb{R})$* , J. Symb.
2563 Log. **79** (2014), no. 4, 1184–1198. MR 3343535
- 2564 28. Steve Jackson, *AD and the projective ordinals*, Cabal Seminar 81–85, Lecture Notes in Math., vol. 1333, Springer, Berlin,
2565 1988, pp. 117–220. MR 960900
- 2566 29. ———, *A computation of δ^1_5* , Mem. Amer. Math. Soc. **140** (1999), no. 670, viii+94. MR 1606035
- 2567 30. ———, *Structural consequences of AD*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1753–1876.
2568 MR 2768700
- 2569 31. Thomas Jech, *Set theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003, The third millennium
2570 edition, revised and expanded. MR 1940513 (2004g:03071)
- 2571 32. A. S. Kechris, *Countable ordinals and the analytical hierarchy. I*, Pacific J. Math. **60** (1975), no. 1, 223–227. MR 387053
- 2572 33. Alexander S. Kechris, *The theory of countable analytical sets*, Trans. Amer. Math. Soc. **202** (1975), 259–297. MR 419235
- 2573 34. ———, *AD and projective ordinals*, Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77), Lecture Notes in
2574 Math., vol. 689, Springer, Berlin, 1978, pp. 91–132. MR 526915
- 2575 35. ———, *The axiom of determinacy implies dependent choices in $L(\mathbb{R})$* , J. Symbolic Logic **49** (1984), no. 1, 161–173.
2576 MR 736611
- 2577 36. ———, *“AD + UNIFORMIZATION” is equivalent to “Half $\text{AD}_{\mathbb{R}}$ ”*, Cabal Seminar 81–85, Lecture Notes in Math., vol.
2578 1333, Springer, Berlin, 1988, pp. 98–102. MR 960897
- 2579 37. Alexander S. Kechris, Eugene M. Kleinberg, Yiannis N. Moschovakis, and W. Hugh Woodin, *The axiom of determinacy,*
2580 *strong partition properties and nonsingular measures*, Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79),
2581 Lecture Notes in Math., vol. 839, Springer, Berlin-New York, 1981, pp. 75–99. MR 611168
- 2582 38. Alexander S. Kechris and W. Hugh Woodin, *Generic codes for uncountable ordinals, partition properties, and elementary*
2583 *embeddings*, Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I, Lect. Notes Log., vol. 31, Assoc. Symbol.
2584 Logic, Chicago, IL, 2008, pp. 379–397. MR 2463619
- 2585 39. Eugene M. Kleinberg, *Infinitary combinatorics and the axiom of determinateness*, Lecture Notes in Mathematics, Vol. 612,
2586 Springer-Verlag, Berlin-New York, 1977. MR 0479903
- 2587 40. Peter Koellner and W. Hugh Woodin, *Large cardinals from determinacy*, Handbook of set theory. Vols. 1, 2, 3, Springer,
2588 Dordrecht, 2010, pp. 1951–2119. MR 2768702
- 2589 41. Kenneth Kunen, *Set theory*, Studies in Logic (London), vol. 34, College Publications, London, 2011. MR 2905394
- 2590 42. Paul B. Larson, *Extensions of the axiom of determinacy*, University Lecture Series, vol. 78, American Mathematical Society,
2591 Providence, RI, 2023. MR 4658673
- 2592 43. Donald A. Martin and John R. Steel, *The extent of scales in $L(\mathbb{R})$* , Cabal seminar 79–81, Lecture Notes in Math., vol. 1019,
2593 Springer, Berlin, 1983, pp. 86–96. MR 730590
- 2594 44. Yiannis N. Moschovakis, *Determinacy and prewellorderings of the continuum*, Mathematical Logic and Foundations of Set
2595 Theory (Proc. Internat. Colloq., Jerusalem, 1968), Stud. Logic Found. Math., North-Holland, Amsterdam-London, 1970,
2596 pp. 24–62. MR 280362
- 2597 45. ———, *Descriptive set theory*, second ed., Mathematical Surveys and Monographs, vol. 155, American Mathematical
2598 Society, Providence, RI, 2009. MR 2526093
- 2599 46. Itay Neeman, *Inner models and ultrafilters in $L(\mathbb{R})$* , Bull. Symbolic Logic **13** (2007), no. 1, 31–53. MR 2300902

- 2600 47. Yinhe Peng and Liang Yu, *TD implies $CC_{\mathbb{R}}$* , Adv. Math. **384** (2021), Paper No. 107755, 7. MR 4246103
- 2601 48. Grigor Sargsyan, *A tale of hybrid mice*, ProQuest LLC, Ann Arbor, MI, 2009, Thesis (Ph.D.)–University of California,
- 2602 Berkeley. MR 2714009
- 2603 49. ———, *Hod mice and the mouse set conjecture*, Mem. Amer. Math. Soc. **236** (2015), no. 1111, viii+172. MR 3362806
- 2604 50. Ralf Schindler, *Set theory*, Universitext, Springer, Cham, 2014, Exploring independence and truth. MR 3243739
- 2605 51. Farmer Schlutzenberg, *Iterability for (transfinite) stacks*, J. Math. Log. **21** (2021), no. 2, Paper No. 2150008, 117.
- 2606 MR 4290497
- 2607 52. Farmer Schlutzenberg and John R. Steel, *Stationary tower free homogeneously Suslin scales*, arXiv e-prints (2024),
- 2608 arXiv:2406.02727.
- 2609 53. David Schritterser and Asger Törnquist, *The Ramsey property implies no mad families*, Proc. Natl. Acad. Sci. USA **116**
- 2610 (2019), no. 38, 18883–18887. MR 4012549
- 2611 54. Robert M. Solovay, *A model of set-theory in which every set of reals is Lebesgue measurable*, Ann. of Math. (2) **92** (1970),
- 2612 1–56. MR 0265151 (42 #64)
- 2613 55. ———, *The independence of DC from AD*, Large cardinals, determinacy and other topics. The Cabal Seminar. Vol. IV,
- 2614 Lect. Notes Log., vol. 49, Cambridge Univ. Press, Cambridge, 2021, pp. 66–95. MR 4567353
- 2615 56. John Steel and Nam Trang, *AD^+ , derived models, and Σ_1 -reflection*, [https://math.berkeley.edu/~steel/papers/AD+](https://math.berkeley.edu/~steel/papers/AD+reflection.pdf)
- 2616 [reflection.pdf](https://math.berkeley.edu/~steel/papers/AD+reflection.pdf).
- 2617 57. John R. Steel, *Scales in $L(\mathbb{R})$* , Cabal seminar 79–81, Lecture Notes in Math., vol. 1019, Springer, Berlin, 1983, pp. 107–156.
- 2618 MR 730592
- 2619 58. ———, *$HOD^{L(\mathbb{R})}$ is a core model below Θ* , Bull. Symbolic Logic **1** (1995), no. 1, 75–84. MR 1324625
- 2620 59. ———, *Derived models associated to mice*, Computational prospects of infinity. Part I. Tutorials, Lect. Notes Ser. Inst.
- 2621 Math. Sci. Natl. Univ. Singap., vol. 14, World Sci. Publ., Hackensack, NJ, 2008, pp. 105–193. MR 2449479
- 2622 60. ———, *An outline of inner model theory*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1595–1684.
- 2623 MR 2768698
- 2624 61. ———, *Ordinal definability in models of determinacy. Introduction to Part V*, Ordinal definability and recursion theory:
- 2625 The Cabal Seminar. Vol. III, Lect. Notes Log., vol. 43, Assoc. Symbol. Logic, Ithaca, NY, 2016, pp. 3–48. MR 3469165
- 2626 62. ———, *A theorem of Woodin on mouse sets*, Ordinal definability and recursion theory: The Cabal Seminar. Vol. III, Lect.
- 2627 Notes Log., vol. 43, Assoc. Symbol. Logic, Ithaca, NY, 2016, pp. 243–256. MR 3469172
- 2628 63. ———, *A comparison process for mouse pairs*, Lecture Notes in Logic, vol. 51, Association for Symbolic Logic, Ithaca,
- 2629 NY; Cambridge University Press, Cambridge, [2023] ©2023. MR 4496338
- 2630 64. John R. Steel and W. Hugh Woodin, ***HOD** as a core model*, Ordinal definability and recursion theory: The Cabal Seminar.
- 2631 Vol. III, Lect. Notes Log., vol. 43, Assoc. Symbol. Logic, Ithaca, NY, 2016, pp. 257–345. MR 3469173
- 2632 65. W. Hugh Woodin, *The cardinals below $|\omega_1|^{<\omega_1}$* , Ann. Pure Appl. Logic **140** (2006), no. 1-3, 161–232. MR 2224057
- 2633 66. ———, *AD and the uniqueness of the supercompact measures on $\mathcal{P}_{\omega_1}(\lambda)$* , Large cardinals, determinacy and other topics.
- 2634 The Cabal Seminar. Vol. IV, Lect. Notes Log., vol. 49, Cambridge Univ. Press, Cambridge, 2021, pp. 136–140. MR 4567357
- 2635 INSTITUTE FOR DISCRETE MATHEMATICS AND GEOMETRY, VIENNA UNIVERSITY OF TECHNOLOGY, VIENNA, AUSTRIA
- 2636 Email address: William.Chan@tuwien.ac.at