

BOLDFACE GCH BELOW THE FIRST UNCOUNTABLE LIMIT CARDINAL

WILLIAM CHAN, STEPHEN JACKSON, AND NAM TRANG

ABSTRACT. If κ is an infinite cardinal, the boldface GCH at κ is the statement that κ^+ does not inject into $\mathcal{P}(\kappa)$. It will be shown here that $\omega_1 \rightarrow (\omega_1)_2^{\omega_1}$ (the strong partition property at ω_1) and $j_{\mu_{\omega_1}}^1(\omega_1) = \omega_2$ (the ultrapower of ω_1 by the club filter on ω_1 is ω_2) imply that the boldface GCH holds at ω_n for all $n < \omega$ using combinatorial arguments. In particular, AD implies the boldface GCH holds at ω_n for all $n < \omega$.

1. INTRODUCTION

This paper will work with the Zermelo-Frankel axiom ZF for set theory (without the axiom of choice, AC). Let κ be an infinite cardinal. There is a cardinal which does not inject into $\mathcal{P}(\kappa)$. What is the smallest cardinal which does not inject into $\mathcal{P}(\kappa)$? Since κ always injects into $\mathcal{P}(\kappa)$, the smallest that this cardinal can be is κ^+ , the cardinal successor of κ . Cantor showed that κ does not surject onto $\mathcal{P}(\kappa)$. Thus $|\kappa| < |\mathcal{P}(\kappa)|$. If the axiom of choice holds, then all sets are wellorderable and one must have that κ^+ injects into $\mathcal{P}(\kappa)$. Assuming the axiom of choice, the smallest cardinal which does not inject into $\mathcal{P}(\kappa)$ must be greater than κ^+ . The usual generalized continuum hypothesis at κ (under AC) is the assertion that $|\mathcal{P}(\kappa)| = 2^\kappa = \kappa^+$. Assuming AC and the generalized continuum hypothesis at κ , one has that κ^{++} is the smallest cardinal which does not inject into $\mathcal{P}(\kappa)$. However, without the axiom of choice, it is potentially possible to have the most elegant answer to the above question: κ^+ is the smallest cardinal that does not inject into $\mathcal{P}(\kappa)$. Steel ([22], Theorem 8.26) calls this phenomenon the boldface GCH at κ which is the assertion that κ^+ does not inject into $\mathcal{P}(\kappa)$. Say that the boldface GCH holds below κ if the boldface GCH holds for all $\delta < \kappa$.

The axiom of determinacy asserts that every infinite two player game on ω has a winning strategy for one of the two players. AD^+ is Woodin's extension of the axiom of determinacy. The axiom of determinacy influences most strongly the sets which are surjective images of $\mathbb{R}$. Let Θ be the supremum of the ordinals onto which $\mathbb{R}$ surjects. Steel ([22] Theorem 8.26) showed that in $L(\mathbb{R})$, the boldface GCH holds below Θ . Woodin ([23] Theorem 2.16) extended these methods to show that AD^+ proves the boldface GCH holds below Θ . Steel and Woodin used an inner model theoretic analysis of HOD-type submodels to establish these results. This paper will prove the boldface GCH below ω_ω using combinatorial hypotheses which follow from AD. The methods here have been adapted to address other combinatorial properties of nonwellorderable set (such as sets of sequences through ω_ω) for which no other techniques are known.

The boldface GCH at ω or the statement that there are no uncountable wellorderable subsets of $\mathbb{R}$ is a very important property of many nice choiceless frameworks for the set theoretic universe. The boldface GCH at ω follows from classical regularity properties. If countable choice for $\mathbb{R}$, $\text{AC}_\omega^\mathbb{R}$, holds and all subsets of $\mathbb{R}$ have the property of Baire, then wellordered unions of meager sets are meager. This implies $\mathbb{R}$ is not wellorderable. If all subsets of $\mathbb{R}$ have the perfect set property, then every uncountable subset of $\mathbb{R}$ is not wellorderable. Thus the boldface GCH at ω holds under $\text{AC}_\omega^\mathbb{R}$ and all subsets of $\mathbb{R}$ have the perfect set property. If ω_1 is measurable (there is a countably complete nonprincipal ultrafilter on ω_1), then also the boldface GCH at ω holds (see Fact 3.2). These properties are all consequences of the axiom of determinacy.

The boldface GCH at ω is very important for the basic theory of determinacy. One important consequence is that if the boldface GCH at ω holds, M is an inner model of ZFC, and $\mathbb{P} \in M$ is a forcing which is countable in the real world, then in the real world, there is a generic $G \subseteq \mathbb{P}$ which is $\mathbb{P}$ -generic over M . The existence of generics for forcings countable in the real world is used in Woodin's analysis of nice models of AD^+ as symmetric extension of their HOD-type submodels using Vopěnka forcing or ordinal definable ∞ -Borel code

June 26, 2026. The first author was supported by the Austrian Science Fund (FWF) 10.55776/Y1498. The third author was supported by NSF grant DMS-1945592.

forcing. The boldface GCH at ω synergizes well with the Baire property. For example, Woodin ([19] Theorem 5.42 Claim 2) showed that $\text{AC}_\omega^\mathbb{R}$, the boldface GCH at ω , and all subsets of $\mathbb{R}$ have the Baire property, then for any set A , if $\mathbb{P} \in \text{HOD}_{\{A\}}$ is a forcing which is countable in the real world, then there is a comeager set of $G \subseteq \mathbb{P}$ which are $\mathbb{P}$ -generic over $\text{HOD}_{\{A\}}$ and moreover $\text{HOD}_{\{A\}}[G] = \text{HOD}_{\{A, G\}}$. Recently, [3] used this observation of Woodin to show the following cardinality computations: Assume $\text{AC}_\omega^\mathbb{R}$, all subsets of $\mathbb{R}$ have the Baire property, and the boldface GCH at ω holds, then $|\omega_{\omega_1}| < |^{<\omega_1}\omega_1|$, ${}^\omega\omega_1$ does not inject into $\mathbb{R} \times \text{ON}$, and S_1 does not inject into ${}^\omega\omega_1$ (where $S_1 = \{f \in [\omega_1]^{<\omega_1} : \sup(f) = \omega_1^{L[f]}\}$).

The boldface GCH at ω and all subsets of $\mathbb{R}$ have the Baire property proves the following result: ([13] Proposition 3.5) For every $\Phi : \mathbb{R} \rightarrow \mathcal{P}(\text{ON})$, there exists a comeager K and countable $\mathcal{E} \subseteq \mathcal{P}(\text{ON})$ so that for all $r \in K$, there exists an $\mathcal{F} \subseteq \mathcal{E}$ so that $\Phi(r) = \bigcup \mathcal{F}$. This result is used to prove some interesting combinatorial results under AD^+ . By [13] Lemma 3.8 and Theorem 4.3, under AD^+ , if $\kappa < \Theta$ is a cardinal of uncountable cofinality, then there are no maximal almost disjoint families $\mathcal{A}$ on κ such that $\neg(|\mathcal{A}| < \text{cof}(\kappa))$. More recently, the above fact was used to obtain large sets with respect to a normal measure or partition filters which are simultaneous homogeneous for many partitions. This is used in [2] to show under AD^+ that there is a four-element basis for linear ordering on $\mathbb{R} \times \kappa$ when $\kappa < \Theta$ is a regular cardinal and there is a twelve-element basis for the linear orderings on $\mathbb{R} \times \kappa$ when $\kappa < \Theta$ is a singular cardinal of uncountable cofinality.

The general boldface GCH plays an important role in the structure of the cardinality of sets which are nonwellorderable but linearly orderable (or equivalently, sets which are in bijection with subsets of the power set of an ordinal). If κ is a cardinal, let $\mathcal{P}_B(\kappa)$ be the set of bounded subsets of κ . By [3] and [8] Theorem 4.8, if the boldface GCH holds below κ , then $\neg(|[\kappa]^{\text{cof}(\kappa)}| \leq |\mathcal{P}_B(\kappa)|)$. If κ is regular cardinal and the boldface GCH at κ holds, then $|[\kappa]^{<\kappa}| < |\mathcal{P}(\kappa)|$. Let $B(\omega, \kappa)$ be the set of all $f : \omega \rightarrow \kappa$ such that $\sup(f) < \kappa$. If $\text{cof}(\kappa) > \omega$, then ${}^\omega\kappa = B(\omega, \kappa)$. However, [3] shows that if the boldface GCH holds below κ , then $|B(\omega, \kappa)| < |{}^\omega\kappa|$ if $\text{cof}(\kappa) = \omega$.

Steel's and Woodin's result that the boldface GCH holds below Θ can be regarded as the first step in classifying the cardinal exponentiations below Θ . Substantial evidence from [7], [8], [12], [11], and [14] suggests that cardinal exponentiation follows a very elegant simple behavior called the ABCD Conjecture: Under AD^+ , for all cardinals $\omega \leq \alpha \leq \beta < \Theta$ and $\omega \leq \gamma \leq \delta < \Theta$, $|\alpha^\beta| \leq |\gamma^\delta|$ if and only if $\alpha \leq \gamma$ and $\beta \leq \delta$. Recently, [3] showed that under AD^+ , if $\omega < \kappa < \Theta$ and $\epsilon < \kappa$, then $\mathcal{P}_B(\kappa)$ does not inject into ${}^\epsilon\text{ON}$, the class of ϵ -length sequences of ordinals. By combining the latter result and the boldface GCH below Θ , [3] proved the ABCD conjecture.

The proof of the boldface GCH below Θ uses the inner model theory analysis of HOD. First, Steel ([21], [24], and [22] Theorem 8.26) showed that if $L(\mathbb{R}) \models \text{AD}$, then $L(\mathbb{R}) \models$ "the boldface GCH below Θ ". To establish this, Steel showed that $\text{HOD}^{L(\mathbb{R})} \restriction \delta_1^2$ is a direct limit of a directed system of certain iterable mice. Woodin (as sketched in [23] Theorem 2.16) generalized this argument to show AD^+ proves the boldface GCH below Θ . To do this, one first applies Suslin-co-Suslin reflection to bring the question of the boldface GCH at some $\kappa < \Theta$ into a nice model of AD^+ . Woodin then showed that a certain HOD-type submodel of this nice AD^+ model has a direct system analysis using hybrid strategy mice.

More recently, many purely combinatorial questions of determinacy have been resolved below ω_ω or the projective ordinals by classical determinacy methods to provide evidence before a general proof using inner model theory is found. The boldface GCH at ω was known by the classical regularity properties or using the fact that ω_1 is measurable. The boldface GCH at ω_1 was known by the fact that ω_2 is measurable since it is a weak partition cardinal as shown by Martin. Remarkably, it seems that Steel established the full boldface GCH below Θ without even knowing that the boldface GCH holds at ω_2 by classical determinacy arguments.

This paper will give a proof that the boldface GCH holds below ω_ω using combinatorial methods of AD . (It should be noted that by the Moschovakis coding lemma, if $\kappa < \Theta^{L(\mathbb{R})}$, the boldface GCH at κ holds in the real world if and only if $L(\mathbb{R}) \models$ "the boldface GCH holds at κ ". Thus Steel's result actually implies that AD proves the boldface GCH below $\Theta^{L(\mathbb{R})}$.) The paper will work with combinatorial principles of ω_1 which are true in AD . $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ is the strong partition relation on ω_1 . Martin showed that AD implies $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$. See Definition 2.10 for the definition of a Kunen function. Essentially, a Kunen function bounding $f : \omega_1 \rightarrow \omega_1$ is a sequence $\langle \varphi_\alpha : \alpha < \omega_1 \rangle$ such that there is a club $C \subseteq \omega_1$ so that for all $\alpha \in C$, φ_α is a surjection of α onto $f(\alpha)$. Let $\star$ denote the following principle. (See Definition 2.10.)

★ For every function $f : \omega_1 \rightarrow \omega_1$, there is a Kunen function $\mathcal{K}$ which bounds f .

Kunen proved that AD implies every function $f : \omega_1 \rightarrow \omega_1$ has a Kunen function bounding it by defining what is known as a Kunen tree. This paper will only use $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and ★. One can show that over $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$, ★ is equivalent to $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$ where $\mu_{\omega_1}^1$ is the club filter on ω_1 . Kleinberg [18] studied the cardinals below ω_ω using the hypothesis that $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$. The principle ★ is much more directly practical than $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$. AD is the only theory in which $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and ★ (or $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$) is known to hold. AD, using the method of good coding system by Martin, is the only known theory that implies the existence of a strong partition cardinal. Radin forcing was used by Mitchell ([20]) to produce a model in which the club filter $\mu_{\omega_1}^1$ is a countably complete ultrafilter and by Woodin to produce a model in which ω_1 is a weak partition cardinal ($\omega_1 \rightarrow_* (\omega_1)_2^\epsilon$ for all $\epsilon < \omega_1$). However, it seems that AD is still the only known theory in which $\mu_{\omega_1}^1$ is a countably complete ultrafilter and $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$.

The main result of the paper is that $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and ★ imply the boldface GCH below ω_ω . The paper is completely self-contained. The combinatorial methods used here can be generalized using Jackson's theory of descriptions ([15]) for the projective ordinals to show the boldface GCH holds below the supremum of the projective ordinals, $\sup\{\delta_n^1 : n \in \omega\}$, and a bit beyond under AD. These methods show the boldface GCH at a level far below Θ . Only inner model theory is known to prove the boldface GCH below Θ assuming AD^+ .

This paper is organized as follows: Section 2 will introduce the basic concepts necessary for the paper. The correct type partition relation and the associated partition filters will be defined. Kunen functions will be introduced. The basic concepts of the Martin's ultrapower analysis below ω_ω while focusing on simple type n functions will be developed. Section 3 will prove the necessary "sliding lemma" which is the combinatorial heart of the main argument. Finally, boldface GCH below ω_ω will be shown.

2. PARTITION RELATIONS AND ULTRAPOWERS BY PARTITION FILTERS

For uncountable cardinal κ , $\epsilon \leq \kappa$ and $\gamma < \kappa$, the correct type partition relation $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ and the associated partition filters μ_κ^ϵ will be defined. Kunen functions for ω_1 will be introduced and used to both show that the ultrapower $j_{\mu_{\omega_1}^n}(\omega_1)$ is wellfounded and less than or equal to ω_{n+1} (Fact 2.19). A result of Martin (Fact 2.22) will be presented which will imply $j_{\mu_{\omega_1}^n}(\omega_1)$ are cardinals. Together these results will show that $j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$. However, for the purpose of analyzing these ultrapowers, one needs to stratify the ordinals below ω_ω by certain "types" suitable for partition arguments. This section will introduce a simple class of types called "type n " for each $1 \leq n < \omega$. Fact 2.37 will show that the ordinals below ω_{n+1} having type n has cardinality ω_{n+1} . Thus for the intended purpose here, one may as well replace ω_{n+1} with the ordinals of type n .

If X is a set and Y is a class, then $^X Y$ is the class of all functions $f : X \rightarrow Y$. If $\epsilon \in \text{ON}$ and $X \subseteq \text{ON}$ is a set, then $[X]^\epsilon$ is the set of all increasing functions $f : \epsilon \rightarrow X$. If κ is a cardinal, $\epsilon \leq \kappa$ and $\gamma < \delta$, then the ordinary partition relation $\kappa \rightarrow (\kappa)_\gamma^\epsilon$ is the assertion that for all $P : [\kappa]^\epsilon \rightarrow \gamma$, there is a $\beta < \gamma$ and an $A \subseteq \kappa$ with $|A| = \kappa$ so that for all $f \in [A]^\epsilon$, $P(f) = \beta$. However one will need the correct type partition relations here since one will be primarily interested in the ultrapowers by the partition measures obtained using the correct type partition relations.

Definition 2.1. Let $\epsilon \in \text{ON}$ and $f : \epsilon \rightarrow \text{ON}$ be a function.

- f is discontinuous everywhere if and only if for all $\alpha < \epsilon$, $\sup(f \upharpoonright \alpha) = \sup\{f(\bar{\alpha}) : \bar{\alpha} < \alpha\} < f(\alpha)$.
- f has uniform cofinality ω if and only if there is a function $F : \epsilon \times \omega \rightarrow \text{ON}$ so that for all $\alpha < \epsilon$ and $n \in \omega$, $F(\alpha, n) < F(\alpha, n+1)$ and $f(\alpha) = \sup\{F(\alpha, n) : n \in \omega\}$.
- f has the correct type if and only if f is both discontinuous everywhere and has uniform cofinality ω .

If $X \subseteq \text{ON}$ and $\epsilon \in \text{ON}$, then let $[X]_*^\epsilon$ denote the set of all increasing function $f : \epsilon \rightarrow X$ of the correct type.

Note that $[\kappa]_*^1$ is the set of ordinals below κ of cofinality ω .

Definition 2.2. Let κ be an uncountable cardinal, $\epsilon \leq \kappa$, and $\gamma < \kappa$. The correct type partition relation $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ is the statement that for all $P : [\kappa]_*^\epsilon \rightarrow \gamma$, there is a $\beta < \gamma$ and a $C \subseteq \kappa$ which is a club subset of κ so that for all $f \in [C]_*^\epsilon$, $P(f) = \beta$.

If κ is an uncountable cardinal, $\epsilon \leq \kappa$, and $\gamma < \kappa$, then $\kappa \rightarrow_* (\kappa)_\gamma^{<\epsilon}$ is the statement that for all $\bar{\epsilon} < \epsilon$, $\kappa \rightarrow_* (\kappa)_\gamma^{\bar{\epsilon}}$. If κ is an uncountable cardinal, $\epsilon \leq \kappa$, and $\gamma \leq \kappa$, then $\kappa \rightarrow_* (\kappa)_{<\gamma}^\epsilon$ is the statement that for all $\bar{\gamma} < \gamma$, $\kappa \rightarrow_* (\kappa)_{\bar{\gamma}}^\epsilon$. If κ is an uncountable cardinal, $\epsilon \leq \kappa$, and $\gamma \leq \kappa$, then $\kappa \rightarrow_* (\kappa)_{<\gamma}^{<\epsilon}$ is the statement that for all $\bar{\epsilon} < \epsilon$ and $\bar{\gamma} < \gamma$, $\kappa \rightarrow_* (\kappa)_{\bar{\gamma}}^{\bar{\epsilon}}$.

If $\kappa \rightarrow_* (\kappa)_2^{<\kappa}$, then κ is called a weak partition cardinal. If $\kappa \rightarrow_* (\kappa)_2^\kappa$, then κ is called a strong partition cardinal. If $\kappa \rightarrow_* (\kappa)_{<\kappa}^\kappa$, then κ is called a very strong partition cardinal.

One can show that $\kappa \rightarrow (\kappa)_\gamma^{\omega \cdot \epsilon}$ implies $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ and $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ implies $\kappa \rightarrow (\kappa)_\gamma^\epsilon$ for all $\epsilon \leq \kappa$ and $\gamma < \kappa$.

Note that every function of uniform cofinality ω must take range among the limit ordinals. Thus for any cardinal κ and $1 \leq \epsilon \leq \kappa$, $[\kappa]_*^\epsilon \neq \emptyset$ requires that κ be an uncountable cardinal. Thus the notions of correct type function and the correct type partition relations are only meaningful for uncountable cardinals. Partition on ω (and notions such as the Ramsey property) can only be expressed using the ordinary partition relation.

Definition 2.3. Let κ be an uncountable cardinal and $\epsilon \leq \kappa$. Define the ϵ -exponent (correct type) partition filter μ_κ^ϵ on $[\kappa]_*^\epsilon$ by $A \in \mu_\kappa^\epsilon$ if and only if there is a club $C \subseteq \kappa$ so that $[C]_*^\epsilon \subseteq A$. Note that μ_κ^1 is the ω -club filter.

If $X \subseteq \text{ON}$, then let $\text{enum}_X : \text{ot}(X) \rightarrow X$ be the increasing enumeration of X . An ordinal γ is indecomposable if and only if for all $\alpha, \beta < \gamma$, $\alpha + \beta < \gamma$ and $\alpha \cdot \beta < \gamma$. If κ is a cardinal, $X \subseteq \kappa$, $\text{ot}(X) = \kappa$, $\alpha < \kappa$, and $\gamma < \kappa$, then let $\text{next}_X^\gamma(\alpha) = \text{enum}_{X \setminus (\alpha+1)}(\gamma)$.

The following results say that if $C \subseteq \kappa$ is a club, then there is a club $D \subseteq C$ which is very thin inside of C . This club is particularly useful for many constructions.

Fact 2.4. Let κ be an uncountable regular cardinal. Let $C \subseteq \kappa$ be a club consisting entirely of indecomposable ordinals. Let $D = \{\alpha \in C : \text{enum}_C(\alpha) = \alpha\}$. Then D is a club subset of C and for any $\epsilon \in D$ and $\alpha, \beta, \gamma, \delta < \epsilon$, $\text{next}_C^{\alpha \cdot \beta + \gamma}(\delta) < \epsilon$.

Proof. D is easily seen to be closed. Let $\alpha < \kappa$. Let $\alpha_0 = \alpha + 1$. If $\alpha_n \in C$ has been defined, then let $\alpha_{n+1} = \text{enum}_C(\alpha_n + 1)$. Let $\alpha_\omega = \sup\{\alpha_n : n \in \omega\}$ and note that $\alpha < \alpha_\omega \in C$ since C is a club. For all $\beta < \alpha_\omega$, there is an $n \in \omega$ so that $\beta < \alpha_n$. Thus $\text{enum}_C(\beta) < \text{enum}_C(\alpha_n) < \text{enum}_C(\alpha_n + 1) = \alpha_{n+1} < \alpha_\omega$. Since $\{\text{enum}_C(\beta) : \beta < \alpha_\omega\} \subseteq \{\gamma \in C : \gamma < \alpha_\omega\}$, $\text{ot}\{\gamma \in C : \gamma < \alpha_\omega\} = \alpha_\omega$. Since $\alpha_\omega \in C$, $\text{enum}_C(\alpha_\omega) = \alpha_\omega$. Thus $\alpha < \alpha_\omega$ and $\alpha_\omega \in D$. This shows that D is unbounded. Thus D is a club. Now suppose $\epsilon \in D$ and $\alpha, \beta, \gamma, \delta < \epsilon$. Since $\epsilon \in D \subseteq C$ and C consists entirely of indecomposable ordinals, ϵ is an indecomposable ordinal. Since ϵ is in particular a limit ordinal and $\epsilon = \text{enum}_C(\epsilon) > \delta$, there is some $\nu < \epsilon$ so that $\delta < \text{enum}_C(\nu) < \text{enum}_C(\epsilon) = \epsilon$. Since ϵ is indecomposable, $\nu + \alpha \cdot \beta + \gamma < \epsilon$. Note that $\text{next}_C^{\alpha \cdot \beta + \gamma}(\delta) < \text{enum}_C(\nu + \alpha \cdot \beta + \gamma) < \text{enum}_C(\epsilon) = \epsilon$. $\square$

Fact 2.5. Let κ be an uncountable cardinal.

- (1) $\kappa \rightarrow_* (\kappa)_2^2$ implies that κ is regular.
- (2) For all $\epsilon \leq \kappa$, $\kappa \rightarrow_* (\kappa)_2^\epsilon$ implies μ_κ^ϵ is an ultrafilter.
- (3) For all $\epsilon \leq \kappa$ and $\gamma < \kappa$, $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$ implies μ_κ^ϵ is a γ^+ -complete ultrafilter.
- (4) If $\epsilon < \kappa$, then $\kappa \rightarrow_* (\kappa)_2^{\epsilon+}$ implies $\kappa \rightarrow_* (\kappa)_{<\kappa}^\epsilon$. Thus $\kappa \rightarrow_* (\kappa)_2^{<\kappa}$ implies $\kappa \rightarrow_* (\kappa)_{<\kappa}^{<\kappa}$.

Proof. (1) Suppose κ is not regular. Let $\delta = \text{cof}(\kappa) < \kappa$ and $\rho : \delta \rightarrow \kappa$ be an increasing cofinal function. Define $P : [\kappa]^2 \rightarrow 2$ by $P(\alpha, \beta) = 0$ if and only if there exists an $\eta < \delta$ so that $\alpha < \rho(\eta) < \beta$. By $\kappa \rightarrow_* (\kappa)_2^2$, let $C \subseteq \kappa$ be a club homogeneous for P . First, suppose C is homogeneous for P taking value 0. For each $\alpha < \kappa$, let $\eta_\alpha = \text{enum}_C(\omega \cdot \alpha + \omega)$. For all $\alpha < \kappa$, $(\eta_\alpha, \eta_{\alpha+1}) \in [C]_*^2$. $P(\eta_\alpha, \eta_{\alpha+1}) = 0$ implies there is a $\xi < \delta$ so that $\eta_\alpha < \rho(\xi) < \eta_{\alpha+1}$. Let ξ_α be the least ξ such that $\eta_\alpha < \rho(\xi) < \eta_{\alpha+1}$. For any $\alpha < \bar{\alpha} < \kappa$, $\rho(\xi_\alpha) < \eta_{\alpha+1} \leq \eta_{\bar{\alpha}} < \rho(\xi_{\bar{\alpha}})$. Since ρ is an increasing function, this implies that $\langle \xi_\alpha : \alpha < \kappa \rangle$ is an increasing function of κ into δ which is impossible since $\delta < \kappa$. Next, suppose C is homogeneous for P taking value 1. Let α be any element of $[C]_*^1$. Since ρ is cofinal, fix $\bar{\xi} < \delta$ so that $\alpha < \rho(\bar{\xi})$. Since C is a club, let β be any element of $[C]_*^1$ so that $\rho(\bar{\xi}) < \beta$. Thus $\alpha < \rho(\bar{\xi}) < \beta$. However, $P(\alpha, \beta) = 0$ implies that there is no $\xi < \delta$ with $\alpha < \rho(\xi) < \beta$ which is contradiction. So C is not homogeneous for P which also a contradiction.

(2) Let $X \subseteq [\kappa]_*^\epsilon$. Define $P_X : [\kappa]^\epsilon \rightarrow 2$ by $P_X(\ell) = 1$ if and only if $\ell \in X$. By $\kappa \rightarrow_* (\kappa)_2^\epsilon$, there is a club C homogeneous for P . If C is homogeneous for P taking value 1, then $[C]_*^\epsilon \subseteq X$ and hence $X \in \mu_\kappa^\epsilon$. If C is homogeneous for P taking value 0, then $[C]_*^\epsilon \subseteq [\kappa]_*^\epsilon \setminus X$ and thus $[\kappa]_*^\epsilon \setminus X \in \mu_\kappa^\epsilon$.

(3) Suppose μ_κ^ϵ is not γ^+ -complete. Let $\delta < \gamma^+$ and $\langle X_\xi : \xi < \delta \rangle$ is a sequence in μ_κ^ϵ such that $\bigcap_{\xi < \delta} X_\xi \notin \mu_\kappa^\epsilon$. Let $\phi : \gamma \rightarrow \delta$ be a surjection. For $\eta < \gamma$, let $Y_\eta = X_{\phi(\eta)}$ and note that $\langle Y_\eta : \eta < \gamma \rangle$ is a sequence in μ_κ^ϵ and $\bigcap_{\eta < \gamma} Y_\eta \notin \mu_\kappa^\epsilon$. Let $C_0 \subseteq \kappa$ be a club so that $[C_0]_*^\epsilon \subseteq [\kappa]_*^\epsilon \setminus \bigcap_{\eta < \gamma} Y_\eta$. Define $P : [C_0]_*^\epsilon \rightarrow \gamma$ by $P(\ell)$ is the least $\eta < \gamma$ so that $\ell \notin Y_\eta$. By $\kappa \rightarrow_* (\kappa)_\gamma^\epsilon$, there is an $\bar{\eta} < \gamma$ and a club $C_1 \subseteq C_0$ so that for all $\ell \in [C_1]_*^\epsilon$, $P(\ell) = \bar{\eta}$. Thus $[C_1]_*^\epsilon \cap Y_{\bar{\eta}} = \emptyset$. Thus $Y_{\bar{\eta}} \notin \mu_\kappa^\epsilon$. Contradiction.

(4) Let $\gamma < \kappa$ and $P : [\kappa]_*^\epsilon \rightarrow \gamma$. If $\ell \in [\kappa]_*^{\epsilon+\epsilon}$, then let $\ell^0, \ell^1 \in [\kappa]^\epsilon$ be defined by $\ell^0 = \ell \restriction \epsilon$ and $\ell^1(\alpha) = \ell(\epsilon + \alpha)$. Define $Q_0 : [\kappa]_*^{\epsilon+\epsilon} \rightarrow 2$ by $Q_0(\ell) = 0$ if and only if $P(\ell^0) = P(\ell^1)$. By $\kappa \rightarrow_* (\kappa)_2^{\epsilon+\epsilon}$, let C_0 be a club homogeneous for Q_0 . Suppose C_0 is homogeneous for Q_0 taking value 1. Define $Q_1 : [\kappa]_*^{\epsilon+\epsilon} \rightarrow 2$ by $Q_1(\ell) = 0$ if and only if $P(\ell^0) < P(\ell^1)$. By $\kappa \rightarrow_* (\kappa)_2^{\epsilon+\epsilon}$, there is a club $C_1 \subseteq C_0$ which is homogeneous for Q_1 . First, suppose C_1 is homogeneous for Q_1 taking value 1. For each $n \in \omega$, let $\iota_n : \epsilon \rightarrow \kappa$ be defined by $\iota_n(\alpha) = \text{enum}_{C_1}((\omega \cdot \epsilon) \cdot n + \omega \cdot \alpha + \omega)$. Let $I_n : \epsilon \times \omega \rightarrow \kappa$ be defined by $I_n(\alpha, k) = \text{enum}_{C_1}((\omega \cdot \epsilon) \cdot n + \omega \cdot \alpha + k)$. For all $n \in \omega$, ι_n is discontinuous and I_n witnesses that ι_n has uniform cofinality ω . Thus $\iota_n \in [C_1]_*^\epsilon$. Note that for all $n < \omega$, $\sup(\iota_n) < \iota_{n+1}(0)$. For each $n \in \omega$, there is an $\ell_n \in [C_1]_*^{\epsilon+\epsilon}$ so that $\ell_n^0 = \iota_n$ and $\ell_n^1 = \iota_{n+1}$. For each $n \in \omega$, $Q_0(\ell_n) = 1$ and $Q_1(\ell_n) = 1$ imply that $P(\iota_n) = P(\ell_n^0) > P(\ell_n^1) = P(\iota_{n+1})$. Thus $\langle P(\iota_n) : n \in \omega \rangle$ is an infinite descending sequence of ordinals which is a contradiction. Now suppose C_1 is homogeneous for Q_1 taking value 0. For each $\xi < \gamma + 1$, let $\tau_\xi(\alpha) = \text{enum}_{C_1}((\omega \cdot \epsilon) \cdot \xi + \omega \cdot \alpha + \omega)$. Let $T_\xi : \epsilon \times \omega \rightarrow \kappa$ be defined by $T_\xi(\alpha, k) = \text{enum}_{C_1}((\omega \cdot \epsilon) \cdot \xi + \omega \cdot \alpha + k)$. For each $\xi < \gamma + 1$, τ_ξ is discontinuous and has uniform cofinality ω as witnessed by T_ξ . For each $\xi_0 < \xi_1 < \gamma + 1$, there is an $\ell_{\xi_0, \xi_1} \in [C_1]_*^{\epsilon+\epsilon}$ so that $\ell_{\xi_0, \xi_1}^0 = \tau_{\xi_0}$ and $\ell_{\xi_0, \xi_1}^1 = \tau_{\xi_1}$. For all $\xi_0 < \xi_1 < \gamma + 1$, $Q_0(\ell_{\xi_0, \xi_1}) = 1$ and $Q_1(\ell_{\xi_0, \xi_1}) = 0$ imply that $P(\tau_{\xi_0}) = P(\ell_{\xi_0, \xi_1}^0) < P(\ell_{\xi_0, \xi_1}^1) = P(\tau_{\xi_1})$. Thus $\langle \tau(\xi) : \xi < \gamma + 1 \rangle$ is an order embedding of $\gamma + 1$ into γ which is impossible. Thus C_0 must have been homogeneous for Q_0 taking value 0. Let $\iota_0, \iota_1 \in [C_0]_*^\epsilon$. Let $\bar{\iota} \in [C_0]_*^\epsilon$ be any element such that $\max\{\sup(\iota_0), \sup(\iota_1)\} < \bar{\iota}(0)$. Then there are $\ell_0, \ell_1 \in [C_0]_*^\epsilon$ so that $\ell_0^0 = \iota_0$, $\ell_1^0 = \iota_1$ and $\ell_0^1 = \bar{\iota}$. Then $Q_0(\ell_0) = 0 = Q_1(\ell_1)$ implies that $P(\iota_0) = P(\ell_0^0) = P(\ell_1^1) = P(\bar{\iota}) = P(\ell_1^0) = P(\iota_1)$. Since $\iota_0, \iota_1 \in [C_0]_*^\epsilon$ were arbitrary, one has that P is constant on $[C_0]_*^\epsilon$. $\square$

Fact 2.6. Let κ be an uncountable cardinal, $1 \leq \epsilon \leq \kappa$, $\delta < \epsilon$, $\kappa \rightarrow_* (\kappa)_2^{\delta+1+(\epsilon-\delta)}$, and $\kappa \rightarrow_* (\kappa)_{<\kappa}^{\epsilon-\delta}$. Let $\Phi : [\kappa]^\epsilon \rightarrow \kappa$ have the property that $\{\iota \in [\kappa]^\epsilon : \Phi(\iota) < \iota(\delta)\} \in \mu_\kappa^\epsilon$. Then there is a club $C \subseteq \kappa$ and a function $\Psi : [C]_*^\delta \rightarrow \kappa$ so that for all $\iota \in [C]_*^\epsilon$, $\Phi(\iota) = \Psi(\iota \restriction \delta)$.

Proof. If $\ell \in [\kappa]_2^{\delta+1+(\epsilon-\delta)}$, then let $\hat{\ell} \in [\omega_1]_*^\epsilon$ be defined by $\hat{\ell}(\alpha) = \ell(\alpha)$ if $\alpha < \delta$ and $\hat{\ell}(\alpha) = \ell(\delta + 1 + (\alpha - \delta))$ if $\delta \leq \alpha < \epsilon$. Let $C_0 \subseteq \kappa$ be a club consisting entirely of indecomposable ordinals so that for all $\iota \in [C_0]_*^\epsilon$, $\Phi(\iota) < \iota(\delta)$. Define $P : [C]_*^{\delta+1+(\epsilon-\delta)} \rightarrow 2$ by $P(\ell) = 0$ if and only if $\Phi(\hat{\ell}) < \ell(\delta)$. By $\kappa \rightarrow_* (\kappa)_2^{\delta+1+(\epsilon-\delta)}$, there is a club $C_1 \subseteq C_0$ which is homogeneous for P . Let $C_2 = \{\alpha \in C_1 : \text{enum}_{C_1}(\alpha) = \alpha\}$. Pick any $\iota \in [C_2]_*^\epsilon$. Since $\Phi(\iota) < \iota(\delta)$ because $\iota \in [C_2]_*^\epsilon \subseteq [C_0]_*^\epsilon$, $\text{next}_{C_1}^\omega(\Phi(\iota)) < \iota(\delta)$ by Fact 2.4. Let $\ell \in [C_1]_*^{\delta+1+(\epsilon-\delta)}$ be such that $\hat{\ell} = \iota$ and $\ell(\delta) = \text{next}_{C_1}^\omega(\Phi(\iota))$ (and note that ℓ has uniform cofinality ω since ι does and $\text{cof}(\text{next}_{C_1}^\omega(\Phi(\iota))) = \omega$). Since $\Phi(\hat{\ell}) = \Phi(\iota) < \text{next}_{C_1}^\omega(\Phi(\iota)) = \ell(\delta)$, one has $P(\ell) = 0$. Thus C_1 is homogeneous for P taking value 0. For any $\sigma \in [C_2]_*^\delta$, let $\Phi_\sigma : [C_2 \setminus (\sup(\sigma) + 1)]_*^{\epsilon-\delta} \rightarrow \kappa$ be defined by $\Phi_\sigma(\tau) = \Phi(\sigma \hat{\tau})$. For any $\tau \in [C_2 \setminus (\sup(\sigma) + 1)]_*^{\epsilon-\delta}$, let $\ell_{\sigma, \tau} = \sigma \hat{\langle \text{next}_{C_1}^\omega(\sup(\sigma)) \rangle \tau}$. Note that $\ell_{\sigma, \tau} \in [C_1]_*^{\delta+1+(\epsilon-\delta)}$, $\hat{\ell}_{\sigma, \tau} = \sigma \hat{\tau}$, and $\ell(\delta) = \text{next}_{C_1}^\omega(\sup(\sigma))$. $P(\ell_{\sigma, \tau}) = 0$ implies that $\Phi_\sigma(\tau) = \Phi(\sigma \hat{\tau}) = \Phi(\hat{\ell}_{\sigma, \tau}) < \ell(\delta) = \text{next}_{C_1}^\omega(\sup(\sigma))$. By $\kappa \rightarrow_* (\kappa)_{<\kappa}^{\epsilon-\delta}$, $\mu_\kappa^{\epsilon-\delta}$ is κ -complete. There is a $\gamma_\sigma < \kappa$ so that for $\mu_\kappa^{\epsilon-\delta}$ -almost all τ , $\Phi_\sigma(\tau) = \gamma_\sigma$. Define $Q : [C_2]_*^\epsilon \rightarrow 2$ by $Q(\iota) = 0$ if and only if $\Phi(\iota) = \gamma_{\iota \restriction \delta}$. By $\kappa \rightarrow_* (\kappa)_2^\epsilon$, there is a club $C_3 \subseteq C_2$ which is homogeneous for Q . Pick any $\sigma \in [C_3]_*^\delta$. There is a club $D \subseteq C_3 \setminus (\sup(\sigma) + 1)$ so that for all $\tau \in [D]_*^{\epsilon-\delta}$, $\Phi_\sigma(\tau) = \gamma_\sigma$. Fix $\tau \in [D]_*^{\epsilon-\delta}$. Let $\iota = \sigma \hat{\tau}$ and note that $\iota \in [C_3]_*^\epsilon$. $\Phi(\iota) = \Phi_{\iota \restriction \delta}(\tau) = \Phi_\sigma(\tau) = \gamma_\sigma = \gamma_{\iota \restriction \delta}$. Thus $Q(\iota) = 0$. This shows that C_3 is homogeneous for Q taking value 0. Define $\Psi : [C_3]_*^\delta \rightarrow \kappa$ by $\Psi(\sigma) = \gamma_\sigma$. For any $\iota \in [C_3]_*^\epsilon$, $Q(\iota) = 0$ implies that $\Phi(\iota) = \gamma_{\iota \restriction \delta} = \Psi(\iota \restriction \delta)$. $\square$

Fact 2.7. Let κ be an uncountable cardinal satisfying $\kappa \rightarrow_* (\kappa)_2^1$. Then μ_κ^1 is normal.

Proof. Note that $\kappa \rightarrow_* (\kappa)_2^1$ implies $\kappa \rightarrow_* (\kappa)_{<\kappa}^1$ by Fact 2.5. This result now follows from Fact 2.6 with $\delta = 0$ and $\epsilon = 1$. $\square$

Fact 2.8. Suppose $\epsilon < \kappa$ and $\kappa \rightarrow_* (\kappa)_2^{\epsilon+1}$. Let $\Phi : [\kappa]^\epsilon \rightarrow \kappa$. Then there is a club $C \subseteq \kappa$ so that for all $f \in [C]_*^\epsilon$, $\Phi(f) < \text{next}_C^\omega(\sup(f))$.

Proof. Define $P : [\kappa]_*^{\epsilon+1} \rightarrow 2$ by $P(g) = 0$ if and only if $\Phi(g \upharpoonright \epsilon) < g(\epsilon)$. By $\kappa \rightarrow_* (\kappa)_2^{\epsilon+1}$, there is a club $C \subseteq \kappa$ which is homogeneous for P . Pick any $f \in [C]_*^\epsilon$. Let $\gamma = \text{next}_C^\omega(\Phi(f))$. Let $g = f^\wedge \langle \gamma \rangle$ and note that $g \in [C]_*^{\epsilon+1}$. Since $\Phi(g \upharpoonright \epsilon) = \Phi(f) < \text{next}_C^\omega(\Phi(f)) = \gamma = g(\epsilon)$, one has that $P(g) = 0$. Since C is homogeneous for P and $g \in [C]_*^{\epsilon+1}$, one has that C is homogeneous for P taking value 0. For any $f \in [C]_*^\epsilon$, let $g_f = f^\wedge \langle \text{next}_C^\omega(\sup(f)) \rangle$. $P(g_f) = 0$ implies that $\Phi(f) = \Phi(g_f \upharpoonright \epsilon) < g_f(\epsilon) = \text{next}_C^\omega(\sup(f))$. $\square$

Note that the ordinary partition relation $\omega \rightarrow (\omega)_2^n$ for $n \in \omega$ is the finite Ramsey theorem. For uncountable cardinals κ , the ordinary partition relation $\kappa \rightarrow (\kappa)_2^2$ is equivalent to the weak compactness of κ which is compatible with the axiom of choice. However, the correct type partition relation $\kappa \rightarrow_* (\kappa)_2^2$ implies μ_κ^1 is normal which can be used to show ${}^\omega \kappa$ is not a wellorderable set. The finite exponent correct type partition relation already seems to imply many of the consequences of the infinite exponent ordinary partition relation. If $\text{AC}_\omega^\mathbb{R}$ holds and $\epsilon < \omega_1$, then a function $f : \epsilon \rightarrow \omega_1$ has uniform cofinality ω if and only if the range of f consists of limit ordinals. However if $\mu_{\omega_1}^1$ is a normal ultrafilter, then one can show that the identity function $\text{id} : \omega_1 \rightarrow \omega_1$ does not have uniform cofinality ω . The notion of a correct type function is a nontrivial concept when handling functions $f : \omega_1 \rightarrow \omega_1$ which will happen frequently in this paper.

Fact 2.9. (*Martin*; [16], [15], [5], [4]) Assume AD. $\omega_1 \rightarrow_* (\omega_1)_{<\omega_1}^1$.

Since AD implies $\omega_1 \rightarrow (\omega_1)_{<\omega_1}^1$, one has that for all $\epsilon \leq \omega_1$, $\mu_{\omega_1}^\epsilon$ are countably complete ultrafilters. Actually, AD implies there are no nonprincipal ultrafilters on ω which can be used to show any ultrafilter on any set is countably complete.

Definition 2.10. Let $\mathfrak{D}_{\omega_1} = \{(\alpha, \beta) : \beta < \alpha < \omega_1\}$. A Kunen function is a function $\mathcal{K} : \mathfrak{D}_{\omega_1} \rightarrow \omega_1$ such that for all $\alpha < \omega_1$, $\{\mathcal{K}(\alpha, \beta) : \beta < \alpha\}$ is an ordinal which will be denote $\chi_\alpha^\mathcal{K}$. Define $\Xi^\mathcal{K} : \omega_1 \rightarrow \omega_1$ by $\Xi^\mathcal{K}(\alpha) = \chi_\alpha^\mathcal{K}$. If $\beta < \omega_1$, then let $\mathcal{K}^\beta : (\omega_1 \setminus \beta + 1) \rightarrow \omega_1$ be defined by $\mathcal{K}^\beta(\alpha) = \mathcal{K}(\alpha, \beta)$.

Let $f : \omega_1 \rightarrow \omega_1$. The Kunen function $\mathcal{K}$ bounds f if and only if $\{\alpha < \omega_1 : f(\alpha) \leq \Xi^\mathcal{K}(\alpha)\} \in \mu_{\omega_1}^1$. The Kunen function $\mathcal{K}$ strictly bounds f if and only if $\{\alpha \in \omega_1 : f(\alpha) < \Xi^\mathcal{K}(\alpha)\} \in \mu_{\omega_1}^1$.

Fact 2.11. (*Kunen*; [15] Lemma 4.1)) AD. For every function $f : \omega_1 \rightarrow \omega_1$, there is a Kunen function $\mathcal{K} : \mathfrak{D}_{\omega_1} \rightarrow \omega_1$ which bounds f .

Definition 2.12. Let $\star$ be the following statement.

- For any function $f : \omega_1 \rightarrow \omega_1$, there is a Kunen function $\mathcal{K} : \mathfrak{D}_{\omega_1} \rightarrow \omega_1$ which bounds f .

Note that $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$ follow from AD by Fact 2.9 and Fact 2.11. The main result of the paper will be proved from the combinatorial principles $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$.

Definition 2.13. Let μ is a measure on a set X . If f and g are two functions on X , then let $f \sim_\mu g$ if and only if $\{x \in X : f(x) = g(x)\} \in \mu$. If $f : X \rightarrow \text{ON}$ and $g : X \rightarrow \text{ON}$, then write $f <_\mu g$ if and only if $\{x \in X : f(x) < g(x)\} \in \mu$. If $f : X \rightarrow \text{ON}$, then let $[f]_\mu$ be the class of all functions g with $g \sim_\mu f$. The ultrapower $\prod_X \text{ON}/\mu$ is the set of $\sim_\mu$ equivalence classes of functions $f : X \rightarrow \text{ON}$. The ultrapower ordering on $\prod_X \text{ON}/\mu$ is defined by $x \prec_\mu y$ if and only if there exists $f, g : X \rightarrow \text{ON}$ so that $x = [f]_\mu$ and $y = [g]_\mu$ and $f <_\mu g$. $j_\mu : \text{ON} \rightarrow \prod_X \text{ON}/\mu$ is defined by $j_\mu(\alpha) = [c_\alpha]_\mu$ where $c_\alpha : X \rightarrow \{\alpha\}$ is the constant function. If $x \in j_\mu(\omega_1)$, then let $\text{init}_\mu(x) = \{y \in j_\mu(\omega_1) : y \prec_\mu x\}$.

Fact 2.14. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$. $j_{\mu_{\omega_1}^1}(\omega_1)$ is wellfounded and $j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$ imply $\star$.

Proof. $\omega_1 \rightarrow_* (\omega_1)_2^2$ implies $\mu_{\omega_1}^1$, the club filter on ω_1 , is a normal ultrafilter on ω_1 by Fact 2.7. Thus $\omega_1 = [\text{id}]_{\mu_{\omega_1}^1}$ where $\text{id} : \omega_1 \rightarrow \omega_1$ is the identity function. Now suppose $j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$. Let $f : \omega_1 \rightarrow \omega_1$ be any function with $[\text{id}]_{\mu_{\omega_1}^1} \leq [f]_{\mu_{\omega_1}^1}$. Thus $\omega_1 = [\text{id}]_{\mu_{\omega_1}^1} \leq [f]_{\mu_{\omega_1}^1} < j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$. Let $\mathfrak{b} : \omega_1 \rightarrow \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1})$ be a bijection. Define a wellordering $\prec$ on ω_1 by $\alpha \prec \beta$ if and only if $\mathfrak{b}(\alpha) < \mathfrak{b}(\beta)$. Let $\mathcal{W} = (\omega_1, \prec)$ and note that $\text{ot}(\mathcal{W}) = [f]_{\mu_{\omega_1}^1}$. For each $\alpha < \omega_1$, let $\mathcal{W}_\alpha = (\alpha, \prec \upharpoonright \alpha)$. If $\beta < \alpha < \omega_1$, then let $\text{ot}(\mathcal{W}_\alpha, \beta)$ be the rank of β in $\mathcal{W}_\alpha$. Define $\mathcal{K} : \mathfrak{D}_{\omega_1} \rightarrow \omega_1$ by $\mathcal{K}(\alpha, \beta) = \text{ot}(\mathcal{W}_\alpha, \beta)$. One seeks to show that $\mathcal{K}$ is a Kunen function for f . It is clear that for all $\alpha \in \omega_1$, $\{\mathcal{K}(\alpha, \beta) : \beta < \alpha\} = \{\text{ot}(\mathcal{W}_\alpha, \beta) : \beta < \alpha\} = \text{ot}(\mathcal{W}_\alpha)$. Thus $\Xi^\mathcal{K}(\alpha) = \text{ot}(\mathcal{W}_\alpha)$. Suppose $\eta < [f]_{\mu_{\omega_1}^1}$. Let $\xi_\eta = \mathfrak{b}^{-1}(\eta)$. Define $g_\eta : \omega_1 \setminus (\xi_\eta + 1) \rightarrow \omega_1$ by $g_\eta(\alpha) = \text{ot}(\mathcal{W}_\alpha, \xi_\eta)$. Note that for all $\alpha \in \omega_1 \setminus (\xi_\eta + 1)$, $g_\eta(\alpha) < \text{ot}(\mathcal{W}_\alpha) = \Xi^\mathcal{K}(\alpha)$. Define $\Psi : \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1}) \rightarrow \text{init}_\mu([\Xi^\mathcal{K}]_{\mu_{\omega_1}^1})$ by $\Psi(\eta) = [g_\eta]_{\mu_{\omega_1}^1}$. Suppose $\eta_0 < \eta_1 < [f]_{\mu_{\omega_1}^1}$. Let $\zeta = \max\{\xi_{\eta_0}, \xi_{\eta_1}\}$. For all $\alpha \in \omega_1 \setminus (\zeta + 1)$, $g_{\eta_0}(\alpha) =$

$\text{ot}(\mathcal{W}_\alpha, \xi_{\eta_0}) < \text{ot}(\mathcal{W}_\alpha, \xi_{\eta_1}) = g_{\eta_1}(\alpha)$ since $\mathbf{b}(\xi_{\eta_0}) = \eta_0 < \eta_1 < \mathbf{b}(\xi_{\eta_1})$. Thus $\Psi(\eta_0) = [g_{\eta_0}]_{\mu_{\omega_1}^1} < [g_{\eta_1}]_{\mu_{\omega_1}^1} = \Psi(\eta_1)$. Ψ is an order embedding of $\text{init}_\mu([f]_{\mu_{\omega_1}^1})$ into $\text{init}_{\mu_{\omega_1}^1}([\Xi^\mathcal{K}])$. Thus $[f]_{\mu_{\omega_1}^1} \leq [\Xi^\mathcal{K}]_{\mu_{\omega_1}^1}$. This shows that $\{\alpha \in \omega_1 : f(\alpha) \leq \Xi^\mathcal{K}(\alpha)\} \in \mu_{\omega_1}^1$. $\mathcal{K}$ is a Kunen function bounding f . $\square$

Fact 2.15. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$. Let $f : \omega_1 \rightarrow \omega_1$ and $\mathcal{K}$ be a Kunen function strictly bounding f . Then there is a $\gamma < \omega_1$ so that $f \sim_{\mu_{\omega_1}^1} \mathcal{K}^\gamma$.

Proof. $\omega_1 \rightarrow_* (\omega_1)_2^2$ implies that $\mu_{\omega_1}^1$ is a normal ultrafilter by Fact 2.7. Let $A = \{\alpha \in \omega_1 : f(\alpha) < \Xi^\mathcal{K}(\alpha)\} \in \mu_{\omega_1}^1$. For each $\alpha \in A$, one has that $f(\alpha) < \Xi^\mathcal{K}(\alpha) = \chi_\alpha^\mathcal{K} = \{\mathcal{K}(\alpha, \beta) : \beta < \alpha\}$. Define $g : A \rightarrow \omega_1$ by $g(\alpha)$ is the least $\beta < \alpha$ so that $\mathcal{K}(\alpha, \beta) = f(\alpha)$. For all $\alpha \in A$, $g(\alpha) < \alpha$. Since $\mu_{\omega_1}^1$ is normal, there is a $\gamma < \omega_1$ and $B \subseteq A$ with $B \in \mu_{\omega_1}^1$ and $g(\alpha) = \gamma$ for all $\alpha \in B$. Thus $\mathcal{K}^\gamma(\alpha) = f(\alpha)$ for all $\alpha \in B$. $\square$

Fact 2.16. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$. Let $f : \omega_1 \rightarrow \omega_1$ and $\mathcal{K}$ be a Kunen function bounding f . Then there is an injection $\Gamma : \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1}) \rightarrow \omega_1$ so that for all $x \in \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1})$, $[\mathcal{K}^{\Gamma(x)}]_{\mu_{\omega_1}^1} = x$.

Proof. Suppose $x \prec_{\mu_{\omega_1}^1} [f]_{\mu_{\omega_1}^1}$. Let $g : \omega_1 \rightarrow \omega_1$ represent x . Then $g <_{\mu_{\omega_1}^1} f$ and hence $\mathcal{K}$ is also a Kunen function bounding g . By Fact 2.15, there is a $\gamma < \omega_1$ so that $\mathcal{K}^\gamma \sim_\mu g$. Let $\Gamma(x)$ be the least γ such that $[\mathcal{K}^\gamma]_{\mu_{\omega_1}^1} = x$. This defines the desired injection $\Gamma : \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1}) \rightarrow \omega_1$. $\square$

Dependent choice implies ultrapowers of ordinals are wellordering. However the existence of Kunen functions bounding functions from ω_1 to ω_1 is sufficient to show that the ultrapower of ω_1 by the finite exponent partition measures on ω_1 are wellorderings.

Fact 2.17. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$ and $\star$. The ultrapower $j_{\mu_{\omega_1}^1}(\omega_1) = \prod_{\omega_1} \omega_1 / \mu_{\omega_1}^1$ is a wellordering.

Proof. Suppose the ultrapower is not wellfounded. There is an $A \subseteq \prod_{\omega_1} \omega_1 / \mu_{\omega_1}^1$ which has no minimal element according to the ultrapower ordering $\prec_{\mu_{\omega_1}^1}$. Pick any element $x \in A$. Let $f : \omega_1 \rightarrow \omega_1$ be a representative for x . Let $\mathcal{K}$ be a Kunen function bounding f . By Fact 2.16, there is an injection $\Gamma : \text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1}) \rightarrow \omega_1$ so that for all $y \prec_{\mu_{\omega_1}^1} [f]_{\mu_{\omega_1}^1}$, $y = [\mathcal{K}^{\Gamma(y)}]_{\mu_{\omega_1}^1}$. Let $B = \Gamma[\text{init}_{\mu_{\omega_1}^1}([f]_{\mu_{\omega_1}^1})]$ be the range of Γ and note B is a set of ordinals. Let δ_0 be the least ordinal $\delta \in B$. Suppose δ_n has been defined. Since $[f]_{\mu_{\omega_1}^1}$ has no $\prec_{\mu_{\omega_1}^1}$ -least element, there is some $\delta \in B$ so that $\mathcal{K}^\delta <_{\mu_{\omega_1}^1} \mathcal{K}^{\delta_n}$. Let δ_{n+1} be the least ordinal $\delta \in B$ so that $\mathcal{K}^\delta <_{\mu_{\omega_1}^1} \mathcal{K}^{\delta_n}$. For each $n \in \omega$, $D_n = \{\alpha \in \omega_1 : \mathcal{K}^{\delta_{n+1}}(\alpha) < \mathcal{K}^{\delta_n}(\alpha)\} \in \mu_{\omega_1}^1$. Since $\mu_{\omega_1}^1$ is countably complete, $D = \bigcap_{n \in \omega} D_n \in \mu_{\omega_1}^1$ and hence nonempty. Let $\bar{\alpha} \in D$. Then $\langle \mathcal{K}^{\delta_n}(\bar{\alpha}) : n \in \omega \rangle$ is an infinite descending sequence of ordinals. Contradiction. $\square$

Fact 2.18. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$ and $\star$. $j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$.

Proof. By Fact 2.17, $j_{\mu_{\omega_1}^1}(\omega_1)$ is a wellordering. By Fact 2.16, each initial segment of $j_{\mu_{\omega_1}^1}(\omega_1)$ injects into ω_1 . Thus $j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$. $\square$

Fact 2.19. Assume $\omega_1 \rightarrow_* (\omega_1)_2^2$. $\star$ and $j_{\mu_{\omega_1}^1}(\omega_1) \leq \omega_2$ are equivalent.

Proof. This follows from Fact 2.14 and Fact 2.18. $\square$

Fact 2.20. Let κ be an uncountable cardinal and μ be a normal ultrafilter on κ containing no bounded subsets of κ . Let $q : \kappa \rightarrow \kappa$ be a function and $A \in \mu$. Then the set $B = \{\alpha \in A : (\forall \alpha' < \alpha)(q(\alpha') < \alpha)\} \in \mu$.

Proof. Suppose not. Then $C = \kappa \setminus B \in \mu$. Let $f : C \rightarrow \kappa$ be defined by $f(\alpha)$ is the least $\alpha' < \alpha$ so that $\alpha \leq q(\alpha')$. Since μ is normal, there is a $D \subseteq C$ with $D \in \mu$ and a $\beta < \kappa$ so that for all $\alpha \in D$, $f(\alpha) = \beta$. Thus for all $\alpha \in D$, $\alpha \leq q(f(\alpha)) = q(\beta)$. This is impossible since D is an unbounded set. $\square$

Fact 2.21. Let κ be an uncountable cardinal and μ be a normal ultrafilter on κ containing no bounded subsets of κ . If $A \in \mu$, then $\{\alpha \in A : \text{enum}_A(\alpha) = \alpha\} \in \mu$.

Proof. By applying Fact 2.20 to enum_A , the set $\bar{A} = \{\alpha \in \kappa : (\forall \alpha' < \alpha)(\text{enum}_A(\alpha') < \alpha)\} \in \mu$. Let $B = A \cap \bar{A} \in \mu$. If $\alpha \in B$, then $\sup(\text{enum}_A \upharpoonright \alpha) = \alpha$ and thus $\text{enum}_A(\alpha) = \alpha$ since $\alpha \in A$. So $B \subseteq \{\alpha \in A : \text{enum}_A(\alpha) = \alpha\}$. $\square$

Fact 2.22. (Martin) Assume κ is an uncountable cardinal satisfying $\kappa \rightarrow_* (\kappa)_2^\kappa$.

- (1) Let μ be an ultrafilter on κ such that $j_\mu(\kappa)$ is a wellordering. Then $j_\mu(\kappa)$ is a cardinal.
- (2) Let μ be a normal ultrafilter on κ which contains no bounded subsets of κ such that $j_\mu(\kappa)$ is a wellordering. Then $j_\mu(\kappa)$ is a regular cardinal.

Proof. (1) First assume μ is an ultrafilter on an uncountable cardinal satisfying $\kappa \rightarrow_* (\kappa)_2^\kappa$ and $j_\mu(\kappa)$ is a wellordering (and thus one may assume $j_\mu(\kappa)$ is an ordinal). For the sake of contradiction, suppose $j_\mu(\kappa)$ is not a cardinal. Then there is a $\lambda < j_\mu(\kappa)$ and an injection $\Phi : j_\mu(\kappa) \rightarrow \lambda$. If $f : \kappa \rightarrow \kappa$ is a function, then let $f^0(\alpha) = f(2 \cdot \alpha)$ and $f^1(\alpha) = f(2 \cdot \alpha + 1)$. Define $P : [\kappa]_*^\kappa \rightarrow 2$ by $P(f) = 0$ if and only if $\Phi([f^0]_\mu) = \Phi([f^1]_\mu)$. By $\kappa \rightarrow_* (\kappa)_2^\kappa$, there is a club $C_0 \subseteq \kappa$ which is homogeneous for P . Take any $f \in [C_0]_*^\kappa$. Note that for all $\alpha < \kappa$, $f^0(\alpha) < f^1(\alpha)$ and hence $[f^0]_\mu < [f^1]_\mu$. Since Φ is injective, $\Phi([f^0]_\mu) \neq \Phi([f^1]_\mu)$ and thus $P(f) = 1$. So C_0 is homogeneous for P taking value 1. Define $Q : [C_0]_*^\kappa \rightarrow 2$ by $Q(f) = 0$ if and only if $\Phi([f^0]_\mu) < \Phi([f^1]_\mu)$. By $\kappa \rightarrow_* (\kappa)_2^\kappa$, there is a club $C_1 \subseteq C_0$ which is homogeneous for Q . First suppose C_1 is homogeneous for Q taking value 1. For each $k \in \omega$ and $\alpha < \kappa$, let $g_k(\alpha) = \text{enum}_{C_1}((\omega \cdot \omega) \cdot \alpha + \omega \cdot k + \omega)$. Note that $g_k \in [C_1]_*^\kappa$. For each $k \in \omega$, there is an $f_k \in [C_1]_*^\kappa$ so that $f_k^0 = g_k$ and $f_k^1 = g_{k+1}$. Then $P(f_k) = 1$ and $Q(f_k) = 1$ imply that $\Phi([g_{k+1}]_\mu) = \Phi([f_k^1]_\mu) < \Phi([f_k^0]_\mu) = \Phi([g_k]_\mu)$. Thus $\langle \Phi([g_k]_\mu) : k \in \omega \rangle$ is an infinite descending sequence in the ordinal λ which is a contradiction. Suppose C_1 is homogeneous for Q taking value 0. Since $\lambda < j_\mu(\kappa)$, let $h : \kappa \rightarrow \kappa$ be such that $[h]_\mu = \lambda$. Define $T = \{(\alpha, \beta) : \alpha < \kappa \wedge \beta < h(\alpha) + 2\}$.¹ Let $<_{\text{lex}}$ be the lexicographic ordering on T . Note that the ordertype of $(T, <_{\text{lex}})$ is κ . Let $\psi : T \rightarrow C_1$ be an order preserving function from $(T, <_{\text{lex}})$ into $(C_1, <)$ of the correct type which means the following two conditions hold:

- For all $x \in T$, $\sup\{\psi(y) : y <_{\text{lex}} x\} < \psi(x)$.
- There is a function $\Psi : T \times \omega \rightarrow \kappa$ so that for all $x \in T$ and $k \in \omega$, $\Psi(x, k) < \Psi(x, k+1)$ and $\psi(x) = \sup\{\Psi(x, k) : k \in \omega\}$.

For any function $g : \kappa \rightarrow \kappa$, define $\hat{g} : \kappa \rightarrow C_1$ by

$$\hat{g}(\alpha) = \begin{cases} \psi(\alpha, g(\alpha)) & g(\alpha) < h(\alpha) + 1 \\ \psi(\alpha, 0) & \text{otherwise} \end{cases}.$$

Define $\hat{G} : \kappa \times \omega \rightarrow \kappa$ by

$$\hat{G}(\alpha, k) = \begin{cases} \Psi((\alpha, g(\alpha)), k) & g(\alpha) < h(\alpha) + 1 \\ \Psi((\alpha, 0), k) & \text{otherwise} \end{cases}.$$

Note that $\hat{G}$ witnesses that $\hat{g}$ has uniform cofinality ω . Since ψ is discontinuous everywhere, $\hat{g}$ is discontinuous everywhere. Thus $\hat{g} : \kappa \rightarrow C_1$ is an increasing function of the correct type and hence $\hat{g} \in [C_1]_*^\kappa$. For any $\eta < \lambda + 1$, let $\delta_\eta = [\hat{g}]_\mu$ for any $g : \kappa \rightarrow \kappa$ such that $[g]_\mu = \eta$. Note that δ_η is independent of the choice of g representing η . Let $\eta_0 < \eta_1 < \lambda + 1$. Let $g_0, g_1 : \kappa \rightarrow \kappa$ be such that $\eta_0 = [g_0]_\mu$ and $\eta_1 = [g_1]_\mu$. For $i \in 2$, let $\tilde{g}_0, \tilde{g}_1 \in [C_1]_*^\kappa$ be defined by

$$\begin{aligned} \tilde{g}_0(\alpha) &= \begin{cases} \psi(\alpha, g_0(\alpha)) & g_0(\alpha) < g_1(\alpha) < h(\alpha) + 1 \\ \psi(\alpha, 0) & \text{otherwise} \end{cases} \\ \tilde{g}_1(\alpha) &= \begin{cases} \psi(\alpha, g_1(\alpha)) & g_0(\alpha) < g_1(\alpha) < h(\alpha) + 1 \\ \psi(\alpha, 1) & \text{otherwise} \end{cases} \end{aligned}$$

Note that for all $\alpha < \kappa$, $\tilde{g}_0(\alpha) < \tilde{g}_1(\alpha)$ by the definitions above and the fact that ψ is order preserving on $(T, <_{\text{lex}})$. For all $\alpha < \kappa$, $\tilde{g}_1(\alpha) < \tilde{g}_0(\alpha + 1)$ since $\tilde{g}_1(\alpha) = \psi(\alpha, \xi)$ for some $\xi < h(\alpha) + 2$, $\tilde{g}_0(\alpha + 1) = \psi(\alpha + 1, \zeta)$ for some $\zeta < h(\alpha + 1) + 2$, and by comparing the first coordinates since ψ is order preserving on $(T, <_{\text{lex}})$. Thus there is an $f \in [C_1]_*^\kappa$ so that $f^0 = \tilde{g}_0$ and $f^1 = \tilde{g}_1$. Since $[g_0]_\mu < [g_1]_\mu < \lambda + 1$, one has that $A = \{\alpha \in \omega_1 : g_0(\alpha) < g_1(\alpha) < h(\alpha) + 1\} \in \mu$. Hence for all $\alpha \in A$, $\hat{g}_0(\alpha) = \tilde{g}_0(\alpha)$ and $\hat{g}_1(\alpha) = \tilde{g}_1(\alpha)$. Thus $\delta_{\eta_0} = [\hat{g}_0]_\mu = [\tilde{g}_0]_\mu = [f^0]_\mu$ and $\delta_{\eta_1} = [\hat{g}_1]_\mu = [\tilde{g}_1]_\mu = [f^1]_\mu$. $P(f) = 1$ and $Q(f) = 0$ imply that $\delta_{\eta_0} = [f^0]_\mu < [f^1]_\mu = \delta_{\eta_1}$. Thus $\langle \delta_\eta : \eta < \lambda + 1 \rangle$ is an order preserving injection of $\lambda + 1$ into λ which is

¹The purpose for adding 2 rather than 1 is to ensure that $(\alpha, 0), (\alpha, 1) \in T$ for all $\alpha < \kappa$.

impossible. (Note that since $j_\mu(\kappa)$ is an ordinal, $\lambda < j_\mu(\kappa)$ is also an ordinal. For ordinals λ , $\lambda + 1$ cannot inject into λ .)

(2) Now suppose μ is a normal ultrafilter on κ which does not contain any bounded subsets of κ and $j_\mu(\kappa)$ is an ordinal. For the sake of contradiction, suppose $j_\mu(\kappa)$ is not regular. Then there is an infinite cardinal $\lambda < j_\mu(\kappa)$ and a cofinal increasing map $\rho : \lambda \rightarrow j_\mu(\kappa)$. Define $V : [\kappa]_\mu^\kappa \rightarrow 2$ by $V(f) = 0$ if and only if there exists a $\xi < \lambda$ so that $[f^0]_\mu < \rho(\xi) < [f^1]_\mu$ (where $f^0, f^1 \in [\kappa]^\kappa$ is obtained from $f \in [\kappa]^\kappa$ as before). By $\kappa \rightarrow_* (\kappa)_2^\kappa$, there is a club C_0 homogeneous for V . First suppose C_0 is homogeneous for V taking value 0. Let $h : \kappa \rightarrow \kappa$ be such that $[h]_\mu = \lambda$. Define $W = \{(\alpha, \beta) : \beta < h(\alpha) + 2\}$. As before, $(W, <_{\text{lex}})$ has ordertype κ . Let $\psi : W \rightarrow C_0$ be an order preserving function from $(W, <_{\text{lex}}) \rightarrow C_0$ of the correct type. For any $g : \kappa \rightarrow \kappa$, define $\check{g}_0, \check{g}_1 : \kappa \rightarrow \kappa$ by

$$\check{g}_0(\alpha) = \begin{cases} \psi(\alpha, g(\alpha)) & g(\alpha + 1) < h(\alpha) + 2 \\ \psi(\alpha, 0) & \text{otherwise} \end{cases}$$

$$\check{g}_1(\alpha) = \begin{cases} \psi(\alpha, g(\alpha) + 1) & g(\alpha + 1) < h(\alpha) + 2 \\ \psi(\alpha, 1) & \text{otherwise} \end{cases}.$$

Note that for all $g : \kappa \rightarrow \kappa$, $\check{g}_0, \check{g}_1 \in [C_0]_\mu^\kappa$ and for all $\alpha < \kappa$, $\check{g}_0(\alpha) < \check{g}_1(\alpha) < \check{g}_0(\alpha + 1)$ by arguments similar to the above. Thus there is some $f \in [C_0]_\mu^\kappa$ so that $f^0 = \check{g}_0$ and $f^1 = \check{g}_1$. Now suppose $\eta < \lambda + 1$. Let $g : \kappa \rightarrow \kappa$ be such that $\eta = [g]_\mu$. Let $f \in [C_0]_\mu^\kappa$ be such that $f^0 = \check{g}_0$ and $f^1 = \check{g}_1$. $V(f) = 0$ implies that there is a $\xi < \lambda$ so that $[\check{g}_0]_\mu = [f^0]_\mu < \rho(\xi) < [f^1]_\mu = [\check{g}_1]_\mu$. Let ξ_η be the least such ξ and note that ξ_η is independent of the choice of g representing η . Now suppose $\eta_0 < \eta_1 < \lambda + 1$. Let $g, p : \kappa \rightarrow \kappa$ be such that $\eta_0 = [g]_\mu$ and $\eta_1 = [p]_\mu$. Note that $B = \{\alpha \in \kappa : g(\alpha) < p(\alpha) < h(\alpha) + 2\} \in \mu$. For all $\alpha \in B$, $\check{g}_1(\alpha) \leq \check{p}_0(\alpha)$. Thus $\rho(\xi_{\eta_0}) < [\check{g}_1]_\mu \leq [\check{p}_0]_\mu < \rho(\xi_{\eta_1})$. Since ρ is an increasing function, one must have that $\xi_{\eta_0} < \xi_{\eta_1}$. Thus $\langle \xi_\eta : \eta < \lambda + 1 \rangle$ is an order preserving injection of $\lambda + 1$ into λ . This is impossible. Now suppose C_0 is homogeneous for V taking value 1. Let $g_0 \in [C_0]_\mu^\kappa$. Since $\rho : \lambda \rightarrow j_\mu(\kappa)$ is cofinal, there is some ξ so that $\rho(\xi) > [g_0]_\mu$. Since $j_\mu(C_0)$ is order isomorphic to $j_\mu(\kappa)$, let $g_1 \in [C_0]_\mu^\kappa$ be so that $[g_1]_\mu > \rho(\xi)$. Since g_0 is an increasing function, for all $\alpha < \kappa$, there is an α' so that $g_1(\alpha) < g_0(\alpha')$. Let $q(\alpha)$ be the least α' so that $g_1(\alpha) < g_0(\alpha')$. Since $[g_0]_\mu < \rho(\xi) < [g_1]_\mu$, let $A_0 \in \mu$ be such that for all $\alpha \in A_0$, $g_0(\alpha) < g_1(\alpha)$. Let $A_1 = \{\alpha \in A_0 : (\forall \alpha' < \alpha)(q(\alpha') < \alpha)\}$ and note that $A_1 \in \mu$ by Fact 2.20. Define $f : \kappa \rightarrow \kappa$ by $f(2 \cdot \alpha) = g_0(\text{enum}_{A_1}(\alpha))$ and $f(2 \cdot \alpha + 1) = g_1(\text{enum}_{A_1}(\alpha))$ for all $\alpha < \kappa$. Note that for all $\alpha < \kappa$, $f(2 \cdot \alpha) = g_0(\text{enum}_{A_1}(\alpha)) < g_1(\text{enum}_{A_1}(\alpha)) = f(2 \cdot \alpha + 1)$ since $g_0(\gamma) < g_1(\gamma)$ for all $\gamma \in A_1$. Note also that for all $\alpha < \kappa$, $f(2 \cdot \alpha + 1) = g_1(\text{enum}_{A_1}(\alpha)) < g_0(q(\text{enum}_{A_1}(\alpha))) < g_0(\text{enum}_{A_1}(\alpha + 1)) = f(2 \cdot (\alpha + 1))$ by the definition of q and A_1 . This shows that $f : \kappa \rightarrow \kappa$ is an increasing function. It is clear that $f \in [C_0]_\mu^\kappa$ since $g_0, g_1 \in [C_0]_\mu^\kappa$. Let $A_2 = \{\alpha \in A_1 : \text{enum}_{A_1}(\alpha) = \alpha\}$ which belongs to μ by Fact 2.21. For all $i \in 2$ and $\alpha \in A_2$, $f^i(\alpha) = g_i(\text{enum}_{A_1}(\alpha)) = g_i(\alpha)$. Thus $[f^i]_\mu = [g_i]_\mu$ for both $i \in 2$. $V(f) = 1$ implies that there is no $\xi < \lambda$ with $[g_0]_\mu = [f^0]_\mu < \rho(\xi) < [f^1]_\mu = [g_1]_\mu$. This is contradiction since $[g_0]_\mu < \rho(\xi) < [g_1]_\mu$. It has been shown that V has no homogeneous club which violates $\kappa \rightarrow_* (\kappa)_2^\kappa$. $\square$

Fact 2.23. (Martin) Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. Then $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$ and ω_2 is a regular cardinal.

Proof. By Fact 2.17, $j_{\mu_{\omega_1}^1}(\omega_1)$ is a wellordering. By Fact 2.16, each initial segment of $j_{\mu_{\omega_1}^1}(\omega_1)$ injects into ω_1 . Thus $\omega_1 = [\text{id}]_{\mu_{\omega_1}^1} < j_{\mu_{\omega_1}^1}(\omega_1) \leq (\omega_1)^+ = \omega_2$. Since Fact 2.22 implies $j_{\mu_{\omega_1}^1}(\omega_1)$ must be a cardinal, one has that $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$. Since $\mu_{\omega_1}^1$ is a normal ultrafilter by Fact 2.7, Fact 2.22 also implies that $\omega_2 = j_{\mu_{\omega_1}^1}(\omega_1)$ is regular. $\square$

Fact 2.24. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$. $\star$ and $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$ are equivalent.

Proof. This follow from Fact 2.14 and Fact 2.23 $\square$

Next, one will show that for all $1 \leq n < \omega$, $j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$ and $\text{cof}(\omega_{n+1}) = \omega_2$ (without assuming any form of dependent choice or even countable choice).

Fact 2.25. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{n+1}$. For each $1 \leq n < \omega$ and $g : \omega_1 \rightarrow \omega_1$, let $\hat{\Sigma}_n(g) : [\omega_1]^n \rightarrow \omega_1$ be defined by $\hat{\Sigma}_n(g)(\iota) = g(\iota(n-1))$. Let $\Sigma_n : j_{\mu_{\omega_1}^1}(\omega_1) \rightarrow j_{\mu_{\omega_1}^n}(\omega_1)$ be defined by $\Sigma_n([g]_{\mu_{\omega_1}^1}) = [\hat{\Sigma}_n(g)]_{\mu_{\omega_1}^n}$. Then $\Sigma_n : j_{\mu_{\omega_1}^1}(\omega_1) \rightarrow j_{\mu_{\omega_1}^n}(\omega_1)$ is cofinal increasing.

Proof. Suppose $x \in j_{\mu_{\omega_1}^n}(\omega_1)$. Let $f : [\omega_1]^n \rightarrow \omega_1$ represent x . Define $P_f : [\omega_1]^{n+1} \rightarrow 2$ by $P_f(\ell) = 0$ if and only if $f(\ell \restriction n) < \ell(n)$. By $\omega_1 \rightarrow_* (\omega_1)_2^{n+1}$, there is a club $C \subseteq \omega_1$ which is homogeneous for P_f . Pick any $\iota \in [C]_*^n$. Let $\gamma \in [C]_*^1$ be such that $f(\iota) < \gamma$. Let $\ell = \iota \hat{\wedge} \langle \gamma \rangle$ and note that $\ell \in [C]_*^{n+1}$. Then $P_f(\ell) = 0$ since $f(\ell \restriction n) = f(\iota) < \gamma = \ell(n)$. This shows that C is homogeneous for P_f taking value 0. Let $g : \omega_1 \rightarrow \omega_1$ be defined by $g(\alpha) = \text{next}_C^\omega(\alpha)$. Let $\iota \in [C]_*^n$ and let $\ell_\iota = \iota \hat{\wedge} \langle g(\iota(n-1)) \rangle$. Note that $\ell_\iota \in [C]_*^{n+1}$ and thus $P_f(\ell_\iota) = 0$. This implies that $f(\iota) = f(\ell_\iota \restriction n) < \ell_\iota(n) = g(\iota(n-1)) = \hat{\Sigma}_n(g)(\iota)$. Since $\iota \in [C]_*^n$ was arbitrary, it has been shown that $x = [f]_{\mu_{\omega_1}^n} \prec_{\mu_{\omega_1}^{n-1}} [\hat{\Sigma}_n(g)]_{\mu_{\omega_1}^n} = \Sigma_n([g]_{\mu_{\omega_1}^1})$. $\square$

Fact 2.26. Assume $1 \leq m < n < \omega$ and $\omega_1 \rightarrow_* (\omega_1)_2^n$. There is an order embedding of $j_{\mu_{\omega_1}^m}(\omega_1)$ into a proper initial segment of $j_{\mu_{\omega_1}^n}(\omega_1)$.

Proof. If $f : [\omega_1]^m \rightarrow \omega_1$, then let $\hat{f} : [\omega_1]^n \rightarrow \omega_1$ be defined by $\hat{f}(\ell) = f(\ell \restriction m)$. Define $\Psi : j_{\mu_{\omega_1}^m}(\omega_1) \rightarrow j_{\mu_{\omega_1}^n}(\omega_1)$ by $\Psi(x) = [\hat{f}]_{\mu_{\omega_1}^n}$ where $f : [\omega_1]^m \rightarrow \omega_1$ represents x . One can check that Ψ is well defined independent of the choice of representative f for x and is an order embedding.

Let $g : [\omega_1]^n \rightarrow \omega_1$ be defined by $g(\ell) = \ell(m)$. The claim is that the range of ψ is below $[g]_{\mu_{\omega_1}^n}$. Let $f : [\omega_1]^m \rightarrow \omega_1$. Let $P_f : [\omega_1]^{m+1} \rightarrow \omega_1$ be defined by $P_f(\sigma) = 0$ if and only if $f(\sigma \restriction m) < \sigma(m)$. By $\omega_1 \rightarrow (\omega_1)_2^{m+1}$, let $C_0 \subseteq \omega_1$ be a club homogeneous for P_f . Pick $\iota \in [C_0]_*^m$. Let $\sigma = \iota \hat{\wedge} \langle \text{next}_{C_0}^\omega(f(\iota)) \rangle$ and note that $\sigma \in [C_0]_*^{m+1}$. Since $f(\sigma \restriction m) = f(\iota) < \text{next}_{C_0}^\omega(f(\iota)) = \sigma(m)$, one has that $P_f(\sigma) = 0$. Thus C_0 is a homogeneous for P_f taking value 0. For any $\iota \in [C_0]_*^m$, let $\sigma_\iota = \iota \hat{\wedge} \langle \text{next}_{C_0}^\omega(\iota(m-1)) \rangle$. $P_f(\sigma_\iota) = 0$ implies that $f(\iota) < \text{next}_{C_0}^\omega(\iota(m-1))$. Let $C_1 = \{\alpha \in C_0 : \text{enum}_{C_0}(\alpha) = \alpha\}$. Let $\ell \in [C_1]^n$. One has $\hat{f}(\ell) = f(\ell \restriction m) < \text{next}_{C_0}^\omega(\ell(m-1)) < \ell(m) = g(\ell)$ since $\ell(m) \in C_1$ and using Fact 2.4. Thus $\Psi([f]_{\mu_{\omega_1}^m}) < [g]_{\mu_{\omega_1}^n}$. This shows that Ψ maps $j_{\mu_{\omega_1}^m}(\omega_1)$ into an initial segment of $j_{\mu_{\omega_1}^n}(\omega_1)$. $\square$

Definition 2.27. Suppose $f : [\omega_1]^n \rightarrow \omega_1$. For each $1 \leq k \leq n$, define $I_f^k : [\omega_1]^k \rightarrow \omega_1$ by $I_f^k(\sigma) = \sup\{f(\tau \hat{\wedge} \sigma) : \tau \in [\omega_1]^{n-k} \wedge \sup(\tau) < \sigma(0)\}$. (Note that $I_f^n = f$.)

Note that if $f, g : [\omega_1]^n \rightarrow \omega_1$ with $[f]_{\mu_{\omega_1}^n} \preceq_{\mu_{\omega_1}^n} [g]_{\mu_{\omega_1}^n}$, then it is not necessarily true that $[I_f^1]_{\mu_{\omega_1}^1} \leq [I_g^1]_{\mu_{\omega_1}^1}$. However, one has the following.

Fact 2.28. Suppose $1 \leq n < \omega_1$, $f, g : [\omega_1]^n \rightarrow \omega_1$ with $[f]_{\mu_{\omega_1}^n} \preceq_{\mu_{\omega_1}^n} [g]_{\mu_{\omega_1}^n}$. Then there is a $\bar{f} : [\omega_1]^n \rightarrow \omega_1$ so that $[\bar{f}]_{\mu_{\omega_1}^n} = [f]_{\mu_{\omega_1}^n}$ and for all $\alpha < \omega_1$, $I_{\bar{f}}^1(\alpha) \leq I_g^1(\alpha)$.

Proof. Since $[f]_{\mu_{\omega_1}^n} \preceq_{\mu_{\omega_1}^n} [g]_{\mu_{\omega_1}^n}$, $A = \{\ell \in [\omega_1]^n : f(\ell) \leq g(\ell)\} \in \mu_{\omega_1}^n$. Define $\bar{f} : [\omega_1]^n \rightarrow \omega_1$ by

$$\bar{f}(\ell) = \begin{cases} f(\ell) & \ell \in A \\ 0 & \ell \notin A \end{cases}.$$

Note that $[f]_{\mu_{\omega_1}^n} = [\bar{f}]_{\mu_{\omega_1}^n}$. For all $\alpha \in \omega_1$, $I_{\bar{f}}^1(\alpha) = \sup\{\bar{f}(\ell) : \ell \in [\omega_1]^n \wedge \ell(n-1) = \alpha\} = \sup\{f(\ell) : \ell \in A \wedge \ell(n-1) = \alpha\} \leq \sup\{g(\ell) : \ell \in A \wedge \ell(n-1) = \alpha\} \leq \sup\{g(\ell) : \ell \in [\omega_1]^n \wedge \ell(n-1) = \alpha\} = I_g^1(\alpha)$ where the first inequality uses the fact that $f(\ell) \leq g(\ell)$ for all $\ell \in A$. $\square$

Definition 2.29. Suppose $1 \leq n < \omega$, $\mathcal{K}$ is a Kunen function, and $h : [\omega_1]^n \rightarrow \omega_1$. Define $\mathcal{K}^{n,h} : [\omega_1]^{n+1} \rightarrow \omega_1$ by $\mathcal{K}^{n,h}(\ell) = \mathcal{K}(\ell(n), h(\ell \restriction n))$ when $h(\ell \restriction n) < \ell(n)$ and $\mathcal{K}^{n,h}(\ell) = 0$ otherwise.

Fact 2.30. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. For all $1 \leq n < \omega$, $j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$ and $\text{cof}(\omega_{n+1}) = \omega_2$.

Proof. Suppose $1 \leq n < \omega$ and the following has been shown:

- (1) $j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$ and $\text{cof}(\omega_{n+1}) = \omega_2$.
- (2) If $A \subseteq \omega_{n+1}$ with $|A| \leq \omega_1$, there is a function Σ so that for all $\alpha \in A$, $\Sigma(\alpha) : [\omega_1]_*^n \rightarrow \omega_1$ and $[\Sigma(\alpha)]_{\mu_{\omega_1}^n} = \alpha$.²

²For this proof, one only needs the result for $|A| \leq \omega$. However, the proof is no different for A with $|A| \leq \omega_1$. There are other applications which require the stronger form.

For $n = 1$, both properties have been shown. (1) is Fact 2.23. To see (2), suppose $A \subseteq \omega_2$ with $|A| \leq \omega_1$. Since ω_2 is regular, $\sup(A) < \omega_2$. Let $f : \omega_1 \rightarrow \omega_1$ represent $\sup(A)$ and $\mathcal{K}$ be a Kunen function bounding f . By Fact 2.16, there is a function Γ so that for all $\alpha < \sup(A)$, $\alpha = [\mathcal{K}^{\Gamma(\alpha)}]_{\mu_{\omega_1}^1}$. For $\alpha \in A$, let $\Sigma(\alpha) = \mathcal{K}^{\Gamma(\alpha)}$.

Now suppose the two properties have been established at n . One seeks to establish the two properties at $n + 1$.

First, one will show that $j_{\mu_{\omega_1}^{n+1}}(\omega_1)$ is wellfounded. Suppose $X \subseteq j_{\mu_{\omega_1}^{n+1}}(\omega_1)$ has no $\prec_{\mu_{\omega_1}^{n+1}}$ -minimal elements. Pick $x \in X$ and let $f : [\omega_1]^{n+1} \rightarrow \omega_1$ represent x . Let $\mathcal{K}$ be a Kunen function bounding $I_f^1 : \omega_1 \rightarrow \omega_1$. For any $y \prec_{\mu_{\omega_1}^{n+1}} x$, use Fact 2.28 to pick a $g : [\omega_1]^{n+1} \rightarrow \omega_1$ which represents y so that for that for all $\alpha < \omega_1$, $I_g^1(\alpha) \leq_{\mu_{\omega_1}^1} I_f^1(\alpha)$. Thus $\mathcal{K}$ is also a Kunen function which bounds I_g^1 . Let $A_g = \{\alpha \in \omega_1 : I_g^1(\alpha) < \Xi^{\mathcal{K}}(\alpha)\}$ and note that $A_g \in \mu_{\omega_1}^1$. For any $\ell \in [A_g]_*^{n+1}$, $g(\ell) \leq I_g^1(\ell(n)) < \Xi^{\mathcal{K}}(\ell(n))$. Let $\hat{h}(\ell)$ be the least ordinal $\gamma < \ell(n)$ so that $g(\ell) = \mathcal{K}(\ell(n), \gamma)$. Since $\hat{h}(\ell) < \ell(n)$ for $\mu_{\omega_1}^{n+1}$ -almost all ℓ , Fact 2.6 implies there is an $h : [\omega_1]^n \rightarrow \omega_1$ so that for $\mu_{\omega_1}^{n+1}$ -almost all ℓ , $\hat{h}(\ell) = h(\ell \upharpoonright n)$. By (1) at n , one has that $[h]_{\mu_{\omega_1}^n} \in j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$. It has been shown that for all $y \prec_{\mu_{\omega_1}^{n+1}} x$, there is an ordinal $\delta < \omega_{n+1}$ so that for any $h : [\omega_1]^n \rightarrow \omega_1$ representing δ , the function $\mathcal{K}^{n,h} : [\omega_1]^{n+1} \rightarrow \omega_1$ defined (in Definition 2.29) by $\mathcal{K}^{n,h}(\ell) = \mathcal{K}(\ell(n), h(\ell \upharpoonright n))$ represents y . Let δ_y be the least such δ for y . Let $B = \{\delta_y : y \in X \wedge y \prec_{\mu_{\omega_1}^{n+1}} x\}$ and note that $B \subseteq \omega_{n+1}$. Let δ_0 be the least member of B according to the usual ordering on ω_{n+1} . Suppose δ_k has been found and let y_k be the element of X represented by $\mathcal{K}^{n,h}$ for any $h : [\omega_1]^n \rightarrow \omega_1$ representing δ_k . Since X has no minimal element, there is some δ so that for any $h : [\omega_1]^n \rightarrow \omega_1$ representing δ , the function $\mathcal{K}^{n,h}$ represents an element of X which is $\prec_{\mu_{\omega_1}^{n+1}}$ below y_k . Let δ_{k+1} be the least such ordinal δ . This defines a sequence $\langle \delta_k : k \in \omega \rangle$ of ordinals below ω_{n+1} and corresponding sequence $\langle y_k : k \in \omega \rangle$ in X . Note that for all $k \in \omega$, $y_{k+1} \prec_{\mu_{\omega_1}^{n+1}} y_k$. Since $|\{\delta_k : k \in \omega\}| = \omega < \omega_1$, property (2) at n gives a sequence $\langle h_k : k \in \omega \rangle$ so that $[h_n]_{\mu_{\omega_1}^n} = \delta_k$. For each $n \in \omega$, let $E_k = \{\ell \in [\omega_1]^{n+1} : \mathcal{K}^{n,h_{k+1}}(\ell) < \mathcal{K}^{n,h_k}(\ell)\}$ and note that $E_k \in \mu_{\omega_1}^{n+1}$ since $y_{k+1} \prec_{\mu_{\omega_1}^{n+1}} y_k$. Since $\omega_1 \rightarrow_* (\omega_1)_2^{2n+2}$ implies $\mu_{\omega_1}^{n+1}$ is countably complete, $E = \bigcap_{k \in \omega} E_k \in \mu_{\omega_1}^{n+1}$ and hence nonempty. Let $\ell \in E$. Then $\langle \mathcal{K}^{n,h_k}(\ell) : k \in \omega \rangle$ is an infinite descending sequence of ordinals which is a contradiction. It has been shown that $j_{\mu_{\omega_1}^{n+1}}(\omega_1)$ is a wellordering.

Next, one will show that $j_{\mu_{\omega_1}^{n+1}}(\omega_1) \leq \omega_{n+2}$. Let $x \in j_{\mu_{\omega_1}^{n+1}}(\omega_1)$, $f : [\omega_1]^{n+1} \rightarrow \omega_1$ represent x , and $\mathcal{K}$ be a Kunen function bounding I_f^1 . The argument above showed that for any $y \prec_{\mu_{\omega_1}^{n+1}} x$, there is an ordinal $\delta < \omega_{n+1}$ so that for any $h : [\omega_1]^n \rightarrow \omega_1$ which represents δ , $\mathcal{K}^{n,h} : [\omega_1]^{n+1} \rightarrow \omega_1$ represents y . Let δ_y be the least such δ for y . The map $\Upsilon : \text{init}_{\mu_{\omega_1}^{n+1}}(x) \rightarrow \omega_{n+1}$ defined by $\Upsilon(y) = \delta_y$ is an injection. So $\text{init}_{\mu_{\omega_1}^{n+1}}(x)$ has cardinality less than or equal to ω_{n+1} . Since $x \in j_{\mu_{\omega_1}^{n+1}}(\omega_1)$ was arbitrary, this shows that $j_{\mu_{\omega_1}^{n+1}}(\omega_1) \leq (\omega_{n+1})^+ = \omega_{n+2}$.

By Fact 2.26, $\omega_{n+1} = j_{\mu_{\omega_1}^n}(\omega_1) < j_{\mu_{\omega_1}^{n+1}}(\omega_1) \leq \omega_{n+2}$. Since Fact 2.22 implies $j_{\mu_{\omega_1}^{n+1}}(\omega_1)$ is a cardinal, one has that $j_{\mu_{\omega_1}^{n+1}}(\omega_1) = \omega_{n+2}$. $\text{cof}(\omega_{n+2}) = j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$ by Fact 2.25. Property (1) at $n + 1$ has been established.

Now to establish property (2) at $n + 1$. Suppose $A \subseteq \omega_{n+2}$ with $|A| \leq \omega_1$. Since $\text{cof}(\omega_{n+2}) = \omega_2$, one has that $\sup(A) < \omega_{n+2}$. Let $f : [\omega_1]^{n+1} \rightarrow \omega_1$ represent $\sup(A)$ and let $\mathcal{K}$ be a Kunen function bounding I_f^1 . As argued above, there is a sequence $\langle \delta_\alpha : \alpha \in A \rangle$ in ω_{n+1} with the property that for all $\alpha \in A$, for any $h : [\omega_1]^n \rightarrow \omega_1$ representing δ_α , $\mathcal{K}^{n,h}$ represents α . The set $\{\delta_\alpha : \alpha \in A\}$ is a subset of ω_{n+1} of cardinality less than or equal to ω_1 . By property (2) at n , there is a sequence $\langle h_\alpha : \alpha \in A \rangle$ so that $h_\alpha : [\omega_1]^n \rightarrow \omega_1$ represents δ_α . Then $\langle \mathcal{K}^{n,h_\alpha} : \alpha \in A \rangle$ has the property that for all $\alpha \in A$, $\mathcal{K}^{n,h_\alpha}$ represents α . This verifies property (2) at $n + 2$.

By induction, this completes the proof. $\square$

As mentioned in the footnote, the proof of Fact 2.30 actually showed that one can find representatives for ω_1 -many elements of $j_{\mu_{\omega_1}^n}(\omega_1)$. Although this is not needed in the proof of Fact 2.30 or anywhere else in this paper, this is a very important instance of choice that is required for many combinatorial results below ω_ω . For example, it is needed to show ω_2 is a weak partition cardinal. This fact proved within the proof of Fact 2.30 is explicitly isolated below.

Fact 2.31. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. Let $1 \leq n < \omega$ and $A \subseteq j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$ with $|A| \leq \omega_1$. There is a function Γ on A so that for all $\alpha \in A$, $\Gamma(\alpha) : [\omega_1]^n \rightarrow \omega_1$ and $\alpha = [\Gamma(\alpha)]_{\mu_{\omega_1}^n}$.

[4] shows that these combinatorial methods using the Kunen functions can show that $j_{\mu_{\omega_1}^\epsilon}(\omega_\omega)$ is wellfounded and even show that $j_{\mu_{\omega_1}^\epsilon}(\omega_\omega) < \omega_{\omega+1}$ for all $\epsilon < \omega_1$. However this seems to be the limit of the purely combinatorial methods. These methods cannot be used to calculate $j_{\mu_{\omega_1}^\epsilon}(\omega_{\omega+1})$ for $\epsilon < \omega_1$. These combinatorial methods have no influence on the ultrapower by the strong partition measure $\mu_{\omega_1}^{\omega_1}$. Using Martin's good coding system for $\omega^\epsilon \omega_1$ for $\epsilon < \omega_1$ to make complexity calculations, [4] showed that $j_{\mu_{\omega_1}^\epsilon}(\omega_{\omega+1}) = \omega_{\omega+1}$ for all $\epsilon < \omega_1$. Calculating the ultrapowers by the strong partition measure on ω_1 is an important question concerning the strong partition property. [4] showed that $j_{\mu_{\omega_1}^{\omega_1}}(\omega_1)$ is wellfounded in AD alone by using Martin's good coding system for $\omega_1 \omega_1$ to bring the ultrapower into $L(\mathbb{R})$ to apply a result of Kechris [17] which states that AD implies $L(\mathbb{R}) \models \text{DC}$. The first step in understanding the ultrapower by the strong partition measure on ω_1 was completed in [4] by answering a question of Goldberg that $j_{\mu_{\omega_1}^{\omega_1}}(\omega_1) < \omega_{\omega+1}$.

Let κ be a cardinal, $\epsilon \leq \kappa$, and $\Phi : [\kappa]^\epsilon \rightarrow \text{ON}$. $\Phi : [\kappa]^\epsilon \rightarrow \text{ON}$ is everywhere monotonic if and only if for all $f, g \in [\kappa]^\epsilon$, if for all $\alpha < \epsilon$, $f(\alpha) \leq g(\alpha)$, then $\Phi(f) \leq \Phi(g)$. Φ is almost everywhere monotonic if and only if there is a club $C \subseteq \kappa$ so that for all $f, g \in [C]_*^\epsilon$, if for all $\alpha < \epsilon$, $f(\alpha) \leq g(\alpha)$, then $\Phi(f) \leq \Phi(g)$. [14] shows that suitable partition relations on κ imply Φ is almost everywhere monotonic. However for functions on $[\kappa]^n$ when n is finite, almost everywhere monotonicity can be shown quite easily.

Fact 2.32. Assume κ is an uncountable cardinal, $1 \leq n < \omega$, and $\kappa \rightarrow_* (\kappa)_2^{n+1}$. For any function $\Phi : [\kappa]^n \rightarrow \text{ON}$, there is a club $C \subseteq \kappa$ so that for all $\ell, \iota \in [C]_*^n$, if for all $k < n$ $\ell(k) \leq \iota(k)$, then $\Phi(\ell) \leq \Phi(\iota)$. There is function $\Psi : [\kappa]^n \rightarrow \text{ON}$ which is everywhere monotonic and $[\Phi]_{\mu_\kappa^n} = [\Psi]_{\mu_\kappa^n}$.

Proof. Let $k < n$. If $\sigma \in [\kappa]^{n+1}$, then for $i \in 2$, define $\sigma^{k,i} \in [\kappa]^n$ by

$$\sigma^{k,i}(j) = \begin{cases} \sigma(j) & j < k \\ \sigma(k+i) & j = k \\ \sigma(j+1) & k < j < n \end{cases}.$$

Define $P_k : [\kappa]^{n+1} \rightarrow 2$ by $P_k(\sigma) = 0$ if and only if $\Phi(\sigma^{k,0}) \leq \Phi(\sigma^{k,1})$. By $\kappa \rightarrow_* (\kappa)_2^{n+1}$, there is a club $C_k \subseteq \kappa$ homogeneous for P_k . Suppose for sake of contradict, C_k is homogeneous for P_k taking value 1. For each $m \in \omega$, define $\tau_m \in [C_k]_*^n$ by

$$\tau_m(j) = \begin{cases} \text{enum}_{[C_k]_*^1}(j) & j < k \\ \text{enum}_{[C_k]_*^1}(\omega + m) & j = k \\ \text{enum}_{[C_k]_*^1}(\omega + \omega + j) & k < j < n \end{cases}.$$

For all $m \in \omega$, there is a unique $\sigma_m \in [C_k]_*^{n+1}$ so that $\sigma_m^{k,0} = \tau_m$ and $\sigma_m^{k,1} = \tau_{m+1}$. For each $m \in \omega$, $P_k(\sigma_m) = 1$ implies that $\Phi(\tau_{m+1}) = \Phi(\sigma_m^{k,1}) < \Phi(\sigma_m^{k,0}) = \Phi(\tau_m)$. Thus $\langle \Phi(\tau_m) : m \in \omega \rangle$ is an infinite descending sequence of ordinals which is a contradiction. It has been shown that C_k must be homogeneous for P_k taking value 0.

Let $C = \bigcap_{k < n} C_k$. Let $\ell, \iota \in [C]_*^n$ be such that $\ell(k) \leq \iota(k)$ for all $k < n$. For $e < n+1$, let $\nu_e \in [C]_*^n$ be defined by

$$\nu_e(k) = \begin{cases} \ell(k) & k < n - e \\ \iota(k) & n - e \leq k < n \end{cases}.$$

Note that $\nu_0 = \ell$ and $\nu_n = \iota$. For each $k < n$, if $\nu_k \neq \nu_{k+1}$, then there is a unique $\sigma_k \in [C]_*^n$ so that $\sigma_k^{n-1-k,0} = \nu_k$ and $\sigma_k^{n-1-k,1} = \nu_{k+1}$. For each $k < n$, if $\nu_k = \nu_{k+1}$, then clearly $\Phi(\nu_k) \leq \Phi(\nu_{k+1})$. For each $k < n$ with $\nu_k \neq \nu_{k+1}$, $P_k(\sigma_k) = 0$ implies that $\Phi(\nu_k) = \Phi(\sigma_k^{n-1-k,0}) \leq \Phi(\sigma_k^{n-1-k,1}) = \Phi(\nu_{k+1})$. Thus $\Phi(\ell) = \Phi(\nu_0) \leq \Phi(\nu_1) \leq \dots \leq \Phi(\nu_{n-1}) \leq \Phi(\nu_n) = \Phi(\iota)$.

Let $D = \{\alpha \in C : \text{enum}_C(\alpha) = \alpha\}$. Define $\Psi(\ell) = \Phi(\text{enum}_C \circ \ell)$. Ψ is everywhere monotonic and $[\Psi]_{\mu_\kappa^n} = [\Phi]_{\mu_\kappa^n}$ since for all $\ell \in [D]_*^n$, $\ell = \text{enum}_C \circ \ell$. $\square$

Definition 2.33. Let $<_{\text{lex}}$ be the lexicographic ordering on n -tuples. If $\iota \in [\omega_1]^n$ (so ι is an increasing function), let $\iota^* \in {}^n \omega_1$ be defined by $\iota^*(k) = \iota(n-1-k)$. Define $\sqsubset_n$ on $[\omega_1]^n$ by $\iota \sqsubset_n \ell$ if and only if $\iota^* <_{\text{lex}} \ell^*$.

(Even more explicitly, let $\alpha_0 < \dots < \alpha_{n-1} < \omega_1$ and $\beta_0 < \dots < \beta_{n-1} < \omega_1$. $(\alpha_0, \dots, \alpha_{n-1}) \sqsubset_n (\beta_0, \dots, \beta_{n-1})$ if and only if $(\alpha_{n-1}, \alpha_{n-2}, \dots, \alpha_0) <_{\text{lex}} (\beta_{n-1}, \beta_{n-2}, \dots, \beta_0)$.)

A function $f : [\omega_1]^n \rightarrow \omega_1$ has type n if and only if the following hold:

- f is order preserving from $([\omega_1]^n, \sqsubset_n)$ into the usual ordinal ordering $(\omega_1, <)$.
- (Discontinuous everywhere) For any $\ell \in [\omega_1]^n$, $\sup\{f(\iota) : \iota \sqsubset_n \ell\} < f(\ell)$.
- (Uniform cofinality ω) There is a function $F : [\omega_1]^n \times \omega \rightarrow \omega_1$ so that for all $\ell \in [\omega_1]^n$ and $k \in \omega$, $F(\ell, k) < F(\ell, k+1)$ and $f(\ell) = \sup\{F(\ell, k) : k \in \omega\}$.

Note that a function $f : \omega_1 \rightarrow \omega_1$ has type 1 if and only if it is an increasing function of the correct type (that is, $f \in [\omega_1]_{\ast}^{\omega_1}$).

Define $\mathfrak{B}_{n+1} \subseteq \omega_{n+1}$ to be the set of $\delta \in \omega_{n+1}$ so that there is a function $f : [\omega_1]^n \rightarrow \omega_1$ of type n with $\delta = [f]_{\mu_{\omega_1}^n}$. If $C \subseteq \omega_1$, then let $\mathfrak{B}_{n+1}^C$ be the set of $\delta \in \omega_{n+1}$ so that there is a function $f : [\omega_1]^n \rightarrow C$ of type n with $\delta = [f]_{\mu_{\omega_1}^n}$.

As mentioned earlier, if $f, g : [\omega_1]^n \rightarrow \omega_1$ and $[f]_{\mu_{\omega_1}^n} = [g]_{\mu_{\omega_1}^n}$, then it is not necessarily true that $[I_f^1]_{\mu_{\omega_1}^1} = [I_g^1]_{\mu_{\omega_1}^1}$. However, this is true if f and g are everywhere monotonic.

Fact 2.34. *Let $1 \leq n < \omega$ and $\omega_1 \rightarrow_{\ast} (\omega_1)_2^n$.*

- (1) *If $f, g : [\omega_1]^n \rightarrow \omega_1$ are everywhere monotonic and $[f]_{\mu_{\omega_1}^n} = [g]_{\mu_{\omega_1}^n}$, then $[I_f^1]_{\mu_{\omega_1}^1} = [I_g^1]_{\mu_{\omega_1}^1}$.*
- (2) *If $f, g : [\omega_1]^n \rightarrow \omega_1$ are order preserving on $\sqsubset_n$ and $[f]_{\mu_{\omega_1}^n} = [g]_{\mu_{\omega_1}^n}$, then for all $k \leq n$, $[I_f^k]_{\mu_{\omega_1}^k} = [I_g^k]_{\mu_{\omega_1}^k}$.*

Proof. The first statement will be shown. The second statement is proved in a similar manner. Suppose $f, g : [\omega_1]^n \rightarrow \omega_1$ are everywhere monotonic and $[f]_{\mu_{\omega_1}^n} = [g]_{\mu_{\omega_1}^n}$. Let $C \subseteq \omega_1$ be a club so that for all $\ell \in [C]_{\ast}^n$, $f(\ell) = g(\ell)$ and assume C consists of indecomposable ordinals. Let $D = \{\alpha \in C : \text{enum}_C(\alpha) = \alpha\}$. Pick $\gamma \in D$. Let $\ell \in [\omega_1]^n$ such that $\ell(n-1) = \gamma$. Let $\iota \in [C]_{\ast}^n$ be defined by $\iota(k) = \text{enum}_C \circ \ell$. By Fact 2.4, $\iota(n-1) = \ell(n-1)$ and for all $k < n$, $\ell(k) \leq \iota(k)$. By everywhere monotonicity, $f(\ell) \leq f(\iota) = g(\iota)$ and $g(\ell) \leq g(\iota) = f(\iota)$. Since $\ell \in [\omega_1]^n$ was arbitrary with $\ell(n-1) = \gamma$, one has that $I_f^1(\gamma) = \sup\{f(\ell) : \ell(n-1) = \gamma\} = \sup\{g(\ell) : \ell(n-1) = \gamma\} = I_g^1(\gamma)$. Since $\gamma \in D$ was arbitrary, this shows $[I_f^1]_{\mu_{\omega_1}^1} = [I_g^1]_{\mu_{\omega_1}^1}$. $\square$

Although in Definition 2.27, I_f^k is made for any function $f : [\omega_1]^n \rightarrow \omega_1$, I_f^k has the intended meaning only for functions f of type n when $1 < k < n$.

Fact 2.35. *Assume $\omega_1 \rightarrow_{\ast} (\omega_1)_2^{\omega_1}$ and $\star$. Let $1 \leq n < \omega$. If $\delta \in \omega_{n+1} \setminus \omega_n$, then there is an everywhere monotonic function $f : [\omega_1]^n \rightarrow \omega_1$ such that $[f]_{\mu_{\omega_1}^n} = \delta$ with the property that for all $\iota_0, \iota_1 \in [\omega_1]^n$, if $\iota_0(n-1) < \iota_1(n-1)$, then $f(\iota_0) < f(\iota_1)$.*

Proof. Using Fact 2.32, let $f_0 : [\omega_1]^n \rightarrow \omega_1$ be an everywhere monotonic representative for δ with respect to $\mu_{\omega_1}^n$. Let $A_0 = \{\iota \in [\omega_1]_{\ast}^n : f(\iota) \geq \iota(n-1)\}$. One must have that $A_0 \in \mu_{\omega_1}^n$ since otherwise Fact 2.6 implies there is a function $g : [\omega_1]^{n-1} \rightarrow \omega_1$ so that for $\mu_{\omega_1}^n$ -almost all ℓ , $f_0(\ell) = g(\ell \upharpoonright n-1)$. This would imply that $\delta = [f_0]_{\mu_{\omega_1}^n} = [g]_{\mu_{\omega_1}^{n-1}} \in j_{\mu_{\omega_1}^{n-1}}(\omega_1) = \omega_n$ which contradicts the assumption that $\delta \in \omega_{n+1} \setminus \omega_n$. Let $C_0 \subseteq \omega_1$ be a club so that $[C_0]_{\ast}^n \subseteq A_0$. Define $P : [C_0]_{\ast}^{n+1} \rightarrow 2$ by $P(\ell) = 0$ if and only if $f_0(\ell \upharpoonright n) < \ell(n)$. Let $C_1 \subseteq C_0$ be homogeneous for P using $\omega_1 \rightarrow_{\ast} (\omega_1)_2^{n+1}$. Pick any $\iota \in [C_1]_{\ast}^n$. Let $\ell = \iota \hat{\ } \langle \text{next}_{C_1}^{\omega}(f_0(\iota)) \rangle$ and note that $\ell \in [C_1]_{\ast}^{n+1}$. Then $f_0(\ell \upharpoonright n) = f_0(\iota) < \text{next}_{C_1}^{\omega}(f_0(\iota)) = \ell(n+1)$. Thus $P(\ell) = 0$ and hence C_1 must be homogeneous for P taking value 0. For any $\iota \in [C_1]_{\ast}^n$, let $\ell_{\iota} = \iota \hat{\ } \langle \text{next}_{C_0}^{\omega}(\iota(n-1)) \rangle$. $P(\ell_{\iota}) = 0$ implies that $f_0(\ell_{\iota}) = f_0(\ell_{\iota} \upharpoonright n) < \ell_{\iota}(n) = \text{next}_{C_1}^{\omega}(\iota(n-1))$. Let $C_2 = \{\alpha \in C_1 : \text{enum}_{C_1}(\alpha) = \alpha\}$. Note that if $\ell \in [\omega_1]_{\ast}^n$, then $\text{enum}_{C_2} \circ \ell \in [C_2]_{\ast}^n$. Define $f_1 : [\omega_1]^n \rightarrow \omega_1$ by $f_1(\iota) = f_0(\text{enum}_{C_2} \circ \iota)$. f_1 is everywhere monotonic. Now suppose $\iota_0, \iota_1 \in [\omega_1]_{\ast}^n$ with $\iota_0(n-1) < \iota_1(n-1)$. By the observations above and the definition of C_2 , one has that $f_1(\iota_0) = f_0(\text{enum}_{C_2} \circ \iota_0) < \text{next}_{C_1}^{\omega}(\text{enum}_{C_2}(\iota_0(n-1))) < (\text{enum}_{C_2} \circ \iota_1)(n-1) \leq f_0(\text{enum}_{C_2} \circ \iota_1) = f_1(\iota_1)$. Let $C_3 = \{\alpha \in C_2 : \text{enum}_{C_2}(\alpha) = \alpha\}$. For all $\iota \in [C_3]_{\ast}^n$, one has that $\text{enum}_{C_2} \circ \iota = \iota$. Thus $\delta = [f_0]_{\mu_{\omega_1}^n} = [f_1]_{\mu_{\omega_1}^n}$. f_1 is the representative of δ with the desired properties. $\square$

Definition 2.36. Let $V_n = \{(\alpha_{n-1}, \dots, \alpha_0, \gamma) \in {}^{n+1}\omega_1 : \alpha_0 < \alpha_1 < \dots < \alpha_{n-1} \wedge \gamma \leq \alpha_{n-1}\}$. Let $\ll$ be the lexicographic ordering on V_n . Let $\mathcal{V}_n = (V_n, \ll)$ which is a wellordering of ordertype ω_1 . A function $\phi : V_n \rightarrow \omega_1$ has the correct type if and only if the following holds:

- ϕ is order preserving from $\mathcal{V}_n$ into $(\omega_1, <)$, the usual ordering on ω_1 .
- ϕ is discontinuous everywhere: for all $x \in V_n$, $\phi(x) > \sup\{\phi(y) : y \ll x\}$.
- ϕ has uniform cofinality ω : there is a function $\Phi : V_n \times \omega \rightarrow \omega_1$ with the property that for all $x \in V$ and $k \in \omega$, $\Phi(x, k) < \Phi(x, k+1)$ and $\phi(x) = \sup\{\Phi(x, k) : k \in \omega\}$.

Fact 2.37. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. Let $C \subseteq \omega_1$ be a club. There is an order embedding of ω_{n+1} into $\mathfrak{B}_{n+1}^C$.

Proof. Let $\phi : V_n \rightarrow C$ be a function of the correct type. Since the order type of $\omega_{n+1} \setminus \omega_n$ is ω_{n+1} , it suffices to define an order embedding of $\omega_{n+1} \setminus \omega_n$ into $\mathfrak{B}_{n+1}^C$. Let $\delta \in \omega_{n+1} \setminus \omega_n$. By Fact 2.35, there is an everywhere monotonic $f : [\omega_1]^n \rightarrow \omega_1$ so that $\delta = [f]_{\mu_{\omega_1}^n}$ and has the property that for all $\iota_0, \iota_1 \in [\omega_1]^n$, if $\iota_0(n-1) < \iota_1(n-1)$, then $f(\iota_0) < f(\iota_1)$. Suppose $\alpha_0 < \alpha_1$ are two limit ordinals. Let γ_0, γ_1 be such that $\alpha_0 < \gamma_0 < \gamma_1 < \alpha_1$. For $i \in 2$, let $\iota_{\gamma_i} = (0, 1, \dots, n-2, \gamma_i)$. By the property of f , one has that $f(\iota_{\gamma_0}) < f(\iota_{\gamma_1})$. Hence $I_f^1(\alpha_0) \leq f(\iota_{\gamma_0}) < f(\iota_{\gamma_1}) < I_f^1(\alpha_1)$. So I_f^1 is an increasing function on the limit ordinals below ω_1 . This implies that $(I_f^1(\omega \cdot \iota(n-1)), I_f^1(\omega \cdot \iota(n-2)), \dots, I_f^1(\omega \cdot \iota(0)), f(\iota)) \in V_n$ when $\iota \in [\omega_1]_*^n$. Let $\hat{f} : [\omega_1]^n \rightarrow \omega_1$ be defined by $\hat{f}(\iota) = \phi(I_f^1(\omega \cdot \iota(n-1)), I_f^1(\omega \cdot \iota(n-2)), \dots, I_f^1(\omega \cdot \iota(0)), f(\iota))$. Define $\Psi : (\omega_{n+1} \setminus \omega_n) \rightarrow \mathfrak{B}_{n+1}^C$ by $\Psi(\delta) = [\hat{f}]_{\mu_{\omega_1}^n}$ for any everywhere monotonic $f : [\omega_1]^n \rightarrow \omega_1$ such that $\delta = [f]_{\mu_{\omega_1}^n}$ and for all $\iota_0, \iota_1 \in [\omega_1]^n$, if $\iota_0(n-1) < \iota_1(n-1)$, then $f(\iota_0) < f(\iota_1)$. Ψ is well defined independent of the choice of such f representing δ using Fact 2.34 (1). Suppose $\iota_0 \sqsubset_n \iota_1$. If $k < n$ is largest such that $\iota_0(k) \neq \iota_1(k)$, then $\iota_0(k) < \iota_1(k)$. By the observation above, $I_f^1(\omega \cdot \iota_0(j)) = I_f^1(\omega \cdot \iota_1(j))$ for all $k < j < n$ and $I_f^1(\omega \cdot \iota_0(k)) < I_f^1(\omega \cdot \iota_1(k))$. Since ϕ is order preserving on $\mathcal{V}_n$, one has that $\hat{f}(\iota_0) < \hat{f}(\iota_1)$. This shows that $\hat{f}$ is order preserving on $([\omega_1]^n, \sqsubset_n)$. $\hat{f}$ is discontinuous everywhere and has uniform cofinality ω since ϕ is discontinuous everywhere and has uniform cofinality ω . So $\hat{f}$ has type n . This shows that Ψ does map into $\mathfrak{B}_{n+1}^C$. Suppose $\delta_0, \delta_1 \in \omega_{n+1} \setminus \omega_n$ and $\delta_0 < \delta_1$. Let $g_0, g_1 : [\omega_1]^n \rightarrow \omega_1$ represent δ_0 and δ_1 , respectively, with the necessary properties stated above. Let $D \subseteq \omega_1$ be a club so that for all $\iota \in [D]_*^n$, $g_0(\iota) < g_1(\iota)$. For $i \in 2$, define $f_i : [\omega_1]^n \rightarrow \omega_1$ by $f_i(\ell) = g_i(\text{enum}_D \circ \ell)$. Let $\tilde{D} = \{\alpha \in D : \text{enum}_D(\alpha) = \alpha\}$. Note that for all $\ell \in \tilde{D}$ and $i \in 2$, $f_i(\ell) = g_i(\ell)$. Thus $\delta_i = [g_i]_{\mu_{\omega_1}^n} = [f_i]_{\mu_{\omega_1}^n}$ and f_i still has the necessary properties. Moreover, for all $\ell \in [\omega_1]^n$, $f_0(\ell) < f_1(\ell)$. Thus for all $\alpha \in [\omega_1]^n$, $I_{f_0}^1(\alpha) \leq I_{f_1}^1(\alpha)$. So for all $\iota \in [\omega_1]^n$, $(I_{f_0}^1(\omega \cdot \iota(n-1)), \dots, I_{f_0}^1(\omega \cdot \iota(0)), f_0(\iota)) \ll (I_{f_1}^1(\omega \cdot \iota(n-1)), \dots, I_{f_1}^1(\omega \cdot \iota(0)), f_1(\iota))$ and therefore $\hat{f}_0(\iota) < \hat{f}_1(\iota)$. It has been shown that $\Psi(\delta_0) = [\hat{f}_0]_{\mu_{\omega_1}^n} < [\hat{f}_1]_{\mu_{\omega_1}^n} = \Psi(\delta_1)$. This shows that $\Psi : (\omega_{n+1} \setminus \omega_n) \rightarrow \mathfrak{B}_{n+1}^C$ is an order preserving injection. $\square$

Definition 2.38. Suppose $1 \leq n < \omega$ and $\delta \in \mathfrak{B}_{n+1}$. For any $1 \leq k \leq n$, let $\mathcal{I}_\delta^k = [I_f^k]_{\mu_{\omega_1}^k}$ for all $f : [\omega_1]^n \rightarrow \omega_1$ of type n such that $[f]_{\mu_{\omega_1}^n} = \delta$. (By Fact 2.34 (2), $\mathcal{I}_\delta^k$ is independent of the choice of f representing δ but f must be a function of type n .)

Fact 2.39. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. Let $1 \leq n < \omega$, $C \subseteq \omega_1$ be a club, and $\phi : V_{n+1} \rightarrow C$ be a function of the correct type. Let $\psi : [\omega_1]^n \rightarrow \omega_1$ be defined by $\psi(\alpha_0, \dots, \alpha_{n-1}) = \sup\{\phi(\alpha_{n-1}, \dots, \alpha_0, \zeta, 0) : \zeta < \alpha_0\}$. Let $\delta = [\psi]_{\mu_{\omega_1}^n}$. Then $|\{\eta \in \mathfrak{B}_{n+2}^C : \mathcal{I}_\eta^n = \delta\}| = |\omega_{n+1}|$.

Proof. Let $A = \{\eta \in \mathfrak{B}_{n+2}^C : \mathcal{I}_\eta^n = \delta\}$. Let $\nu \in \omega_{n+1}$. Let $f : [\omega_1]^n \rightarrow \omega_1$ be such that $[f]_{\mu_{\omega_1}^n} = \nu$. By Fact 2.8, there is a club $D_0 \subseteq \omega_1$ so that for all $\ell \in [D]_*^n$, $f(\ell) < \text{next}_{D_0}^\omega(\ell(n-1))$. Let $D_1 = \{\alpha \in D_0 : \text{enum}_{D_0}(\alpha) = \alpha\}$. For all $\ell \in [D_1]^{n+1}$, $f(\ell \upharpoonright n) < \text{next}_{D_0}^\omega(\ell(n-1)) < \ell(n)$ using Fact 2.4. Define a function $\hat{f} : [D_1]^{n+1} \rightarrow C$ by

$$\hat{f}(\alpha_0, \dots, \alpha_n) = \phi(\alpha_n, \dots, \alpha_0, f(\alpha_0, \dots, \alpha_{n-1})).$$

Note that this is well defined since $(\alpha_n, \dots, \alpha_0, f(\alpha_0, \dots, \alpha_{n-1})) \in V_{n+1}$ by the property of D_1 . Let $\tilde{f} : [\omega_1]^{n+1} \rightarrow C$ be defined by $\tilde{f}(\ell) = \hat{f}(\text{enum}_{D_1} \circ \ell)$. Let $D_2 = \{\alpha \in D_1 : \text{enum}_{D_1}(\alpha) = \alpha\}$. Note that for all $\ell \in [D_2]_*^{n+1}$, $\ell = \text{enum}_{D_1} \circ \ell$. Thus $[\tilde{f}]_{\mu_{\omega_1}^{n+1}} = [\hat{f}]_{\mu_{\omega_1}^{n+1}}$. Note that $\tilde{f}$ has type $n+1$ since ϕ has the correct type from $\mathcal{V}_{n+1}$ into $(C, <)$. So $[\tilde{f}]_{\mu_{\omega_1}^{n+1}} \in \mathfrak{B}_{n+2}^C$. Observe that for all $(\alpha_0, \dots, \alpha_{n-1}) \in [D_2]_*^n$,

$$\begin{aligned} I_{\tilde{f}}^n(\alpha_0, \dots, \alpha_{n-1}) &= \sup\{\tilde{f}(\zeta, \alpha_0, \dots, \alpha_{n-1}) : \zeta < \alpha_0\} = \sup\{\hat{f}(\zeta, \alpha_0, \dots, \alpha_{n-1}) : \zeta < \alpha_0\} \\ &= \sup\{\phi(\alpha_{n-1}, \dots, \alpha_0, \zeta, f(\alpha_0, \dots, \alpha_{n-1})) : \zeta < \alpha_0\} = \sup\{\phi(\alpha_{n-1}, \dots, \alpha_0, \zeta, 0) : \zeta < \alpha_0\} = \psi(\alpha_0, \dots, \alpha_{n-1}) \end{aligned}$$

Let $\Upsilon(\nu) = [\tilde{f}]_{\mu_{\omega_1}^{n+1}} = [\hat{f}]_{\mu_{\omega_1}^{n+1}}$. By the above discussion, $\Upsilon(\nu) \in \mathfrak{B}_{n+2}^C$ and $\mathcal{I}_{\Upsilon(\nu)}^n = [I_{\tilde{f}}^n]_{\mu_{\omega_1}^n} = [\psi]_{\mu_{\omega_1}^n} = \delta$. Thus $\Upsilon : \omega_{n+1} \rightarrow A$. Suppose $\eta_0 < \eta_1$. Let $f_0, f_1 : [\omega_1]^n \rightarrow \omega_1$ be such that $\eta_0 = [f_0]_{\mu_{\omega_1}^n}$ and $\eta_1 = [f_1]_{\mu_{\omega_1}^n}$. For $\mu_{\omega_1}^n$ -almost all ℓ , $f_0(\ell) < f_1(\ell)$. Thus for $\mu_{\omega_1}^{n+1}$ -almost all ι , $\hat{f}_0(\iota) < \hat{f}_1(\iota)$. Thus $\Upsilon : \omega_{n+1} \rightarrow A$ is order preserving and hence an injection. This shows that $|\omega_{n+1}| \leq |A|$. Since for every $\eta \in A$, $\mathcal{I}_{\eta}^n = \delta$, one has that A is a bounded subset of ω_{n+2} . Therefore $|A| \leq |\omega_{n+1}|$. It has been shown that $|A| = |\omega_{n+1}|$. $\square$

Fact 2.40. Suppose $1 \leq n < \omega$ and $g : [\omega_1]^n \rightarrow \omega_1$ is order preserving on $\sqsubset_n$. Then $[I_g^1]_{\mu_{\omega_1}^1} = [I_{I_g^{n-1}}^1]_{\mu_{\omega_1}^1}$. Thus if $\eta = [g]_{\mu_{\omega_1}^n}$ and $\chi = [I_g^{n-1}]_{\mu_{\omega_1}^{n-1}}$, then $\mathcal{I}_{\eta}^1 = \mathcal{I}_{\chi}^1$.

Proof. Fix $\omega \leq \alpha < \omega_1$. Let $\ell \in [\omega_1]^n$ be such that $\ell(n-1) = \alpha$. Let $\iota \in [\omega_1]^{n-1}$ be such that $\iota(i) = \ell(i+1)$ for all $i < n-1$. Then $I_g^{n-1}(\iota) \geq g(\ell)$. This shows that

$$I_{I_g^{n-1}}^1(\alpha) = \sup\{I_g^{n-1}(\iota) : \iota \in [\omega_1]^{n-1} \wedge \iota(n-2) = \alpha\} \geq g(\ell).$$

Since $\ell \in [\omega_1]_n^n$ so that $\ell(n-1) = \alpha$ was arbitrary, one has that

$$I_{I_g^{n-1}}^1(\alpha) \geq \sup\{g(\ell) : \ell \in [\omega_1]^n \wedge \ell(n-1) = \alpha\} = I_g^1(\alpha).$$

This shows that $I_{I_g^{n-1}}^1(\alpha) \geq I_g^1(\alpha)$ for all $\omega \leq \alpha < \omega_1$.

Fix a limit ordinal $\alpha < \omega_1$. Suppose $\iota \in [\omega_1]^{n-1}$ with $\iota(n-2) = \alpha$ and $\iota(0) > 0$. Let $\tilde{\iota} \in [\omega_1]^{n-1}$ be defined by $\tilde{\iota}(i) = \iota(i) + 1$ if $i < n-2$ and $\tilde{\iota}(n-2) = \alpha$. Let $\tilde{\ell} \in [\omega_1]^n$ be defined by $\tilde{\ell} = \langle 0 \rangle \hat{\sim} \tilde{\iota}$. Suppose $\ell \in [\omega_1]^n$ is such that for all $i \in \omega$, $\iota(i) = \ell(i+1)$, then $\ell \sqsubset_n \tilde{\ell}$. Since g is order preserving on $\sqsubset_n$, $g(\ell) < g(\tilde{\ell})$. Thus

$$I_g^{n-1}(\iota) = \sup\{g(\ell) : (\forall i < n-1)(\ell(i+1) = \iota(i))\} \leq g(\tilde{\ell})$$

Then

$$I_g^{n-1}(\iota) \leq \sup\{g(\ell) : \ell \in [\omega_1]^n \wedge \ell(n-1) = \alpha\} = I_g^1(\alpha)$$

Since $\iota \in [\omega_1]^{n-2}$ with $\iota(n-2) = \alpha$ was arbitrary,

$$I_{I_g^{n-1}}^1(\alpha) = \sup\{I_g^{n-1}(\iota) : \iota \in [\omega_1]^{n-1} \wedge \iota(n-2) = \alpha\} \leq I_g^1(\alpha).$$

It has been shown that for all $\omega \leq \alpha < \omega_1$, $I_{I_g^{n-1}}^1(\alpha) \leq I_g^1(\alpha)$.

Thus for all limit $\alpha < \omega_1$, $I_g^1(\alpha) = I_{I_g^{n-1}}^1(\alpha)$. $\square$

3. BOLDFACE GCH BELOW ω_ω

This section will prove the main result that the boldface GCH holds below ω_ω (Theorem 3.10). The key combinatorial lemma for this argument is Lemma 3.9 which is frequently called a “sliding lemma”.

Definition 3.1. Let κ be a cardinal. The boldface GCH holds at κ if and only if there is no injection of κ^+ into $\mathcal{P}(\kappa)$. The boldface GCH below κ is the statement that for all $\delta < \kappa$, the boldface GCH holds at δ .

Fact 3.2. Let κ be a cardinal and $\delta < \kappa$. If there is a δ^+ -complete nonprincipal ultrafilter on κ , then there is no injection of κ into $\mathcal{P}(\delta)$.

Proof. Let μ be a δ^+ -complete nonprincipal ultrafilter on κ . Suppose $\langle A_\alpha : \alpha < \kappa \rangle$ is an injection of κ into $\mathcal{P}(\delta)$. For each $\xi < \delta$, let $E_\xi^0 = \{\alpha \in \kappa : \xi \notin A_\alpha\}$ and $E_\xi^1 = \{\alpha \in \kappa : \xi \in A_\alpha\}$. Since μ is an ultrafilter, there is a unique $i_\xi \in 2$ so that $E_\xi^{i_\xi} \in \mu$. Let $E = \bigcap_{\xi < \delta} E_\xi^{i_\xi}$ and note that $E \in \mu$ since μ is δ^+ -complete. Since μ is nonprincipal, let $\alpha_0, \alpha_1 \in E$ with $\alpha_0 \neq \alpha_1$. For all $\xi < \delta$, since $\alpha_0, \alpha_1 \in E \subseteq E_\xi^{i_\xi}$, $\xi \in A_{\alpha_0}$ if and only if $i_\xi = 1$ if and only if $\xi \in A_{\alpha_1}$. Thus $A_{\alpha_0} = A_{\alpha_1}$. This contradicts the injectiveness of $\langle A_\alpha : \alpha < \kappa \rangle$. $\square$

Fact 3.3. If $\kappa \rightarrow_* (\kappa)_2^2$, then there is no injection of κ into $\mathcal{P}(\delta)$ for any $\delta < \kappa$.

Proof. $\kappa \rightarrow_* (\kappa)_2^2$ implies that μ_κ^1 is a κ -complete ultrafilter by Fact 2.5. The result follow from Fact 3.2. $\square$

Fact 3.4. $\omega_1 \rightarrow_* (\omega_1)_2^2$ implies the boldface GCH at ω .

Martin [16] (and Kleinberg [18]) (also see [5]) showed that $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$ imply that ω_2 is a weak partition cardinal (satisfies $\omega_2 \rightarrow (\omega_2)_2^{<\omega_2}$). Thus under these assumptions, Fact 3.3 implies that the boldface GCH holds at ω_1 . Although the proof of the weak partition property on ω_2 can be shown by similar techniques used here, it is very enlightening to see a direct proof of the boldface GCH at ω_1 to motivate the proof of the boldface GCH at ω_n for $2 \leq n < \omega$.

Definition 3.5. Let $U_1 = \{(0, 0)\} \cup \{(1, \alpha, i) : \alpha < \omega_1 \wedge i < 2\}$. Let $\ll_1$ be the lexicographic ordering on U_1 . Let $\mathcal{U}_1 = (U_1, \ll_1)$. Note that $(0, 0)$ is the minimal element of $\mathcal{U}_1$. Note that the ordertype of $\mathcal{U}_1$ is ω_1 . Suppose $F : U_1 \rightarrow \omega_1$. Define $F^0, F^1 : \omega_1 \rightarrow \omega_1$ and $F^2 \in \omega_1$ by $F^0(\alpha) = F(1, \alpha, 0)$, $F^1(\alpha) = F(1, \alpha, 1)$, and $F^2 = F(0, 0)$. (Note that F^2 is just a countable ordinal.) A function $F : U_1 \rightarrow \omega_1$ has the correct type if and only if the following two conditions hold.

- F is discontinuous everywhere: For all $x \in U_1$, $\sup\{F(y) : y \ll_1 x\} < F(x)$.
- F has uniform cofinality ω : There is a function $\mathcal{F} : U_1 \times \omega \rightarrow \omega$ so that for all $x \in U_1$ and $k \in \omega$, $\mathcal{F}(x, k) < \mathcal{F}(x, k+1)$ and $F(x) = \sup\{\mathcal{F}(x, k) : k \in \omega\}$.

If $F : U_1 \rightarrow \omega_1$ has the correct type, then $F^0, F^1 : \omega_1 \rightarrow \omega_1$ are functions of the correct type and $\text{cof}(F^2) = \omega$. If $X \subseteq \omega_1$, then let $[X]_*^{\mathcal{U}_1}$ be the set of all $F : U_1 \rightarrow X$ of the correct type and order preserving between $\mathcal{U}_1$ and $(X, <)$, where $<$ is the usual ordinal ordering.

Lemma 3.6. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{n+1}$. Let $f_0, f_1 : \omega_1 \rightarrow \omega_1$ be functions of type 1 such that $[f_0]_{\mu_{\omega_1}^1} < [f_1]_{\mu_{\omega_1}^1}$ and let $\delta \in [\omega_1]_*^1$ (i.e. is a limit ordinal). Then there is an $F \in [\omega_1]_*^{\mathcal{U}_1}$ so that $[F^0]_{\mu_{\omega_1}^1} = [f_0]_{\mu_{\omega_1}^1}$, $[F^1]_{\mu_{\omega_1}^1} = [f_1]_{\mu_{\omega_1}^1}$, $F^2 = \delta$, $F^0[\omega_1] \subseteq f_0[\omega_1]$, and $F^1[\omega_1] \subseteq f_1[\omega_1]$.

Proof. Since f_0 and f_1 are of type 1, they are both increasing, discontinuous, and have uniform cofinality ω . Let $G_0 : \omega_1 \times \omega \rightarrow \omega_1$ witness that f_0 has uniform cofinality ω and let $G_1 : \omega_1 \times \omega \rightarrow \omega_1$ witness that f_1 has uniform cofinality ω . Since $\delta < \omega_1$ is a limit ordinal, let $\rho : \omega \rightarrow \delta$ be an increasing cofinal function. For each $\alpha < \omega_1$, let $h(\alpha)$ be the least element $\bar{\alpha} \in \omega_1$ so that $f_1(\alpha) < f_0(\bar{\alpha})$. Since $[f_0]_{\mu_{\omega_1}^1} < [f_1]_{\mu_{\omega_1}^1}$, there is a club $C_0 \subseteq \omega_1$ so that for all $\alpha \in [C_0]_*^1$, $f_0(\alpha) < f_1(\alpha)$. Let $C_1 = \{\alpha \in C_0 : (\forall \alpha' < \alpha)(h(\alpha') < \alpha)\}$. C_1 is a club subset of C_0 . One may assume $\delta < \min(C_1)$. For notational simplicity, let $\epsilon = \text{enum}_{[C_1]_*^1}$. Define $F : U_1 \rightarrow \omega_1$ by $F(0, 0) = \delta$, and $F(1, \alpha, i) = f_i(\epsilon(\alpha))$ for $i < 2$ and $\alpha < \omega_1$. Fix $\alpha < \omega_1$. Note that $F(1, \alpha, 0) = f_0(\epsilon(\alpha)) < f_1(\epsilon(\alpha)) = F(1, \alpha, 1)$ by the property of the club C_0 and the fact that $\epsilon(\alpha) \in [C_1]_*^1$. $F(1, \alpha, 1) = f_1(\epsilon(\alpha)) < f_0(h(\epsilon(\alpha))) < f_0(\epsilon(\alpha + 1)) = F(1, \alpha + 1, 0)$ since the first inequality comes from the property of h and the second inequality comes from the property of C_1 . This shows that F is order preserving from $\mathcal{U}_1$ into the usual ordering on ω_1 . The elements of $\mathcal{U}_1$ of limit rank take the form $(1, \alpha, 0)$ where α is a limit ordinal. Note that $\sup\{F(x) : x \ll_1 (1, \alpha, 0)\} = \sup\{F(1, \alpha', 0) : \alpha' < \alpha\} = \sup\{f_0 \upharpoonright \epsilon(\alpha) < f_0(\epsilon(\alpha)) = F(1, \alpha, 0)$ using the discontinuity of f_0 . This shows that F is discontinuous everywhere. Let $\mathcal{F} : U_1 \times \omega \rightarrow \omega_1$ be defined by

$$\mathcal{F}(x, k) = \begin{cases} \rho(k) & x = (0, 0) \\ G_i(\epsilon(\alpha), k) & x = (1, \alpha, i) \end{cases}$$

$\mathcal{F}$ witnesses that F has uniform cofinality ω . Thus F has the correct type. By construction, it is clear that $F^0[\omega_1] \subseteq f_0[\omega_1]$ and $F^1[\omega_1] \subseteq f_1[\omega_1]$. Let $C_2 = \{\alpha \in C_1 : \text{enum}_{C_1}(\alpha) = \alpha\}$ which is a club subset of C_1 . For all $\alpha \in C_2$ and $i \in 2$, $F(1, \alpha, i) = f_i(\epsilon(\alpha)) = f_i(\text{enum}_{[C_1]_*^1}(\alpha)) = f_i(\alpha)$ since $\text{enum}_{C_1}(\alpha) = \alpha$ implies that $\text{ot}(\{\bar{\alpha} \in C_1 : \bar{\alpha} < \alpha \wedge \text{cof}(\bar{\alpha}) = \omega\}) = \alpha$ and thus $\text{enum}_{[C_1]_*^1}(\alpha) = \alpha$. This shows that for all $i < 2$, $[F^i]_{\mu_{\omega_1}^1} = [f_i]_{\mu_{\omega_1}^1}$. $\square$

Theorem 3.7. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $\star$. Then the boldface GCH holds at ω_1 .

Proof. By Fact 3.4, one has that the boldface GCH holds at ω . Suppose the boldface GCH at ω_1 fails. Let $\langle A_\eta : \eta < \omega_2 \rangle$ be an injection of ω_2 into $\mathcal{P}(\omega_1)$. Define $P : [\omega_1]_*^{\mathcal{U}_1} \rightarrow 2$ by $P(F) = 0$ if and only if $\min(A_{[F^0]_{\mu_{\omega_1}^1}} \Delta A_{[F^1]_{\mu_{\omega_1}^1}}) < F^2$ (where Δ refers to symmetric difference). Since $\mathcal{U}_1$ has ordertype ω_1 , $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ implies there is a club $C \subseteq \omega_1$ homogeneous for P . Pick any $f_0, f_1 : \omega_1 \rightarrow C$ of type 1 so that $[f_0]_{\mu_{\omega_1}^1} < [f_1]_{\mu_{\omega_1}^1}$. Since $\langle A_\eta : \eta < \omega_2 \rangle$ is an injection, $A_{[f_0]_{\mu_{\omega_1}^1}} \neq A_{[f_1]_{\mu_{\omega_1}^1}}$ and thus $A_{[f_0]_{\mu_{\omega_1}^1}} \Delta A_{[f_1]_{\mu_{\omega_1}^1}} \neq \emptyset$. Let $\delta \in [C]_*^1$ be such that $\min(A_{[f_0]_{\mu_{\omega_1}^1}} \Delta A_{[f_1]_{\mu_{\omega_1}^1}}) < \delta$. By Lemma 3.6, there is an $F \in [\omega_1]_*^{\mathcal{U}_1}$ so that

$[F^0]_{\mu_{\omega_1}^1} = [f_0]_{\mu_{\omega_1}^1}$, $[F^1]_{\mu_{\omega_1}^1} = [f_1]_{\mu_{\omega_1}^1}$, $F^2 = \delta$, $F^0[\omega_1] \subseteq f_0[\omega_1] \subseteq C$, and $F^1[\omega_1] \subseteq f_1[\omega_1] \subseteq C$. Thus $F \in [C]_*^{\mathcal{U}_1}$. Thus $\min(A_{[F^0]_{\mu_{\omega_1}^1}} \triangle A_{[F^1]_{\mu_{\omega_1}^1}}) < F^2$ implies that $P(F) = 0$. This shows that C must be homogeneous for P taking value 0. Fix a $\delta \in [C]_*^1$. By Fact 2.37, ω_2 order embeds into $\mathfrak{B}_2^C$. Pick any $\nu \in \mathfrak{B}_2^C$ so that $\mathfrak{B}_2^C \upharpoonright \nu = \{\eta \in \mathfrak{B}_2^C : \eta < \nu\}$ has cardinality ω_1 . Let $\eta_0, \eta_1 \in \mathfrak{B}_2^C \upharpoonright \nu$ with $\eta_0 \neq \eta_1$. Without loss of generality, suppose $\eta_0 < \eta_1$. Let $f_0, f_1 : \omega_1 \rightarrow C$ be functions of type 1 so that $\eta_0 = [f_0]_{\mu_{\omega_1}^1}$ and $\eta_1 = [f_1]_{\mu_{\omega_1}^1}$. By Lemma 3.6, there is an $F \in [C]_*^{\mathcal{U}_1}$ so that $[F^0]_{\mu_{\omega_1}^1} = [f_0]_{\mu_{\omega_1}^1} = \eta_0$, $[F^1]_{\mu_{\omega_1}^1} = [f_1]_{\mu_{\omega_1}^1} = \eta_1$, and $F^2 = \delta$. Thus $P(F) = 0$ implies that $\min(A_{\eta_0} \triangle A_{\eta_1}) < \delta$. This implies that the function $\Phi : \mathfrak{B}_2^C \upharpoonright \nu \rightarrow \mathcal{P}(\delta)$ defined by $\Phi(\eta) = A_\eta \cap \delta$ is an injection. Since $\delta < \omega_1$ implies $|\mathcal{P}(\delta)| = |\mathcal{P}(\omega)|$ and $|\mathfrak{B}_2^C \upharpoonright \nu| = \omega_1$, one has an injection of ω_1 into $\mathcal{P}(\omega)$. This violates the boldface GCH at ω . $\square$

Definition 3.8. Let $2 \leq n < \omega$. Let $U_n = \{(\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i) : \alpha_0 < \dots < \alpha_{n-1} < \omega_1 \wedge i < 2\} \cup \{(\alpha, 1) : n-1 \leq \alpha < \omega_1\}$. Let $\ll_n$ be the lexicographic ordering on U_n . Let $\mathcal{U}_n = (U_n, \ll_n)$. Observe that $\text{ot}(\mathcal{U}_n) = \omega_1$. Suppose $F : U_n \rightarrow \omega_1$. Define $F^0, F^1 : [\omega_1]^n \rightarrow \omega_1$ and $F^2 : \omega_1 \setminus (n-1) \rightarrow \omega_1$ by $F^0(\iota) = F(\iota(n-1), 0, \iota(n-2), \dots, \iota(0), 0)$, $F^1(\iota) = F(\iota(n-1), 0, \iota(n-2), \dots, \iota(0), 1)$, and $F^2(\alpha) = F(\alpha, 1)$. A function $F : U_n \rightarrow \omega_1$ has the correct type if and only if the following two conditions hold:

- F is discontinuous everywhere: For all $x \in U_n$, $\sup\{F(y) : y \ll_n x\} < F(x)$.
- F has uniform cofinality ω : There is a function $\mathcal{F} : U_n \times \omega \rightarrow \omega_1$ so that for all $x \in U_n$ and $k \in \omega$, $\mathcal{F}(x, k) < \mathcal{F}(x, k+1)$ and $F(x) = \sup\{\mathcal{F}(x, k) : k \in \omega\}$.

Note that if $F : U_n \rightarrow \omega_1$ has the correct type, then $F^0, F^1 : [\omega_1]^n \rightarrow \omega_1$ has type n , $[F^0]_{\mu_{\omega_1}^n} < [F^1]_{\mu_{\omega_1}^n}$, and $F^2 : \omega_1 \setminus (n-1) \rightarrow \omega_1$ has the correct type. If $X \subseteq \omega_1$, then let $[X]_*^{\mathcal{U}_n}$ be the set of all $F : U_n \rightarrow X$ of the correct type and order preserving between $\mathcal{U}_n$ and $(X, <)$ with the usual ordering.

Lemma 3.9. Suppose $2 \leq n < \omega$. Assume $\omega_1 \rightarrow_* (\omega_1)_2^{n+1}$. Let $f_0, f_1 : [\omega_1]^n \rightarrow \omega_1$ and $f_2 : \omega_1 \rightarrow \omega_1$ be functions with the following properties.

- f_0 and f_1 have type n . f_2 has type 1.
- $[I_{f_0}^{n-1}]_{\mu_{\omega_1}^{n-1}} = [I_{f_1}^{n-1}]_{\mu_{\omega_1}^{n-1}}$ and $[f_0]_{\mu_{\omega_1}^n} < [f_1]_{\mu_{\omega_1}^n}$.
- $[I_{f_0}^1]_{\mu_{\omega_1}^1} = [I_{f_1}^1]_{\mu_{\omega_1}^1} < [f_2]_{\mu_{\omega_1}^1}$.

Then there is an $F \in [\omega_1]_*^{\mathcal{U}_n}$ with the following properties.

- $[F^0]_{\mu_{\omega_1}^n} = [f_0]_{\mu_{\omega_1}^n}$, $[F^1]_{\mu_{\omega_1}^n} = [f_1]_{\mu_{\omega_1}^n}$, and $[F^2]_{\mu_{\omega_1}^1} = [f_2]_{\mu_{\omega_1}^1}$,
- $F^0[[\omega_1]^n] \subseteq f_0[[\omega_1]^n]$, $F^1[[\omega_1]^n] \subseteq f_1[[\omega_1]^n]$, and $F^2[\omega_1] \subseteq f_2[\omega_1]$.

Proof. Let $C_0 \subseteq \omega_1$ be a club with the following properties:

- (1) For all $\ell \in [C_0]_*^n$, $I_{f_0}^{n-1}(\ell) = I_{f_1}^{n-1}(\ell)$.
- (2) For all $\ell \in [C_0]_*^n$, $f_0(\ell) < f_1(\ell)$.
- (3) For all $\alpha \in C_0$, $I_{f_0}^1(\alpha) = I_{f_1}^1(\alpha) < f_2(\alpha)$.

If $\ell \in [\omega_1]^{n+1}$, let $\ell^0, \ell^1 \in [\omega_1]^n$ be defined by $\ell^1(k) = \ell(k+1)$ and

$$\ell^0(k) = \begin{cases} \ell(0) & k = 0 \\ \ell(k+1) & 0 < k < n \end{cases}.$$

Define $P : [C_0]^{n+1} \rightarrow 2$ by $P(\ell) = 0$ if and only if $f_1(\ell^0) < f_0(\ell^1)$. By $\omega_1 \rightarrow_* (\omega_1)_2^{n+1}$, there is a club $C_1 \subseteq C_0$ which is homogeneous for P . Let $C_2 = \{\alpha \in C_1 : \text{enum}_{C_1}(\alpha) = \alpha\}$. Pick $(\alpha_0, \alpha_1, \dots, \alpha_{n-1}) \in [C_2]_*^n$. Since α_1 is a limit point of C_1 and $I_{f_0}^{n-1}(\alpha_1, \dots, \alpha_{n-1}) = I_{f_1}^{n-1}(\alpha_1, \dots, \alpha_{n-1})$ because $(\alpha_1, \dots, \alpha_{n-1}) \in [C_2]_*^{n-1} \subseteq [C_0]_*^{n-1}$, there must be some $\gamma \in C_1$ so that $\alpha_0 < \gamma < \alpha_1$ and $f_1(\alpha_0, \dots, \alpha_{n-1}) < f_0(\gamma, \alpha_1, \dots, \alpha_{n-1})$. Pick $\ell \in [C_1]_*^{n+1}$ so that $\ell^0 = (\alpha_0, \alpha_1, \dots, \alpha_{n-1})$ and $\ell^1 = (\gamma, \alpha_1, \dots, \alpha_{n-1})$. Then $P(\ell) = 0$ since $f_1(\ell^0) = f_1(\alpha_0, \dots, \alpha_{n-1}) < f_0(\gamma, \alpha_1, \dots, \alpha_{n-1}) = f_0(\ell^1)$. This shows that C_1 is homogeneous for P taking value 0. Since f_0 and f_1 have type n , $I_{f_0}^1$ and $I_{f_1}^1$ are increasing functions. For any $\alpha \in \omega_1$, there is some $\gamma \in C_1$ so that $f_2(\alpha) < I_{f_0}^1(\gamma) = I_{f_1}^1(\gamma)$. Let $h : \omega_1 \rightarrow C_1$ be defined by $h(\alpha)$ is the least such $\gamma \in C_1$. Let $C_3 = \{\alpha \in C_1 : \text{enum}_{C_1}(\alpha) = \alpha \wedge (\forall \alpha' < \alpha)(h(\alpha') < \alpha)\}$ which is a club subset of C_1 .

For notational simplicity, let $\mathfrak{e} = \text{enum}_{[C_3]_*^1}$. Define $F : U_n \rightarrow \omega_1$ by $F(\alpha_{n-1}, 0, \dots, \alpha_0, i) = f_i(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1}))$ for all $(\alpha_0, \dots, \alpha_{n-1}) \in [\omega_1]^n$ and $i \in 2$ and $F(\alpha, 1) = f_2(\mathfrak{e}(\alpha))$ for all $\alpha \in \omega_1 \setminus (n-1)$.

First, one will show that F is an order preserving map from $\mathcal{U}_n$ into the usual ordering on ω_1 . Suppose $x, y \in U_n$ and $x \ll_n y$. One seeks to show $F(x) < F(y)$.

- Suppose $x = (\alpha, 1)$ and $y = (\beta, 1)$ with $\alpha < \beta$:

$$F(x) = F(\alpha, 1) = f_2(\mathfrak{e}(\alpha)) < f_2(\mathfrak{e}(\beta)) = F(\beta, 1) = F(y)$$

since f_2 has type 1.

- Suppose $x = (\alpha, 1)$ and $y = (\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, i)$ for some $i < 2$ and $\beta_0 < \dots < \beta_{n-1}$ with $\alpha < \beta_{n-1}$: Note that

$$\begin{aligned} F(x) &= F(\alpha, 1) = f_2(\mathfrak{e}(\alpha)) < I_{f_i}^1(h(\mathfrak{e}(\alpha))) \\ &< f_i(\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1})) = F(\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, i) = F(y). \end{aligned}$$

The first inequality comes from the definition of h , the second inequality comes from $h(\mathfrak{e}(\alpha)) < \mathfrak{e}(\beta_{n-1})$ by the definition of $\mathfrak{e}(\beta_{n-1}) \in C_3$, and the last inequality comes from f_i having type n .

- Suppose $x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i)$ and $y = (\beta, 1)$ for some $i < 2$ and $\alpha_0 < \dots < \alpha_{n-1}$ with $\alpha_{n-1} \leq \beta$:

$$\begin{aligned} F(x) &= F(\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i) = f_i(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) \leq I_{f_i}^1(\mathfrak{e}(\alpha_{n-1})) \\ &< f_2(\mathfrak{e}(\alpha_{n-1})) \leq f_2(\mathfrak{e}(\beta)) = F(\beta, 1) = F(y). \end{aligned}$$

The first inequality comes from the definition of $I_{f_i}^1$, the second inequality comes from property (3) of the club C_0 , and the third inequality comes from the fact that f_2 has type 1.

- Suppose $x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i)$ and $y = (\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, j)$ for some $i, j \in 2$, $\alpha_0 < \dots < \alpha_{n-1}$, $\beta_0 < \dots < \beta_{n-1}$, and there is some $k > 0$ so that $\alpha_k < \beta_k$ and for all $k < k' < n$, $\alpha_{k'} = \beta_{k'}$:

$$\begin{aligned} F(x) &= F(\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i) = f_i(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) \leq I_{f_i}^{n-k}(\mathfrak{e}(\alpha_k), \dots, \mathfrak{e}(\alpha_{n-1})) \\ &= I_{f_j}^{n-k}(\mathfrak{e}(\alpha_k), \dots, \mathfrak{e}(\alpha_{n-1})) < I_{f_j}^{n-k}(\text{next}_{C_1}(\mathfrak{e}(\alpha_k)), \mathfrak{e}(\alpha_{k+1}), \dots, \mathfrak{e}(\alpha_{n-1})) \\ &= I_{f_j}^{n-k}(\text{next}_{C_1}(\mathfrak{e}(\alpha_k)), \mathfrak{e}(\beta_{k+1}), \dots, \mathfrak{e}(\beta_{n-1})) < f_j(\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1})) \\ &= F(\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, j) = F(y). \end{aligned}$$

- Suppose $x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, 0)$ and $y = (\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, 1)$ for $\alpha_0 < \dots < \alpha_{n-1}$ and $\beta_0 < \dots < \beta_{n-1}$ such that for all $0 < k < n$, $\alpha_k = \beta_k$ and $\alpha_0 \leq \beta_0$:

$$\begin{aligned} F(x) &= F(\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, 0) = f_0(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) < f_1(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) \\ &\leq f_1(\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1})) = F(\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, 1) = F(y). \end{aligned}$$

- Suppose $x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, 1)$ and $y = (\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, 0)$ for $\alpha_0 < \dots < \alpha_{n-1}$ and $\beta_0 < \dots < \beta_{n-1}$ such that for all $0 < k < n$, $\alpha_k = \beta_k$ and $\alpha_0 < \beta_0$: Let $\ell = (\mathfrak{e}(\alpha_0), \mathfrak{e}(\beta_0), \mathfrak{e}(\beta_1), \dots, \mathfrak{e}(\beta_{n-1})) = (\mathfrak{e}(\alpha_0), \mathfrak{e}(\beta_0), \mathfrak{e}(\alpha_1), \dots, \mathfrak{e}(\alpha_{n-1}))$. In the notation above, $\ell^0 = (\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1}))$ and $\ell^1 = (\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1}))$. $P(\ell) = 0$ implies that $f_1(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) = f_1(\ell^0) < f_0(\ell^1) = f_0(\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1}))$. So we have the following.

$$\begin{aligned} F(x) &= F(\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, 1) = f_1(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) \\ &< f_0(\mathfrak{e}(\beta_0), \dots, \mathfrak{e}(\beta_{n-1})) = F(\beta_{n-1}, 0, \beta_{n-2}, \dots, \beta_0, 0) = F(y) \end{aligned}$$

This shows that F is order preserving.

Next one will show that F is discontinuous everywhere. Suppose $x \in U_n$ has limit rank in $\ll_n$.

- Suppose $x = (\alpha, 1)$ for some $\alpha \in \omega_1$. Then $\sup(F \upharpoonright x) = \sup\{F(\alpha, 0, \alpha_{n-2}, \dots, \alpha_0, i) : i \in 2 \wedge \alpha_0 < \dots < \alpha_{n-2} < \alpha\} = \sup\{F(\alpha, 0, \alpha_{n-2}, \dots, \alpha_0, 0) : \alpha_0 < \dots < \alpha_{n-2} < \alpha\} = \sup\{f_0(\mathfrak{e}(\alpha_0), \mathfrak{e}(\alpha_1), \dots, \mathfrak{e}(\alpha_{n-2}), \mathfrak{e}(\alpha)) : \alpha_0 < \alpha_1 < \dots < \alpha_{n-2} < \alpha\} = I_{f_0}^1(\mathfrak{e}(\alpha)) < f_2(\mathfrak{e}(\alpha)) = F(x)$.
- Suppose $x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, 0)$ and has limit rank. Then $\sup(F \upharpoonright x) = \sup\{f_0(\mathfrak{e} \circ \ell) : \ell \sqsubset_n (\alpha_0, \dots, \alpha_{n-1})\} < f_0(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) = F(x)$ using the discontinuity of f_0 .

This shows that F is discontinuous everywhere.

Let $G_0, G_1 : [\omega_1]^n \times \omega \rightarrow \omega_1$ witness that f_0 and f_1 have uniform cofinality ω . Let $G_2 : \omega_1 \times \omega \rightarrow \omega_1$ witness that f_2 has uniform cofinality ω . Define $\mathcal{F} : U_n \times \omega \rightarrow \omega_1$ be defined as follows.

$$\mathcal{F}(x, k) = \begin{cases} G_2(\mathfrak{e}(\alpha), k) & x = (\alpha, 1) \\ G_i((\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})), k) & x = (\alpha_{n-1}, 0, \alpha_{n-2}, \dots, \alpha_0, i) \wedge i \in 2 \end{cases}$$

$\mathcal{F}$ witness that F has uniform cofinality ω . It has been shown that F is a function of the correct type.

It is clear from the construction that $F^0[[\omega_1]^n] \subseteq f_0[[\omega_1]^n]$, $F^1[[\omega_1]^n] \subseteq f_1[[\omega_1]^n]$, and $F^2[\omega_1] \subseteq f_2[\omega_1]$. Let $C_4 = \{\alpha \in C_3 : \text{enum}_{C_3}(\alpha) = \alpha\}$ which is a club subset of C_3 . For all $\alpha \in [C_4]_*^1$, $\alpha = \text{enum}_{C_3}(\alpha) = \text{enum}_{[C_3]_*^1}(\alpha) = \mathfrak{e}(\alpha)$. For all $(\alpha_0, \dots, \alpha_{n-1}) \in [C_4]^n$ and $i \in 2$, $F^i(\alpha_0, \dots, \alpha_{n-1}) = f_i(\mathfrak{e}(\alpha_0), \dots, \mathfrak{e}(\alpha_{n-1})) = f_i(\alpha_0, \dots, \alpha_{n-1})$. For all $\alpha \in [C_4]_*^1$, $F^2(\alpha) = f_2(\mathfrak{e}(\alpha)) = f_2(\alpha)$. This shows that $[F^1]_{\mu_{\omega_1}^n} = [f_0]_{\mu_{\omega_1}^n}$, $[F^1]_{\mu_{\omega_1}^n} = [f_1]_{\mu_{\omega_1}^n}$, and $[F^2]_{\mu_{\omega_1}^1} = [f_2]_{\mu_{\omega_1}^1}$. $\square$

Theorem 3.10. Assume $\omega_1 \rightarrow_* (\omega_1)_{2^1}^{\omega_1}$ and $\star$. The boldface GCH holds below ω_ω .

Proof. The boldface GCH at ω_n for all $n < \omega$ will be shown by induction. For $n = 0$, the boldface GCH at ω has already been shown by Fact 3.4. For $n = 1$, the boldface GCH at ω_1 has already been shown by Theorem 3.7. Suppose $n > 1$ and the boldface GCH has been shown at ω_{n-1} . Suppose for the sake of contradiction, the boldface GCH at ω_n fails. Let $\langle A_\eta : \eta < \omega_{n+1} \rangle$ be an injection of ω_{n+1} into $\mathcal{P}(\omega_n)$. Recall $\mathcal{U}_n = (U_n, \ll_n)$ from Definition 3.8. By Fact 2.30, $\text{cof}(\omega_{n+1}) = \omega_2$ for all $1 \leq n < \omega$. Fix $\rho : \omega_2 \rightarrow \omega_n$ be an increasing cofinal map. Define $P : [\omega_1]^{\mathcal{U}_n} \rightarrow 2$ by $P(F) = 0$ if and only if $\min(A_{[F^0]_{\mu_{\omega_1}^n}} \Delta A_{[F^1]_{\mu_{\omega_1}^n}}) < \rho([F^2]_{\mu_{\omega_1}^1})$, where Δ refers to the symmetric difference. (Note that here one is using the fact that $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$ and $j_{\mu_{\omega_1}^n}(\omega_1) = \omega_{n+1}$ established in Fact 2.30.) Since $\text{ot}(\mathcal{U}_n) = \omega_1$, $\omega_1 \rightarrow_* (\omega_1)_{2^1}^{\omega_1}$ implies there is a club $C \subseteq \omega_1$ which is homogeneous for P . Let $f_0, f_1 : [\omega_1]^n \rightarrow C$ be any two function of type n with $[f_0]_{\mu_{\omega_1}^n} < [f_1]_{\mu_{\omega_1}^n}$ and $[I_{f_0}^{n-1}]_{\mu_{\omega_1}^{n-1}} = [I_{f_1}^{n-1}]_{\mu_{\omega_1}^{n-1}}$. Since $\langle A_\eta : \eta < \omega_{n+1} \rangle$ is an injection, $A_{[f_0]_{\mu_{\omega_1}^n}} \neq A_{[f_1]_{\mu_{\omega_1}^n}}$ and thus $A_{[f_0]_{\mu_{\omega_1}^n}} \Delta A_{[f_1]_{\mu_{\omega_1}^n}} \neq \emptyset$. Let $f_2 : \omega_1 \rightarrow C$ be any function of type 1 so that $\rho([f_2]_{\mu_{\omega_1}^1}) > \min(A_{[f_0]_{\mu_{\omega_1}^n}} \Delta A_{[f_1]_{\mu_{\omega_1}^n}})$ and $[f_2]_{\mu_{\omega_1}^1} > [I_{f_0}^1]_{\mu_{\omega_1}^1} = [I_{f_1}^1]_{\mu_{\omega_1}^1}$. By Lemma 3.9, there is an $F \in [\omega_1]^{\mathcal{U}_n}$ so that $[F^0]_{\mu_{\omega_1}^n} = [f_0]_{\mu_{\omega_1}^n}$, $[F^1]_{\mu_{\omega_1}^n} = [f_1]_{\mu_{\omega_1}^n}$, $[F^2]_{\mu_{\omega_1}^1} = [f_2]_{\mu_{\omega_1}^1}$, $F^0[[\omega_1]^n] \subseteq f_0[[\omega_1]^n] \subseteq C$, $F^1[[\omega_1]^n] \subseteq f_1[[\omega_1]^n] \subseteq C$, and $F^2[\omega_1] \subseteq f_2[\omega_1] \subseteq C$. Thus $F \in [C]_*^{\mathcal{U}_n}$. Then $\rho([F^2]_{\mu_{\omega_1}^1}) > \min(A_{[F^0]_{\mu_{\omega_1}^n}} \Delta A_{[F^1]_{\mu_{\omega_1}^n}})$ implies that $P(F) = 0$. This shows that C must be homogeneous for P taking value 0. Pick any $\phi : V_n \rightarrow C$ of the correct type from $\mathcal{V}_n$ into $(C, <)$ (where recall V_n is defined in Definition 2.36). By Fact 2.39, there is a $\chi < \omega_n$ so that $E_\chi = \{\eta \in \mathfrak{B}_{n+1}^C : I_\eta^{n-1} = \chi\}$ has cardinality ω_n . Let $g : \omega_1 \rightarrow C$ be any function of type 1 so that $[g]_{\mu_{\omega_1}^1} > \mathcal{I}_\chi^1$. Let $\epsilon = [g]_{\mu_{\omega_1}^1}$. Suppose $\eta_0, \eta_1 \in E_\chi$ and $\eta_0 \neq \eta_1$. Without loss of generality, suppose $\eta_0 < \eta_1$. Let $f_0, f_1 : [\omega_1]^n \rightarrow C$ be functions of type n so that $[f_0]_{\mu_{\omega_1}^n} = \eta_0$ and $[f_1]_{\mu_{\omega_1}^n} = \eta_1$. By definition of $\eta_0, \eta_1 \in E_\chi$, $[I_{f_0}^{n-1}]_{\mu_{\omega_1}^{n-1}} = \chi$. Note that $\mathcal{I}_\chi^1 = \mathcal{I}_{\eta_i}^1$ for both $i \in 2$ by Fact 2.40. Thus $[I_{f_i}^1]_{\mu_{\omega_1}^1} = \mathcal{I}_{\eta_i}^1 = \mathcal{I}_\chi^1 < [g]_{\mu_{\omega_1}^1} = \epsilon$ for both $i \in 2$. By Lemma 3.9, there is an $F \in [C]_*^{\mathcal{U}_n}$ so that $[F^0]_{\mu_{\omega_1}^n} = [f_0]_{\mu_{\omega_1}^n} = \eta_0$, $[F^1]_{\mu_{\omega_1}^n} = [f_1]_{\mu_{\omega_1}^n} = \eta_1$, and $[F^2]_{\mu_{\omega_1}^1} = [g]_{\mu_{\omega_1}^1} = \epsilon$. By $P(F) = 0$, one has that $\min(A_{\eta_0} \Delta A_{\eta_1}) < \rho(\epsilon)$. This shows that the function $\Upsilon : E_\chi \rightarrow \mathcal{P}(\rho(\epsilon))$ defined by $\Upsilon(\eta) = A_\eta \cap \rho(\epsilon)$ is an injection. Since $|E_\chi| = \omega_n$ and $|\mathcal{P}(\rho(\epsilon))| = |\mathcal{P}(\omega_{n-1})|$ because $\rho(\epsilon) < \omega_n$, Υ induces an injection of ω_n into $\mathcal{P}(\omega_{n-1})$ which violates the inductive assumption that the boldface GCH holds at ω_{n-1} . $\square$

Under AD, $\omega_{\omega+1} = \delta_3^1$ and there is a $\omega_{\omega+1}$ -complete nonprincipal ultrafilter on $\omega_{\omega+1}$. Thus the boldface GCH holds at ω_ω by Fact 3.2. The combinatorial methods used here can be generalized with Jackson's theory of descriptions ([15]) for the projective ordinals to show that the boldface GCH holds below the supremum of the projective ordinals, $\sup\{\delta_n^1 : n \in \omega\}$, assuming AD. Jackson's theory can go slightly beyond the projective ordinals but not all the way through Θ . The inner model theoretic techniques of Steel and Woodin are the only known methods to prove the boldface GCH below Θ under AD^+ .

The methods used here have been expanded to address other combinatorial questions concerning important subsets of $\mathcal{P}(\omega_\omega)$ for which there are currently no other known methods. A brief survey of these new results will be provided below. This paper is a good introduction to these other results which require a substantially more challenging analysis of sequences of ordinals below ω_ω . Here, the Kunen functions were used merely

to establish the basic theory of Martin’s analysis of the ultrapower below ω_ω . In these new results, the Kunen functions will appear very prominently and will be used to make selections of representatives for ω_1 -many ordinals below ω_ω . In each of these new results, an appropriate “sliding lemma” captures the core combinatorics of the proof. These new “sliding lemmas” will be much more elaborate. The relatively simple “sliding lemma” of Lemma 3.9 is a good warm-up.

Let X and Y be sets. X is said to be Y -regular if and only if for every function $\Phi : X \rightarrow Y$, there is a $y \in Y$ so that $|\Phi^{-1}[\{y\}]| = |X|$. In [10], this concept is used to form a theory of cofinality for nonwellorderable sets which extends the usual theory of cofinality for ordinals.

Theorem 3.11. ([10]) *Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$.*

- $\omega(\omega_n)$, ${}^{\omega_1}(\omega_n)$, and ${}^{<\omega_2}(\omega_n)$ are $\mathcal{P}(\omega_1)$ -regular for all $2 \leq n < \omega$.
- $\omega(\omega_\omega)$ is $\mathcal{P}(\omega_1)$ -regular.

A set X is prime if and only if for all sets A and B , if $|X| \leq |A \times B|$, then $|X| \leq |A|$ or $|X| \leq |B|$.

Theorem 3.12. ([9]) *Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$, $j_{\mu_{\omega_1}^1}(\omega) = \omega_2$, and $\text{AC}_\omega^{\mathcal{P}(\omega_1)}$.*

- For all $n \in \omega$, $\omega(\omega_n)$ is prime.
- $\omega(\omega_\omega)$ is prime.
- For all $n < \omega$, $\omega(\omega_n)$ is $\mathcal{P}(\omega_n)$ -regular.

If $X \subseteq \omega_\omega$, then let $\text{Bl}_{\omega_\omega}(<\omega_1, X)$ be the set of countable subsets of X bounded below ω_ω . ω_ω is said to be a Magidor cardinal if and only if for every function $\Phi : \text{Bl}_{\omega_\omega}(<\omega_1, \omega_\omega) \rightarrow \omega_\omega$, there is an $X \subseteq \omega_\omega$ so that $|X| = |\omega_\omega|$ and $\Phi[\text{Bl}_{\omega_\omega}(<\omega_1, X)] \neq \omega_\omega$. Ben-Neria and Garti ([1]) asked whether ω_ω is a Magidor cardinal under AD. The following gives a positive answer:

Theorem 3.13. ([6]) *Assume $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$ and $j_{\mu_{\omega_1}^1}(\omega_1) = \omega_2$. ω_ω is a Magidor cardinal.*

REFERENCES

1. Omer Ben-Neria and Shimon Garti, *Jonsson and Magidor filters*, arXiv preprint arXiv:2503.13744 (2025).
2. William Chan, *Basis for uncountable linear orders*, In preparation.
3. ———, *Cardinality of the set of bounded subsets of a cardinal*, In preparation.
4. ———, *Definable combinatorics of strong partition cardinals*, In preparation.
5. William Chan, *An introduction to combinatorics of determinacy*, Trends in Set Theory, Contemp. Math., vol. 752, Amer. Math. Soc., Providence, RI, 2020, pp. 21–75. MR 4132099
6. ———, *Coding and antcoding of a cardinal by bounded subsets of the cardinal*, J. Lond. Math. Soc. (2) **113** (2026), no. 3, Paper No. e70501, 51. MR 5042535
7. William Chan and Stephen Jackson, *Definable combinatorics at the first uncountable cardinal*, Trans. Amer. Math. Soc. **374** (2021), no. 3, 2035–2056. MR 4216731
8. ———, *Applications of infinity-Borel codes to definability and definable cardinals*, Fund. Math. **265** (2024), no. 3, 215–258. MR 4771874
9. William Chan, Stephen Jackson, and Nam Trang, *Primeness property for cardinalities*, In preparation.
10. ———, *A search for the cofinality of the power set of the first uncountable cardinal*, In preparation.
11. ———, *The size of the class of countable sequences of ordinals*, Trans. Amer. Math. Soc. **375** (2022), no. 3, 1725–1743. MR 4378077
12. ———, *More definable combinatorics around the first and second uncountable cardinals*, J. Math. Log. **23** (2023), no. 3, Paper No. 2250029, 31. MR 4603918
13. ———, *Almost disjoint families under determinacy*, Adv. Math. **437** (2024), Paper No. 109410, 34. MR 4669334
14. ———, *Almost Everywhere Behavior of Functions According to Partition Measures*, Forum Math. Sigma **12** (2024), Paper No. e16. MR 4696011
15. Steve Jackson, *Structural consequences of AD*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1753–1876. MR 2768700
16. Alexander S. Kechris, *AD and projective ordinals*, Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77), Lecture Notes in Math., vol. 689, Springer, Berlin, 1978, pp. 91–132. MR 526915
17. ———, *The axiom of determinacy implies dependent choices in $L(\mathbb{R})$* , J. Symbolic Logic **49** (1984), no. 1, 161–173. MR 736611
18. Eugene M. Kleinberg, *Infinitary combinatorics and the axiom of determinateness*, Lecture Notes in Mathematics, Vol. 612, Springer-Verlag, Berlin-New York, 1977. MR 0479903
19. Peter Koellner and W. Hugh Woodin, *Large cardinals from determinacy*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1951–2119. MR 2768702

20. William Mitchell, *How weak is a closed unbounded ultrafilter?*, Logic Colloquium '80 (Prague, 1980), Studies in Logic and the Foundations of Mathematics, vol. 108, North-Holland, Amsterdam-New York, 1982, pp. 209–230. MR 673794
21. John R. Steel, $\text{HOD}^{L(\mathbb{R})}$ is a core model below Θ , Bull. Symbolic Logic **1** (1995), no. 1, 75–84. MR 1324625
22. ———, *An outline of inner model theory*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1595–1684. MR 2768698
23. ———, *Ordinal definability in models of determinacy. Introduction to Part V*, Ordinal definability and recursion theory: The Cabal Seminar. Vol. III, Lect. Notes Log., vol. 43, Assoc. Symbol. Logic, Ithaca, NY, 2016, pp. 3–48. MR 3469165
24. John R. Steel and W. Hugh Woodin, **HOD** as a core model, Ordinal definability and recursion theory: The Cabal Seminar. Vol. III, Lect. Notes Log., vol. 43, Assoc. Symbol. Logic, Ithaca, NY, 2016, pp. 257–345. MR 3469173

INSTITUTE FOR DISCRETE MATHEMATICS AND GEOMETRY, VIENNA UNIVERSITY OF TECHNOLOGY, VIENNA, AUSTRIA
Email address: `William.Chan@tuwien.ac.at`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203
Email address: `Stephen.Jackson@unt.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203
Email address: `Nam.Trang@unt.edu`