

INTERVAL ENFORCEABLE PROPERTIES OF FINITE GROUPS

WILLIAM DEMEO

ABSTRACT. We propose a classification of group properties according to whether they can be deduced from the assumption that a group’s subgroup lattice contains an interval isomorphic to some lattice. Suppose \mathcal{P} is a group property and suppose there exists a lattice L such that if G is a group with L isomorphic to an interval $[[H, G]]$ in $\text{Sub}(G)$, with H core-free, then G has property \mathcal{P} . We call such \mathcal{P} *core-free interval enforceable*. Among other things we show that if both a property and its negation could be proved core-free interval enforceable, this would solve an important problem in universal algebra.

1. INTRODUCTION

The study of subgroup lattices has a long history, starting with Richard Dedekind [8] and Ada Rottlaender [26], and later a number of important contributions by Reinhold Baer, Øystein Ore, Michio Suzuki, Roland Schmidt, and many others (see Schmidt [27]). Much of this work focuses on the problem of inferring properties of a group G based on the structure of its lattice of subgroups, $\text{Sub}(G)$, or, conversely, inferring lattice theoretical properties of $\text{Sub}(G)$ from properties of G .

Historically, less attention was paid to the local structure of the subgroup lattice of a finite group, perhaps because it seemed that very little about G could be inferred from knowledge of, say, an *upper interval* $[[H, G]] = \{K \mid H \leq K \leq G\}$ in the subgroup lattice of G . Recently, however, this topic has attracted more attention (e.g., [2, 3, 5, 7, 16, 18, 20, 21]), mostly owing to its connection to the most important open problem in universal algebra, the *Finite Lattice Representation Problem* (FLRP). This is the problem of characterizing the lattices that are (isomorphic to) congruence lattices of finite algebras (see [6, 10, 21, 22]). There is a remarkable theorem relating this problem to intervals in subgroup lattices of finite groups.

Theorem 1.1 (Pálffy and Pudlák [23]). *The following statements are equivalent:*

- (A) *Every finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (B) *Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

If these statements are true (resp., false), then we say the FLRP has a positive (resp., negative) answer. Thus, if we can find a finite lattice L for which it can be

2010 *Mathematics Subject Classification.* Primary 20D30; Secondary 06B15, 08A30.

Key words and phrases. subgroup lattice; congruence lattice; group properties.

The author wishes to thank the anonymous referee for many excellent suggestions that greatly improved the presentation of the paper.

proved that there is no finite group G with $L \cong \llbracket H, G \rrbracket$ for some $H < G$, then the FLRP has a negative answer.

In this paper we propose a new classification of group properties according to whether or not they can be deduced from the assumption that $\text{Sub}(G)$ has an upper interval isomorphic to some finite lattice. We believe that discovering which group properties can (or cannot) be connected in this way to the local structure of a subgroup lattice is itself a worthwhile endeavor, but we will also describe how such a classification could be used to solve the FLRP.

Suppose \mathcal{P} is a *group theoretical property*¹ and suppose there exists a finite lattice L such that if G is a finite group with $L \cong \llbracket H, G \rrbracket$ for some $H \leq G$, then G has property \mathcal{P} . We call such a property \mathcal{P} *interval enforceable (IE) by L* . An *interval enforceable class of groups* is a class of all groups having property \mathcal{P} , for some interval enforceable property \mathcal{P} .

Although it depends on the lattice L , generally speaking it is difficult to deduce very much about a group G from the assumption that an upper interval in $\text{Sub}(G)$ is isomorphic to L . It gets easier easier if, in addition to the hypothesis $L \cong \llbracket H, G \rrbracket$, we assume that the subgroup H is *core-free* in G ; that is, H contains no nontrivial normal subgroup of G . Properties of G that can be deduced from these assumptions are what we call *core-free interval enforceable (cf-IE)*.

Extending this idea, we consider finite collections \mathcal{L} of finite lattices and ask what can be proved about a group G if one assumes that each $L_i \in \mathcal{L}$ is isomorphic to an upper interval $\llbracket H_i, G \rrbracket \leq \text{Sub}(G)$, with each H_i core-free in G . Clearly, if $\text{Sub}(G)$ has such upper intervals, and if corresponding to each $L_i \in \mathcal{L}$ there is a property \mathcal{P}_i that is cf-IE by L_i , then G must have all of the properties \mathcal{P}_i . A related question is the following: Given a set \mathcal{P} of cf-IE properties, is the conjunction $\bigwedge \mathcal{P}$ cf-IE? Corollary 3.11 answers this question affirmatively.

In this paper, we will identify some group properties that are cf-IE, and others that are not. We will see that the cf-IE properties that we have found thus far are negations of common group properties (for example, “not solvable,” “not almost simple,” “not alternating,” “not symmetric”). Moreover, we prove that in these special cases the corresponding group properties (“solvable,” “almost simple,” “alternating,” “symmetric”) that are not cf-IE. This and other considerations suggest that a group property and its negation cannot both be cf-IE. As yet, we are unable to prove this. A related question is whether we should expect that, for every group property \mathcal{P} , either \mathcal{P} is cf-IE or $\neg\mathcal{P}$ is cf-IE.

One of our main results (Theorem 3.9) connects the foregoing ideas with the FLRP, as follows:

Statement (B) of Theorem 1.1 is equivalent to each of the following statements:

- (C) *For every finite lattice L , for every finite collection $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ of cf-IE classes of groups, there exists a finite group $G \in \bigcap_{i=1}^n \mathfrak{X}_i$ such that $L \cong \llbracket H, G \rrbracket$ for some subgroup H that is core-free in G .*
- (D) *For every finite collection \mathcal{L} of finite lattices, there exists a finite group G such that for each $L \in \mathcal{L}$ we have $L \cong \llbracket H, G \rrbracket$ for some subgroup H that is core-free in G .*

¹This and other italicized terms in the introduction will be defined more formally in Section 2.

In fact, the arguments proving the equivalence of these statements are easily combined to show that the following is also equivalent to statement (B):

- (E) *For every finite collection \mathcal{L} of finite lattices, for every finite collection $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ of cf-IE classes of groups, there exists a finite group $G \in \bigcap_{i=1}^n \mathfrak{X}_i$ such that each $L_i \in \mathcal{L}$ is isomorphic to an upper interval $\llbracket H_i, G \rrbracket$ with H core-free in G .*

Core-free interval enforceable properties are intimately related to permutation representations of groups. If H is a core-free subgroup of G , then G has a faithful permutation representation $\varphi : G \hookrightarrow \text{Sym}(G/H)$. Let $\langle G/H, \varphi(G) \rangle$ denote the algebra comprised of the right cosets G/H acted upon by right multiplication by elements of G ; that is, $\varphi(g) : Hx \mapsto Hxg$. It is well known that the congruence lattice of this algebra (i.e., the lattice of systems of imprimitivity) is isomorphic to the interval $\llbracket H, G \rrbracket$ in the subgroup lattice of G .² This puts statement (E) into perspective. If the FLRP has a positive answer, then no matter what we take as our finite collection \mathcal{L} —for example, we might take \mathcal{L} to be *all* finite lattices with at most N elements for some large $N < \omega$ —we can always find a *single* finite group G such that every lattice in \mathcal{L} is isomorphic to the interval in $\text{Sub}(G)$ above a core-free subgroup. As a result, this group G must have so many faithful representations $G \hookrightarrow \text{Sym}(G/H_i)$ with systems of imprimitivity isomorphic to L_i , one such representation for each distinct $L_i \in \mathcal{L}$. Moreover, the group G having this property can be chosen from the class $\bigcap_{i=1}^n \mathfrak{X}_i$, where $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ is an arbitrary collection of cf-IE classes of groups.

2. NOTATION AND DEFINITIONS

In this paper, *all groups and lattices are finite*. We use \mathfrak{G} to denote the class of all finite groups. Given a group G , we denote the set of subgroups of G by $\text{Sub}(G)$. The algebra $\langle \text{Sub}(G), \wedge, \vee \rangle$ is a lattice where the \wedge (“meet”) and \vee (“join”) operations are defined for all H and K in $\text{Sub}(G)$ by $H \wedge K = H \cap K$ and $H \vee K = \langle H, K \rangle =$ the smallest subgroup of G containing both H and K . We will refer to the set $\text{Sub}(G)$ as a lattice, without explicitly mentioning the \wedge and \vee operations.

By $H \leq G$ (resp., $H < G$) we mean H is a subgroup (resp., proper subgroup) of G . For $H \leq G$, the *core of H in G* , denoted by $\text{core}_G(H)$, is the largest normal subgroup of G contained in H . If $\text{core}_G(H) = 1$, then we say that H is *core-free in G* . For $H \leq G$, by the *interval $\llbracket H, G \rrbracket$* we mean the set $\{K \mid H \leq K \leq G\}$, which is a sublattice of $\text{Sub}(G)$. That is, $\llbracket H, G \rrbracket$ is the lattice of those subgroups of G that contain H . With this notation, $\text{Sub}(G) = \llbracket 1, G \rrbracket$. When viewing $\llbracket H, G \rrbracket$ as a sublattice of $\text{Sub}(G)$, we sometimes refer to it as an *upper interval*. Given an abstract lattice L , if there is no mention of specific groups H and G , then the expression $L \cong \llbracket H, G \rrbracket$ means “there exist (finite) groups $H \leq G$ such that L is isomorphic to the interval $\{K \mid H \leq K \leq G\}$ in the subgroup lattice of G .”

By a *group theoretical class*, or *class of groups*, we mean a collection \mathfrak{X} of groups that is closed under isomorphism: if $G_0 \in \mathfrak{X}$ and $G_1 \cong G_0$, then $G_1 \in \mathfrak{X}$. A *group theoretical property*, or simply *property of groups*, is a property \mathcal{P} such that if a

²See [19, Lemma 4.20] or [13, Theorem 1.5A].

group G_0 has property \mathcal{P} and $G_1 \cong G_0$, then G_1 has property \mathcal{P} .³ Thus if $\mathfrak{X}_{\mathcal{P}}$ denotes the collection of all groups having the group property \mathcal{P} , then $\mathfrak{X}_{\mathcal{P}}$ is a class of groups, and belonging to a particular class of groups is a group theoretical property. Therefore, we need not distinguish between a property of groups and the class of groups that possess this property. A group in the class \mathfrak{X} is called a \mathfrak{X} -group, and we sometimes write $G \models \mathcal{P}$ to indicate that G has property \mathcal{P} .

If \mathcal{K} is a class of algebras (e.g., a class of groups), then we say that \mathcal{K} is *closed under homomorphic images* and we write $\mathbf{H}(\mathcal{K}) = \mathcal{K}$ provided $\varphi(G) \in \mathcal{K}$ whenever $G \in \mathcal{K}$ and φ is a homomorphism of G . By the first isomorphism theorem for groups, this is equivalent to: $G/N \in \mathcal{K}$ whenever $G \in \mathcal{K}$ and $N \trianglelefteq G$. For algebras, $\mathbf{H}(\mathcal{K}) = \mathcal{K}$ holds if and only if $\mathbf{A}/\theta \in \mathcal{K}$ for all $\mathbf{A} \in \mathcal{K}$ and all $\theta \in \text{Con } \mathbf{A}$, where $\text{Con } \mathbf{A}$ denotes the lattice of congruence relations of \mathbf{A} . Apart from possible notational differences, the foregoing terminology is standard.

We now introduce some new terminology that we find useful.⁴ Let \mathfrak{L} denote the class of all finite lattices, and \mathfrak{G} the class of all finite groups. Let \mathcal{P} be a given group theoretical property and $\mathfrak{X}_{\mathcal{P}}$ the associated class of all groups with property \mathcal{P} . We call \mathcal{P} (and $\mathfrak{X}_{\mathcal{P}}$)

- *interval enforceable* (IE) provided

$$(\exists L \in \mathfrak{L}) (\forall G \in \mathfrak{G}) (L \cong \llbracket H, G \rrbracket \longrightarrow G \in \mathfrak{X}_{\mathcal{P}})$$

- *core-free interval enforceable* (cf-IE) provided

$$(\exists L \in \mathfrak{L}) (\forall G \in \mathfrak{G}) ((L \cong \llbracket H, G \rrbracket \bigwedge \text{core}_G(H) = 1) \longrightarrow G \in \mathfrak{X}_{\mathcal{P}})$$

- *minimal interval enforceable* (min-IE) provided there exists $L \in \mathfrak{L}$ such that if $L \cong \llbracket H, G \rrbracket$ for some group $G \in \mathfrak{G}$ of minimal order (with respect to $L \cong \llbracket H, G \rrbracket$), then $G \in \mathfrak{X}_{\mathcal{P}}$.

In this paper we will have little to say about min-IE properties. Nonetheless, we include this class in our list of new definitions because properties of this type arise often (see, e.g., [18]), and a primary aim of this paper is to formalize various notions of interval enforceability that we believe are useful in applications.

3. RESULTS

Clearly, if \mathcal{P} is an interval enforceable property, then it is also core-free interval enforceable. There is an easy sufficient condition under which the converse holds. Suppose \mathcal{P} is a group property, let $\mathfrak{X}_{\mathcal{P}}$ denote the class of all groups with property \mathcal{P} , and let $\mathfrak{X}_{\mathcal{P}}^c$ denote the class of all groups that do not have property \mathcal{P} .

Lemma 3.1. *Suppose \mathcal{P} is a core-free interval enforceable property. If $\mathbf{H}(\mathfrak{X}_{\mathcal{P}}^c) = \mathfrak{X}_{\mathcal{P}}^c$, then \mathcal{P} is an interval enforceable property.*

Proof. Since \mathcal{P} is cf-IE there is a lattice L such that

$$(3.1) \quad (L \cong \llbracket H, G \rrbracket \bigwedge \text{core}_G(H) = 1) \longrightarrow G \in \mathfrak{X}_{\mathcal{P}}.$$

³It seems there is no single standard definition of *group theoretical class*. While some authors (e.g., [14], [4]) use the same definition we use here, others (e.g. [24], [25]) require that every group theoretical class contains the one element group. In the sequel we consider negations of group properties, and we would like these to qualify as group properties. Therefore, we don't require that every group theoretical class contains the one element group.

⁴The author thanks Bjørn Kjos-Hanssen and David Ross for suggesting improvements to the wording of these definitions.

Under the assumption $\mathbf{H}(\mathfrak{X}_{\mathcal{P}}^c) = \mathfrak{X}_{\mathcal{P}}^c$ we prove

$$(3.2) \quad L \cong \llbracket H, G \rrbracket \longrightarrow G \in \mathfrak{X}_{\mathcal{P}}.$$

If (3.2) fails, then there is a group $G \in \mathfrak{X}_{\mathcal{P}}^c$ with $L \cong \llbracket H, G \rrbracket$. Let $N = \text{core}_G(H)$. Then $L \cong \llbracket H/N, G/N \rrbracket$ and H/N is core-free in G/N so, by hypothesis (3.1), $G/N \in \mathfrak{X}_{\mathcal{P}}$. But $G/N \in \mathfrak{X}_{\mathcal{P}}^c$, since $\mathfrak{X}_{\mathcal{P}}^c$ is closed under homomorphic images. \square

In [21], Péter Pálffy gives an example of a lattice that cannot occur as an upper interval in the subgroup lattice finite solvable group. (We give other examples in §3.3 and §??.) In his Ph.D. thesis [5], Alberto Basile proves that if G is an alternating or symmetric group, then there are certain lattices that cannot occur as upper intervals in $\text{Sub}(G)$. Another class of lattices with this property is described by Aschbacher and Shareshian in [2]. Thus, two classes of groups that are known to be at least cf-IE are the following:

- $\mathfrak{X}_0 = \mathfrak{S}^c = \text{nonsolvable finite groups};$
- $\mathfrak{X}_1 = \{G \in \mathfrak{G} \mid (\forall n < \omega) (G \neq A_n \wedge G \neq S_n)\},$

where A_n and S_n denote, respectively, the alternating and symmetric groups on n letters. Note that both classes \mathfrak{X}_0 and \mathfrak{X}_1 satisfy the hypothesis of 3.1. Explicitly, $\mathfrak{X}_0^c = \mathfrak{S}$, the class of solvable groups, is closed under homomorphic images, as is the class \mathfrak{X}_1^c of alternating and symmetric groups. Therefore, by Lemma 3.1, \mathfrak{X}_0 and \mathfrak{X}_1 are IE classes. By contrast, suppose there exists a finite lattice L such that

$$(L \cong \llbracket H, G \rrbracket \bigwedge \text{core}_G(H) = 1) \longrightarrow G \text{ is subdirectly irreducible.}$$

Lemma 3.1 does not apply in this case since the class of subdirectly reducible groups is obviously not closed under homomorphic images.⁵ In Sections 3.3 and ?? below we describe lattices with which we can prove that the following classes are at least cf-IE:

- $\mathfrak{X}_2 = \text{the subdirectly irreducible groups};$
- $\mathfrak{X}_3 = \text{the groups having no nontrivial abelian normal subgroups};$
- $\mathfrak{X}_4 = \{G \in \mathfrak{G} \mid C_G(M) = 1 \text{ for all } 1 \neq M \trianglelefteq G\}.$

We noted above that \mathfrak{X}_2 fails to satisfy the hypothesis of 3.1. The same can be said of \mathfrak{X}_3 and \mathfrak{X}_4 . That is, $\mathbf{H}(\mathfrak{X}_i^c) \neq \mathfrak{X}_i^c$ for $i = 2, 3, 4$. To verify this take $H \in \mathfrak{X}_i$, $K \in \mathfrak{X}_i^c$, and consider $H \times K$. In each case ($i = 2, 3, 4$) we see that $H \times K$ belongs to \mathfrak{X}_i^c , but the homomorphic image $(H \times K)/(1 \times K) \cong H$ does not.

3.1. Negations of interval enforceable properties. The following definition is useful: if a lattice L is isomorphic to an interval in the subgroup lattice of a finite group, then we call L *group representable*. Recall, Theorem 1.1 says that the FLRP has a negative answer if we can find a finite lattice that is not group representable.

Suppose there exists a property \mathcal{P} such that both \mathcal{P} and its negation $\neg\mathcal{P}$ are interval enforceable by the lattices L and L_c , respectively. That is $L \cong \llbracket H, G \rrbracket$ implies $G \in \mathfrak{X}_{\mathcal{P}}$ and $L_c \cong \llbracket H_c, G_c \rrbracket$ implies $G_c \in \mathfrak{X}_{\mathcal{P}}^c$. Then clearly the lattice in Figure 1 could not be group representable. As the next result shows, however, if a group property and its negation are interval enforceable by the lattices L and L_c , then already at least one of these lattices is not group representable.

⁵Recall, for groups *subdirectly irreducible* is equivalent to having a unique minimal normal subgroup. Every algebra, in particular every group G , has a subdirect decomposition into subdirectly irreducibles, say, $G \hookrightarrow G/N_1 \times \cdots \times G/N_n$, so there are always subdirectly irreducible homomorphic images.




FIGURE 1.

Lemma 3.2. *If \mathcal{P} is a group property that is interval enforceable by a group representable lattice, then $\neg\mathcal{P}$ is not interval enforceable by a group representable lattice.*

Proof. Assume the contrary. Then both \mathcal{P} and its negation $\neg\mathcal{P}$ are interval enforceable by group representable lattices L and L_c , respectively. Let G and G_c be groups for which $L \cong \llbracket H, G \rrbracket$ and $L_c \cong \llbracket H_c, G_c \rrbracket$ for some $H \leq G$ and $H_c \leq G_c$. Then the group $G \times G_c$ has upper intervals $L \cong \llbracket H \times G_c, G \times G_c \rrbracket$ and $L_c \cong \llbracket G \times H_c, G \times G_c \rrbracket$. Thus, by the interval enforceability assumptions, the group $G \times G_c$ both is and is not a $\mathfrak{X}_{\mathcal{P}}$ -group. \square

To take a concrete example, insolubility is IE. However, solubility is obviously not IE. For, if $L \cong \llbracket H, G \rrbracket$ then for any nonsolvable group K we have $L \cong \llbracket H \times K, G \times K \rrbracket$, and of course $G \times K$ is nonsolvable. Note that here (and in the proof of Lemma 3.2) the group $H \times K$ at the bottom of the interval is not core-free. So a more interesting question is whether a property and its negation can both be cf-IE. Again, if such a property were found, a lattice of the form in Figure 1 would give a negative answer to the FLRP, though this requires additional justification to address the core-free aspect (see Section 3.3). We suspect the answer is no, as suggested by

Conjecture 3.1. *If \mathcal{P} is core-free interval enforceable by a group representable lattice, then $\neg\mathcal{P}$ is not core-free interval enforceable by a group representable lattice.*

We will confirm a few special cases of the foregoing conjecture—in particular, the cases when \mathcal{P} means “not solvable” or “not almost simple.” Indeed, Lemma 3.4 below implies that the class of solvable groups, and more generally any class of groups that omits certain wreath products, cannot be core-free interval enforceable by a group representable lattice. Before stating and proving this lemma, however, we pause to state a result that gives further support of for the conjecture above.

Proposition 3.3. *The following statements are equivalent...*

Lemma 3.4. *Suppose \mathcal{P} is core-free interval enforceable by a group representable lattice. Then, for any finite nonabelian simple group S , there exists a wreath product group of the form $W = S \wr \bar{U}$ that has property \mathcal{P} .*

Proof. Let L be a group representable lattice such that if $L \cong \llbracket H, G \rrbracket$ and $\text{core}_G(H) = 1$ then $G \models \mathcal{P}$. Since L is group representable, there exists a \mathcal{P} -group G with $L \cong \llbracket H, G \rrbracket$. We apply an idea of Hans Kurzweil (see [17]) twice. Fix a finite nonabelian simple group S . Suppose the index of H in G is $|G : H| = n$. Then the action of G on the cosets of H induces an automorphism of the group S^n by permutation of coordinates. Denote this representation by $\varphi : G \rightarrow \text{Aut}(S^n)$, and

let the image of G be $\varphi(G) = \bar{G} \leq \text{Aut}(S^n)$. The wreath product under this action is the group

$$U := S \wr_{\varphi} G = S^n \rtimes_{\varphi} G = S^n \rtimes \bar{G},$$

with multiplication given by

$$(s_1, \dots, s_n, x)(t_1, \dots, t_n, y) = (s_1 t_{x(1)}, \dots, s_n t_{x(n)}, xy),$$

for $s_i, t_i \in S$ and $x, y \in \bar{G}$. (For the remainder of the proof, we suppress the semidirect product symbol and write, for example, $S^n \bar{G}$ instead of $S^n \rtimes \bar{G}$.)

An illustration of the subgroup lattice of such a wreath product appears in Figure 2. Note that the interval $[[D, S^n]]$, where D denotes the diagonal subgroup of S^n , is isomorphic to $\text{Eq}(n)'$, the dual of the lattice of partitions of an n -element set. The dual lattice L' is an upper interval of $\text{Sub}(U)$, namely, $L' \cong [[D\bar{G}, U]]$.⁶




FIGURE 2. Hasse diagram illustrating some features of the subgroup lattice of the wreath product U .

It is important to note (and we prove below) that if H is core-free in G – equivalently, if $\ker \varphi = 1$ – then the foregoing construction results in the subgroup $D\bar{G}$ being core-free in U . Therefore, by repeating the foregoing procedure, with $H_1 = D\bar{G}$ denoting the (core-free) subgroup of U such that $L' \cong [[H_1, U]]$, we find that $L = L'' \cong [[D_1 \bar{U}, S^m \bar{U}]]$, where $m = |U : H_1|$, and D_1 denotes the diagonal subgroup of S^m . Since $D_1 \bar{U}$ will be core-free in $S^m \bar{U}$ then, it follows by the original hypothesis that $S^m \bar{U} = S \wr \bar{U}$ must have property \mathcal{P} .

To complete the proof, we check that starting with a core-free subgroup $H \leq G$ in the Kurzweil construction just described results in a core-free subgroup $D\bar{G} \leq U$. Let $N = \text{core}_U(D\bar{G})$. Then, for all $w = (d, \dots, d, x) \in N$ and for all $u = (t_1, \dots, t_n, g) \in U$, we have $uwu^{-1} \in N$. Fix $w = (d, \dots, d, x) \in N$. We will choose

⁶These facts, which were proved by Kurzweil in [17], are discussed in greater detail in [10, Section 2.2].

$u \in U$ so that the condition $uwu^{-1} \in N$ implies x acts trivially on $\{1, \dots, n\}$. First note that if $u = (t_1, \dots, t_n, 1)$, then

$$\begin{aligned} uwu^{-1} &= (t_1, \dots, t_n, 1)(d, \dots, d, x)(t_1^{-1}, \dots, t_n^{-1}, 1) \\ &= (t_1 d t_{x(1)}^{-1}, \dots, t_n d t_{x(n)}^{-1}, 1) \in N, \end{aligned}$$

and this implies that $t_1 d t_{x(1)}^{-1} = t_2 d t_{x(2)}^{-1} = \dots = t_n d t_{x(n)}^{-1}$. Suppose by way of contradiction that $x(1) = j \neq 1$. Then, since x is a permutation (hence, one-to-one), $x(k) \neq j$ for each $k \in \{2, 3, \dots, n\}$. Pick one such k other than j . (This is possible since $n = |G : H| > 2$; for otherwise $H \trianglelefteq G$ contradicting $\text{core}_G(H) = 1$.) Since $u \in U$ is arbitrary, we may assume $t_1 = t_k$ and $t_{x(1)} = t_j \neq t_{x(k)}$. But this contradicts $t_1 d t_{x(1)}^{-1} = t_k d t_{x(k)}^{-1}$. Therefore, $x(1) = 1$. The same argument shows that $x(i) = i$ for each $1 \leq i \leq n$, and we see that $w = (d, \dots, d, x) \in N$ implies $x \in \ker \varphi = 1$. This puts N below D , and the only normal subgroup of U that lies below D is the trivial group. \square

By the foregoing result we conclude that a class of groups that does not include wreath products of the form $S \wr G$, where S is an arbitrary finite nonabelian simple group, is not a core-free interval enforceable class. The class of solvable groups is an example.

3.2. Dedekind's rule and its consequences. When A and B are subgroups of a group G , by AB we mean the set $\{ab \mid a \in A, b \in B\}$, and we write $A \vee B$ or $\langle A, B \rangle$ to denote the subgroup of G generated by A and B . Clearly $AB \subseteq \langle A, B \rangle$; equality holds if and only if A and B *permute*, by which we mean $AB = BA$.

We will need the following well known result:⁷

Theorem 3.5 (Dedekind's rule). *Let G be a group and let A, B and C be subgroups of G with $A \leq B$. Then,*

$$(3.3) \quad A(C \cap B) = AC \cap B, \quad \text{and}$$

$$(3.4) \quad (C \cap B)A = CA \cap B.$$

For $A \in \llbracket H, G \rrbracket$ we let $A^{\perp(H, G)}$ denote the set of complements of A in the interval $\llbracket H, G \rrbracket$. That is, $A^{\perp(H, G)} = \{B \in \llbracket H, G \rrbracket \mid A \cap B = H, \langle A, B \rangle = G\}$. Clearly $H^{\perp(H, G)} = \{G\}$ and $G^{\perp(H, G)} = \{H\}$. Recall that an *antichain* of a partially ordered set is a subset of pairwise incomparable elements.

Corollary 3.6. *If $A \in \llbracket H, G \rrbracket$ and if A permutes with each subgroup in $A^{\perp(H, G)}$, then $A^{\perp(H, G)}$ is an antichain or empty.*

Proof. If $A^{\perp(H, G)}$ contains fewer than two elements, the result holds trivially. Let B_1 and B_2 denote two distinct elements of $A^{\perp(H, G)}$. We will prove $\neg(B_1 \leq B_2)$. Indeed, if $B_1 \leq B_2$, then Theorem 3.5 implies

$$B_1 = B_1 H = B_1 (A \cap B_2) = B_1 A \cap B_2 = G \cap B_2 = B_2,$$

which is a contradiction. The penultimate equality holds by the hypothesis that A permutes with B_1 . \square

For the present work, the most important consequence of Dedekind's rule is the following:

⁷See [25, p. 122], for example.

Corollary 3.7. *If $H \leq G$ and $N \trianglelefteq G$, then $(HN)^{\perp(H,G)}$ is an antichain.*

Proof. Note that HN permutes with each subgroup in $\llbracket H, G \rrbracket$. Indeed, if $H \leq A \leq G$, then $HNA = HAN = HN = AHN$. Therefore, by Corollary 3.6, the result holds. \square

Our next lemma (Lemma 3.8) generalizes a standard result. We find this generalization useful for proving that certain properties are core-free interval enforceable. To state Lemma 3.8, we need some new notation. Let U and H be subgroups of a group, let $U_0 = U \cap H$, and consider the interval $\llbracket U_0, U \rrbracket = \{V \mid U_0 \leq V \leq U\}$. It will be helpful to visualize part of the subgroup lattice of $\langle U, H \rangle$, as shown in Figure 3.




FIGURE 3. Some intervals in a subgroup lattice.

Recall that the usual isomorphism theorem for groups implies that if H is a normal subgroup of $\langle U, H \rangle$, then the interval $\llbracket H, \langle U, H \rangle \rrbracket$ is isomorphic to the interval $\llbracket U \cap H, U \rrbracket$. The purpose of the next lemma is to relate these two intervals in cases where we drop the assumption $H \trianglelefteq \langle U, H \rangle$ and add the assumption $UH = \langle U, H \rangle$.

If the two subgroups U and H permute, then we define

$$(3.5) \quad \llbracket U_0, U \rrbracket^H = \{V \in \llbracket U_0, U \rrbracket \mid VH = HV\},$$

which consists of those subgroups in $\llbracket U_0, U \rrbracket$ that permute with H .

If H normalizes U (which implies $UH = HU$), then we define

$$(3.6) \quad \llbracket U_0, U \rrbracket_H = \{V \in \llbracket U_0, U \rrbracket \mid H \leq N_{UH}(V)\}.$$

This is the set consisting of those subgroups in $\llbracket U_0, U \rrbracket$ that are normalized by H (sometimes called *H-invariant subgroups* in $\llbracket U_0, U \rrbracket$). Notice that to even define $\llbracket U_0, U \rrbracket_H$ we must have $H \leq N_{UH}(U)$, and in this case, as we will see below, the sublattices $\llbracket U_0, U \rrbracket_H$ and $\llbracket U_0, U \rrbracket^H$ coincide.

We are now ready to state the main result relating the sets defined in (3.5) and (3.6) to the interval $\llbracket H, UH \rrbracket$.

Lemma 3.8. *Suppose U and H are permuting subgroups of a group. Let $U_0 = U \cap H$. Then*

- (i) $\llbracket H, UH \rrbracket \cong \llbracket U_0, U \rrbracket^H \leq \llbracket U_0, U \rrbracket$.
- (ii) If $U \trianglelefteq UH$, then $\llbracket U_0, U \rrbracket_H = \llbracket U_0, U \rrbracket^H \leq \llbracket U_0, U \rrbracket$.
- (iii) If $H \trianglelefteq UH$, then $\llbracket U_0, U \rrbracket_H = \llbracket U_0, U \rrbracket^H = \llbracket U_0, U \rrbracket$.

Remarks. The expression $\llbracket H, UH \rrbracket \cong \llbracket U_0, U \rrbracket^H$ means that $\llbracket H, UH \rrbracket$ and $\llbracket U_0, U \rrbracket^H$ are isomorphic as lattices. Thus, when two subgroups permute, we can identify the interval above either one of them with the set of subgroups below the other that permute with the first. The expression $\llbracket U_0, U \rrbracket^H \leq \llbracket U_0, U \rrbracket$ means that the lattice $\llbracket U_0, U \rrbracket^H$ is, in fact, a sublattice of $\llbracket U_0, U \rrbracket$. That is, meets and joins in $\llbracket U_0, U \rrbracket^H$ coincide with those in $\llbracket U_0, U \rrbracket$. Part (ii) of Lemma 3.8 is a standard result (see, e.g., [7] or [15]). Since $G = UH$ is a group, the hypothesis of (ii) is equivalent to $H \leq N_G(U)$, and the hypothesis of (iii) is equivalent to $U \leq N_G(H)$. Once we have proved (i), the proof of (iii) follows trivially from the standard isomorphism theorem for groups, so we omit the details.

Proof. To prove (i), we first show that the following maps are inverse order isomorphisms:

$$(3.7) \quad \begin{aligned} \varphi : \llbracket H, UH \rrbracket \ni X &\mapsto U \cap X \in \llbracket U_0, U \rrbracket^H \\ \psi : \llbracket U_0, U \rrbracket^H \ni V &\mapsto VH \in \llbracket H, UH \rrbracket. \end{aligned}$$

Then we show that $\llbracket U_0, U \rrbracket^H$ is a sublattice of $\llbracket U_0, U \rrbracket$, that is, $\llbracket U_0, U \rrbracket^H \leq \llbracket U_0, U \rrbracket$. Fix $X \in \llbracket H, UH \rrbracket$. We claim that $U \cap X \in \llbracket U_0, U \rrbracket^H$. Indeed,

$$(U \cap X)H = UH \cap X = HU \cap X = H(U \cap X).$$

The first equality holds by (3.4) since $H \leq X$, the second holds by assumption, and the third by (3.3). This proves $U \cap X \in \llbracket U_0, U \rrbracket^H$. Moreover, by the first equality, $\psi \circ \varphi(X) = (U \cap X)H = UH \cap X = X$, so $\psi \circ \varphi$ is the identity on $\llbracket H, UH \rrbracket$.

If $V \in \llbracket U_0, U \rrbracket^H$, then $VH = HV$ implies $VH \in \llbracket H, UH \rrbracket$. Also, $\varphi \circ \psi$ is the identity on $\llbracket U_0, U \rrbracket^H$, since $\varphi \circ \psi(V) = VH \cap U = V(H \cap U) = VU_0 = V$, by (3.3). This proves that φ and ψ are inverses of each other on the sets indicated, and it's easy to see that they are order preserving: $X \leq Y$ implies $U \cap X \leq U \cap Y$, and $V \leq W$ implies $VH \leq WH$. Therefore, φ and ψ are inverse order isomorphisms.


To complete the proof of (i), we show that $\llbracket U_0, U \rrbracket^H$ is a sublattice of $\llbracket U_0, U \rrbracket$. Suppose V_1 and V_2 are subgroups in $\llbracket U_0, U \rrbracket$ that permute with H . It is easy to see that their join $V_1 \vee V_2 = \langle V_1, V_2 \rangle$ also permutes with H , so we just check that their intersection permutes with H . Fix $x \in V_1 \cap V_2$ and $h \in H$. We show $xh = h'x'$ for some $h' \in H$, $x' \in V_1 \cap V_2$. Since V_1 and V_2 permute with H , we have $xh = h_1v_1$ and $xh = h_2v_2$ for some $h_1, h_2 \in H$, $v_1 \in V_1$, $v_2 \in V_2$. Therefore, $h_1v_1 = h_2v_2$, which implies $v_1 = h_1^{-1}h_2v_2 \in HV_2$, so v_1 belongs to $V_1 \cap HV_2$. Note that $V_1 \cap HV_2$ is below both V_1 and $U \cap HV_2 = \varphi\psi(V_2) = V_2$. Therefore, $v_1 \in V_1 \cap HV_2 \leq V_1 \cap V_2$, and we have proved that $xh = h_1v_1$ for $h_1 \in H$ and $v_1 \in V_1 \cap V_2$, as desired.

To prove (ii), assuming $U \trianglelefteq G := UH$, we show that if $U_0 \leq V \leq U$, then $VH = HV$ if and only if $H \leq N_G(V)$. If $H \leq N_G(V)$, then $VH = HV$ (even when $U \not\trianglelefteq G$). Suppose $VH = HV$. We must show $(\forall v \in V)(\forall h \in H) hvh^{-1} \in V$. Fix $v \in V$, $h \in H$. Then, $hv = v'h'$ for some $v' \in V$, $h' \in H$, since $VH = HV$. Therefore, $v'h'h^{-1} = hvh^{-1} = u$ for some $u \in U$, since $H \leq N_G(U)$. This proves that $hvh^{-1} \in VH \cap U = V(H \cap U) = VU_0 = V$, as desired. \square

3.3. Parachute lattices. We now prove the equivalence of statements (B), (C), and (D) mentioned in Section 1. (That statement (E) of Section 1 is also equivalent to (B) follows easily from the arguments given below, so we omit the details.)

Theorem 3.9. *The following statements are equivalent:*

FIGURE 4. The parachute construction.



- (B) Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.
- (C) For every finite lattice L , for every finite collection $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ of cf-IE classes of groups, there exists a finite group $G \in \bigcap_{i=1}^n \mathfrak{X}_i$ such that $L \cong \llbracket H, G \rrbracket$ for some core-free subgroup $H \leq G$.
- (D) For every finite collection \mathcal{L} of finite lattices, there exists a finite group G such that each $L_i \in \mathcal{L}$ is isomorphic to $\llbracket H_i, G \rrbracket$ for some core-free subgroup $H_i \leq G$.
- (E) For every finite collection \mathcal{L} of finite lattices, for every finite collection $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ of cf-IE classes of groups, there exists a finite group $G \in \bigcap_{i=1}^n \mathfrak{X}_i$ such that for each $L_i \in \mathcal{L}$ we have $L_i \cong \llbracket H_i, G \rrbracket$ for some subgroup H_i that is core-free in G .

Remark. By (C), the FLRP would have a negative answer if we could find a collection $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ of cf-IE classes such that $\bigcap_{i=1}^n \mathfrak{X}_i$ is empty.

Proof. We prove the equivalence of (B) and (C). Obviously (C) implies (B). Assume (B) holds and let L be any finite lattice. Suppose $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ is a collection of cf-IE enforceable classes of groups. Then there exist finite lattices L_1, \dots, L_n such that $L_i \cong \llbracket H_i, G_i \rrbracket$ with $\text{core}_G(H_i) = 1$ implies $G_i \in \mathfrak{X}_i$. Construct a new lattice, denoted $\mathcal{P} = \mathcal{P}(L, L_1, \dots, L_n)$, as shown in the Hasse diagram of Figure 4 (a). Note that the bottoms of the L_i sublattices are atoms in \mathcal{P} . By (B), there exist groups $H < G$ with $\mathcal{P} \cong \llbracket H, G \rrbracket$. We can assume H is a core-free subgroup of G . (If not, replace G and H with G/N and H/N , where $N = \text{core}_G(H)$, and note that $\mathcal{P} \cong \llbracket H, G \rrbracket \cong \llbracket H/N, G/N \rrbracket$.) Let K, K_1, \dots, K_n be the subgroups of G in which H is maximal and for which $L \cong \llbracket K, G \rrbracket$ and $L_i \cong \llbracket K_i, G \rrbracket$, $1 \leq i \leq n$ (Figure 4

(b)). If we can prove for each $1 \leq i \leq n$ that $\text{core}_G(K_i) = 1$, then $G \in \mathfrak{X}_i$ by hypothesis, and it will follow that $G \in \bigcap_{i=1}^n \mathfrak{X}_i$, proving that (B) implies (C).

If $L \cong \mathbf{2}$, the two element lattice, or if $L_j \cong \mathbf{2}$ for all $1 \leq j \leq n$, then the consequent of the implication in (C) holds trivially. So we can assume without loss of generality that $L \not\cong \mathbf{2}$ and that there is at least one $1 \leq j \leq n$ for which $L_j \not\cong \mathbf{2}$. Let N be a minimal normal subgroup of G and suppose $N \leq K_i$. Since H is core-free, $K_i = NH$. Suppose $K_i \neq K$. Note that the groups K and K_i permute:

$$KK_i = KNH = NKH = NHK = K_iK.$$

Therefore, by Lemma 3.8 (i) we see that the interval $\llbracket K, G \rrbracket \cong L$ must be isomorphic to a sublattice of the interval $\llbracket H, K_i \rrbracket \cong \mathbf{2}$, but this contradicts $L \not\cong \mathbf{2}$. Suppose instead that $K = K_i$. Note that $K \neq K_j$, and since $K = NH$, we see that K and K_j permute. Therefore, by Lemma 3.8 (i) again, the lattice $L_j \not\cong \mathbf{2}$ is isomorphic to a sublattice of $\llbracket H, K \rrbracket \cong \mathbf{2}$, which is impossible. This proves that $NH = G$ for all $N \trianglelefteq G$, so each K_i is core-free.

The proof that statements (B) and (D) are equivalent follows by a similar construction. Roughly, if $\mathcal{L} = \{L_1, \dots, L_n\}$, we form the lattice $\mathcal{P} = \mathcal{P}(L_1, \dots, L_n)$. If (B) holds, then there exists a group G with $\mathcal{P} \cong \llbracket H, G \rrbracket$ and $\text{core}_G(H) = 1$. The proof that each K_i is core-free, where $L_i \cong \llbracket K_i, G \rrbracket$, is similar to the argument above. \square

By a *parachute lattice*, denoted $\mathcal{P}(L_1, \dots, L_m)$, we mean a lattice just like the one illustrated in Figure 4 (a), but with the lattices L_1, \dots, L_m appearing as the upper intervals.

Next we prove that any group that has a nontrivial parachute lattice as an upper interval in its subgroup lattice must have some rather special properties.

Lemma 3.10. *Let $\mathcal{P} = \mathcal{P}(L_1, \dots, L_n)$ with $n \geq 2$ and $|L_i| > 2$ for all i , and suppose $\mathcal{P} \cong \llbracket H, G \rrbracket$, with H core-free in G .*

- (i) *If $1 \neq N \trianglelefteq G$, then $NH = G$ and $C_G(N) = 1$.*
- (ii) *G is subdirectly irreducible and nonsolvable.*

Remark. If a subgroup $N \leq G$ is abelian, then $N \leq C_G(N)$, so (i) implies that every nontrivial normal subgroup of G is nonabelian.

Proof. (i) Let $1 \neq N \trianglelefteq G$. Then $N \not\leq H$, since H is core-free in G . Therefore, $H < NH$. As in Section 3.3, we let K_i denote the subgroups of G corresponding to the atoms of \mathcal{P} . Then H is covered by each K_i , so $K_j \leq NH$ for some $1 \leq j \leq n$. Suppose, by way of contradiction, that $NH < G$. By assumption, $n \geq 2$ and $|L_i| > 2$. Thus for any $i \neq j$ we have $K_i \leq Y < Z < G$ for some subgroups Y and Z which satisfy $(NH) \cap Z = H$ and $(NH) \vee Y = G$. Also, $(NH)Y = NY$ is a group, so $(NH)Y = NH \vee Y = G$. But then, by Dedekind's rule, we have

$$Y = HY = ((NH) \cap Z)Y = (NH)Y \cap Z = G \cap Z = Z,$$

contrary to $Y < Z$. This contradiction proves that $NH = G$.

To prove that $C_G(N) = 1$, let M be a minimal normal subgroup of G contained in N . It suffices to prove $C_G(M) = 1$. Assume the contrary. Then, since $C_G(M) \trianglelefteq N_G(M) = G$, it follows by what we just proved that $C_G(M)H = G$. Consider any $H < K < G$. Then $1 < M \cap K < M$ (strictly, by Lemma 3.8). Now $M \cap K$

is normalized by H and centralized (hence normalized) by $C_G(M)$. Therefore, $M \cap K \trianglelefteq C_G(M)H = G$, contradicting the minimality of M .

To prove (ii) we first show that G has a unique minimal normal subgroup. Let M be a minimal normal subgroup of G and let $N \trianglelefteq G$ be any normal subgroup not containing M . We show that $N = 1$. Since both subgroups are normal, the commutator subgroup $[M, N]$ lies in the intersection $M \cap N$, which is trivial by the minimality of M . Thus, M and N centralize each other. In particular, $N \leq C_G(M) = 1$, by (i).

Finally, since G has a unique minimal normal subgroup that is nonabelian (see the remark preceding the proof), G is nonsolvable. \square

To summarize what we have thus far, the lemmas above imply that (B) holds if and only if every finite lattice is an interval $\llbracket H, G \rrbracket$, with H core-free in G , where

- (i) G is nonsolvable, not alternating, and not symmetric;
- (ii) G has a unique minimal normal subgroup M which satisfies $MH = G$ and $C_G(M) = 1$; in particular, M is nonabelian and $\text{core}_G(X) = 1$ for all $H \leq X < G$.

We conclude this section by formalizing the remarks of the previous sentence. Given two group theoretical properties $\mathcal{P}_1, \mathcal{P}_2$, we write $\mathcal{P}_1 \rightarrow \mathcal{P}_2$ to denote that property \mathcal{P}_1 implies property \mathcal{P}_2 . In other words, $G \models \mathcal{P}_1$ only if $G \models \mathcal{P}_2$. Thus \rightarrow provides a natural partial order on any given set of properties, as follows:

$$\mathcal{P}_1 \leq \mathcal{P}_2 \iff \mathcal{P}_1 \rightarrow \mathcal{P}_2 \iff \mathfrak{X}_{\mathcal{P}_1} \subseteq \mathfrak{X}_{\mathcal{P}_2},$$

where $\mathfrak{X}_{\mathcal{P}_i} = \{G \in \mathfrak{G} \mid G \models \mathcal{P}_i\}$. The following is an immediate corollary of the parachute construction described above.

Corollary 3.11. *If $\mathcal{P} = \{\mathcal{P}_i \mid i \in \mathcal{I}\}$ is a collection of (cf-)IE properties, then $\bigwedge \mathcal{P}$ is (cf-)IE.*

The conjunction $\bigwedge \mathcal{P}$ corresponds to the class $\bigcap_{i \in \mathcal{I}} \mathfrak{X}_{\mathcal{P}_i} = \{G \in \mathfrak{G} \mid (\forall i \in \mathcal{I}) G \models \mathcal{P}_i\}$.

REFERENCES

- [1] Michael Aschbacher. On intervals in subgroup lattices of finite groups. *J. Amer. Math. Soc.*, 21(3):809–830, 2008. doi:10.1090/S0894-0347-08-00602-4.
- [2] Michael Aschbacher and John Shareshian. Restrictions on the structure of subgroup lattices of finite alternating and symmetric groups. *J. Algebra*, 322(7):2449–2463, 2009. URL: <http://dx.doi.org/10.1016/j.jalgebra.2009.05.042>.
- [3] Robert Baddeley and Andrea Lucchini. On representing finite lattices as intervals in subgroup lattices of finite groups. *J. Algebra*, 196(1):1–100, 1997. URL: <http://dx.doi.org/10.1006/jabr.1997.7069>.
- [4] Adolfo Ballester-Bolinches and Luis M. Ezquerro. *Classes of finite groups*, volume 584 of *Mathematics and Its Applications (Springer)*. Springer, Dordrecht, 2006.
- [5] Alberto Basile. *Second maximal subgroups of the finite alternating and symmetric groups*. PhD thesis, Australian National University, Canberra, April 2001.
- [6] Joel Berman. *Congruence lattices of finite universal algebras*. PhD thesis, University of Washington, 1970. URL: <http://db.tt/mXUVtzSr>.
- [7] Ferdinand Börner. A remark on the finite lattice representation problem. In *Contributions to general algebra, 11 (Olomouc/Velké Karlovice, 1998)*, pages 5–38, Klagenfurt, 1999. Heyn.
- [8] Richard Dedekind. Über die Anzahl der Ideal-classes in den verschiedenen Ordnungen eines endlichen Körpers. In *Festschrift zur Saecularfeier des Geburtstages von C. F. Gauss*, pages 1–55. Vieweg, Braunschweig, 1877. see Ges. Werke, Band I, 1930, 105–157.

- [9] William DeMeo. Given a lattice L with n elements, are there finite groups $H < G$ such that $L \cong$ the lattice of subgroups between H and G ? MathOverflow. (accessed 2012-05-08). URL: <http://mathoverflow.net/questions/85724>.
- [10] William DeMeo. *Congruence lattices of finite algebras*. PhD thesis, University of Hawai'i at Mānoa, Honolulu, HI, May 2012. [arXiv:1204.4305](https://arxiv.org/abs/1204.4305).
- [11] William DeMeo. Expansions of finite algebras and their congruence lattices. *Algebra Universalis*, 69:257–278, 2013. Available from: github.com/williamdemeo/Overalgebras. doi: 10.1007/s00012-013-0226-3.
- [12] William DeMeo and Ralph Freese. Congruence lattices of intransitive G -sets. *Preprint*, 2012.
- [13] John D. Dixon and Brian Mortimer. *Permutation groups*, volume 163 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1996.
- [14] Klaus Doerk and Trevor Hawkes. *Finite soluble groups*, volume 4 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1992.
- [15] I. Martin Isaacs. *Finite group theory*, volume 92 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [16] Peter Köhler. M_7 as an interval in a subgroup lattice. *Algebra Universalis*, 17(3):263–266, 1983. URL: <http://dx.doi.org/10.1007/BF01194535>.
- [17] Hans Kurzweil. Endliche Gruppen mit vielen Untergruppen. *J. Reine Angew. Math.*, 356:140–160, 1985. URL: <http://dx.doi.org/10.1515/crll.1985.356.140>.
- [18] Andrea Lucchini. Intervals in subgroup lattices of finite groups. *Comm. Algebra*, 22(2):529–549, 1994. doi:10.1080/00927879408824862.
- [19] Ralph N. McKenzie, George F. McNulty, and Walter F. Taylor. *Algebras, lattices, varieties. Vol. I*. Wadsworth & Brooks/Cole, Monterey, CA, 1987.
- [20] Péter Pál Pálffy. On Feit's examples of intervals in subgroup lattices. *J. Algebra*, 116(2):471–479, 1988. doi:10.1016/0021-8693(88)90230-x.
- [21] Péter Pál Pálffy. Intervals in subgroup lattices of finite groups. In *Groups '93 Galway/St. Andrews, Vol. 2*, volume 212 of *London Math. Soc. Lecture Note Ser.*, pages 482–494. Cambridge Univ. Press, Cambridge, 1995. doi:10.1017/CB09780511629297.014.
- [22] Péter Pál Pálffy. Groups and lattices. In *Groups St. Andrews 2001 in Oxford. Vol. II*, volume 305 of *London Math. Soc. Lecture Note Ser.*, pages 428–454, Cambridge, 2003. Cambridge Univ. Press. doi:10.1017/CB09780511542787.014.
- [23] Péter Pál Pálffy and Pavel Pudlák. Congruence lattices of finite algebras and intervals in subgroup lattices of finite groups. *Algebra Universalis*, 11(1):22–27, 1980. doi:10.1007/BF02483080.
- [24] Derek J. S. Robinson. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.
- [25] John S. Rose. *A course on group theory*. Dover Publications Inc., New York, 1994. Reprint of the 1978 original [Cambridge University Press; MR0498810 (58 #16847)].
- [26] Ada Rottlaender. Nachweis der Existenz nicht-isomorpher Gruppen von gleicher Situation der Untergruppen. *Math. Z.*, 28(1):641–653, 1928. URL: <http://dx.doi.org/10.1007/BF01181188>.
- [27] Roland Schmidt. *Subgroup lattices of groups*, volume 14 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [28] Yasuo Watatani. Lattices of intermediate subfactors. *J. Funct. Anal.*, 140(2):312–334, 1996. URL: <http://dx.doi.org/10.1006/jfan.1996.0110>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC 29208,
UNITED STATES

E-mail address: williamdemeo@gmail.com

URL: <http://williamdemeo.org>