

Homework 2

Matthew Corley and Melissa Murphy

Problem 1 (Golan 56). Is it possible to define on $\mathbb{Z}/(4)$ the structure of a vector space over $\text{GF}(2)$ in such a way that the vector addition is the usual addition in $\mathbb{Z}/(4)$?

[*Hints*: Recall that (n) denotes the set $\{\dots, -2n, -n, 0, n, 2n, 3n, \dots\}$, which we denoted in lecture by $n\mathbb{Z}$. This is the ideal generated by n in the ring \mathbb{Z} , but don't worry about that for now. Just take $\mathbb{Z}/(n)$ to be the abelian group of integers $\{0, 1, 2, \dots, n-1\}$ with addition modulo n . In lecture, we used $\mathbb{Z}/n\mathbb{Z}$ to denote $\mathbb{Z}/(n)$. Use whichever notation you prefer.]

Solution. Assume toward a contradiction that $\mathbb{Z}/(4)$ is a vector space over $\text{GF}(2)$, with vector addition defined as the usual modular addition. Then for any $v \in \mathbb{Z}/(4)$,

$$\begin{aligned} 0 &= (0)v \\ &= (1+1)v \\ &= 1 \cdot v + 1 \cdot v \\ &= v + v \end{aligned}$$

Now, let $v = 3$ to see that $3 + 3 = 2 \pmod{4} \neq 0$. Thus we have a contradiction. The answer is no, it is not possible with the usual modular vector addition.

Problem 2 (Golan 60). Let $V = C(0, 1)$. Define an operation \boxplus on V by setting $f \boxplus g : x \mapsto \max\{f(x), g(x)\}$. Does this operation of vector addition, together with the usual scalar multiplication make V into a vector space over \mathbb{R} ?

Solution. We must show that the \boxplus operation is closed in V , that is, $f \boxplus g \in C(0, 1)$. We will prove that $f \boxplus g$ is continuous by the "delta-epsilon" method from real analysis.

Let $\epsilon > 0$. As f, g continuous on $(0, 1)$, there exist $\delta_f > 0$ and $\delta_g > 0$ such that

$$\begin{aligned} |x - x_0| < \delta_f &\implies |f(x) - f(x_0)| < \frac{\epsilon}{2} \\ |x - x_0| < \delta_g &\implies |g(x) - g(x_0)| < \frac{\epsilon}{2} \end{aligned}$$

Let $h(x) = \max\{f(x), g(x)\}$. Let $\delta_h = \min\{\delta_f, \delta_g\}$. Then

$$\begin{aligned}
 |x - x_0| < \delta_h &\implies |h(x) - h(x_0)| \\
 &= \left| \frac{f(x) + g(x) + |f(x) - g(x)| - f(x_0) - g(x_0) - |f(x_0) - g(x_0)|}{2} \right| \\
 &\leq \left| \frac{f(x) - f(x_0)}{2} \right| + \left| \frac{g(x) - g(x_0)}{2} \right| + \left| \frac{f(x) - f(x_0) - (g(x) - g(x_0))}{2} \right| \\
 &< \frac{\epsilon}{2} + \left| \frac{f(x) - f(x_0) - (g(x) - g(x_0))}{2} \right| \\
 &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon
 \end{aligned}$$

Therefore $f \boxplus g$ is continuous for $x_0 \in (0, 1)$, so vector addition is closed in V . With the usual scalar multiplication, V is a vector space over \mathbb{R} .

Problem 3 (Golan 63). Let $V = \{i \in \mathbb{Z} \mid 0 \leq i < 2^n\}$ for some fixed positive integer n . Define operations of vector addition and scalar multiplication on V in such a way as to turn it into a vector space over $\text{GF}(2)$.

[*Hints*: Recall that $\text{GF}(2)$ denotes the Galois field with two elements, $\{0, 1\}$, with addition mod 2 and the usual multiplication. Other than this field, the only restriction given in the problem is that V must have 2^n elements. Do you know of any sets of this size?]

Solution. Let us define vector addition in a "bitwise xor" fashion such that $v + v = 0$ and $v + w = 1$ for all $v, w \in V, w \neq v$. Furthermore, let us define scalar multiplication in the natural way such that $1 \cdot v = v$ and $0 \cdot v = 0$. Then we can see that vector addition is closed as $0, 1 \in V$, as well as being associative and commutative. And every v has an additive inverse, namely v . Scalar multiplication is also closed in V , as the product is always $0 \in V$ or $v \in V$. So V is a vector space over $\text{GF}(2)$.

Problem 4 (Golan 70). Show that \mathbb{Z} is not a vector space over any field.

Solution. Let $\mathbb{F} = \text{GF}(2)$. Assume \mathbb{Z} is a vector space over \mathbb{F} . Then

$$\begin{aligned} 0 &= 1_Z(1_F + 1_F) \\ &= 1_Z(1_F) + 1_Z(1_F) \\ &= 2 \end{aligned}$$

But $0 \neq 2$, so \mathbb{Z} is not a vector space over $\text{GF}(2)$. Now let \mathbb{F} be a field with characteristic greater than 2. Assume \mathbb{Z} a vector space over \mathbb{F} . Then we know

$$1_F + 1_F = 2_F \implies 1_F = \frac{1}{2_F} + \frac{1}{2_F}$$

Therefore

$$\begin{aligned} 1_Z &= 1_F(1_Z) \\ &= \left(\frac{1}{2_F} + \frac{1}{2_F}\right)1_Z \\ &= \frac{1}{2_F}(1_Z) + \frac{1}{2_F}(1_Z) \end{aligned}$$

Now let $\frac{1}{2_F}(1_Z) = n \in \mathbb{Z}$. But there is no element in $n \in \mathbb{Z}$ that satisfies $n + n = 1$. So \mathbb{Z} is not a vector space over any field with characteristic greater than 2. Thus \mathbb{Z} is not a vector space over any field.

Problem 5 (Golan 76). Let $V = \mathbb{R}^{\mathbb{R}}$ and let W be the subset of V containing the constant function $x \mapsto 0$ and all of those functions $f \in V$ satisfying the following condition: $f(a) = 0$ for at most finitely many real numbers a . Is W a subspace of V .

[Hint: It's easy.]

Solution. Let us assume toward a contradiction that W is a subspace of V . Let $p(x), g(x), h(x)$ be distinct functions in W , such that $p(x) = g(x) + h(x)$ and $g(x) = 0$ for $x = b_0, \dots, b_l \in \mathbb{R}$ and $h(x) = 0$ for $x = c_0, \dots, c_m \in \mathbb{R}$. Then $p(x) = 0$ when $g(x) = h(x) = 0$ or when $g(x) = -h(x)$. The former happens when $b_i = c_j$ for $1 \leq i \leq l, 1 \leq j \leq m$. We can see this is a finite set of points. The latter, however, could happen for an infinite number of points (e.g. define $g(x) = -h(x)$ for $x > \max(b_l, c_m) \in \mathbb{R}$). In that case, $p(a) = 0$ for an infinitely many real numbers a , but it is not the constant function $x \mapsto 0$. So vector addition is not closed and therefore W is not a subspace in V .

Problem 6 (Golan 79). A function $f \in \mathbb{R}^{\mathbb{R}}$ is *piecewise constant* if and only if it is a constant function $x \mapsto c$ or there exist $a_1 < a_2 < \dots < a_n$ and $c_0 < c_1 < \dots < c_n$ in \mathbb{R} such that

$$f : x \mapsto \begin{cases} c_0 & \text{if } x < a_1, \\ c_i & \text{if } a_i \leq x < a_{i+1} \text{ for } 1 \leq i < n, \\ c_n & \text{if } a_n \leq x. \end{cases}$$

Does the set of all piecewise constant functions form a subspace of the vector space $\mathbb{R}^{\mathbb{R}}$ over \mathbb{R} ?

Solution. Let W denote the set in question. W is clearly a subset of $\mathbb{R}^{\mathbb{R}}$. Let $r \in \mathbb{R}$ and $k_i = rc_i$. Then

$$rf : x \mapsto \begin{cases} k_0 & \text{if } x < a_1, \\ k_i & \text{if } a_i \leq x < a_{i+1} \text{ for } 1 \leq i < n, \\ k_n & \text{if } a_n \leq x. \end{cases}$$

So W is closed under scalar multiplication.

Let us describe $f(x), g(x) \in W$ using the characteristic function χ :

$$f(x) = \sum_{i=1}^n f(a_i) \chi_{[a_i, a_{i+1}]}(x)$$

$$g(x) = \sum_{i=1}^n g(b_i) \chi_{[b_i, b_{i+1}]}(x)$$

Now let $\{z_0, z_1, \dots\}$ be the reordering of $\{a_0, a_1, \dots, a_n, b_0, b_1, \dots, b_n\}$. Then we can describe the sum $f(x) + g(x)$ as follows:

$$(f + g)(x) = \sum_{i=1}^n (f(z_i) + g(z_i)) \chi_{[z_i, z_{i+1}]}(x)$$

Thus $f(x) + g(x)$ is a piecewise constant function and so W is closed under vector addition. Therefore W is a subspace of $\mathbb{R}^{\mathbb{R}}$.

Problem 7 (Golan 81). Let W be the subspace of $V = \text{GF}(2)^5$ consisting of all vectors (a_1, \dots, a_5) satisfying $\sum_{i=1}^5 a_i = 0$. Is W a subspace of V ?

Solution. (We assume the problem means "subset" not "subspace" in the first line. Otherwise trivial.)

We know W is a subset of V and $O_V \in V$. Let $x \in V, x = \langle a_1, a_2, a_3, a_4, a_5 \rangle$ and $\sum_{i=1}^5 a_i = 0$. Let $b \in F$. Then

$$bx = \langle b(a_1), b(a_2), b(a_3), b(a_4), b(a_5) \rangle$$

And

$$\sum_{i=1}^5 (ba_i) = b \left(\sum_{i=1}^5 a_i \right) = b(0) = 0$$

So W is closed under scalar multiplication.

Let $y \in V, y = \langle c_1, c_2, c_3, c_4, c_5 \rangle$ such that $\sum_{i=1}^5 c_i = 0$. Then

$$x + y = \langle a_1 + c_1, a_2 + c_2, a_3 + c_3, a_4 + c_4, a_5 + c_5 \rangle$$

Now let $d_i = a_i + c_i$.

$$\begin{aligned} \sum_{i=1}^5 d_i &= \sum_{i=1}^5 (a_i + c_i) \\ &= \sum_{i=1}^5 a_i + \sum_{i=1}^5 c_i \\ &= 0 + 0 = 0 \end{aligned}$$

Therefore W is closed under vector addition and scalar multiplication so it is a subspace.

Problem 8 (Golan 85). Let $V = \mathbb{R}^{\mathbb{R}}$ and let W be the subset of V consisting of all functions f satisfying the following condition: there exists $c \in \mathbb{R}$ (that depends on f) such that $|f(a)| \leq c|a|$ for all $a \in \mathbb{R}$. Is W a subspace of V ?

Solution. We know W is a subset of V . Let $g(x) \in W$. There exists a $c_1 \in \mathbb{R}$ such that $\forall x \in \mathbb{R} |h(x)| \leq c|x|$. Let $b \in \mathbb{R}$. Let $h(x) = b(g(x))$. Then

$$\begin{aligned} |h(x)| &= |b(g(x))| \\ &\leq |b||g(x)| \\ &\leq |b|(c_1|x|) \\ &= c|x| \end{aligned}$$

Where $c = |b|c_1$. So $\exists c \in \mathbb{R}$ such that $\forall x \in \mathbb{R}$ we have

$$|h(x)| \leq c|x|$$

Therefore W is closed under scalar multiplication.

Let $p(x), g(x) \in W$. There exist $c_1, c_2 \in \mathbb{R}$ such that $\forall x \in \mathbb{R}$

$$|p(x)| \leq c_1|x|$$

$$|q(x)| \leq c_2|x|$$

Let $t(x) = p(x) + g(x)$. Then

$$\begin{aligned} t(x) &= |p(x) + g(x)| \\ &\leq |p(x)| + |g(x)| \\ &\leq c_1|x| + c_2|x| \\ &= (c_1 + c_2)|x| \\ &= c_3|x| \end{aligned}$$

where $c_3 = c_1 + c_2$. Therefore $\exists c \in \mathbb{R}$ such that $\forall x \in \mathbb{R}$ we have

$$t(x) \leq c|x|$$

Thus W is closed under addition and scalar multiplication so W is a subspace of V .

Problem 9 (Golan 93). Let V be a vector space over a field F and let P be the collection of all subsets of V , which we know is a vector space over $\text{GF}(2)$. Is the collection of all subspaces of V a subspace of P ?

Solution. P is the collection of all subset of V . Let U be the collection of all subspaces of V . As all subspaces of V must be subsets of V , we know that $U \subseteq V$. We want to show that U is a subspace of P .

Let $X \in U$. Then $X \in P$. Let $a \in F$. Let $y, z \in X$. Then $ay, az \in aX$. $ay + az = a(x + y)$. As X is a subspace, $(x + y) \in X$ so $a(x + y) \in aX$. Thus aX is closed under addition. Let $b \in F$.

$$\begin{aligned} b(ay) &= (ba)y \\ &= (ab)y \\ &= a(by) \end{aligned}$$

So $by \in X \implies a(by) \in aX$. aX is closed under scalar multiplication. And $X \subseteq V \implies aX \subseteq V$. Then aX is a subspace, so U is closed under scalar multiplication.

Let $Y \in U, x \in X, y \in Y$. $X + Y = \mathbb{F}\{x, y\}$. Let $p, q \in X + Y$ such that for $x_1, x_2 \in X, y_1, y_2 \in Y, a, b \in F$

$$p = ax_1 + by_1$$

$$q = cx_2 + dy_2$$

Now we add $p + q$

$$p + q = (ax_1 + by_1) + (cx_2 + dy_2) \quad (1)$$

$$= (ax_1 + cx_2) + (by_1 + dy_2) \quad (2)$$

Since we know $(ax_1 + cx_2) \in X$ and $(by_1 + dy_2) \in Y$, we can deduce that $p + q \in X + Y$. Then U is closed under addition. So U is a subspace of P .

Problem 10 (Golan 105). Let V be a vector space over a field F and let $0_V \neq w \in V$. Given a vector $v \in V \setminus Fw$, find the set G of all scalars $a \in F$ satisfying $F\{v, w\} = F\{v, aw\}$.

Solution.

$$F\{v, w\} = \{bv + cw \mid b, c \in F\}$$

$$F\{v, aw\} = \{dv + e(aw) \mid d, e \in F\}$$

F is a field, so it is closed under multiplication. v is not a scalar multiple of w . We know $1 \in G$. Thus $F\{v, aw\} \leq F\{v, w\} \forall a \in F$. Let $x \in F\{v, w\}$. Then for $s, t \in F$

$$\begin{aligned} x &= sv + tw \\ &= sv + \frac{t}{a}(aw) \end{aligned}$$

So $\frac{t}{a} = ta^{-1}$, and $t \in F, a^{-1} \in F$ so $ta^{-1} \in F$ as long as $a \neq 0$. So $G = F \setminus 0_V$.