

THE FINITE LATTICE REPRESENTATION PROBLEM  
AND  
INTERVALS IN SUBGROUP LATTICES  
PART I

William DeMeo

`williamdemeo@gmail.com`

joint work with

Ralph Freese, Peter Jipsen, Bill Lampe, J.B. Nation

University of South Carolina

Algebra & Logic Seminar

September 21, 2012


*These slides and other resources are available at*

<http://www.math.sc.edu/~demeow/FLRP.html> →



# CONGRUENCE DECOMPOSITIONS


CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

# CONGRUENCE DECOMPOSITIONS

CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

# THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

## THEOREM (GRÄTZER-SCHMIDT, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

What if the algebra is finite?

**Problem:** Given a finite lattice  $L$ , does there exist a *finite* algebra  $A$  such that  $\text{Con } A \cong L$ ?

## DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

# THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

## THEOREM (GRÄTZER-SCHMIDT, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

What if the algebra is finite?

**Problem:** Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{Con A} \cong \mathbf{L}$ ?

## DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

# THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

## THEOREM (GRÄTZER-SCHMIDT, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

What if the algebra is finite?

**Problem:** Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{Con A} \cong \mathbf{L}$ ?

## DEFINITION

We call a finite lattice **representable** if it is isomorphic to the congruence lattice of a finite algebra.

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .



## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .
- Does equality hold in each case?

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$ 
  - (Birkhoff, 1935) Is every lattice  $\cong$  a sublattice of a partition lattice?
  - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
  - (Pudlák and Tůma, 1980) Yes,  $\mathcal{L}_0 = \mathcal{L}_1$ .

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$ 
  - (Birkhoff, 1935) Is every lattice  $\cong$  a sublattice of a partition lattice?
  - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
  - (Pudlák and Tůma, 1980) Yes,  $\mathcal{L}_0 = \mathcal{L}_1$ .
- **Main problem:** Is  $\mathcal{L}_0 = \mathcal{L}_3$  true?

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$ 
  - (Birkhoff, 1935) Is every lattice  $\cong$  a sublattice of a partition lattice?
  - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
  - (Pudlák and Tůma, 1980) Yes,  $\mathcal{L}_0 = \mathcal{L}_1$ .
- **Main problem:** Is  $\mathcal{L}_0 = \mathcal{L}_3$  true?
- (Pálffy and Pudlák, 1980)  $\mathcal{L}_0 = \mathcal{L}_3$  if and only if  $\mathcal{L}_0 = \mathcal{L}_4$ .

## SOME IMPORTANT CLASSES OF FINITE LATTICES

- $\mathcal{L}_0$  = all finite lattices
- $\mathcal{L}_1$  = lattices isomorphic to sublattices of finite partition lattices
- $\mathcal{L}_2$  = ...strong congruence lattices of finite partial algebras
- $\mathcal{L}_3$  = ...congruence lattices of finite algebras
- $\mathcal{L}_4$  = ...intervals in subgroup lattices of finite groups
- $\mathcal{L}_5$  = ...subgroup lattices of finite groups

- Fact:  $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$ .
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$ 
  - (Birkhoff, 1935) Is every lattice  $\cong$  a sublattice of a partition lattice?
  - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
  - (Pudlák and Tůma, 1980) Yes,  $\mathcal{L}_0 = \mathcal{L}_1$ .
- **Main problem:** Is  $\mathcal{L}_0 = \mathcal{L}_3$  true?
- (Pálffy and Pudlák, 1980)  $\mathcal{L}_0 = \mathcal{L}_3$  if and only if  $\mathcal{L}_0 = \mathcal{L}_4$ .

This does **not** say  $\mathcal{L}_3 = \mathcal{L}_4$ . It's possible that  $\mathcal{L}_0 \subsetneq \mathcal{L}_3 \subsetneq \mathcal{L}_4$ .

## RECAP

### THEOREM (PUDLÁK AND TŮMA, 1980)

*Every finite lattice can be embedded in  $\text{Eq}(X)$  with  $X$  finite.*

In other words,  $\mathcal{L}_0 = \mathcal{L}_1$ .

### THEOREM (PÁLFY AND PUDLÁK, 1980)

*The following statements are equivalent:*

- (I) *Every finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (II) *Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

In other words,  $\mathcal{L}_0 = \mathcal{L}_3$  if and only if  $\mathcal{L}_0 = \mathcal{L}_4$ .

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 1 (USE CLOSURE PROPERTIES)

The class  $\mathcal{L}_3$  is closed under the following operations:

- lattice duals (Kurzweil and Netter, 1986)
- interval sublattices (follows from Kurzweil-Netter)
- direct products (Tůma, 1986)
- ordinal sums (McKenzie, 1984; Snow, 2000)
- parallel sums (Snow, 2000)
- certain sublattices of lattices in  $\mathcal{L}_3$  (Snow, 2000)  
(namely, those obtained as a union of a filter and ideal)

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix  $\theta \subseteq X \times X$ ,  $f : X^n \rightarrow X$ .

Say that  $f$  **respects**  $\theta$  and write  $f(\theta) \subseteq \theta$  provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$



# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix  $\theta \subseteq X \times X$ ,  $f : X^n \rightarrow X$ .

Say that  $f$  **respects**  $\theta$  and write  $f(\theta) \subseteq \theta$  provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For  $L \subseteq \text{Eq}(X)$  define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on  $X$  which respect all relations in  $L$ .

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix  $\theta \subseteq X \times X$ ,  $f : X^n \rightarrow X$ .

Say that  $f$  **respects**  $\theta$  and write  $f(\theta) \subseteq \theta$  provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For  $L \subseteq \text{Eq}(X)$  define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on  $X$  which respect all relations in  $L$ .

- For  $F \subseteq X^X$  define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on  $X$  respected by all  $f \in F$ .

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix  $\theta \subseteq X \times X$ ,  $f : X^n \rightarrow X$ .

Say that  $f$  **respects**  $\theta$  and write  $f(\theta) \subseteq \theta$  provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For  $L \subseteq \text{Eq}(X)$  define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on  $X$  which respect all relations in  $L$ .

- For  $F \subseteq X^X$  define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on  $X$  respected by all  $f \in F$ .

- Then  $L \subseteq \rho\lambda(L)$  and  $\rho\lambda$  is a *closure operator* on  $\text{Sub}[\text{Eq}(X)]$ .  
(idempotent, extensive, order preserving)

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix  $\theta \subseteq X \times X$ ,  $f : X^n \rightarrow X$ .

Say that  $f$  **respects**  $\theta$  and write  $f(\theta) \subseteq \theta$  provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For  $L \subseteq \text{Eq}(X)$  define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on  $X$  which respect all relations in  $L$ .

- For  $F \subseteq X^X$  define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on  $X$  respected by all  $f \in F$ .

- Then  $L \subseteq \rho\lambda(L)$  and  $\rho\lambda$  is a *closure operator* on  $\text{Sub}[\text{Eq}(X)]$ .  
(idempotent, extensive, order preserving)
- If a lattice  $L \leq \text{Eq}(X)$  is *closed*, i.e.  $\rho\lambda(L) = L$ , then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 3 (SUBGROUP LATTICE INTERVAL)

Find  $L$  as an interval in a subgroup lattice of a finite group.

If  $H \leq G$  are finite groups, then the interval above  $H$  in  $\text{Sub}(G)$ ,

$$[H, G] := \{K \mid H \leq K \leq G\},$$


is isomorphic to  $\text{Con} \langle G/H, G \rangle$ .

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 4 (FILTER+IDEAL)

Find  $L$  as the union of a filter and ideal in a representable lattice.

Suppose  $L_0 \cong \text{Con } \langle A, F \rangle$ ,  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ .



# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE


## METHOD 4 (FILTER+IDEAL)

Find  $L$  as the union of a filter and ideal in a representable lattice.

Suppose  $L_0 \cong \text{Con } \langle A, F \rangle$ ,  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ .

Consider  $L = \alpha^\uparrow \cup \beta^\downarrow$ .

$$L \leq L_0$$



# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 4 (FILTER+IDEAL)

Find  $L$  as the union of a filter and ideal in a representable lattice.


Suppose  $L_0 \cong \text{Con} \langle A, F \rangle$ ,  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ .

Consider  $L = \alpha^\uparrow \cup \beta^\downarrow$ .

Then there exists a set  $F' \subset A^A$  such that


$$L \cong \text{Con} \langle A, F \cup F' \rangle.$$

$$L \leq L_0$$





# LATTICES WITH AT MOST 6 ELEMENTS ARE REPRESENTABLE.




Watatani (1996) *J. Funct. Anal.*


Aschbacher (2008) *JAMS*

**Theorem:** *Every lattice with at most 6 elements is an interval in the subgroup lattice of a finite group.*


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$




$L_{20}$




$L_{17}$




$L_{13}$



$L_{11}$




$L_9$




$L_7$


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$




$L_{20}$




$L_{17}$




$L_{13}$



$L_{11}$





$L_9$



$L_7$


# FINDING REPRESENTATIONS...

...AS INTERVALS IN SUBGROUP LATTICES



# FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



`SmallGroup(288,1025)`


$$|G : H| = 48$$

- The group  $G = (A_4 \times A_4) \rtimes C_2$  has a subgroup  $H \cong S_3$  such that  $[H, G] \cong L_{17}$ .
- ...so the dual  $L_{16}$  is also representable.



# FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$

- The group  $G = (A_4 \times A_4) \rtimes C_2$  has a subgroup  $H \cong S_3$  such that  $[H, G] \cong L_{17}$ .
- ...so the dual  $L_{16}$  is also representable.




SmallGroup(960,11358)

$$|G : H| = 80$$

- The group  $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$  has a subgroup  $H \cong A_4$  such that  $[H, G] \cong L_{13}$ .

## FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.



# FINDING REPRESENTATIONS...

...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)




- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .



# FINDING REPRESENTATIONS...

...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)




- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$

# FINDING REPRESENTATIONS...



...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)



- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$




- $\text{Sub}(A_4)$  is a congruence lattice (of  $A_4$  acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓




$L_9$




$L_7$


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓




$L_9$  ✓




$L_7$


# SEVEN ELEMENT LATTICES: SUMMARY




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓




$L_9$  ✓




$L_7$


## SEVEN ELEMENT LATTICES: SUMMARY




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓



$L_9$  ✓



$L_7$


# HAS ANYONE SEEN THIS LATTICE?

[login](#)[faq](#)[how to ask](#)[meta](#)**mathoverflow**[Questions](#)[Tags](#)[Users](#)[Badges](#)[Unanswered](#)[Ask Question](#)

Given a lattice  $L$  with  $n$  elements, are there finite groups  $H < G$  such that  $L \cong$  the lattice of subgroups between  $H$  and  $G$ ?

▲  
13  
▼

If there is no restriction on  $n$ , this is a famous [open problem](#). I'm wondering if any recent work has been done for small  $n > 6$ . I believe the question is answered (positively) for  $n = 6$  by Watatani (1996) [MR1409040](#) and Aschbacher (2008) [MR2393428](#). I also believe we can answer it for  $n = 7$ , with one possible exception. The exceptional case is shown below.



So my two questions are these:

- 1) Does anyone know of recent work on this special case of the problem (specifically for  $n = 7$  or  $n = 8$ )?
- 2) Has anyone found a finite group  $G$  with a subgroup  $H$  such that the interval  $[H, G] = \{K : H \leq K \leq G\}$

is the lattice shown above?

## Welcome to MathOverflow

A place for mathematicians to ask and answer questions.

tagged

[finite-groups](#) × 343

[open-problem](#) × 216

[lattices](#) × 157

[universal-algebra](#) × 55

[congruences](#) × 7

asked


**8 months ago**

viewed

**468 times**

## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES


- $L_7$  cannot be obtained using the overalgebra construction.






## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- $L_7$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_7$  must come from a transitive  $G$ -set.




## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- $L_7$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_7$  must come from a transitive  $G$ -set.
- Suppose  $L_7 \cong [H, G]$  for some finite groups  $H < G$ .  
What can we say about the group  $G$ ?




## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- $L_7$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_7$  must come from a transitive  $G$ -set.
- Suppose  $L_7 \cong [H, G]$  for some finite groups  $H < G$ .  
What can we say about the group  $G$ ?
- If we prove  $G$  must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- $L_7$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_7$  must come from a transitive  $G$ -set.
- Suppose  $L_7 \cong [H, G]$  for some finite groups  $H < G$ .  
What can we say about the group  $G$ ?
- If we prove  $G$  must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




### THEOREM

Suppose  $H < G$ ,  $\text{core}_G(H) = 1$ ,  $L_7 \cong [H, G]$ .

## INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- $L_7$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_7$  must come from a transitive  $G$ -set.
- Suppose  $L_7 \cong [H, G]$  for some finite groups  $H < G$ .  
What can we say about the group  $G$ ?
- If we prove  $G$  must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



### THEOREM

Suppose  $H < G$ ,  $\text{core}_G(H) = 1$ ,  $L_7 \cong [H, G]$ .

- (I)  $G$  is a primitive permutation group.
- (II) If  $N \triangleleft G$ , then  $C_G(N) = 1$ .
- (III)  $G$  contains no non-trivial abelian normal subgroup.
- (IV)  $G$  is not solvable.
- (V)  $G$  is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, all proper subgroups in the interval  $[H, G]$  are core-free.

## SUBGROUP LATTICE BASICS

Let  $U$  and  $H$  be subgroups of a finite group.

- By  $UH$  we mean the set  $\{uh \mid u \in U, h \in H\}$ .

## SUBGROUP LATTICE BASICS

Let  $U$  and  $H$  be subgroups of a finite group.

- By  $UH$  we mean the set  $\{uh \mid u \in U, h \in H\}$ .
- $U \vee V = \langle U, H \rangle$  means the group generated by  $U$  and  $H$ .

## SUBGROUP LATTICE BASICS

Let  $U$  and  $H$  be subgroups of a finite group.

- By  $UH$  we mean the set  $\{uh \mid u \in U, h \in H\}$ .
- $U \vee V = \langle U, H \rangle$  means the group generated by  $U$  and  $H$ .
- $UH \subseteq \langle U, H \rangle$  and equality holds iff  $U$  and  $H$  permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$




## SUBGROUP LATTICE BASICS

Let  $U$  and  $H$  be subgroups of a finite group.

- By  $UH$  we mean the set  $\{uh \mid u \in U, h \in H\}$ .
- $U \vee V = \langle U, H \rangle$  means the group generated by  $U$  and  $H$ .
- $UH \subseteq \langle U, H \rangle$  and equality holds iff  $U$  and  $H$  permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$




## SUBGROUP LATTICE BASICS

Let  $U$  and  $H$  be subgroups of a finite group.

- By  $UH$  we mean the set  $\{uh \mid u \in U, h \in H\}$ .
- $U \vee H = \langle U, H \rangle$  means the group generated by  $U$  and  $H$ .
- $UH \subseteq \langle U, H \rangle$  and equality holds iff  $U$  and  $H$  permute:


$$UH = \langle U, H \rangle \iff UH = HU.$$




$$[U_0, U] := \{V \mid U_0 \leq V \leq U\}$$

## INTERVAL ISOMORPHISMS

- If  $H \trianglelefteq \langle U, H \rangle$ , then  $UH = \langle U, H \rangle$  and  $[U_0, U] \cong [H, UH]$ .



## INTERVAL ISOMORPHISMS




- If  $H \trianglelefteq \langle U, H \rangle$ , then  $UH = \langle U, H \rangle$  and  $[U_0, U] \cong [H, UH]$ .
- Instead of  $H \trianglelefteq \langle U, H \rangle$ , assume only  $UH = \langle U, H \rangle$  and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

## INTERVAL ISOMORPHISMS



- If  $H \trianglelefteq \langle U, H \rangle$ , then  $UH = \langle U, H \rangle$  and  $[U_0, U] \cong [H, UH]$ .
- Instead of  $H \trianglelefteq \langle U, H \rangle$ , assume only  $UH = \langle U, H \rangle$  and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$


the ***H-permuting subgroups***.

- If  $U \trianglelefteq UH$ , define

$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***:  $V^h = V$  ( $\forall h \in H$ ).

# INTERVAL ISOMORPHISMS



- If  $H \trianglelefteq \langle U, H \rangle$ , then  $UH = \langle U, H \rangle$  and  $[U_0, U] \cong [H, UH]$ .
- Instead of  $H \trianglelefteq \langle U, H \rangle$ , assume only  $UH = \langle U, H \rangle$  and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

- If  $U \trianglelefteq UH$ , define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***:  $V^h = V$  ( $\forall h \in H$ ).

## LEMMA

- 1  $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$
- 2 If  $U \trianglelefteq UH$ , then  $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$ .
- 3 If  $H \trianglelefteq UH$ , then  $[U_0, U]_H = [U_0, U]^H = [U_0, U]$ .

# INTERVAL ISOMORPHISMS



- If  $H \trianglelefteq \langle U, H \rangle$ , then  $UH = \langle U, H \rangle$  and  $[U_0, U] \cong [H, UH]$ .
- Instead of  $H \trianglelefteq \langle U, H \rangle$ , assume only  $UH = \langle U, H \rangle$  and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

- If  $U \trianglelefteq UH$ , define

$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***:  $V^h = V$  ( $\forall h \in H$ ).

## LEMMA

- 1  $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$
- 2 If  $U \trianglelefteq UH$ , then  $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$ .
- 3 If  $H \trianglelefteq UH$ , then  $[U_0, U]_H = [U_0, U]^H = [U_0, U]$ .

## EXAMPLE 1

- Consider  $G \cong C_3 \times S_3$ , say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$



## EXAMPLE 1


- Consider  $G \cong C_3 \times S_3$ , say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ( $UH = HU$ ) but neither one normalizes the other.



## EXAMPLE 1


- Consider  $G \cong C_3 \times S_3$ , say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ( $UH = HU$ ) but neither one normalizes the other.



## EXAMPLE 1


- Consider  $G \cong C_3 \times S_3$ , say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ( $UH = HU$ ) but neither one normalizes the other.




- Three of the four subgroups of  $U$  permute with  $H$ .  
As the lemma predicts,  $U \cap \langle b, c \rangle = \langle b \rangle$ .

## EXAMPLE 2

- The group  $S_4$  has subgroups  $U \cong D_8$  and  $H \cong C_3$  that permute but neither one normalizes the other.

## EXAMPLE 2


- The group  $S_4$  has subgroups  $U \cong D_8$  and  $H \cong C_3$  that permute but neither one normalizes the other.



- Only four subgroups of  $U$  permute with  $H$


## EXAMPLE 2

- The group  $S_4$  has subgroups  $U \cong D_8$  and  $H \cong C_3$  that permute but neither one normalizes the other.



- Only four subgroups of  $U$  permute with  $H$ , including

$$U \cap A_4 \cong C_2 \times C_2, \quad U \cap S_3 \cong C_2.$$




## THEOREM

Suppose  $H < G$ ,  $\text{core}_G(H) = 1$ , and  $L_7 \cong [H, G]$ . Then

- (I)  $G$  is a primitive permutation group.
- (II) If  $N \triangleleft G$ , then  $C_G(N) = 1$ .
- (III)  $G$  contains no non-trivial abelian normal subgroup.
- (IV)  $G$  is not solvable.
- (V)  $G$  is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup,  $M_1$  or  $M_2$ , all proper subgroups in the interval  $[H, G]$  are core-free.

## IDEA OF THE PROOF




**Claim:**  $J_1$  and  $J_2$  are core-free subgroups of  $G$ .

**Proof:**



## IDEA OF THE PROOF




**Claim:**  $J_1$  and  $J_2$  are core-free subgroups of  $G$ .

**Proof:**

- If  $N \triangleleft G$  then  $NH$  permutes with each subgroup containing  $H$ .

## IDEA OF THE PROOF




**Claim:**  $J_1$  and  $J_2$  are core-free subgroups of  $G$ .

**Proof:**

- If  $N \triangleleft G$  then  $NH$  permutes with each subgroup containing  $H$ .
- If  $1 \neq N \leq J_1$ , then  $NH = J_1$ , so  $J_1$  and  $K$  permute.

## IDEA OF THE PROOF




**Claim:**  $J_1$  and  $J_2$  are core-free subgroups of  $G$ .

**Proof:**

- If  $N \triangleleft G$  then  $NH$  permutes with each subgroup containing  $H$ .
- If  $1 \neq N \leq J_1$ , then  $NH = J_1$ , so  $J_1$  and  $K$  permute.
- Since  $J_1K = G$  and  $J_1 \cap K = H$ , our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1X = XJ_1\}.$$

## IDEA OF THE PROOF



**Claim:**  $J_1$  and  $J_2$  are core-free subgroups of  $G$ .

**Proof:**

- If  $N \triangleleft G$  then  $NH$  permutes with each subgroup containing  $H$ .
- If  $1 \neq N \leq J_1$ , then  $NH = J_1$ , so  $J_1$  and  $K$  permute.
- Since  $J_1K = G$  and  $J_1 \cap K = H$ , our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1X = XJ_1\}.$$

Impossible!



# ASCHBACHER-O'NAN-SCOTT THEOREM

Let  $G$  be a primitive permutation group of degree  $d$ , and let  $N := \text{Soc}(G) \cong T^m$  with  $m \geq 1$ . Then one of the following holds.

- ①  $N$  is regular and
  - **(Affine type)**  $T$  is cyclic of order  $p$ , so  $|N| = p^m$ . Then  $d = p^m$  and  $G$  is permutation isomorphic to a subgroup of the affine general linear group  $\text{AGL}(m, p)$ .
  - **(Twisted wreath product type)**  $m \geq 6$ , the group  $T$  is nonabelian and  $G$  is a group of *twisted wreath product type*, with  $d = |T|^m$ .
- ②  $N$  is non-regular, non-abelian, and
  - **(Almost simple type)**  $m = 1$  and  $T \leq G \leq \text{Aut}(T)$ .
  - **(Product action type)**  $m \geq 2$  and  $G$  is permutation isomorphic to a subgroup of the product action wreath product  $P \wr_{m/l} S_m$  of degree  $d = nm/l$ . The group  $P$  is primitive of type 2.(a) or 2.(c),  $P$  has degree  $n$  and  $\text{Soc}(P) \cong T^l$ , where  $l \geq 1$  divides  $m$ .
  - **(Diagonal type)**  $m \geq 2$  and  $T^m \leq G \leq T^m \cdot (\text{Out}(T) \times S_m)$ , with the diagonal action. The degree  $d = |T|^{m-1}$ .

## ASCHBACHER-O'NAN-SCOTT THEOREM

For some interesting history, see Peter Cameron's blog at

<http://cameroncounts.wordpress.com/tag/onan-scott-theorem/>



## CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)



## CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.

## CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).

## CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).
- The lattice  $L_7$  puts many restrictions on the class of groups which could possibly represent it. Perhaps this lattice, or a modified version of it, will enable us to restrict to the class of almost simple groups.

## CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).
- The lattice  $L_7$  puts many restrictions on the class of groups which could possibly represent it. Perhaps this lattice, or a modified version of it, will enable us to restrict to the class of almost simple groups.
- Future work: Explore “interval enforceable properties of finite groups” and try to restrict to almost simple groups. Then solve the problem using the CFSG Theorem.

