

# The Finite Lattice Representation Problem

William J. DeMeo

University of Hawai'i

KMS-AMS Joint Meeting  
December 18, 2009

## Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

## Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

In particular, this shows there is no lattice-theoretic condition stronger than algebraicity satisfied by all congruence lattices.

## Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

In particular, this shows there is no lattice-theoretic condition stronger than algebraicity satisfied by all congruence lattices.

What if the lattice is finite?

## Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

In particular, this shows there is no lattice-theoretic condition stronger than algebraicity satisfied by all congruence lattices.

## What if the lattice is finite?

Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{ConA} \cong \mathbf{L}$ ?

## Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

In particular, this shows there is no lattice-theoretic condition stronger than algebraicity satisfied by all congruence lattices.

## What if the lattice is finite?

Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{ConA} \cong \mathbf{L}$ ?

status: open

age: 45+ years

# What is an algebra?

## Definition (algebra)

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where  
     $A$  is a nonempty set, called the *universe* of  $\mathbf{A}$   
     $F$  is a family of finitary operations acting on  $\mathbf{A}$

# What is an algebra?

## Definition (algebra)

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

- $A$  is a nonempty set, called the *universe* of  $\mathbf{A}$
- $F$  is a family of finitary operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is **finite** if  $|A|$  is finite.



# What is an algebra?

## Definition (algebra)

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of finitary operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is **finite** if  $|A|$  is finite.

- Examples: semigroups, groups, rings, modules, lattices,  $\ast$ -algebras (with a vector space reduct, ring reduct, and unary  $\ast$ )

# What is an algebra?

## Definition (algebra)

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of finitary operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is **finite** if  $|A|$  is finite.

- Examples: semigroups, groups, rings, modules, lattices,  $*$ -algebras (with a vector space reduct, ring reduct, and unary  $*$ )
- A **variety**  $\mathcal{K}$  of algebras is a class of (similar) algebras defined by equations. They are closed under homomorphic images, subalgebras and direct products, and in fact

$$\mathbf{V}(\mathcal{K}) = \mathbf{HSP}(\mathcal{K})$$

is the variety generated by a class  $\mathcal{K}$  of algebras.

# Examples

- A **group** is an algebra  $\mathbf{G} = \langle G, \cdot, ^{-1}, 1 \rangle$  with binary, unary, and nullary operations satisfying,  $\forall x, y, z \in G$ ,

G1:  $x \cdot (y \cdot z) \approx (x \cdot y) \cdot z$

G2:  $x \cdot 1 \approx 1 \cdot x \approx x$

G3:  $x \cdot x^{-1} \approx x^{-1} \cdot x \approx 1$


# Examples

- A **group** is an algebra  $\mathbf{G} = \langle G, \cdot, ^{-1}, 1 \rangle$  with binary, unary, and nullary operations satisfying,  $\forall x, y, z \in G$ ,
  - G1:  $x \cdot (y \cdot z) \approx (x \cdot y) \cdot z$
  - G2:  $x \cdot 1 \approx 1 \cdot x \approx x$
  - G3:  $x \cdot x^{-1} \approx x^{-1} \cdot x \approx 1$
- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$ , a partially ordered set, and binary operations:
  - $x \wedge y = \text{g.l.b.}(x, y)$  the “meet” of  $x$  and  $y$
  - $x \vee y = \text{l.u.b.}(x, y)$  the “join” of  $x$  and  $y$


# Examples

- A **group** is an algebra  $\mathbf{G} = \langle G, \cdot, ^{-1}, 1 \rangle$  with binary, unary, and nullary operations satisfying,  $\forall x, y, z \in G$ ,
  - G1:  $x \cdot (y \cdot z) \approx (x \cdot y) \cdot z$
  - G2:  $x \cdot 1 \approx 1 \cdot x \approx x$
  - G3:  $x \cdot x^{-1} \approx x^{-1} \cdot x \approx 1$
- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$ , a partially ordered set, and binary operations:
  - $x \wedge y = \text{g.l.b.}(x, y)$  the “meet” of  $x$  and  $y$
  - $x \vee y = \text{l.u.b.}(x, y)$  the “join” of  $x$  and  $y$
- *Examples of lattices:*
  - subsets of a set
  - closed subsets of a topology
  - subgroups of a group, normal subgroups of a group
  - ideals of a ring
  - submodules of a module
  - invariant subspaces of an operator or operator algebra


# Lattices



# Lattices




# Lattices




$$v \leq w$$




# Lattices



# Lattices




# Lattices



$x, y, z$  generate

# Lattices



$$v = y \wedge (x \vee z)$$

# Distributivity and Modularity

- A lattice is **distributive** if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

# Distributivity and Modularity

- A lattice is **distributive** if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

- It is **modular** if

$$x \wedge (y \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$$

# Distributivity and Modularity

- A lattice is **distributive** if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

- It is **modular** if

$$x \wedge (y \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$$

Equivalently,

$$z \leq x \quad \Rightarrow \quad x \wedge (y \vee z) = (x \wedge y) \vee z$$

# Distributivity and Modularity

- A lattice is **distributive** if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$


- It is **modular** if

$$x \wedge (y \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$$

Equivalently,

$$z \leq x \quad \Rightarrow \quad x \wedge (y \vee z) = (x \wedge y) \vee z$$

- **M<sub>3</sub>** is modular but not distributive.





# Distributivity and Modularity

- A lattice is **distributive** if

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$



- It is **modular** if

$$x \wedge (y \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$$

Equivalently,

$$z \leq x \Rightarrow x \wedge (y \vee z) = (x \wedge y) \vee z$$

- M**<sub>3</sub> is modular but not distributive.
- N**<sub>5</sub> is not even modular.



# The subspaces of a NLS

- Let  $\mathbf{X}$  be a normed linear space and let  $\text{CSub}[\mathbf{X}]$  denote the set of closed linear manifolds (subspaces).

# The subspaces of a NLS

- Let  $\mathbf{X}$  be a normed linear space and let  $\text{CSub}[\mathbf{X}]$  denote the set of closed linear manifolds (subspaces).
- Define meet to be set intersection and join to be the norm-closure of the span:

$$V_1 \wedge V_2 = V_1 \cap V_2, \quad V_1 \vee V_2 = \overline{V_1 + V_2}$$

# The subspaces of a NLS

- Let  $\mathbf{X}$  be a normed linear space and let  $\text{CSub}[\mathbf{X}]$  denote the set of closed linear manifolds (subspaces).
- Define meet to be set intersection and join to be the norm-closure of the span:

$$V_1 \wedge V_2 = V_1 \cap V_2, \quad V_1 \vee V_2 = \overline{V_1 + V_2}$$

- Then  $\mathbf{CSub}[\mathbf{X}] = \langle \text{CSub}[\mathbf{X}], \wedge, \vee \rangle$  is a lattice.

# The subspaces of a NLS

- Let  $\mathbf{X}$  be a normed linear space and let  $\mathbf{CSub}[\mathbf{X}]$  denote the set of closed linear manifolds (subspaces).
- Define meet to be set intersection and join to be the norm-closure of the span:

$$V_1 \wedge V_2 = V_1 \cap V_2, \quad V_1 \vee V_2 = \overline{V_1 + V_2}$$

- Then  $\mathbf{CSub}[\mathbf{X}] = \langle \mathbf{CSub}[\mathbf{X}], \wedge, \vee \rangle$  is a lattice.
- It is *modular* if and only if  $\mathbf{X}$  is finite dimensional.
- It is *distributive* if and only if  $\mathbf{X}$  has dimension 0 or 1.

(See e.g. Halmos, “A Hilbert Space Problem Book,” Springer, 1984.)

## Example: **Sub**[**G**]

- The **lattice of subgroups** of a group **G**

$$\mathbf{Sub}[\mathbf{G}] = \langle \mathbf{Sub}[\mathbf{G}], \subseteq \rangle = \langle \mathbf{Sub}[\mathbf{G}], \wedge, \vee \rangle$$

has universe  $\mathbf{Sub}[\mathbf{G}]$ , the set of subgroups of **G**.

# Example: **Sub[G]**

- The **lattice of subgroups** of a group **G**

$$\mathbf{Sub[G]} = \langle \mathbf{Sub[G]}, \subseteq \rangle = \langle \mathbf{Sub[G]}, \wedge, \vee \rangle$$

has universe  $\mathbf{Sub[G]}$ , the set of subgroups of **G**.


- For subgroups  $H, K \in \mathbf{Sub[G]}$ ,  
**meet** is set intersection:

$$H \wedge K = H \cap K$$

**join** is the subgroup generated by set union:

$$H \vee K = \bigcap \{J \in \mathbf{Sub[G]} \mid H \cup K \subseteq J\}$$

## Example: Hasse diagram of $\text{Sub}[D_4]$



The lattice of subgroups of the dihedral group  $D_4$ , represented as groups of rotations and reflections of a plane figure.



...ok, but is it useful?

## ...ok, but is it useful?

Lattice-theoretic information (about **Sub**[**G**]) can be used to obtain group-theoretic information (about **G**).

# ...ok, but is it useful?

Lattice-theoretic information (about **Sub**[**G**]) can be used to obtain group-theoretic information (about **G**).

## Examples:

- **G** is locally cyclic if and only if **Sub**[**G**] is distributive.
- Øystein Ore, “Structures and group theory,” *Duke Math. J.* (1937)

# ...ok, but is it useful?

Lattice-theoretic information (about **Sub**[**G**]) can be used to obtain group-theoretic information (about **G**).

## Examples:

- **G** is locally cyclic if and only if **Sub**[**G**] is distributive.  
Øystein Ore, “Structures and group theory,” *Duke Math. J.* (1937)
- Similar lattice-theoretic characterizations exist for solvable and perfect groups.  
Michio Suzuki, “On the lattice of subgroups of finite groups,”  
*Trans. AMS* (1951)  
    , “Structure of a group and the structure of its  
    lattice of subgroups,” *Springer* (1956)

## Example: equivalence relations

- The set of equivalence relations on a set is a 0-1 lattice:

$$\mathbf{Eq}(A) = \langle \text{Eq}(A), \subseteq \rangle = \langle \text{Eq}(A), \wedge, \vee \rangle$$

# Example: equivalence relations

- The set of equivalence relations on a set is a 0-1 lattice:

$$\mathbf{Eq}(A) = \langle \mathbf{Eq}(A), \subseteq \rangle = \langle \mathbf{Eq}(A), \wedge, \vee \rangle$$

- $\mathbf{Eq}(A) \subset \mathcal{P}(A \times A)$  and meet is set intersection, while join is the equivalence generated by set union:

$$\alpha \wedge \beta = \alpha \cap \beta$$

$$\alpha \vee \beta = \bigcap \{ \theta \in \mathbf{Eq}(A) \mid \alpha \leq \theta, \beta \leq \theta \}$$

# Example: equivalence relations

- The set of equivalence relations on a set is a 0-1 lattice:

$$\mathbf{Eq}(A) = \langle \text{Eq}(A), \subseteq \rangle = \langle \text{Eq}(A), \wedge, \vee \rangle$$

- $\text{Eq}(A) \subset \mathcal{P}(A \times A)$  and meet is set intersection, while join is the equivalence generated by set union:

$$\alpha \wedge \beta = \alpha \cap \beta$$

$$\alpha \vee \beta = \bigcap \{ \theta \in \text{Eq}(A) \mid \alpha \leq \theta, \beta \leq \theta \}$$


- The greatest equivalence is the *all* relation:

$$\nabla = A \times A$$

- The least equivalence is the *diagonal* relation:

$$\Delta = \{ (x, y) \in A \times A \mid x = y \}$$

# Example: Eq(4)



The lattice of equivalence relations on the set of four elements.



# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $\mathbf{A}^2$ .

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $\mathbf{A}^2$ .
- The set  $\text{Con}(\mathbf{A})$  of congruences of  $\mathbf{A}$  is a sublattice of  $\mathbf{Eq}(A)$ ,

$$\mathbf{ConA} = \langle \text{Con}(\mathbf{A}), \wedge, \vee \rangle$$

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $\mathbf{A}^2$ .
- The set  $\text{Con}(\mathbf{A})$  of congruences of  $\mathbf{A}$  is a sublattice of  $\mathbf{Eq}(A)$ ,

$$\mathbf{ConA} = \langle \text{Con}(\mathbf{A}), \wedge, \vee \rangle$$

- For groups this is the same as the lattice of normal subgroups.

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $A^2$ .
- The set  $\text{Con}(\mathbf{A})$  of congruences of  $\mathbf{A}$  is a sublattice of  $\mathbf{Eq}(A)$ ,

$$\mathbf{ConA} = \langle \text{Con}(\mathbf{A}), \wedge, \vee \rangle$$

- For groups this is the same as the lattice of normal subgroups.
- For rings this is the same as the lattice of ideals.

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $A^2$ .
- The set  $\text{Con}(\mathbf{A})$  of congruences of  $\mathbf{A}$  is a sublattice of  $\mathbf{Eq}(A)$ ,

$$\mathbf{ConA} = \langle \text{Con}(\mathbf{A}), \wedge, \vee \rangle$$

- For groups this is the same as the lattice of normal subgroups.
- For rings this is the same as the lattice of ideals.
- For most classical algebras,  $\mathbf{ConA}$  is modular.

# Congruence Lattices

- If  $f : A \rightarrow B$  is a mapping of one set to another, the relation  $\theta$  on  $A$ ,

$$x \theta y \iff f(x) = f(y)$$

is an equivalence relation.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are algebras and  $f \in \text{Hom}(\mathbf{A}, \mathbf{B})$  an algebra hom,

$$\theta = \ker f = \{(x, y) \in A^2 \mid f(x) = f(y)\}$$

is called a **congruence relation** on  $\mathbf{A}$ .

- A relation on  $\mathbf{A} = \langle A, F \rangle$  is a congruence relation iff it is an equivalence relation which is a subalgebra of  $A^2$ .
- The set  $\text{Con}(\mathbf{A})$  of congruences of  $\mathbf{A}$  is a sublattice of  $\mathbf{Eq}(A)$ ,

$$\mathbf{ConA} = \langle \text{Con}(\mathbf{A}), \wedge, \vee \rangle$$

- For groups this is the same as the lattice of normal subgroups.
- For rings this is the same as the lattice of ideals.
- For most classical algebras,  $\mathbf{ConA}$  is modular.
- For lattices, and the algebras of logic,  $\mathbf{ConA}$  is distributive.



# The finite lattice representation problem

## Definition (representable lattice)

Call a finite lattice **representable** if it is (isomorphic to) the congruence lattice of a finite algebra.

# The finite lattice representation problem

## Definition (representable lattice)

Call a finite lattice **representable** if it is (isomorphic to) the congruence lattice of a finite algebra.

## The $\leq \$1m$ question

Is every finite lattice representable?

# The finite lattice representation problem

## Definition (representable lattice)

Call a finite lattice **representable** if it is (isomorphic to) the congruence lattice of a finite algebra.

## The $\leq \$1m$ question

Is every finite lattice representable?

Equivalently, given a finite lattice **L**, does there exist a finite algebra **A** such that **ConA**  $\cong$  **L**?

# An equivalent problem in group theory

Theorem (Pálffy and Pudlák, AU 11, 1980)

*The following statements are equivalent:*

- (i) Any finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (ii) Any finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

# An equivalent problem in group theory

Theorem (Pálffy and Pudlák, AU 11, 1980)

*The following statements are equivalent:*

- (i) Any finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (ii) Any finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

A quote from MathSciNet reviews

*"In AU 11, Pálffy and Pudlák proved that...a finite lattice is representable if and only if it occurs as an interval in the subgroup lattice of a finite group."*

# An equivalent problem in group theory

Theorem (Pálffy and Pudlák, AU 11, 1980)

*The following statements are equivalent:*

- (i) Any finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (ii) Any finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

A quote from MathSciNet reviews

*"In AU 11, Pálffy and Pudlák proved that...a finite lattice is representable if and only if it occurs as an interval in the subgroup lattice of a finite group."*

False!

# Concrete representation

Theorem (Pudlák and Tůma, AU 10, 1980)

*A finite lattice can be embedded in  $Eq(X)$ , for some finite  $X$ .*

Theorem (Pudlák and Tůma, AU 10, 1980)

*A finite lattice can be embedded in  $\mathbf{Eq}(X)$ , for some finite  $X$ .*

- That is, if  $\mathbf{L}$  is any finite lattice, there exists a finite set  $X$  with

$$\mathbf{L} \cong \mathbf{L}' \leq \mathbf{Eq}(X)$$



# Concrete representation

Theorem (Pudlák and Tůma, AU 10, 1980)

*A finite lattice can be embedded in  $\mathbf{Eq}(X)$ , for some finite  $X$ .*

- That is, if  $\mathbf{L}$  is any finite lattice, there exists a finite set  $X$  with

$$\mathbf{L} \cong \mathbf{L}' \leq \mathbf{Eq}(X)$$

- Assume  $\mathbf{L}$  is itself *concretely represented* as  $\mathbf{L} \leq \mathbf{Eq}(X)$

# Concrete representation

Theorem (Pudlák and Tůma, AU 10, 1980)

*A finite lattice can be embedded in  $\mathbf{Eq}(X)$ , for some finite  $X$ .*

- That is, if  $\mathbf{L}$  is any finite lattice, there exists a finite set  $X$  with

$$\mathbf{L} \cong \mathbf{L}' \leq \mathbf{Eq}(X)$$

- Assume  $\mathbf{L}$  is itself *concretely represented* as  $\mathbf{L} \leq \mathbf{Eq}(X)$
- Define a relation  $R$  on  $X^X \times \mathbf{Eq}(X)$  as follows:

$$h R \theta \quad \Leftrightarrow \quad h(x) \theta h(y) \text{ whenever } x \theta y$$

# Concrete representation

Theorem (Pudlák and Tůma, AU 10, 1980)

*A finite lattice can be embedded in  $\mathbf{Eq}(X)$ , for some finite  $X$ .*

- That is, if  $\mathbf{L}$  is any finite lattice, there exists a finite set  $X$  with

$$\mathbf{L} \cong \mathbf{L}' \leq \mathbf{Eq}(X)$$

- Assume  $\mathbf{L}$  is itself *concretely represented* as  $\mathbf{L} \leq \mathbf{Eq}(X)$
- Define a relation  $R$  on  $X^X \times \mathbf{Eq}(X)$  as follows:

$$h R \theta \quad \Leftrightarrow \quad h(x) \theta h(y) \text{ whenever } x \theta y$$

- If  $h R \theta$  we say “ $h$  respects  $\theta$ ” or “ $\theta$  admits  $h$ ”

- Let  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$  and  $\mathcal{H} = \mathcal{P}(X^X)$  be po'd by set inclusion.

- Let  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$  and  $\mathcal{H} = \mathcal{P}(X^X)$  be po'd by set inclusion.
- Define maps  $\lambda : \mathcal{E} \rightarrow \mathcal{H}$  and  $\rho : \mathcal{H} \rightarrow \mathcal{E}$  by

$$\lambda(E) = \{h \in X^X \mid h R \theta \text{ for all } \theta \in E\} \quad (E \in \mathcal{E})$$

$$\rho(H) = \{\theta \in \text{Eq}(X) \mid h R \theta \text{ for all } h \in H\} \quad (H \in \mathcal{H})$$

- Let  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$  and  $\mathcal{H} = \mathcal{P}(X^X)$  be po'd by set inclusion.
- Define maps  $\lambda : \mathcal{E} \rightarrow \mathcal{H}$  and  $\rho : \mathcal{H} \rightarrow \mathcal{E}$  by

$$\lambda(E) = \{h \in X^X \mid h R \theta \text{ for all } \theta \in E\} \quad (E \in \mathcal{E})$$

$$\rho(H) = \{\theta \in \text{Eq}(X) \mid h R \theta \text{ for all } h \in H\} \quad (H \in \mathcal{H})$$

- $(\lambda, \rho)$  is a pair of antitone maps with

$$\rho\lambda \geq \text{id}_{\mathcal{E}} \quad \text{and} \quad \lambda\rho \geq \text{id}_{\mathcal{H}}$$

- Let  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$  and  $\mathcal{H} = \mathcal{P}(X^X)$  be po'd by set inclusion.
- Define maps  $\lambda : \mathcal{E} \rightarrow \mathcal{H}$  and  $\rho : \mathcal{H} \rightarrow \mathcal{E}$  by

$$\lambda(E) = \{h \in X^X \mid h R \theta \text{ for all } \theta \in E\} \quad (E \in \mathcal{E})$$

$$\rho(H) = \{\theta \in \text{Eq}(X) \mid h R \theta \text{ for all } h \in H\} \quad (H \in \mathcal{H})$$

- $(\lambda, \rho)$  is a pair of antitone maps with

$$\rho\lambda \geq \text{id}_{\mathcal{E}} \quad \text{and} \quad \lambda\rho \geq \text{id}_{\mathcal{H}}$$

- $(\lambda, \rho)$  is a *Galois correspondence* between  $\text{Eq}(X)$  and  $X^X$

- Let  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$  and  $\mathcal{H} = \mathcal{P}(X^X)$  be po'd by set inclusion.
- Define maps  $\lambda : \mathcal{E} \rightarrow \mathcal{H}$  and  $\rho : \mathcal{H} \rightarrow \mathcal{E}$  by

$$\lambda(E) = \{h \in X^X \mid h R \theta \text{ for all } \theta \in E\} \quad (E \in \mathcal{E})$$

$$\rho(H) = \{\theta \in \text{Eq}(X) \mid h R \theta \text{ for all } h \in H\} \quad (H \in \mathcal{H})$$

- $(\lambda, \rho)$  is a pair of antitone maps with

$$\rho\lambda \geq \text{id}_{\mathcal{E}} \quad \text{and} \quad \lambda\rho \geq \text{id}_{\mathcal{H}}$$

- $(\lambda, \rho)$  is a *Galois correspondence* between  $\text{Eq}(X)$  and  $X^X$
- Easy consequences:  
 $\rho\lambda$  and  $\lambda\rho$  are idempotent;  $\rho\lambda\rho = \rho$  and  $\lambda\rho\lambda = \lambda$ ;  
 $F \subseteq \rho\lambda(F)$ , for any set  $F \in \mathcal{E}$ .



# Closure operator

- The map  $\rho\lambda$  is idempotent, extensive, and order preserving; i.e.

$\rho\lambda$  is a *closure operator* on  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$

# Closure operator

- The map  $\rho\lambda$  is idempotent, extensive, and order preserving; i.e.

$\rho\lambda$  is a *closure operator* on  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$

- Call a set  $F \in \mathcal{E}$  **closed** provided  $\rho\lambda(F) = F$ .

# Closure operator

- The map  $\rho\lambda$  is idempotent, extensive, and order preserving; i.e.

$\rho\lambda$  is a *closure operator* on  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$

- Call a set  $F \in \mathcal{E}$  **closed** provided  $\rho\lambda(F) = F$ .
- To reiterate, for  $F \subseteq \text{Eq}(X)$ , we have

$$F \subseteq \rho\lambda(F) \subseteq \text{Eq}(X)$$

and  $F$  is **closed** iff  $\rho\lambda(F) = F$ .

We call  $F$  **dense** iff  $\rho\lambda(F) = \text{Eq}(X)$

# Closure operator

- The map  $\rho\lambda$  is idempotent, extensive, and order preserving; i.e.

$\rho\lambda$  is a *closure operator* on  $\mathcal{E} = \mathcal{P}(\text{Eq}(X))$

- Call a set  $F \in \mathcal{E}$  **closed** provided  $\rho\lambda(F) = F$ .
- To reiterate, for  $F \subseteq \text{Eq}(X)$ , we have

$$F \subseteq \rho\lambda(F) \subseteq \text{Eq}(X)$$

and  $F$  is **closed** iff  $\rho\lambda(F) = F$ .

We call  $F$  **dense** iff  $\rho\lambda(F) = \text{Eq}(X)$

- If  $\mathbf{L} \cong \mathbf{L}' \leq \mathbf{Eq}(X)$  and if  $\rho\lambda(L') = \text{Eq}(X)$ , then we say  $\mathbf{L}$  can be **densely embedded** in  $\mathbf{Eq}(X)$ .

# A density result

## Theorem

*If  $\mathbf{L} \leq \mathbf{Eq}(X)$ , then  $\mathbf{L} = \mathbf{ConA}$  for some algebra  $\mathbf{A} = \langle X, F \rangle$  if and only if  $\mathbf{L}$  is closed; that is, iff  $\rho\lambda(L) = L$ .*

# A density result

## Theorem

*If  $\mathbf{L} \leq \mathbf{Eq}(X)$ , then  $\mathbf{L} = \mathbf{ConA}$  for some algebra  $\mathbf{A} = \langle X, F \rangle$  if and only if  $\mathbf{L}$  is closed; that is, iff  $\rho\lambda(L) = L$ .*

- Much research has focused on the height two lattices  $\mathbf{M}_n$ , which are perceived as crucial for the general representation problem.

# A density result

## Theorem

*If  $\mathbf{L} \leq \mathbf{Eq}(X)$ , then  $\mathbf{L} = \mathbf{ConA}$  for some algebra  $\mathbf{A} = \langle X, F \rangle$  if and only if  $\mathbf{L}$  is closed; that is, iff  $\rho\lambda(L) = L$ .*

- Much research has focused on the height two lattices  $\mathbf{M}_n$ , which are perceived as crucial for the general representation problem.
- J.B. Nation asked how “bad” can concrete representations of  $\mathbf{M}_3$  be in terms of non-closure, and do there exist “superbad” or dense  $\mathbf{M}_3$ ’s in  $\mathbf{Eq}(X)$ ?

# A density result

## Theorem

*If  $\mathbf{L} \leq \mathbf{Eq}(X)$ , then  $\mathbf{L} = \mathbf{ConA}$  for some algebra  $\mathbf{A} = \langle X, F \rangle$  if and only if  $\mathbf{L}$  is closed; that is, iff  $\rho\lambda(L) = L$ .*

- Much research has focused on the height two lattices  $\mathbf{M}_n$ , which are perceived as crucial for the general representation problem.
- J.B. Nation asked how “bad” can concrete representations of  $\mathbf{M}_3$  be in terms of non-closure, and do there exist “superbad” or dense  $\mathbf{M}_3$ ’s in  $\mathbf{Eq}(X)$ ?

## Theorem (Snow 2009)

*The lattice  $\mathbf{Eq}(X)$  contains a proper dense  $\mathbf{M}_3$  if and only if  $|X| \geq 5$ .*



# A density result

## Theorem

If  $\mathbf{L} \leq \mathbf{Eq}(X)$ , then  $\mathbf{L} = \mathbf{ConA}$  for some algebra  $\mathbf{A} = \langle X, F \rangle$  if and only if  $\mathbf{L}$  is closed; that is, iff  $\rho\lambda(L) = L$ .

- Much research has focused on the height two lattices  $\mathbf{M}_n$ , which are perceived as crucial for the general representation problem.
- J.B. Nation asked how “bad” can concrete representations of  $\mathbf{M}_3$  be in terms of non-closure, and do there exist “superbad” or dense  $\mathbf{M}_3$ ’s in  $\mathbf{Eq}(X)$ ?

## Theorem (Snow 2009)

The lattice  $\mathbf{Eq}(X)$  contains a proper dense  $\mathbf{M}_3$  if and only if  $|X| \geq 5$ .

**Idea of proof:** Find an  $\mathbf{L} \cong \mathbf{M}_3$  in  $\mathbf{Eq}(X)$  such that every non-trivial operation in  $X^X$  violates some equivalence in the universe  $L$  of  $\mathbf{L}$ . Then  $\lambda(L)$  is trivial, so the closure  $\rho\lambda(L)$  is all of  $\mathbf{Eq}(X)$ . John Snow proved this for  $|X|$  odd.

# Another density result

Snow's result can be generalized to  $\mathbf{M}_n$  as follows:

Let  $\text{Eq}(n)$  denote the set of equivalences on an  $n$ -element set.

# Another density result

Snow's result can be generalized to  $\mathbf{M}_n$  as follows:

Let  $\text{Eq}(n)$  denote the set of equivalences on an  $n$ -element set.

**Theorem (Snow-wjd 2009)**

*For  $n \geq 1$ , the lattice  $\mathbf{Eq}(2n + 1)$  contains a dense  $\mathbf{M}_{n+2}$ .*

# Another density result

Snow's result can be generalized to  $\mathbf{M}_n$  as follows:

Let  $\text{Eq}(n)$  denote the set of equivalences on an  $n$ -element set.

**Theorem (Snow-wjd 2009)**

*For  $n \geq 1$ , the lattice  $\mathbf{Eq}(2n + 1)$  contains a dense  $\mathbf{M}_{n+2}$ .*

So, for any  $n \geq 3$ ,  $\mathbf{M}_n$  can be densely embedded in  $\mathbf{Eq}(X)$ , for some finite set  $X$ .

# A non-density result

On the other hand, we noticed that certain lattices, like  $\mathbf{N}_5$ , are never densely embedded.

# A non-density result

On the other hand, we noticed that certain lattices, like  $\mathbf{N}_5$ , are never densely embedded.

## Lemma

*Suppose  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  is a complete 0, 1-lattice. TFAE*

- (i) There is an element  $\alpha \in L \setminus \{0_L\}$  such that  $\bigvee \{\gamma \in L : \gamma \not\leq \alpha\} < 1_L$*
- (ii) There is an element  $\alpha \in L \setminus \{1_L\}$  such that  $\bigwedge \{\gamma \in L : \gamma \not\leq \alpha\} > 0_L$ .*
- (iii)  $\mathbf{L}$  is the union of a proper ideal and a proper filter.*

# A non-density result

On the other hand, we noticed that certain lattices, like  $\mathbf{N}_5$ , are never densely embedded.

## Lemma

*Suppose  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  is a complete 0, 1-lattice. TFAE*

- (i) There is an element  $\alpha \in L \setminus \{0_L\}$  such that  $\bigvee \{\gamma \in L : \gamma \not\leq \alpha\} < 1_L$*
- (ii) There is an element  $\alpha \in L \setminus \{1_L\}$  such that  $\bigwedge \{\gamma \in L : \gamma \not\leq \alpha\} > 0_L$ .*
- (iii)  $\mathbf{L}$  is the union of a proper ideal and a proper filter.*

## Theorem (wjd 2009)

*If  $\mathbf{L} \not\cong \mathbf{2}$  is a sublattice of  $\mathbf{Eq}(X)$  satisfying conditions of the lemma, then  $\lambda(L)$  contains a non-trivial unary function.*

# A non-density result

On the other hand, we noticed that certain lattices, like  $\mathbf{N}_5$ , are never densely embedded.

## Lemma

*Suppose  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  is a complete 0, 1-lattice. TFAE*

- (i) There is an element  $\alpha \in L \setminus \{0_L\}$  such that  $\bigvee \{\gamma \in L : \gamma \not\leq \alpha\} < 1_L$*
- (ii) There is an element  $\alpha \in L \setminus \{1_L\}$  such that  $\bigwedge \{\gamma \in L : \gamma \not\leq \alpha\} > 0_L$ .*
- (iii)  $\mathbf{L}$  is the union of a proper ideal and a proper filter.*

## Theorem (wjd 2009)

*If  $\mathbf{L} \not\cong \mathbf{2}$  is a sublattice of  $\mathbf{Eq}(X)$  satisfying conditions of the lemma, then  $\lambda(L)$  contains a non-trivial unary function.*

## Corollary

*If  $\mathbf{L} \not\cong \mathbf{2}$  is a lattice satisfying conditions of the lemma, then  $\mathbf{L}$  cannot be densely embedded in  $\mathbf{Eq}(X)$ .*



# More non-density consequences...

## Corollary

*If  $\mathbf{L} \not\cong \mathbf{2}$  is a finite lattice with a prime element and  $X$  is any set, then  $\mathbf{L}$  cannot be densely embedded in  $\mathbf{Eq}(X)$ .*

## Corollary

*If  $\mathbf{L} \in SD_{\wedge}$  is a finite semi-distributive lattice with  $\mathbf{L} \not\cong \mathbf{2}$ , and  $X$  is any set, then  $\mathbf{L}$  cannot be densely embedded in  $\mathbf{Eq}(X)$ .*

# Finally, a closure result

## Theorem (Snow 2009)

*Suppose  $\mathbf{L} \leq \mathbf{Eq}(X)$  is a closed sublattice and  $\mathbf{L}' \leq \mathbf{L}$  is a sublattice with universe  $A \cup B$ , where  $A = \{x \in L \mid x \leq \alpha\}$  and  $B = \{x \in L \mid x \leq \beta\}$  for some  $\alpha, \beta \in L$ . Then  $\mathbf{L}'$  is closed.*

# Finally, a closure result

## Theorem (Snow 2009)

*Suppose  $\mathbf{L} \leq \mathbf{Eq}(X)$  is a closed sublattice and  $\mathbf{L}' \leq \mathbf{L}$  is a sublattice with universe  $A \cup B$ , where  $A = \{x \in L \mid x \leq \alpha\}$  and  $B = \{x \in L \mid x \leq \beta\}$  for some  $\alpha, \beta \in L$ . Then  $\mathbf{L}'$  is closed.*

- This is another recent result of John Snow, which he proved using *primitive positive formulas*.

# Finally, a closure result

## Theorem (Snow 2009)

*Suppose  $\mathbf{L} \leq \mathbf{Eq}(X)$  is a closed sublattice and  $\mathbf{L}' \leq \mathbf{L}$  is a sublattice with universe  $A \cup B$ , where  $A = \{x \in L \mid x \leq \alpha\}$  and  $B = \{x \in L \mid x \leq \beta\}$  for some  $\alpha, \beta \in L$ . Then  $\mathbf{L}'$  is closed.*

- This is another recent result of John Snow, which he proved using *primitive positive formulas*.
- An easy consequence is that all hexagons are congruence hereditary. That is, if a hexagon is closed, so are its sublattices.

- Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a finite algebra  $\mathbf{A}$  such that  $\mathbf{L} \cong \mathbf{ConA}$ ?

# Summary

- Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a finite algebra  $\mathbf{A}$  such that  $\mathbf{L} \cong \mathbf{ConA}$ ?
- It is generally believed the answer is no.

# Summary

- Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a finite algebra  $\mathbf{A}$  such that  $\mathbf{L} \cong \mathbf{ConA}$ ?
- It is generally believed the answer is no.
- Pálffy and Pudlák translated it into one for the group theorists, but still no answer...

- Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a finite algebra  $\mathbf{A}$  such that  $\mathbf{L} \cong \mathbf{ConA}$ ?
- It is generally believed the answer is no.
- Pálffy and Pudlák translated it into one for the group theorists, but still no answer...
- The problem can be stated very concretely in terms of partitions of a set allowing us to analyze many concrete examples with the computer and locate specific representable lattices.



- Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a finite algebra  $\mathbf{A}$  such that  $\mathbf{L} \cong \mathbf{ConA}$ ?
- It is generally believed the answer is no.
- Pálffy and Pudlák translated it into one for the group theorists, but still no answer...
- The problem can be stated very concretely in terms of partitions of a set allowing us to analyze many concrete examples with the computer and locate specific representable lattices.
- In recent years, the partial results have gathered significant momentum, and there is some hope that the full solution is forthcoming.

감사합니다

 Thank You