

OVERALGEBRAS: EXPANSIONS AND EXTENSIONS OF FINITE ALGEBRAS


William DeMeo
williamdemeo@gmail.com

Iowa State University
Algebra & Combinatorics Seminar

22 Feb 2016


These slides and other resources are available at
<https://github.com/williamdemeo/Talks>

REPRESENTING LATTICES



Question: Is this a congruence lattice?


REPRESENTING LATTICES



Question: Is this a congruence lattice?


Answer: Yes! ...of which algebra?

REPRESENTING LATTICES




Question: Is this a congruence lattice?

Answer: Yes! ...of which algebra?




Question: Is this a congruence lattice?

REPRESENTING LATTICES



Question: Is this a congruence lattice?

Answer: Yes! ...of which algebra?





Question: Is this a congruence lattice?

Answer: Yes! ...but this one is harder.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Find a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.


There are infinitely many but here's an easy one:



CONSTRUCTION OF AN ALGEBRA \mathbf{A} WITH $\text{Con } \mathbf{A} \cong L_9$.


STEP 1 Find a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

There are infinitely many but here's an easy one:




- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 .
- the right regular action of S_3 on itself has generators $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ has elements

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$



CONSTRUCTION OF AN ALGEBRA \mathbf{A} WITH $\text{Con } \mathbf{A} \cong L_9$.




STEP 1 Find a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

There are infinitely many but here's an easy one:

- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 .
- the right regular action of S_3 on itself has generators $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ has elements

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$



STEP 2 Define $\mathbf{A} = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$


$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.


STEP 1 Find a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

There are infinitely many but here's an easy one:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 .
- the right regular action of S_3 on itself has generators $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ has elements

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$



STEP 2 Define $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$


$$B_2 = \{11, 12, 13, 14, 15\}.$$

STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.


STEP 1 Find a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

There are infinitely many but here's an easy one:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 .
- the right regular action of S_3 on itself has generators $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ has elements

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$



STEP 2 Define $A = B_0 \cup B_1 \cup B_2$ where


$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA \mathbf{A} WITH $\text{Con } \mathbf{A} \cong L_9$.




$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con } \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$


$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

CONSTRUCTION OF AN ALGEBRA \mathbf{A} WITH $\text{Con } \mathbf{A} \cong L_9$.




$\text{Con} \langle B, \{g_0, g_1\} \rangle$

$$\alpha = [0, 1, 2 | 3, 4, 5]$$

$$\beta = [0, 3 | 1, 4 | 2, 5]$$

$$\gamma = [0, 4 | 1, 5 | 2, 3]$$

$$\delta = [0, 5 | 1, 3 | 2, 4]$$



$\text{Con} \langle A, F_A \rangle$

$$\hat{\alpha} = [0, 1, 2, 6, 7, 11, 12 | 3, 4, 5 | 8, 9, 10, 13, 14, 15]$$

$$\alpha^* = [0, 1, 2, 6, 7, 11, 12 | 3, 4, 5 | 8, 9, 10 | 13, 14, 15]$$

$$\beta^* = [0, 3, 8 | 1, 4 | 2, 5, 15 | 6, 9 | 7, 10 | 11, 13 | 12, 14]$$

$$\gamma^* = [0, 4, 9 | 1, 5 | 2, 3, 13 | 6, 10 | 7, 8 | 11, 14 | 12, 15]$$

$$\delta^* = [0, 5, 10 | 1, 3 | 2, 4, 14 | 6, 8 | 7, 9, 11, 15 | 12, 13]$$

$$\alpha = \alpha^* \cap B^2 = \hat{\alpha} \cap B^2, \quad \beta = \beta^* \cap B^2, \quad \dots$$

EXPANSION & EXTENSION

Expanded Universe

$A = B_0 \cup B_1 \cup B_2$ where

$B_0 = \{0, 1, 2, 3, 4, 5\}$

$B_1 = \{0, 6, 7, 8, 9, 10\}$

$B_2 = \{11, 12, 13, 14, 15\}$.

New Operations

$e_0 : A \rightarrow B_0$


$e_1 : A \rightarrow B_1$

$e_2 : A \rightarrow B_2$

$s : A \rightarrow B_0,$

$ge_0 : A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0,$

for each $g \in F^B$.



WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

*Why don't β classes
of B_1 , B_2 mix?*

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$



$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

VARIATIONS ON THE SAME EXAMPLE...

- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.



VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = \llbracket \beta^*, \widehat{\beta} \rrbracket$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



THE P^5 LEMMA

LEMMA (PÁLFY-PUDLÁK, 1980)

Let $\mathbf{A} = \langle A, F \rangle$ be a unary algebra where F is a monoid.

Suppose $e \in F$ satisfies $e \circ e = e$.

Define $\mathbf{B} = \langle B, G \rangle$

$$B = e(A) \quad \text{and} \quad G = \{ef|_B \mid f \in F\}.$$

Let $|_B : \text{Con}(\mathbf{A}) \rightarrow \text{Con}(\mathbf{B})$ be the restriction mapping:

$$\theta|_B = \theta \cap B^2$$

Then $|_B$ is a surjective homomorphism (even for arbitrary meets and joins).

Péter Pál Pálfi and Pavel Pudlák: *Congruence lattices of finite algebras* AU (1980).

THE STAR MAP AND HAT MAP



star map



hat map

THE STAR MAP AND HAT MAP



star-map star chart



hat-map map hat

THE STAR MAP AND HAT MAP



STAR MAP $*$: $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is congruence generation:

$$\beta^* = \text{Cg}^{\mathbf{A}}(\beta) \quad (\forall \beta \in \text{Con } \mathbf{B})$$

HAT MAP $\hat{}$: $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is

$$\hat{\beta} = \{(x, y) \in A^2 \mid (ef(x), ef(y)) \in \beta, \forall f \in \text{Pol}_1(\mathbf{A})\}.$$

THE STAR MAP AND HAT MAP



STAR MAP $*$: $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is congruence generation:

$$\beta^* = \text{Cg}^{\mathbf{A}}(\beta) \quad (\forall \beta \in \text{Con } \mathbf{B})$$

HAT MAP $\hat{}$: $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is

$$\hat{\beta} = \{(x, y) \in A^2 \mid (ef(x), ef(y)) \in \beta, \forall f \in \text{Pol}_1(\mathbf{A})\}.$$

The hat map appears in McKenzie's "Finite Forbidden Lattices" paper (Puebla, 1982) where he gives an alternative proof of the P^5 Lemma.

RESIDUATION LEMMA

A lemma relating the three maps * , $|_B$ and $\hat{}$.

LEMMA

- (I) $^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is a **residuated mapping** with **residual** $|_B$.
- (II) $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$ is a **residuated mapping** with **residual** $\hat{}$.
- (III) For all $\alpha \in \text{Con } \mathbf{A}$, $\beta \in \text{Con } \mathbf{B}$,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \hat{\beta}.$$

In particular, $\beta^*|_B = \beta = \hat{\beta}|_B$.

ADJUNCTION LEMMA

New version of the lemma:

$$* \dashv \mid_B \dashv \widehat{}$$

That is,

LEMMA

- (I) $*$: $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is *left adjoint* to \mid_B .
- (II) \mid_B : $\text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$ is *left adjoint* to $\widehat{}$.
- (III) For all $\alpha \in \text{Con } \mathbf{A}$, $\beta \in \text{Con } \mathbf{B}$,

$$\beta = \alpha \mid_B \Leftrightarrow \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular, $\beta^* \mid_B = \beta = \widehat{\beta} \mid_B$.

PROOF OF THE P^5 LEMMA

LEMMA (PÁLFY-PUDLÁK, 1980)

The restriction mapping

$$\text{Con } \mathbf{A} \ni \alpha \mapsto \alpha|_B = \alpha \cap B^2 \in \text{Con } \mathbf{B}$$

is a complete lattice epimorphism.

NEW PROOF

Recall, for $f : X \rightarrow Y$ a monotone function on preorders X and Y , if f has a right (left) adjoint, then f preserves all joins (meets) that exist in X .

By the lemma $|_B$ has both a left and right adjoint.