

# OVERALGEBRAS: EXPANSIONS AND EXTENSIONS OF FINITE ALGEBRAS



William DeMeo  
williamdemeo@gmail.com

Iowa State University  
Algebra & Combinatorics Seminar

22 Feb 2016



*These slides and other resources are available at*  
<https://github.com/williamdemeo/Talks>

CONSTRUCTION OF AN ALGEBRA  $\mathbf{A}$  WITH  $\text{Con } \mathbf{A} \cong L_9$ .



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .


**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .


**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .

Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences


$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$


$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where


$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .




$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con } \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$


$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .




$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con } \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

$$\alpha = \alpha^* \cap B^2 = \hat{\alpha} \cap B^2, \quad \beta = \beta^* \cap B^2, \quad \dots$$

## EXTENSION & EXPANSION

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

$B_0$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \Downarrow & \Downarrow & \Downarrow & \Downarrow & \Downarrow & \Downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \Uparrow & \Uparrow & \Uparrow & \Uparrow & \Uparrow & \Uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

$B_0$

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

	$B_1$	2	5
10	9	8	4
7	6	0	3

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

$B_1$

10	9	8
7	6	0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## EXTENSION & EXPANSION


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## EXTENSION & EXPANSION

Con  $\langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

- $A = B_0 \cup B_1 \cup B_2$

- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con } \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con } \langle A, F_A \rangle$$

*Why don't the  $\beta$  classes  
of  $B_1$  and  $B_2$  mix?*

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$



$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$



## VARIATIONS ON THE SAME EXAMPLE...

- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$ .



## VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$





## THE $P^5$ LEMMA

### LEMMA (PÁLFY AND PUDLÁK)

Let  $\mathbf{A} = \langle A, F \rangle$  be a unary algebra with  $e^2 = e \in F$ .

Define  $\mathbf{B} = \langle B, G \rangle$  with

$$B = e(A) \quad \text{and} \quad G = \{ef|_B : f \in F\}.$$

Then

$$\text{Con } \mathbf{A} \ni \theta \mapsto \theta \cap B^2 \in \text{Con } \mathbf{B}$$

is a lattice epimorphism.

## RESIDUATION LEMMA

- Define  $\hat{\phantom{\beta}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\hat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

## RESIDUATION LEMMA

- Define  $\hat{\phantom{\beta}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\hat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to  $\text{Con } \mathbf{B}$ .

## RESIDUATION LEMMA

- Define  $\widehat{\phantom{x}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to  $\text{Con } \mathbf{B}$ .

### LEMMA


- (I)  $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is a residuated mapping with residual  $|_B$ .
- (II)  $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is a residuated mapping with residual  $\widehat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \widehat{\beta}|_B$ .


## THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .



## THE STRUCTURE OF THE INTERVAL $[\beta^*, \widehat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .




*More generally...*

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

## THE STRUCTURE OF THE INTERVAL $[\beta^*, \widehat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .



More generally...

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :


$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

$$\text{Then } [\beta^*, \widehat{\beta}] = \{\theta \in \mathbf{Eq(A)} : \beta^* \subseteq \theta \subseteq \widehat{\beta}\} \cong \prod_{r=1}^m (\mathbf{Eq} | T_r|)^{m-1}.$$


## SLIGHTLY MORE GENERAL EXAMPLES...

Returning to our original example, the base algebra **B** is the right regular  $S_3$ -set, and the nontrivial relations in  $\text{Con } \mathbf{B}$  are


$$\alpha = |0, 1, 2|3, 4, 5| \quad \beta = |0, 3|1, 4|2, 5| \quad \gamma = |0, 4|1, 5|2, 3| \quad \delta = |0, 5|1, 3|2, 4|$$




$$T = \{0, 1\}$$




$$T = \{0, 1, 2\}$$



$$T = \{0, 2, 3\}$$



$$T = \{0, 1, 2, 3\}$$



$$T = \{0, 2, 3, 5\}$$

$$L \cong 2^2 \times 2^2$$



## LIMITATIONS

Two limitations of the foregoing construction:

- 1 The sizes  $|T_r|$  of the partition lattice factors in

$$[\beta^*, \hat{\beta}] \cong \prod_{r=1}^m (\text{Eq}|T_r|)^{m-1}$$

are limited by the size of the blocks of  $\beta$ .

- 2 If  $\beta$  is not principal,  $[\theta^*, \hat{\theta}]$  may be non-trivial for some  $\theta \not\leq \beta$ .

## A GENERALIZATION

### THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose

$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .

# A GENERALIZATION

## THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose


$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .



## THE $P^5$ LEMMA

LEMMA (PÁLFY-PUDLÁK, 1980)

Let  $\mathbf{A} = \langle A, F \rangle$  be a unary algebra where  $F$  is a monoid.

Suppose  $e \in F$  satisfies  $e \circ e = e$ .

Define  $\mathbf{B} = \langle B, G \rangle$

$$B = e(A) \quad \text{and} \quad G = \{ef|_B \mid f \in F\}.$$

Let  $|_B : \text{Con}(\mathbf{A}) \rightarrow \text{Con}(\mathbf{B})$  be the restriction mapping:

$$\theta|_B = \theta \cap B^2$$

Then  $|_B$  is a surjective homomorphism (even for arbitrary meets and joins).



Péter Pál Pálfi and Pavel Pudlák: *Congruence lattices of finite algebras and intervals in subgroup lattices of finite groups.*

*Algebra Universalis* **11**(1), 22–27 (1980).

<http://dx.doi.org/10.1007/BF02483080>

## STAR MAP AND HAT MAP

**STAR MAP**  $^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is the congruence generation operator restricted to the set  $\text{Con } \mathbf{B}$ :

$$\beta^* = \text{Cg}^{\mathbf{A}}(\beta) \quad (\forall \beta \in \text{Con } \mathbf{B})$$

## STAR MAP AND HAT MAP

**STAR MAP**  $^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is the congruence generation operator restricted to the set  $\text{Con } \mathbf{B}$ :

$$\beta^* = \text{Cg}^{\mathbf{A}}(\beta) \quad (\forall \beta \in \text{Con } \mathbf{B})$$

**HAT MAP**  $\hat{\phantom{x}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is

$$\hat{\beta} = \{(x, y) \in A^2 \mid (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

(Used by McKenzie (1982) in an alternative proof of the  $P^5$  Lemma.)



Ralph McKenzie: *Finite forbidden lattices*.

In: Universal algebra and lattice theory (Puebla, 1982),  
*Lecture Notes in Math.*, vol. 1004, pp. 176–205. Springer, Berlin (1983).

<http://dx.doi.org/10.1007/BFb0063438>

## RESIDUATION LEMMA

A little lemma relating the three maps  $^*$ ,  $|_B$  and  $\hat{\phantom{x}}$ .

### LEMMA

- (I)  $^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is a **residuated mapping** with **residual**  $|_B$ .
- (II)  $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is a **residuated mapping** with **residual**  $\hat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \hat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \hat{\beta}|_B$ .

## ADJUNCTION LEMMA

New version (of the little lemma):

$$* \dashv \mid_B \dashv \widehat{\phantom{x}}$$

### LEMMA

- (I)  $*$  :  $\text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is **left adjoint** to  $\mid_B$ .
- (II)  $\mid_B$  :  $\text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is **left adjoint** to  $\widehat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \iff \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \widehat{\beta}|_B$ .



## PROOF OF THE $P^5$ LEMMA

### LEMMA (PÁLFY-PUDLÁK, 1980)

*The restriction mapping*

$$\text{Con } \mathbf{A} \ni \alpha \mapsto \alpha|_B = \alpha \cap B^2 \in \text{Con } \mathbf{B}$$

*is a complete lattice epimorphism.*

### PROOF.


Recall, for  $f : X \rightarrow Y$  a monotone function on preorders  $X, Y$ , if  $f$  has a right (left) adjoint, then  $f$  preserves all joins (meets) existing in  $X$ .

By the little lemma  $|_B$  has both a left and right adjoint.




## THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .



# THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .




*More generally...*

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

## THE STRUCTURE OF THE INTERVAL $[\beta^*, \widehat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .



More generally...

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :


$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

$$\text{Then } [\beta^*, \widehat{\beta}] = \{\theta \in \mathbf{Eq(A)} : \beta^* \subseteq \theta \subseteq \widehat{\beta}\} \cong \prod_{r=1}^m (\mathbf{Eq} | T_r|)^{m-1}.$$


## SLIGHTLY MORE GENERAL EXAMPLES...

Returning to our original example, the base algebra **B** is the right regular  $S_3$ -set, and the nontrivial relations in  $\text{Con } \mathbf{B}$  are


$$\alpha = |0, 1, 2|3, 4, 5| \quad \beta = |0, 3|1, 4|2, 5| \quad \gamma = |0, 4|1, 5|2, 3| \quad \delta = |0, 5|1, 3|2, 4|$$




$$T = \{0, 1\}$$




$$T = \{0, 1, 2\}$$



$$T = \{0, 2, 3\}$$



$$T = \{0, 1, 2, 3\}$$



$$T = \{0, 2, 3, 5\}$$

$$L \cong 2^2 \times 2^2$$

## LIMITATIONS

Two limitations of the foregoing construction:

- 1 The sizes  $|T_r|$  of the partition lattice factors in

$$[\beta^*, \hat{\beta}] \cong \prod_{r=1}^m (\text{Eq}|T_r|)^{m-1}$$

are limited by the size of the blocks of  $\beta$ .

- 2 If  $\beta$  is not principal,  $[\theta^*, \hat{\theta}]$  may be non-trivial for some  $\theta \not\leq \beta$ .

## A GENERALIZATION

### THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose

$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .

# A GENERALIZATION

## THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose

$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .

