

# The Finite Lattice Representation Problem

William DeMeo

joint work with

Ralph Freese, Peter Jipsen, Bill Lampe, J.B. Nation

University of Hawai'i at Mānoa

ARCS Presentation

April 23, 2011

I am sorry...

I am sorry...  
...this talk is about math.

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where  
     $A$  is a nonempty set, called the *universe* of  $\mathbf{A}$   
     $F$  is a family of operations acting on  $\mathbf{A}$

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is called **finite** if  $|A|$  is finite.

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is called **finite** if  $|A|$  is finite.

- **Example:** The set of integers

$$A = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is called **finite** if  $|A|$  is finite.

- **Example:** The set of integers

$$A = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

along with the operations  $F = \{+, -, \times\}$ .



# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is called **finite** if  $|A|$  is finite.

- **Example:** The set of integers

$$A = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

along with the operations  $F = \{+, -, \times\}$ .

*Question:* Is  $\div$  an operation on this set,  $A$ ?

# What is an algebra?

...an utterly fundamental object in mathematics.

## Definition

An **algebra**  $\mathbf{A}$  is an ordered pair  $\mathbf{A} = \langle A, F \rangle$  where

$A$  is a nonempty set, called the *universe* of  $\mathbf{A}$

$F$  is a family of operations acting on  $\mathbf{A}$

An algebra  $\langle A, F \rangle$  is called **finite** if  $|A|$  is finite.

- **Example:** The set of integers

$$A = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

along with the operations  $F = \{+, -, \times\}$ .

- **Other Examples:** semigroups, groups, quasigroups, rings, modules, lattices, Boolean algebras,  $*$ -algebras, etc.

# Examples

- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$  and binary operations:

# Examples

- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$  and binary operations:

$x \wedge y = \text{g.l.b.}(x, y)$     the “meet” of  $x$  and  $y$

$x \vee y = \text{l.u.b.}(x, y)$     the “join” of  $x$  and  $y$

# Examples

- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$  and binary operations:

$x \wedge y = \text{g.l.b.}(x, y)$     the “meet” of  $x$  and  $y$

$x \vee y = \text{l.u.b.}(x, y)$     the “join” of  $x$  and  $y$

- *Examples of lattices:*

- subsets of a set
- closed subsets of a topology
- subgroups of a group, normal subgroups of a group
- ideals of a ring
- submodules of a module
- invariant subspaces of an operator or operator algebra


# Examples

- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$  and binary operations:

$x \wedge y = \text{g.l.b.}(x, y)$  the “meet” of  $x$  and  $y$

$x \vee y = \text{l.u.b.}(x, y)$  the “join” of  $x$  and  $y$

- Examples of lattices:*




# Examples

- A **lattice** is an algebra  $\mathbf{L} = \langle L, \wedge, \vee \rangle$  with universe  $L$  and binary operations:


$x \wedge y = \text{g.l.b.}(x, y)$  the “meet” of  $x$  and  $y$

$x \vee y = \text{l.u.b.}(x, y)$  the “join” of  $x$  and  $y$

- Examples of lattices:*




# Lattices






# Lattices




# Lattices




$$v \leq w$$


# Lattices



# Lattices




# Lattices



$x, y, z$  generate

# Lattices



$$v = y \wedge (x \vee z)$$

# How do we study algebras?

## The Key Features

Given an algebra  $\mathbf{A}$ , there are three other important algebras which enable us to characterize and understand  $\mathbf{A}$ . These are...

# How do we study algebras?

## The Key Features

Given an algebra  $\mathbf{A}$ , there are three other important algebras which enable us to characterize and understand  $\mathbf{A}$ . These are...

- **SubA**, the lattice of [subalgebras](#) of  $\mathbf{A}$ .



# How do we study algebras?

## The Key Features

Given an algebra  $\mathbf{A}$ , there are three other important algebras which enable us to characterize and understand  $\mathbf{A}$ . These are...

- **SubA**, the lattice of [subalgebras](#) of  $\mathbf{A}$ .
- **EndA** and **AutA**, the [endomorphisms](#) and [automorphisms](#) of  $\mathbf{A}$ .

# How do we study algebras?

## The Key Features

Given an algebra  $\mathbf{A}$ , there are three other important algebras which enable us to characterize and understand  $\mathbf{A}$ . These are...

- **SubA**, the lattice of **subalgebras** of  $\mathbf{A}$ .
- **EndA** and **AutA**, the **endomorphisms** and **automorphisms** of  $\mathbf{A}$ .
- **ConA**, the lattice of **congruence relations** of  $\mathbf{A}$ .

# How do we study algebras?

## The Key Features

Given an algebra  $\mathbf{A}$ , there are three other important algebras which enable us to characterize and understand  $\mathbf{A}$ . These are...

- **SubA**, the lattice of **subalgebras** of  $\mathbf{A}$ .
- **EndA** and **AutA**, the **endomorphisms** and **automorphisms** of  $\mathbf{A}$ .
- **ConA**, the lattice of **congruence relations** of  $\mathbf{A}$ .


...okay, there are four.

# Congruence Relations

A **congruence** of an algebra  $\mathbf{A} = \langle A, F \rangle$  is a partition of the set  $A$  into blocks that are preserved by the operations in  $F$ .

# Congruence Relations


A **congruence** of an algebra  $\mathbf{A} = \langle A, F \rangle$  is a partition of the set  $A$  into blocks that are preserved by the operations in  $F$ .



$$\theta_1 = (a_0, a_1, \dots | a_2, a_8, \dots | a_3 \dots)$$

# Congruence Relations


A **congruence** of an algebra  $\mathbf{A} = \langle A, F \rangle$  is a partition of the set  $A$  into blocks that are preserved by the operations in  $F$ .



$$\theta_1 = (a_0, a_1, \dots | a_2, a_8, \dots | a_3 \dots)$$

# Congruence Relations


A **congruence** of an algebra  $\mathbf{A} = \langle A, F \rangle$  is a partition of the set  $A$  into blocks that are preserved by the operations in  $F$ .



$$\theta_1 = (a_0, a_1, \dots | a_2, a_8, \dots | a_3 \dots)$$

# Congruence Relations

A **congruence** of an algebra  $\mathbf{A} = \langle A, F \rangle$  is a partition of the set  $A$  into blocks that are preserved by the operations in  $F$ .



$$\theta_2 = (a_0, a_1, a_2, \dots | a_3, \dots | \dots), \quad \theta_1 = (a_0, a_1, \dots | a_2, a_8, \dots | a_3 \dots)$$




# Congruence Decompositions

We know an algebra by the congruences it keeps.


**A**

The shape/structure  
of **ConA** tells us  
how the algebra can  
be decomposed.



# Congruence Decompositions

We know an algebra by the congruences it keeps.



# The Problem

What are the possible shapes of congruence lattices?

# The Problem

What are the possible shapes of congruence lattices?

Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

# The Problem

What are the possible shapes of congruence lattices?

Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

Thus, there is essentially no restriction on the shape of a congruence lattice of an *infinite* algebra.

# The Problem

What are the possible shapes of congruence lattices?

Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

Thus, there is essentially no restriction on the shape of a congruence lattice of an *infinite* algebra.

What if the algebra is finite?

Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{ConA} \cong \mathbf{L}$ ?

# The Problem

What are the possible shapes of congruence lattices?

Theorem (Grätzer-Schmidt, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

Thus, there is essentially no restriction on the shape of a congruence lattice of an *infinite* algebra.


What if the algebra is finite?

Problem: Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{ConA} \cong \mathbf{L}$ ?

status: open


age: 50+ years

# Known Results: lattices of size $\leq 6$ are congruence lattices.





# Known Results: lattices of size $\leq 6$ are congruence lattices.



Aschbacher recently found the last two of these as intervals in subgroup lattices.

**Theorem:** *Every lattice with at most 6 elements is a congruence lattice of a finite algebra.*

# Recent Results


LATTICES OF SIZE  $\leq 7$  NOT YET KNOWN TO BE CONGRUENCE  
LATTICES OF FINITE ALGEBRAS



9\*



10\*



11\*



13



16\*



17\*



18\*



19\*



20\*



21\*

Figure courtesy of Peter Jipsen.

# Recent Results


LATTICES OF SIZE  $\leq 7$  NOT YET KNOWN TO BE CONGRUENCE  
LATTICES OF FINITE ALGEBRAS




9\*




10\*




11\*



13



16\*



17\*



18\*



19\*



20\*



21\*

# Recent Results


LATTICES OF SIZE  $\leq 7$  NOT YET KNOWN TO BE CONGRUENCE  
LATTICES OF FINITE ALGEBRAS




9\*




10\*




11\*



13




16\*




17\*




18\*



19\*



20\*



21\*

# Recent Results


LATTICES OF SIZE  $\leq 7$  NOT YET KNOWN TO BE CONGRUENCE  
LATTICES OF FINITE ALGEBRAS




9\*




10\*




11\*



13




16\*



17\*




18\*



19\*



20\*



21\*

# Recent Results


LATTICES OF SIZE  $\leq 7$  NOT YET KNOWN TO BE CONGRUENCE  
LATTICES OF FINITE ALGEBRAS



9\*



10\*




11\*




13




16\*



17\*



18\*



19\*



20\*



21\*

감사합니다

 Thank You

