# THE FINITE LATTICE REPRESENTATION PROBLEM AND INTERVALS IN SUBGROUP LATTICES PART II

#### William DeMeo

williamdemeo@gmail.com

joint work with

Ralph Freese, Peter Jipsen, Bill Lampe, J.B. Nation


University of South Carolina

Algebra & Logic Seminar September 28, 2012

These slides and other resources are available at  $\label{eq:http://www.math.sc.edu/~demeow/FLRP.html} \longrightarrow$ 




## Contruction of an algebra ${\bf A}$ with Con ${\bf A}\cong L_9$ .





## Contruction of an algebra $\mathbf{A}$ with Con $\mathbf{A} \cong \mathbf{L}_9$ .

STEP 1 Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\operatorname{Con} \mathbf{B} \cong M_4$ .






## Contruction of an algebra $\mathbf{A}$ with Con $\mathbf{A} \cong L_9$ .

STEP 1 Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice Con  $\mathbf{B} \cong M_4$ .

#### Example:

- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0=(0,4)(1,3)(2,5)$  and  $g_1=(0,1,2)(3,4,5)$  generate this action group, the image of  $S_3\hookrightarrow S_6$ .
- $\operatorname{Con} \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \ \beta = |03|14|25|, \gamma = |04|15|23|, \ \delta = |05|13|24|.$$






## Contruction of an algebra **A** with Con $\mathbf{A} \cong L_9$ .

STEP 1 Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\operatorname{Con} \mathbf{B} \cong M_4$ .

#### Example:





• 
$$g_0 = (0, 4)(1, 3)(2, 5)$$
 and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .

• Con  $\langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |{\tt 012|345|}, \; \beta = |{\tt 03|14|25|}, \gamma = |{\tt 04|15|23|}, \; \delta = |{\tt 05|13|24|}.$$

**Goal:** expand **B** to an algebra **A** that has  $\alpha$  "doubled" in Con **A**.






## Contruction of an algebra $\mathbf{A}$ with Con $\mathbf{A} \cong L_9$ .

STEP 1 Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\operatorname{Con} \mathbf{B} \cong M_4$ .

#### Example:




• 
$$g_0 = (0, 4)(1, 3)(2, 5)$$
 and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .


• Con  $\langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences


$$\alpha = |{\tt 012|345|}, \; \beta = |{\tt 03|14|25|}, \gamma = |{\tt 04|15|23|}, \; \delta = |{\tt 05|13|24|}.$$





$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$
 
$$B_1 = \{0, 6, 7, 8, 9, 10\}$$
 
$$B_2 = \{11, 12, 2, 13, 14, 15\}.$$






## Contruction of an algebra $\mathbf{A}$ with Con $\mathbf{A} \cong L_9$ .

STEP 1 Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\operatorname{Con} \mathbf{B} \cong M_4$ .

#### Example:



• 
$$g_0 = (0, 4)(1, 3)(2, 5)$$
 and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .

• Con  $\langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \ \beta = |03|14|25|, \gamma = |04|15|23|, \ \delta = |05|13|24|.$$





Con B

Con A

STEP 2 Since  $\alpha = Cg^{B}(0,2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where


$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 2, 13, 14, 15\}.$$

STEP 3 Define unary operations  $e_0$ ,  $e_1$ ,  $e_2$ , s,  $g_0e_0$ , and  $g_1e_0$ .

## Contruction of an algebra **A** with Con $\mathbf{A} \cong L_9$ .




$$\operatorname{Con}\left\langle B,\left\{ g_{0},g_{1}\right\} \right\rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



 $\operatorname{Con} \langle A, F_A \rangle$ 

$$\widehat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$


$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## Contruction of an algebra **A** with Con $\mathbf{A} \cong L_9$ .



$$\begin{array}{c}
\widehat{\alpha} \\
\alpha^* \\
\alpha^*
\end{array}$$

$$\begin{array}{c}
\beta^* \\
0_A
\end{array}$$

$$\operatorname{Con}\left\langle B,\left\{ g_{0},g_{1}\right\} \right\rangle$$

 $\alpha = [0, 1, 2|3, 4, 5]$ 

Con 
$$\langle A, F_A \rangle$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0,4|1,5|2,3|$$

$$\delta = |0,5|1,3|2,4|$$

$$\widehat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

$$\alpha = \alpha^* \cap B^2 = \widehat{\alpha} \cap B^2, \quad \beta = \beta^* \cap B^2, \quad \dots$$

$$\text{Con}\,\langle B,\{g_0,g_1\}\rangle$$

$$lpha = |0,1,2|3,4,5|$$
 $eta = |0,3|1,4|2,5|$ 
 $\gamma = |0,4|1,5|2,3|$ 
 $\delta = |0,5|1,3|2,4|$ 

$$\operatorname{Con}\left\langle B,\left\{ g_{0},g_{1}\right\} \right\rangle$$


$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con}\,\langle \textbf{\textit{B}},\{\textbf{\textit{g}}_0,\textbf{\textit{g}}_1\}\rangle$$


$$\begin{split} \alpha &= |0,1,2|3,4,5| \\ \beta &= |0,3|1,4|2,5| \\ \gamma &= |0,4|1,5|2,3| \\ \delta &= |0,5|1,3|2,4| \end{split}$$

$$\beta = \begin{bmatrix} 0 & 1 & 2 \\ 3 & 4 & 5 \end{bmatrix}$$

$$\operatorname{Con}\left\langle B,\left\{ g_{0},g_{1}\right\} \right\rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$
 $\beta = |0, 3|1, 4|2, 5|$ 
 $\gamma = |0, 4|1, 5|2, 3|$ 


$$\delta=|0,5|1,3|2,4|$$



$$\operatorname{Con}\left\langle B,\left\{ g_{0},g_{1}\right\} \right\rangle$$

$$lpha = |0, 1, 2|3, 4, 5|$$
 $eta = |0, 3|1, 4|2, 5|$ 
 $\gamma = |0, 4|1, 5|2, 3|$ 

$$\delta = |0, 5|1, 3|2, 4|$$



$$\operatorname{Con}\langle B,\{g_0,g_1\}\rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$


$$\delta = |0, 5|1, 3|2, 4|$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$


$$\alpha = |0, 1, 2|3, 4, 5|$$
 
$$\beta = |0, 3|1, 4|2, 5|$$
 
$$\gamma = |0, 4|1, 5|2, 3|$$
 
$$\delta = |0, 5|1, 3|2, 4|$$

$$\bullet \ A=B_0\cup B_1\cup B_2$$



$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



10


 $B_2$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

IIIII

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$
 $B_0 = \{ 1 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$s: A \rightarrow B_0$$

$$B_2$$

$$A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

 $B_1 = \{ 0 \ 6 \ 7 \ 8 \ 9 \ 10 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$


$$S: A \rightarrow B_0$$

$$D_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$D_1 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$S: A \rightarrow B_0$$

$$D_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$D_1 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow$$


$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$ 

 $\delta = [0, 5|1, 3|2, 4]$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$


$$\downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$\begin{aligned} &\operatorname{Con} \left\langle B, \{g_0, g_1\} \right\rangle \\ &\alpha = |0, 1, 2|3, 4, 5| \\ &\beta = |0, 3|1, 4|2, 5| \\ &\gamma = |0, 4|1, 5|2, 3| \\ &\delta = |0, 5|1, 3|2, 4| \end{aligned}$$



 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $S$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$B_2 = \{ \, 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \, \}$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$lpha = |0,1,2|3,4,5|$$
 $eta = |0,3|1,4|2,5|$ 
 $\gamma = |0,4|1,5|2,3|$ 

 $\delta = [0, 5|1, 3|2, 4]$ 

Con  $\langle B, \{g_0, g_1\} \rangle$ 

 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$S: A \rightarrow B_0$$

$$D_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$D_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$s: A \rightarrow B_0$$

$$B_2:$$
 $ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$ 

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\begin{split} \alpha &= |0,1,2|3,4,5| \\ \beta &= |0,3|1,4|2,5| \\ \gamma &= |0,4|1,5|2,3| \\ \delta &= |0,5|1,3|2,4| \end{split}$$



- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$


$$\delta = |0, 5|1, 3|2, 4|$$

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



10 }

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 


 $ae_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$ 

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \}$ 

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ae_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$


Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



# • $A = B_0 \cup B_1 \cup B_2$ Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0\colon A\stackrel{e_0}{\twoheadrightarrow} B_0\stackrel{g}{\to} B_0$$




$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

 $B_1 = \{ 0 \ 6 \ 7 \ 8 \ 9 \ 10 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations


$$e_0$$
:  $A \rightarrow B_0$   
 $e_1$ :  $A \rightarrow B_1$   
 $e_2$ :  $A \rightarrow B_2$   
 $s$ :  $A \rightarrow B_0$ 

 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \}$ 

10 }

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

 $B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0$$
:  $A woheadrightarrow B_0$   
 $e_1$ :  $A woheadrightarrow B_1$   
 $e_2$ :  $A woheadrightarrow B_2$   
 $s$ :  $A woheadrightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 




- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$




- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

 $\delta = [0, 5|1, 3|2, 4]$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$ 

 $\gamma = |0, 4|1, 5|2, 3|$   $\delta = |0, 5|1, 3|2, 4|$ 


- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

 $\delta = [0, 5|1, 3|2, 4]$ 



- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$lpha = |0, 1, 2|3, 4, 5|$$
  
 $eta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$ 


 $\delta = [0, 5|1, 3|2, 4]$ 

• 
$$A = B_0 \cup B_1 \cup B_2$$

Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$



$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

 $\delta = [0, 5|1, 3|2, 4]$ 

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

 $ae_0: A \stackrel{e_0}{\rightarrow} B_0 \stackrel{g}{\rightarrow} B_0$ 

$$B_2$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

 $\delta = [0, 5|1, 3|2, 4]$ 

- B<sub>1</sub>
- 10 9 8 7 6 0

- $\bullet \ A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \ \}$$

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$

Con 
$$\langle B, \{g_0, g_1\} \rangle$$
  
 $\alpha = |0, 1, 2|3, 4, 5|$   
 $\beta = |0, 3|1, 4|2, 5|$   
 $\gamma = |0, 4|1, 5|2, 3|$   
 $\delta = |0, 5|1, 3|2, 4|$ 



- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
 $e_1: A \rightarrow B_1$ 
 $e_2: A \rightarrow B_2$ 
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $\bullet \ A=B_0\cup B_1\cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$
  
 $e_1: A \rightarrow B_1$   
 $e_2: A \rightarrow B_2$   
 $s: A \rightarrow B_0$ 

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$
 $B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$ 
 $B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$ 

$$ge_0: A \stackrel{e_0}{\twoheadrightarrow} B_0 \stackrel{g}{\rightarrow} B_0$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = [0, 1, 2|3, 4, 5]$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$


$$\operatorname{Con} \left\langle B, \left\{ g_0, g_1 \right\} \right\rangle$$

$$\alpha = |\mathbf{0}, \mathbf{1}, \mathbf{2}|\mathbf{3}, \mathbf{4}, \mathbf{5}|$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta=|0,5|1,3|2,4|$$



$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$


$$\text{Con}\,\langle \textbf{\textit{B}}, \{\textbf{\textit{g}}_0,\textbf{\textit{g}}_1\}\rangle$$

$$\alpha = |\mathbf{0}, \mathbf{1}, \mathbf{2}|\mathbf{3}, \mathbf{4}, \mathbf{5}|$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$

$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$
 $\beta = |0, 3|1, 4|2, 5|$ 
 $\gamma = |0, 4|1, 5|2, 3|$ 

 $\delta = |0, 5|1, 3|2, 4|$ 

Con 
$$\langle A, F_A \rangle$$


$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$

$$\text{Con}\,\langle \textbf{\textit{B}}, \{\textbf{\textit{g}}_0,\textbf{\textit{g}}_1\}\rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$
  
 $\beta = |0, 3|1, 4|2, 5|$ 

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



Con 
$$\langle A, F_A \rangle$$

$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15| \\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15| \\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14| \\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15| \\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$


$$\operatorname{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |\mathbf{0}, \mathbf{1}, \mathbf{2}|\mathbf{3}, \mathbf{4}, \mathbf{5}|$$

$$\beta = |0,3|1,4|2,5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$





Con 
$$\langle A, F_A \rangle$$

Why don't the  $\beta$  classes of  $B_1$  and  $B_2$  mix?

$$\begin{split} \widehat{\alpha} &= |0,1,2,6,7,11,12|3,4,5|8,9,10,13,14,15|\\ \alpha^* &= |0,1,2,6,7,11,12|3,4,5|8,9,10|13,14,15|\\ \beta^* &= |0,3,8|1,4|2,5,15|6,9|7,10|11,13|12,14|\\ \gamma^* &= |0,4,9|1,5|2,3,13|6,10|7,8|11,14|12,15|\\ \delta^* &= |0,5,10|1,3|2,4,14|6,8|7,9,11,15|12,13| \end{split}$$

• Suppose we want  $\beta = \mathrm{Cg^B}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$ 






- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:


$$A = B_0 \cup B_1 \cup B_2$$
 where


$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$



- Suppose we want  $\beta = \operatorname{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$
 
$$B_0 = \{0, 1, 2, 3, 4, 5\}$$
 
$$B_1 = \{0, 6, 7, 8, 9, 10\}$$
 
$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$








- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2$$
 where

$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$










- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2$$
 where

$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$










- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2$$
 where

$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$










- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2$$
 where

$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$










- Suppose we want  $\beta = \mathrm{Cg}^{\mathbf{B}}(0,3) = |0,3|2,5|1,4|$  to have non-trivial inverse image  $\beta|_{\mathcal{B}}^{-1} = [\beta^*,\widehat{\beta}].$
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2$$
 where

$$\begin{split} B_0 &= \{0,1,2,3,4,5\} \\ B_1 &= \{0,6,7,8,9,10\} \\ B_2 &= \{11,12,13,3,14,15\}. \end{split}$$







### RESIDUATION LEMMA

• Define  $\widehat{\phantom{a}}$  : Con  ${\bf B} \to {\rm Con}\, {\bf A}$  by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

#### RESIDUATION LEMMA

• Define  $\widehat{\ }$  : Con  ${f B} 
ightarrow {
m Con}\, {f A}$  by

$$\widehat{\beta} = \{(x,y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in Pol_1(\mathbf{A})\}.$$

• For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$^*$$
: Con  ${f B} 
ightarrow$  Con  ${f A}$ 

is the congruence generation operator restricted to  $\operatorname{Con} \boldsymbol{B}$ .

#### RESIDUATION LEMMA

• Define  $\widehat{\phantom{a}}$  : Con  ${\bf B} \to {\rm Con}\, {\bf A}$  by

$$\widehat{\beta} = \{(x,y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

• For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$^*$$
: Con  ${f B} 
ightarrow$  Con  ${f A}$ 

is the congruence generation operator restricted to  $\operatorname{Con} \boldsymbol{B}$ .

#### LEMMA


- (I) \* : Con  $\mathbf{B} \to \operatorname{Con} \mathbf{A}$  is a residuated mapping with residual  $|_{\mathcal{B}}$ .
- (II)  $|_{B} : \operatorname{Con} \mathbf{A} \to \operatorname{Con} \mathbf{B}$  is a residuated mapping with residual  $\hat{\ }$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}, \ \beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_{\mathcal{B}} \quad \Leftrightarrow \quad \beta^* \leqslant \alpha \leqslant \widehat{\beta}.$$

In particular,  $\beta^*|_{B} = \beta = \widehat{\beta}|_{B}$ .


# The structure of the interval $[\beta^*, \widehat{\beta}] \leq \mathbf{Con} \mathbf{A}$ .

• If  $\beta \in \operatorname{Con} \mathbf{B}$  is a coatom of  $\operatorname{Con} \mathbf{B}$  with m congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\operatorname{Con} \mathbf{A}$  is  $\mathbf{2}^{m-1}$ .



# The structure of the interval $[\beta^*, \widehat{\beta}] \leq \mathbf{Con} \mathbf{A}$ .

• If  $\beta \in \operatorname{Con} \mathbf{B}$  is a coatom of  $\operatorname{Con} \mathbf{B}$  with m congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\operatorname{Con} \mathbf{A}$  is  $\mathbf{2}^{m-1}$ .




More generally...

- Suppose  $\beta \in \text{Con } \mathbf{B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by T<sub>r</sub> the set of intersection points in the r-th block of β:

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta$$

# The structure of the interval $[\beta^*, \widehat{\beta}] \leqslant \mathbf{Con} \mathbf{A}$ .

• If  $\beta \in \operatorname{Con} \mathbf{B}$  is a coatom of  $\operatorname{Con} \mathbf{B}$  with m congruence classes then the interval  $[\beta^*, \widehat{\beta}]$  in  $\operatorname{Con} \mathbf{A}$  is  $\mathbf{2}^{m-1}$ .



More generally...

- Suppose  $\beta \in \text{Con } \mathbf{B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the r-th block of  $\beta$ :


$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta$$

Then 
$$[\beta^*, \widehat{\beta}] = \{\theta \in \text{Eq}(A) : \beta^* \subseteq \theta \subseteq \widehat{\beta}\} \cong \prod_{r=1}^m (\text{Eq}|T_r|)^{m-1}.$$

#### SLIGHTLY MORE GENERAL EXAMPLES...

Returning to our original example, the base algebra  ${\bf B}$  is the right regular  $S_3$ -set, and the nontrivial relations in Con  ${\bf B}$  are

$$\alpha = [0, 1, 2|3, 4, 5]$$
  $\beta = [0, 3|1, 4|2, 5]$   $\gamma = [0, 4|1, 5|2, 3]$   $\delta = [0, 5|1, 3|2, 4]$ 



#### LIMITATIONS

Two limitations of the foregoing construction:

• The sizes  $|T_r|$  of the partition lattice factors in

$$[\beta^*,\widehat{\beta}]\cong\prod_{r=1}^m(\mathrm{Eq}|T_r|)^{m-1}$$

are limited by the size of the blocks of  $\beta$ .

**4** If  $\beta$  is not principal,  $[\theta^*, \hat{\theta}]$  may be non-trivial for some  $\theta \not \geq \beta$ .

## A GENERALIZATION

#### **THEOREM**

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose

$$\beta = \mathrm{Cg}^{\mathbf{B}}((a_1, b_1), \ldots, (a_{K-1}, b_{K-1}))$$

has m blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leqslant \operatorname{Con} \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \ngeq \beta$  in Con **A**.

## A GENERALIZATION

#### **THEOREM**

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose

$$\beta = \mathrm{Cg}^{\mathbf{B}}((a_1, b_1), \ldots, (a_{K-1}, b_{K-1}))$$

has m blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leqslant \operatorname{Con} \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not \geqslant \beta$  in Con **A**.

