

Constraint Satisfaction Problems, and Universal Algebra

Cliff Bergman and William DeMeo
Iowa State University
`williamdemeo@gmail.com`

April 15, 2016

What is a CSP?

Informally, a **C**onstraint **S**atisfaction **P**roblem consists of

- a list of variables ranging over a finite domain and
- a set of constraints on those variables.

Problem: can we assign values to all the variables so that all of the constraints are satisfied?

Examples

A system of linear equations is a CSP

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m$$

Also, a system of nonlinear equations is a CSP


$$a_{11}x_1^2x_3 + a_{12}x_2x_3x_7 + \cdots + a_{1n}x_4x_n^3 = b_1$$

$$a_{21}x_2x_5 + a_{22}x_2 \quad + \cdots + a_{2n}x_4^3 = b_2$$

$$\vdots$$

$$a_{m1}x_3x_5x_8 + a_{m2}x_2 \quad + \cdots + a_{mn}x_n = b_m$$

For a fixed k , determining whether a graph is k -colorable is a CSP



Given a propositional formula $\varphi(x_1, \dots, x_n)$, determine whether φ is satisfiable

$$\varphi(x, y, z) = (x \vee y \vee z') \wedge (x' \vee y \vee z')$$

then

$$\varphi(0, 0, 1) = 1$$

Algorithms

There is an efficient algorithm (Gaussian elimination) for solving any linear system. That is

There is an algorithm that accepts as **input** a linear system and decides whether that system has a solution. The running time of the algorithm is bounded above by $f(s)$ where f is a polynomial and s is the size of the system.

Algorithms

There is an efficient algorithm (Gaussian elimination) for solving any linear system. That is

There is an algorithm that accepts as **input** a linear system and decides whether that system has a solution. The running time of the algorithm is bounded above by $f(s)$ where f is a polynomial and s is the size of the system.

The particular system is an **instance** of the **problem**
LINEAR SYSTEM

Similarly

There is an algorithm that accepts as input a graph and decides whether the graph is 2-colorable.

Running time bounded by $f(s)$, a polynomial in size s .

The graph is an **instance** of the **problem**
2-COLORABILITY.

Similarly

There is an algorithm that accepts as input a graph and decides whether the graph is 2-colorable.

Running time bounded by $f(s)$, a polynomial in size s .

The graph is an **instance** of the **problem** 2-COLORABILITY.

There is an algorithm that accepts as input a formula, $\varphi = \varphi_1 \wedge \varphi_2 \wedge \cdots \wedge \varphi_k$, each φ_i bijunctive, and decides whether φ is satisfiable.

Running time bounded by $f(k)$, a polynomial in size k .

φ is an **instance** of the **problem** 2-SAT.

We say these algorithms run in **polynomial time**.

No polynomial-time algorithm is known for, NONLINEAR SYSTEM, 3-COLORABILITY, or 3-SAT.

No polynomial-time algorithm is known for, NONLINEAR SYSTEM, 3-COLORABILITY, or 3-SAT.

However, any candidate solution to either of these problems can be checked in polynomial-time.

No polynomial-time algorithm is known for, NONLINEAR SYSTEM, 3-COLORABILITY, or 3-SAT.

However, any candidate solution to either of these problems can be checked in polynomial-time.

Thus these problems are solvable in **nondeterministic polynomial time**.

Let X and Y be two problems. We write $X \leq_p Y$ to indicate that Y is at least as hard as X .

Let X and Y be two problems. We write $X \leq_p Y$ to indicate that Y is at least as hard as X .

Somewhat more precisely: any algorithm for solving Y can be transformed into an algorithm for X without drastically increasing its running time.

Let X and Y be two problems. We write $X \leq_p Y$ to indicate that Y is at least as hard as X .

Somewhat more precisely: any algorithm for solving Y can be transformed into an algorithm for X without drastically increasing its running time.

It is possible for $X \leq_p Y \leq_p X$. In that case, write $X \equiv_p Y$.

\mathbb{P} is the class of all problems solvable in polynomial time.
Its members are called **tractable**.

NP is the class of problems solvable in nondeterministic polynomial time.

\mathbb{P} is the class of all problems solvable in polynomial time.
Its members are called **tractable**.

\mathbb{NP} is the class of problems solvable in nondeterministic polynomial time.

- $\mathbb{P} \subseteq \mathbb{NP}$
- Both \mathbb{P} and \mathbb{NP} are downsets, i.e.,
 $Y \in \mathbb{P} \ \& \ X \leq_p Y \implies X \in \mathbb{P}$

\mathbb{P} is the class of all problems solvable in polynomial time. Its members are called **tractable**.

\mathbb{NP} is the class of problems solvable in nondeterministic polynomial time.

- $\mathbb{P} \subseteq \mathbb{NP}$
- Both \mathbb{P} and \mathbb{NP} are downsets, i.e.,
$$Y \in \mathbb{P} \ \& \ X \leq_p Y \implies X \in \mathbb{P}$$

The maximal members of \mathbb{NP} are called **\mathbb{NP} -complete**.

3-COLORABILITY, NONLINEAR SYSTEM, and 3-SAT are known to be \mathbb{NP} -complete.

\$2²⁰ question: $\mathbb{P} \stackrel{?}{=} \text{NP}$.

\$2²⁰ question: $\mathbb{P} \stackrel{?}{=} \text{NP}$.

If $\mathbb{P} = \text{NP}$ then all of the above distinctions go away.
Almost every problem that mathematicians actually care about can be solved efficiently. Just build bigger computers.

\$2²⁰ question: $\mathbb{P} \stackrel{?}{=} \text{NP}$.

If $\mathbb{P} = \text{NP}$ then all of the above distinctions go away. Almost every problem that mathematicians actually care about can be solved efficiently. Just build bigger computers.

In particular, this talk becomes pointless. So assume $\mathbb{P} \neq \text{NP}$.


\$2²⁰ question: $\mathbb{P} \stackrel{?}{=} \text{NP}$.

If $\mathbb{P} = \text{NP}$ then all of the above distinctions go away. Almost every problem that mathematicians actually care about can be solved efficiently. Just build bigger computers.

In particular, this talk becomes pointless. So assume $\mathbb{P} \neq \text{NP}$.

Theorem (Ladner, 1975)

If $\mathbb{P} \neq \text{NP}$ then there are problems in $\text{NP} - \mathbb{P}$ that are not NP-complete.



If $P \neq NP$ then
the pink area is
nonempty.

Formal Definition of CSP

Let D be a set, n a positive integer

An n -ary relation on D is a subset of D^n

Formal Definition of CSP

Let D be a set, n a positive integer

An n -ary relation on D is a subset of D^n

$\text{Rel}_n(D)$ denotes the set of all n -ary relations on D

$$\text{Rel}(D) = \bigcup_{n>0} \text{Rel}_n(D)$$

Let D be a finite set and $\Delta \subseteq \text{Rel}(D)$

$\text{CSP}(\langle D, \Delta \rangle)$ is the following decision problem:

Instance. A finite set $V = \{v_1, \dots, v_n\}$ of **variables** and a finite set $\{C_1, \dots, C_m\}$ of **constraints**;

each constraint C_i is a pair $(\langle x_{i1}, \dots, x_{ip_i} \rangle, \delta_i)$ in which $x_{i1}, \dots, x_{ip_i} \in V$ and $\delta_i \in \Delta$

Let D be a finite set and $\Delta \subseteq \text{Rel}(D)$

$\text{CSP}(\langle D, \Delta \rangle)$ is the following decision problem:

Instance. A finite set $V = \{v_1, \dots, v_n\}$ of **variables** and a finite set $\{C_1, \dots, C_m\}$ of **constraints**;

each constraint C_i is a pair $(\langle x_{i1}, \dots, x_{ip_i} \rangle, \delta_i)$ in which $x_{i1}, \dots, x_{ip_i} \in V$ and $\delta_i \in \Delta$

Question. Does there exist a **solution**, that is, a “context” $\rho: V \rightarrow D$, such that for all $i \leq m$, $\langle \rho(x_{i1}), \dots, \rho(x_{ip_i}) \rangle \in \delta_i$?

Let D be a finite set and $\Delta \subseteq \text{Rel}(D)$

$\text{CSP}(\langle D, \Delta \rangle)$ is the following decision problem:

Instance. A finite set $V = \{v_1, \dots, v_n\}$ of **variables** and a finite set $\{C_1, \dots, C_m\}$ of **constraints**;

each constraint C_i is a pair $(\langle x_{i1}, \dots, x_{ip_i} \rangle, \delta_i)$ in which $x_{i1}, \dots, x_{ip_i} \in V$ and $\delta_i \in \Delta$

Question. Does there exist a **solution**, that is, a “context” $\rho: V \rightarrow D$, such that for all $i \leq m$, $\langle \rho(x_{i1}), \dots, \rho(x_{ip_i}) \rangle \in \delta_i$?


$\text{CSP}(\langle D, \Delta \rangle)$ always lies in NP .

Example: 3-colorability

$$D = \{\textcolor{red}{r}, \textcolor{green}{g}, \textcolor{blue}{b}\}, \quad \Delta = \{\kappa_3\}$$

$$\kappa_3 = \{ (x, y) \in D : x \neq y \}$$

Then $\text{CSP}(\langle D, \Delta \rangle)$ is the 3-colorability problem



$$V = \{v_1, \dots, v_6\}$$

$$\langle v_1, v_2 \rangle \in \kappa$$

$$\langle v_1, v_3 \rangle \in \kappa$$

$$\langle v_1, v_4 \rangle \in \kappa$$

$$\langle v_2, v_4 \rangle \in \kappa$$

$$\vdots$$

$$\langle v_5, v_6 \rangle \in \kappa$$

Two Motivating Questions

1 Dichotomy Conjecture

Every $\text{CSP}(\langle D, \Delta \rangle)$ either lies in \mathbb{P} or is NP -complete.

Two Motivating Questions

1 Dichotomy Conjecture

Every $\text{CSP}(\langle D, \Delta \rangle)$ either lies in \mathbb{P} or is NP -complete.

2 Tractability Problem

Characterize those CSPs that lie in \mathbb{P} .

Two Motivating Questions

1 Dichotomy Conjecture

Every $\text{CSP}(\langle D, \Delta \rangle)$ either lies in \mathbb{P} or is NP -complete.

2 Tractability Problem

Characterize those CSPs that lie in \mathbb{P} .

What would a characterization look like? What language could we use?

Polymorphisms

Definition

Let $\delta \in \text{Rel}_k(D)$ and $f: D^n \rightarrow D$. We say f preserves δ if

$$(a_{11}, \dots, a_{1k}), \dots, (a_{n1}, \dots, a_{nk}) \in \delta \implies \\ (f(a_{11}, \dots, a_{n1}), \dots, f(a_{1k}, \dots, a_{nk})) \in \delta$$

Polymorphisms

Definition

Let $\delta \in \text{Rel}_k(D)$ and $f: D^n \rightarrow D$. We say f preserves δ if

$$(a_{11}, \dots, a_{1k}), \dots, (a_{n1}, \dots, a_{nk}) \in \delta \implies \\ (f(a_{11}, \dots, a_{n1}), \dots, f(a_{1k}, \dots, a_{nk})) \in \delta$$

f is an n -ary operation on D .

Polymorphisms

Definition

Let $\delta \in \text{Rel}_k(D)$ and $f: D^n \rightarrow D$. We say f preserves δ if

$$(a_{11}, \dots, a_{1k}), \dots, (a_{n1}, \dots, a_{nk}) \in \delta \implies \\ (f(a_{11}, \dots, a_{n1}), \dots, f(a_{1k}, \dots, a_{nk})) \in \delta$$

$$\begin{array}{ccccccc} a_{11} & a_{12} & \dots & a_{1k} & \in & \delta \\ a_{21} & a_{22} & \dots & a_{2k} & \in & \delta \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nk} & \in & \delta \\ \downarrow f & \downarrow f & & \downarrow f & & \\ \star & \star & \dots & \star & \in & \delta \end{array}$$

Definition

Let Δ be a set of relations on D . Then $\text{Pol}(\Delta)$ denotes the set of all operations preserving all members of Δ . These are the **polymorphisms** of Δ .

Definition

Let Δ be a set of relations on D . Then $\text{Pol}(\Delta)$ denotes the set of all operations preserving all members of Δ . These are the **polymorphisms** of Δ .

Let F be a set of operations on D . Then $\text{Inv}(F)$ denotes the set of all relations preserved by all operations in F .

Definition

Let Δ be a set of relations on D . Then $\text{Pol}(\Delta)$ denotes the set of all operations preserving all members of Δ . These are the **polymorphisms** of Δ .

Let F be a set of operations on D . Then $\text{Inv}(F)$ denotes the set of all relations preserved by all operations in F .

Important point: $\langle D, \text{Pol}(\Delta) \rangle$ is an algebraic structure

Theorem

Let $\Gamma, \Delta \subseteq \text{Rel}(D)$. Then

$$\text{Pol}(\Gamma) \subseteq \text{Pol}(\Delta) \implies \text{CSP}(\Delta) \leq_p \text{CSP}(\Gamma).$$

Theorem

Let $\Gamma, \Delta \subseteq \text{Rel}(D)$. Then

$$\text{Pol}(\Gamma) \subseteq \text{Pol}(\Delta) \implies \text{CSP}(\Delta) \leq_p \text{CSP}(\Gamma).$$

Thus, the richer the algebraic structure, the easier the corresponding CSP

One can go back and forth between relational and algebraic structures

$$\begin{array}{ccc} \textbf{Relational} & & \textbf{Algebraic} \\ \langle D, \Delta \rangle & \longrightarrow & \langle D, \text{Pol}(\Delta) \rangle \\ \langle D, \text{Inv}(F) \rangle & \longleftarrow & \langle D, F \rangle \end{array}$$

$$\text{CSP} \langle D, \Delta \rangle \equiv_p \text{CSP} \langle D, \text{Inv}(\text{Pol}(\Delta)) \rangle$$

One can go back and forth between relational and algebraic structures

$$\begin{array}{ccc} \textbf{Relational} & & \textbf{Algebraic} \\ \langle D, \Delta \rangle & \longrightarrow & \langle D, \text{Pol}(\Delta) \rangle \\ \langle D, \text{Inv}(F) \rangle & \longleftarrow & \langle D, F \rangle \end{array}$$

$$\text{CSP}\langle D, \Delta \rangle \equiv_p \text{CSP}\langle D, \text{Inv}(\text{Pol}(\Delta)) \rangle$$

Perhaps the expressive power of algebra can be used to classify CSPs.

Algebraic Facts

Let \mathbf{A} and \mathbf{B} be algebras

\mathbf{B} a subalgebra of $\mathbf{A} \implies \text{CSP}(\mathbf{B}) \leq_p \text{CSP}(\mathbf{A})$.

\mathbf{B} a homomorphic image of $\mathbf{A} \implies \text{CSP}(\mathbf{B}) \leq_p \text{CSP}(\mathbf{A})$.

$\text{CSP}(\mathbf{A}^n) \equiv_p \text{CSP}(\mathbf{A})$

Theorem (Bulatov, Jeavons, Krokhin, 2000)

If $\langle D, \Delta \rangle$ is a core and every polymorphism is essentially unary, then $\text{CSP}(\Delta)$ is NP -complete.

f is **essentially unary** if $f(x_1, \dots, x_n) = g(x_j)$ for some unary g and some $j \leq n$.

Theorem (Bulatov, Jeavons, Krokhin, 2000)

If $\langle D, \Delta \rangle$ is a core and every polymorphism is essentially unary, then $\text{CSP}(\Delta)$ is NP -complete.

f is **essentially unary** if $f(x_1, \dots, x_n) = g(x_j)$ for some unary g and some $j \leq n$.

Corollary

3-COLORABILITY, NONLINEAR SYSTEM, and 3-SAT are NP -complete.

Informal reformulation of the dichotomy conjecture

If \mathbf{A} has some kind of decent algebraic structure then $\text{CSP}(\mathbf{A}) \in \mathbb{P}$ otherwise $\text{CSP}(\mathbf{A})$ is NP -complete.