

CSPs OF FINITE COMMUTATIVE IDEMPOTENT BINARS

William DeMeo

williamdemeo@gmail.com

Iowa State University

joint work with

Cliff Bergman

Jiali Li

Shanks Workshop

Vanderbilt University

May 30, 2015

General Problem: Find Maltsev conditions that characterize the complexity of CSPs of universal algebras.

CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra \mathbf{A} ...

$\text{CSP}(\mathbf{A})$ is tractable $\iff \mathbf{A}$ has a weak-nu term operation

General Problem: Find Maltsev conditions that characterize the complexity of CSPs of universal algebras.

CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra \mathbf{A} ...

$\text{CSP}(\mathbf{A})$ is tractable $\implies \mathbf{A}$ has a weak-nu term operation ✓

The left-to-right direction is known.

General Problem: Find Maltsev conditions that characterize the complexity of CSPs of universal algebras.

CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra \mathbf{A} ...

$\text{CSP}(\mathbf{A})$ is tractable $\iff \mathbf{A}$ has a weak-nu term operation (?)

The right-to-left direction is open.

General Problem: Find Maltsev conditions that characterize the complexity of CSPs of universal algebras.

CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra \mathbf{A} ...

$\text{CSP}(\mathbf{A})$ is tractable $\iff \mathbf{A}$ has a weak-nu term operation (?)

A **weak near unanimity** (weak-nu) term operation is one that satisfies

$$t(x, x, \dots, x) \approx x \quad (\text{idempotent})$$

$$t(y, x, \dots, x) \approx t(x, y, \dots, x) \approx \dots \approx t(x, x, \dots, y)$$

General Problem: Find Maltsev conditions that characterize the complexity of CSPs of universal algebras.

CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra \mathbf{A} ...

$\text{CSP}(\mathbf{A})$ is tractable $\iff \mathbf{A}$ has a weak-nu term operation (?)

A **weak near unanimity** (weak-nu) term operation is one that satisfies

$$t(x, x, \dots, x) \approx x \quad (\text{idempotent})$$

$$t(y, x, \dots, x) \approx t(x, y, \dots, x) \approx \dots \approx t(x, x, \dots, y)$$

A *binary* operation $t(x, y)$ is weak-nu if

$$t(x, x) \approx x \quad (\text{idempotent})$$

$$t(y, x) \approx t(x, y) \quad (\text{commutative})$$

So let's try to prove (?) for **commutative idempotent binars**.

COMMUTATIVE IDEMPOTENT BINARS

A **CIB** is an algebra $\mathbf{A} = \langle A, \cdot \rangle$ satisfying $x \cdot y \approx y \cdot x$ and $x \cdot x \approx x$.

COMMUTATIVE IDEMPOTENT BINARS

A **CIB** is an algebra $\mathbf{A} = \langle A, \cdot \rangle$ satisfying $x \cdot y \approx y \cdot x$ and $x \cdot x \approx x$.

QUESTION

Is every finite commutative idempotent binar tractable?

COMMUTATIVE IDEMPOTENT BINARS

A **CIB** is an algebra $\mathbf{A} = \langle A, \cdot \rangle$ satisfying $x \cdot y \approx y \cdot x$ and $x \cdot x \approx x$.

QUESTION

Is every finite commutative idempotent binar tractable?

First Example: a semilattice is an associative CIB.

Semilattices are tractable (in fact, they have *finite width*).

COMMUTATIVE IDEMPOTENT BINARS

A **CIB** is an algebra $\mathbf{A} = \langle A, \cdot \rangle$ satisfying $x \cdot y \approx y \cdot x$ and $x \cdot x \approx x$.

QUESTION

Is every finite commutative idempotent binar tractable?

First Example: a semilattice is an associative CIB.

Semilattices are tractable (in fact, they have *finite width*).

Pause to consider more general case for a minute...

GENERAL CASE

SOME WELL KNOWN FACTS

Let \mathbf{A} be a finite idempotent algebra. Let \mathbf{S}_2 be the 2-elt semilattice.

$$\mathbf{V}(\mathbf{A}) \text{ is CP} \iff \mathbf{A} \text{ has Malcev term}$$

GENERAL CASE

SOME WELL KNOWN FACTS

Let \mathbf{A} be a finite idempotent algebra. Let \mathbf{S}_2 be the 2-elt semilattice.

$$V(\mathbf{A}) \text{ is CP} \iff \mathbf{A} \text{ has Malcev term}$$




GENERAL CASE

SOME WELL KNOWN FACTS

Let \mathbf{A} be a finite idempotent algebra. Let \mathbf{S}_2 be the 2-elt semilattice.

$$\begin{aligned} \mathbf{V}(\mathbf{A}) \text{ is CP} &\iff \mathbf{A} \text{ has Malcev term} \\ &\implies \mathbf{A} \text{ has cube term} \end{aligned}$$



GENERAL CASE


SOME WELL KNOWN FACTS

Let \mathbf{A} be a finite idempotent algebra. Let \mathbf{S}_2 be the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has Malcev term

\implies \mathbf{A} has cube term

$\implies V(\mathbf{A})$ is CM



GENERAL CASE

SOME WELL KNOWN FACTS


Let \mathbf{A} be a finite idempotent algebra. Let \mathbf{S}_2 be the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has Malcev term

\implies \mathbf{A} has cube term

$\implies V(\mathbf{A})$ is CM

$\implies \mathbf{S}_2$ is not in $V(\mathbf{A})$



SOME WELL KNOWN FACTS

\mathbf{A} = a finite idempotent algebra


\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term

\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$



SOME WELL KNOWN FACTS

\mathbf{A} = a finite idempotent algebra


\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term

\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$



■ cube term \implies CM

(Berman, Idziak, Marković, McKenzie, Valeriote, Willard 2010)

■ CM \implies \mathbf{S}_2 is not in $V(\mathbf{A})$

Proof: $\mathbf{S}_2 \in V(\mathbf{A}) \Rightarrow \mathbf{S}_2^2 \in V(\mathbf{A})$;

$\text{Con}(\mathbf{S}_2^2)$ is not modular.

SOME WELL KNOWN FACTS

\mathbf{A} = a finite **CIB**


\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term

\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$



CIB case

- 1st reduction by cube-term blockers.

SOME WELL KNOWN FACTS

\mathbf{A} = a finite **CIB**

\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term


\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$

\implies \mathbf{A} has a cube term

CIB case

- 1st reduction by cube-term blockers.



SOME WELL KNOWN FACTS

\mathbf{A} = a finite **CIB**

\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term


\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$

\implies \mathbf{A} has a cube term

CIB case

- 1st reduction by cube-term blockers.
- 2nd reduction by Kearnes-Tschantz.



SOME WELL KNOWN FACTS

\mathbf{A} = a finite **CIB**

\mathbf{S}_2 = the 2-elt semilattice.

$V(\mathbf{A})$ is CP \iff \mathbf{A} has a Malcev term

\implies \mathbf{A} has a cube term

\implies $V(\mathbf{A})$ is CM

\implies \mathbf{S}_2 is not in $V(\mathbf{A})$

\implies \mathbf{A} has a cube term

\implies $V(\mathbf{A})$ is CP

CIB case

- 1st reduction by cube-term blockers.
- 2nd reduction by Kearnes-Tschantz.



FIRST REDUCTION

BY CUBE-TERM BLOCKERS

Marković, M. Maróti, McKenzie (M^4)

“Finitely related clones and algebras with cube terms” (2012)

A **cube-term blocker** (CTB) is a pair (C, B) of subuniverses satisfying $\emptyset < C < B \leq A$ and for every $t(x_1, \dots, x_n)$ there is an index $i \in [n]$ with

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

FIRST REDUCTION

BY CUBE-TERM BLOCKERS

Marković, M. Maróti, McKenzie (M^4)

“Finitely related clones and algebras with cube terms” (2012)

A **cube-term blocker** (CTB) is a pair (C, B) of subuniverses satisfying $\emptyset < C < B \leq A$ and for every $t(x_1, \dots, x_n)$ there is an index $i \in [n]$ with

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

M^4 prove a finite idempotent algebra has a cube term iff it has no CTB.

FIRST REDUCTION

BY CUBE-TERM BLOCKERS

Marković, M. Maróti, McKenzie (M^4)

“Finitely related clones and algebras with cube terms” (2012)

A **cube-term blocker** (CTB) is a pair (C, B) of subuniverses satisfying $\emptyset < C < B \leq A$ and for every $t(x_1, \dots, x_n)$ there is an index $i \in [n]$ with

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

M^4 prove a finite idempotent algebra has a cube term iff it has no CTB.

LEMMA

A finite CIB \mathbf{A} has a CTB if and only if $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$.

FIRST REDUCTION

BY CUBE-TERM BLOCKERS

Marković, M. Maróti, McKenzie (M^4)

“Finitely related clones and algebras with cube terms” (2012)

A **cube-term blocker** (CTB) is a pair (C, B) of subuniverses satisfying $\emptyset < C < B \leq A$ and for every $t(x_1, \dots, x_n)$ there is an index $i \in [n]$ with

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

M^4 prove a finite idempotent algebra has a cube term iff it has no CTB.

LEMMA

A finite CIB \mathbf{A} has a CTB if and only if $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$.

PROOF.

(C, B) a CTB implies $\theta = C^2 \cup (B - C)^2$ a congruence with $\mathbf{B}/\theta \cong \mathbf{S}_2$.

Conversely, suppose $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$, and \mathbf{B} is a subalgebra of \mathbf{A} with \mathbf{B}/θ a meet-SL for some θ . Let C/θ be the bottom of \mathbf{B}/θ , then (C, B) is a CTB. □

SECOND REDUCTION

Kearnes and Tschantz

“Automorphism groups of squares and of free algebras” (2007)

LEMMA

If V is an idempotent variety that is not congruence permutable, then there are subuniverses U and W of $\mathbf{F} := \mathbf{F}_V\{x, y\}$ satisfying

1. $x \in U \cap W$
2. $y \in U^c \cap W^c$
3. $(U \times F) \cup (F \times W) \leq \mathbf{F}^2$

SECOND REDUCTION

Kearnes and Tschantz

“Automorphism groups of squares and of free algebras” (2007)

LEMMA

If V is an idempotent variety that is not congruence permutable, then there are subuniverses U and W of $\mathbf{F} := \mathbf{F}_V\{x, y\}$ satisfying

1. $x \in U \cap W$
2. $y \in U^c \cap W^c$
3. $(U \times F) \cup (F \times W) \leq \mathbf{F}^2$

For CIB's, either U or W will be an ideal.

This implies a CTB and a semilattice.

REMAINING QUESTIONS FOR FINITE CIBs

CONCLUSION

Let \mathbf{A} be a finite CIB. Then

$\mathbf{S}_2 \notin \text{HS}(\mathbf{A})$ if and only if $\mathbf{V}(\mathbf{A})$ is congruence permutable.

REMAINING QUESTIONS FOR FINITE CIBS

CONCLUSION

Let \mathbf{A} be a finite CIB. Then

$\mathbf{S}_2 \notin \text{HS}(\mathbf{A})$ if and only if $\mathbf{V}(\mathbf{A})$ is congruence permutable.

(so $\text{CSP}(\mathbf{A})$ tractable in this case)

REMAINING QUESTIONS FOR FINITE CIBs

CONCLUSION

Let \mathbf{A} be a finite CIB. Then

$S_2 \notin \text{HS}(\mathbf{A})$ if and only if $V(\mathbf{A})$ is congruence permutable.

(so $\text{CSP}(\mathbf{A})$ tractable in this case)

OPEN QUESTION

Let \mathbf{A} be a finite CIB with S_2 in $\text{HS}(\mathbf{A})$. Is $\text{CSP}(\mathbf{A})$ tractable?

REMAINING QUESTIONS FOR FINITE CIBS

CONCLUSION

Let \mathbf{A} be a finite CIB. Then

$S_2 \notin \text{HS}(\mathbf{A})$ if and only if $V(\mathbf{A})$ is congruence permutable.

(so $\text{CSP}(\mathbf{A})$ tractable in this case)

OPEN QUESTION

Let \mathbf{A} be a finite CIB with S_2 in $\text{HS}(\mathbf{A})$. Is $\text{CSP}(\mathbf{A})$ tractable?

Recall, if $V(\mathbf{A})$ is SD_\wedge , then $\text{CSP}(\mathbf{A})$ is tractable.

REMAINING QUESTIONS FOR FINITE CIBS

CONCLUSION

Let \mathbf{A} be a finite CIB. Then

$S_2 \notin \text{HS}(\mathbf{A})$ if and only if $V(\mathbf{A})$ is congruence permutable.

(so $\text{CSP}(\mathbf{A})$ tractable in this case)

OPEN QUESTION

Let \mathbf{A} be a finite CIB with S_2 in $\text{HS}(\mathbf{A})$. Is $\text{CSP}(\mathbf{A})$ tractable?

Recall, if $V(\mathbf{A})$ is SD_\wedge , then $\text{CSP}(\mathbf{A})$ is tractable.

REVISED QUESTION

Let \mathbf{A} be a finite CIB with S_2 in $\text{HS}(\mathbf{A})$, and $V(\mathbf{A})$ not SD_\wedge .

Is $\text{CSP}(\mathbf{A})$ tractable?

EXAMPLES

\cdot	0	1	2	3
0	0	0	0	1
1	0	1	3	2
2	0	3	2	1
3	1	2	1	3

$*$	0	1	2	3
0	0	0	1	1
1	0	1	3	2
2	1	3	2	1
3	1	2	1	3

\circ	0	1	2	3
0	0	0	2	1
1	0	1	3	2
2	2	3	2	1
3	1	2	1	3

Maroti's idea:

Bergman's idea: replace basic binary operation with a term from $\text{Clo}(\mathbf{A})$, say $t(x, y) = (x * y) * x$.

If $\langle A, t \rangle$ tractable, then so is $\langle A, * \rangle$