

THE FINITE LATTICE REPRESENTATION PROBLEM
AND
INTERVALS IN SUBGROUP LATTICES
PART I

William DeMeo

`williamdemeo@gmail.com`

joint work with

Ralph Freese, Peter Jipsen, Bill Lampe, J.B. Nation

University of South Carolina

Algebra & Logic Seminar

September 21, 2012


These slides and other resources are available at

<http://www.math.sc.edu/~demeow/FLRP.html> →



CONGRUENCE DECOMPOSITIONS


CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

CONGRUENCE DECOMPOSITIONS

CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice L , does there exist a *finite* algebra A such that $\text{Con } A \cong L$?

DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice \mathbf{L} , does there exist a *finite* algebra \mathbf{A} such that $\mathbf{Con A} \cong \mathbf{L}$?

DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice \mathbf{L} , does there exist a *finite* algebra \mathbf{A} such that $\mathbf{Con A} \cong \mathbf{L}$?

DEFINITION

We call a finite lattice **representable** if it is isomorphic to the congruence lattice of a finite algebra.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) Yes, $\mathcal{L}_0 = \mathcal{L}_1$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) Yes, $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) Yes, $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Fact: $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) Yes, $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \subsetneq \mathcal{L}_3 \subsetneq \mathcal{L}_4$.

RECAP

THEOREM (PUDLÁK AND TŮMA, 1980)

Every finite lattice can be embedded in $\text{Eq}(X)$ with X finite.

In other words, $\mathcal{L}_0 = \mathcal{L}_1$.

THEOREM (PÁLFY AND PUDLÁK, 1980)

The following statements are equivalent:

- (I) *Every finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (II) *Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

In other words, $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 1 (USE CLOSURE PROPERTIES)

The class \mathcal{L}_3 is closed under the following operations:

- lattice duals (Kurzweil and Netter, 1986)
- interval sublattices (follows from Kurzweil-Netter)
- direct products (Tůma, 1986)
- ordinal sums (McKenzie, 1984; Snow, 2000)
- parallel sums (Snow, 2000)
- certain sublattices of lattices in \mathcal{L}_3 (Snow, 2000)
(namely, those obtained as a union of a filter and ideal)

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 3 (SUBGROUP LATTICE INTERVAL)

Find L as an interval in a subgroup lattice of a finite group.

If $H \leq G$ are finite groups, then the interval above H in $Sub(G)$,

$$[H, G] := \{K \mid H \leq K \leq G\},$$


is isomorphic to $\text{Con} \langle G/H, G \rangle$.

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.

Suppose $L_0 \cong \text{Con } \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.



HOW TO FIND A REPRESENTATION OF A FINITE LATTICE


METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.

Suppose $L_0 \cong \text{Con } \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.

Consider $L = \alpha^\uparrow \cup \beta^\downarrow$.

$$L \leq L_0$$



HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.


Suppose $L_0 \cong \text{Con} \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.

Consider $L = \alpha^\uparrow \cup \beta^\downarrow$.


Then there exists a set $F' \subset A^A$ such that

$$L \cong \text{Con} \langle A, F \cup F' \rangle.$$

$$L \leq L_0$$



LATTICES WITH AT MOST 6 ELEMENTS ARE REPRESENTABLE.




Watatani (1996) *J. Funct. Anal.*


Aschbacher (2008) *JAMS*

Theorem: *Every lattice with at most 6 elements is an interval in the subgroup lattice of a finite group.*


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19}




L_{20}




L_{17}




L_{13}



L_{11}




L_9




L_7


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19}




L_{20}




L_{17}




L_{13}



L_{11}





L_9



L_7


FINDING REPRESENTATIONS...

...AS INTERVALS IN SUBGROUP LATTICES



FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



`SmallGroup(288,1025)`


$$|G : H| = 48$$

- The group $G = (A_4 \times A_4) \rtimes C_2$ has a subgroup $H \cong S_3$ such that $[H, G] \cong L_{17}$.
- ...so the dual L_{16} is also representable.



FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$

- The group $G = (A_4 \times A_4) \rtimes C_2$ has a subgroup $H \cong S_3$ such that $[H, G] \cong L_{17}$.
- ...so the dual L_{16} is also representable.




SmallGroup(960,11358)

$$|G : H| = 80$$

- The group $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$ has a subgroup $H \cong A_4$ such that $[H, G] \cong L_{13}$.

FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.



FINDING REPRESENTATIONS...

...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)




- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.

FINDING REPRESENTATIONS...

...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)




- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$

FINDING REPRESENTATIONS...



...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)



- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$




- $\text{Sub}(A_4)$ is a congruence lattice (of A_4 acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓




L_9




L_7


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓




L_9 ✓




L_7


SEVEN ELEMENT LATTICES: SUMMARY




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓




L_9 ✓




L_7


SEVEN ELEMENT LATTICES: SUMMARY




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓



L_9 ✓



L_7


HAS ANYONE SEEN THIS LATTICE?

[login](#)[faq](#)[how to ask](#)[meta](#)**mathoverflow**[Questions](#)[Tags](#)[Users](#)[Badges](#)[Unanswered](#)[Ask Question](#)

Given a lattice L with n elements, are there finite groups $H < G$ such that $L \cong$ the lattice of subgroups between H and G ?

▲
13
▼

If there is no restriction on n , this is a famous [open problem](#). I'm wondering if any recent work has been done for small $n > 6$. I believe the question is answered (positively) for $n = 6$ by Watatani (1996) [MR1409040](#) and Aschbacher (2008) [MR2393428](#). I also believe we can answer it for $n = 7$, with one possible exception. The exceptional case is shown below.



So my two questions are these:

- 1) Does anyone know of recent work on this special case of the problem (specifically for $n = 7$ or $n = 8$)?
- 2) Has anyone found a finite group G with a subgroup H such that the interval $[H, G] = \{K : H \leq K \leq G\}$

is the lattice shown above?

Welcome to MathOverflow

A place for mathematicians to ask and answer questions.

tagged

[finite-groups](#) × 343

[open-problem](#) × 216

[lattices](#) × 157

[universal-algebra](#) × 55

[congruences](#) × 7

asked


8 months ago

viewed

468 times


INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, $L_7 \cong [H, G]$.

INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$. What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, $L_7 \cong [H, G]$.

- (I) G is a primitive permutation group.
- (II) If $N \triangleleft G$, then $C_G(N) = 1$.
- (III) G contains no non-trivial abelian normal subgroup.
- (IV) G is not solvable.
- (V) G is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, all proper subgroups in the interval $[H, G]$ are core-free.

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:


$$UH = \langle U, H \rangle \iff UH = HU.$$

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$




SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee H = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:


$$UH = \langle U, H \rangle \iff UH = HU.$$




$$[U_0, U] := \{V \mid U_0 \leq V \leq U\}$$

INTERVAL ISOMORPHISMS

- If $H \trianglelefteq \langle U, H \rangle$, then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



INTERVAL ISOMORPHISMS




- If $H \trianglelefteq \langle U, H \rangle$, then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.
- Instead of $H \trianglelefteq \langle U, H \rangle$, assume only $UH = \langle U, H \rangle$ and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

INTERVAL ISOMORPHISMS



- If $H \trianglelefteq \langle U, H \rangle$, then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.
- Instead of $H \trianglelefteq \langle U, H \rangle$, assume only $UH = \langle U, H \rangle$ and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$


the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define

$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $V^h = V$ ($\forall h \in H$).

INTERVAL ISOMORPHISMS



- If $H \trianglelefteq \langle U, H \rangle$, then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.
- Instead of $H \trianglelefteq \langle U, H \rangle$, assume only $UH = \langle U, H \rangle$ and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $V^h = V$ ($\forall h \in H$).

LEMMA

- 1 $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$
- 2 If $U \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$.
- 3 If $H \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H = [U_0, U]$.

INTERVAL ISOMORPHISMS



- If $H \trianglelefteq \langle U, H \rangle$, then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.
- Instead of $H \trianglelefteq \langle U, H \rangle$, assume only $UH = \langle U, H \rangle$ and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\},$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define

$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $V^h = V$ ($\forall h \in H$).

LEMMA

- 1 $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$
- 2 If $U \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$.
- 3 If $H \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H = [U_0, U]$.

EXAMPLE 1

- Consider $G \cong C_3 \times S_3$, say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

EXAMPLE 1


- Consider $G \cong C_3 \times S_3$, say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ($UH = HU$) but neither one normalizes the other.



EXAMPLE 1


- Consider $G \cong C_3 \times S_3$, say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ($UH = HU$) but neither one normalizes the other.



EXAMPLE 1


- Consider $G \cong C_3 \times S_3$, say,

$$G = \langle a, b, c \mid a^2, b^3, c^3, [b, a], [c, b], c^{-1} a^{-1} a^c \rangle$$

- The subgroups

$$U = \langle a, b \rangle \cong C_6, \quad H = \langle bc \rangle \cong C_3$$

permute ($UH = HU$) but neither one normalizes the other.




- Three of the four subgroups of U permute with H .
As the lemma predicts, $U \cap \langle b, c \rangle = \langle b \rangle$.

EXAMPLE 2

- The group S_4 has subgroups $U \cong D_8$ and $H \cong C_3$ that permute but neither one normalizes the other.

EXAMPLE 2


- The group S_4 has subgroups $U \cong D_8$ and $H \cong C_3$ that permute but neither one normalizes the other.



- Only four subgroups of U permute with H


EXAMPLE 2

- The group S_4 has subgroups $U \cong D_8$ and $H \cong C_3$ that permute but neither one normalizes the other.



- Only four subgroups of U permute with H , including

$$U \cap A_4 \cong C_2 \times C_2, \quad U \cap S_3 \cong C_2.$$




THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, and $L_7 \cong [H, G]$. Then

- (I) G is a primitive permutation group.
- (II) If $N \triangleleft G$, then $C_G(N) = 1$.
- (III) G contains no non-trivial abelian normal subgroup.
- (IV) G is not solvable.
- (V) G is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, M_1 or M_2 , all proper subgroups in the interval $[H, G]$ are core-free.


IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

IDEA OF THE PROOF




Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .

IDEA OF THE PROOF




Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.

IDEA OF THE PROOF




Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1X = XJ_1\}.$$

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!



ASCHBACHER-O'NAN-SCOTT THEOREM

Let G be a primitive permutation group of degree d , and let $N := \text{Soc}(G) \cong T^m$ with $m \geq 1$. Then one of the following holds.

- ① N is regular and
 - **(Affine type)** T is cyclic of order p , so $|N| = p^m$. Then $d = p^m$ and G is permutation isomorphic to a subgroup of the affine general linear group $\text{AGL}(m, p)$.
 - **(Twisted wreath product type)** $m \geq 6$, the group T is nonabelian and G is a group of *twisted wreath product type*, with $d = |T|^m$.
- ② N is non-regular, non-abelian, and
 - **(Almost simple type)** $m = 1$ and $T \leq G \leq \text{Aut}(T)$.
 - **(Product action type)** $m \geq 2$ and G is permutation isomorphic to a subgroup of the product action wreath product $P \wr_{m/l} S_{m/l}$ of degree $d = nm/l$. The group P is primitive of type 2.(a) or 2.(c), P has degree n and $\text{Soc}(P) \cong T^l$, where $l \geq 1$ divides m .
 - **(Diagonal type)** $m \geq 2$ and $T^m \leq G \leq T^m \cdot (\text{Out}(T) \times S_m)$, with the diagonal action. The degree $d = |T|^{m-1}$.

ASCHBACHER-O'NAN-SCOTT THEOREM

For some interesting history, see Peter Cameron's blog at

<http://cameroncounts.wordpress.com/tag/onan-scott-theorem/>



CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).
- The lattice L_7 puts many restrictions on the class of groups which could possibly represent it. Perhaps this lattice, or a modified version of it, will enable us to restrict to the class of almost simple groups.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways. (Tune in next week!)
- We have representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We want to further explore the idea of finding restrictions on groups that have certain lattices as intervals in their subgroup lattices (next week).
- The lattice L_7 puts many restrictions on the class of groups which could possibly represent it. Perhaps this lattice, or a modified version of it, will enable us to restrict to the class of almost simple groups.
- Future work: Explore “interval enforceable properties of finite groups” and try to restrict to almost simple groups. Then solve the problem using the CFSG Theorem.

