

# CSP THEORY OF COMMUTATIVE IDEMPOTENT BINARS

William DeMeo

`williamdemeo@gmail.com`

Iowa State University

joint work with

Cliff Bergman

Jiali Li

May 25, 2015

**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation

**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation

The left-to-right direction is known, the converse is open.

**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation

The left-to-right direction is known, the converse is open.

**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation

A term  $t(x_1, \dots, x_n)$  is a **weak near unanimity** term operation if it satisfies

$$t(x, x, \dots, x) \approx x \quad (\text{idempotent})$$


$$t(y, x, \dots, x) \approx t(x, y, \dots, x) \approx \dots \approx t(x, x, \dots, y).$$

**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation




**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation




**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation






**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation




**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation




**General Problem:** Find Maltsev conditions that characterize complexity of CSPs of universal algebras.

### CSP DICHOTOMY CONJECTURE

For a (finite, idempotent) algebra  $\mathbf{A}$ ...

$\text{CSP}(\mathbf{A})$  is tractable  $\iff \mathbf{A}$  has a wnu term operation



## COMMUTATIVE IDEMPOTENT BINARS

Some more definitions.

- A set  $A$  together with a single binary operation is called a **binar**.
- A **commutative idempotent binar** is an algebra  $A = \langle A, \cdot \rangle$  satisfying  $x \cdot y \approx y \cdot x$  and  $x \cdot x \approx x$ .
- A binary operation  $x \cdot y = t(x, y)$  is a WNU term if and only if it is idempotent and commutative. This suggests the following

## COMMUTATIVE IDEMPOTENT BINARS

Some more definitions.

- A set  $A$  together with a single binary operation is called a **binar**.
- A **commutative idempotent binar** is an algebra  $A = \langle A, \cdot \rangle$  satisfying  $x \cdot y \approx y \cdot x$  and  $x \cdot x \approx x$ .
- A binary operation  $x \cdot y = t(x, y)$  is a WNU term if and only if it is idempotent and commutative. This suggests the following

### QUESTION

Is every finite commutative idempotent binar tractable?

If the dichotomy conjecture is to hold, then the answer must be “yes.”

## COMMUTATIVE IDEMPOTENT BINARS

Some more definitions.

- A set  $A$  together with a single binary operation is called a **binar**.
- A **commutative idempotent binar** is an algebra  $A = \langle A, \cdot \rangle$  satisfying  $x \cdot y \approx y \cdot x$  and  $x \cdot x \approx x$ .
- A binary operation  $x \cdot y = t(x, y)$  is a WNU term if and only if it is idempotent and commutative. This suggests the following

### QUESTION

Is every finite commutative idempotent binar tractable?

If the dichotomy conjecture is to hold, then the answer must be “yes.”

A semilattice is an associative CIB.

Semilattices are tractable (in fact, they have *finite width*).

## SOME WELL KNOWN FACTS

Let  $\mathbf{A}$  be a finite idempotent algebra. Let  $S_2$  be the 2-elt semilattice.

$$V(\mathbf{A}) \text{ is CP} \iff \mathbf{A} \text{ has Malcev term}$$

## SOME WELL KNOWN FACTS

Let  $\mathbf{A}$  be a finite idempotent algebra. Let  $S_2$  be the 2-elt semilattice.

$$V(\mathbf{A}) \text{ is CP} \iff \mathbf{A} \text{ has Malcev term}$$






## SOME WELL KNOWN FACTS

Let  $\mathbf{A}$  be a finite idempotent algebra. Let  $S_2$  be the 2-elt semilattice.

$$\begin{aligned} V(\mathbf{A}) \text{ is CP} &\iff \mathbf{A} \text{ has Malcev term} \\ &\implies \mathbf{A} \text{ has cube term} \end{aligned}$$




## SOME WELL KNOWN FACTS

Let  $\mathbf{A}$  be a finite idempotent algebra. Let  $S_2$  be the 2-elt semilattice.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has Malcev term

$\implies$   $\mathbf{A}$  has cube term

$\implies$   $V(\mathbf{A})$  is CM



## SOME WELL KNOWN FACTS


Let  $\mathbf{A}$  be a finite idempotent algebra. Let  $\mathbf{S}_2$  be the 2-elt semilattice.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has Malcev term

$\implies$   $\mathbf{A}$  has cube term

$\implies V(\mathbf{A})$  is CM

$\implies \mathbf{S}_2$  is not in  $V(\mathbf{A})$



## SOME WELL KNOWN FACTS

$\mathbf{A}$  = a finite idempotent algebra


$\mathbf{S}_2$  = the 2-elt semilattice.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has a Malcev term

$\implies$   $\mathbf{A}$  has a cube term

$\implies$   $V(\mathbf{A})$  is CM

$\implies$   $\mathbf{S}_2$  is not in  $V(\mathbf{A})$



## SOME WELL KNOWN FACTS

$\mathbf{A}$  = a finite idempotent algebra

$\mathbf{S}_2$  = the 2-elt semilattice.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has a Malcev term

$\implies$   $\mathbf{A}$  has a cube term


$\implies$   $V(\mathbf{A})$  is CM

$\implies$   $\mathbf{S}_2$  is not in  $V(\mathbf{A})$

■ cube term  $\implies$  CM

*Proof:* few subalgebras of powers

Berman, Idziak, Marković, McKenzie,  
Valeriote, Willard (BIMMVW) 2010.



## SOME WELL KNOWN FACTS

$\mathbf{A}$  = a finite idempotent algebra


$\mathbf{S}_2$  = the 2-elt semilattice.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has a Malcev term

$\implies$   $\mathbf{A}$  has a cube term

$\implies$   $V(\mathbf{A})$  is CM

$\implies$   $\mathbf{S}_2$  is not in  $V(\mathbf{A})$



■ cube term  $\implies$  CM

*Proof:* few subalgebras of powers

Berman, Idziak, Marković, McKenzie, Valeriote, Willard (BIMMVW) 2010.

■ CM  $\implies$   $\mathbf{S}_2$  is not in  $V(\mathbf{A})$

*Proof:*  $\mathbf{S}_2 \in V(\mathbf{A}) \Rightarrow \mathbf{S}_2^2 \in V(\mathbf{A})$ ;

$\text{Con}(\mathbf{S}_2^2)$  is not modular.

# COMMUTATIVE IDEMPOTENT BINARS (CIBs)


Let  $\mathbf{A}$  be a CIB.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has Malcev term

$\implies$   $\mathbf{A}$  has cube term

$\implies V(\mathbf{A})$  is CM

$\implies \mathbf{S}_2$  is not in  $V(\mathbf{A})$



# COMMUTATIVE IDEMPOTENT BINARS (CIBs)


Let  $\mathbf{A}$  be a CIB.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has Malcev term

$\implies$   $\mathbf{A}$  has cube term

$\implies$   $V(\mathbf{A})$  is CM

$\implies$   $\mathbf{S}_2$  is not in  $V(\mathbf{A})$





## COMMUTATIVE IDEMPOTENT BINARS (CIBs)


Let  $\mathbf{A}$  be a CIB.

$V(\mathbf{A})$  is CP  $\iff$   $\mathbf{A}$  has Malcev term

$\implies$   $\mathbf{A}$  has cube term

$\implies V(\mathbf{A})$  is CM

$\implies \mathbf{S}_2$  is not in  $V(\mathbf{A})$



## CUBE TERMS

A **cube operation** is a function  $c : A^n \rightarrow A$  satisfying for each  $1 \leq i \leq n$   $c(w_1, \dots, w_n) = x$  where  $\{w_1, \dots, w_n\} \subseteq \{x, y\}$  and  $w_i = y$ .

Here  $x$  and  $y$  are distinct variables.

An algebra **A** is said to have a **cube term** if its clone of term operations contains a cube operation.

## CUBE TERMS

A **cube operation** is a function  $c : A^n \rightarrow A$  satisfying for each  $1 \leq i \leq n$   $c(w_1, \dots, w_n) = x$  where  $\{w_1, \dots, w_n\} \subseteq \{x, y\}$  and  $w_i = y$ .

Here  $x$  and  $y$  are distinct variables.

An algebra **A** is said to have a **cube term** if its clone of term operations contains a cube operation.

Cube terms were introduced in... ?

Berman, Idziak, Marković, McKenzie, Valeriote, Willard,  
"Varieties with few subalgebras of powers," 2010.

Marković, Maróti, McKenzie,  
"Finitely related clones & algebras with cube terms," 2012.

## CUBE TERM BLOCKERS

A **cube term blocker** (CTB) for  $\mathbf{A}$  is a pair  $(C, B)$  of subuniverses of  $\mathbf{A}$  satisfying  $\emptyset < C < B \leq A$  and for every term  $t(x_1, \dots, x_n)$  of  $\mathbf{A}$  there is an index  $i \in [n]$  such that

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

Marković, Maróti, McKenzie proved that a finite idempotent algebra has a cube term iff it possesses no CTB.

## CUBE TERM BLOCKERS

A **cube term blocker** (CTB) for  $\mathbf{A}$  is a pair  $(C, B)$  of subuniverses of  $\mathbf{A}$  satisfying  $\emptyset < C < B \leq A$  and for every term  $t(x_1, \dots, x_n)$  of  $\mathbf{A}$  there is an index  $i \in [n]$  such that

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

Marković, Maróti, McKenzie proved that a finite idempotent algebra has a cube term iff it possesses no CTB.

### LEMMA

*A finite CIB  $\mathbf{A} = \langle A, \cdot \rangle$  has a CTB if and only if  $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$ .*

## CUBE TERM BLOCKERS

A **cube term blocker** (CTB) for  $\mathbf{A}$  is a pair  $(C, B)$  of subuniverses of  $\mathbf{A}$  satisfying  $\emptyset < C < B \leq A$  and for every term  $t(x_1, \dots, x_n)$  of  $\mathbf{A}$  there is an index  $i \in [n]$  such that

$$(\forall (b_1, \dots, b_n) \in B^n)(b_i \in C \longrightarrow t(b_1, \dots, b_n) \in C).$$

Marković, Maróti, McKenzie proved that a finite idempotent algebra has a cube term iff it possesses no CTB.

### LEMMA

*A finite CIB  $\mathbf{A} = \langle A, \cdot \rangle$  has a CTB if and only if  $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$ .*

### PROOF.

If  $(C, B)$  is a CTB, then  $\theta = C^2 \cup (B - C)^2$  is a congruence of  $\mathbf{B} = \langle B, \cdot \rangle$  and  $\mathbf{B}/\theta \cong \mathbf{S}_2$ .

Conversely, suppose  $\mathbf{S}_2 \in \text{HS}(\mathbf{A})$ , and  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$  with  $\mathbf{B}/\theta$  a meet-SL for some  $\theta$ . Let  $C/\theta$  be the bottom of  $\mathbf{B}/\theta$ , then  $(C, B)$  is a CTB. □

## COLLAPSE FOR CIBS

Kearnes and Tschantz, “Automorphism groups of squares and of free algebras,” 2007.

### LEMMA

*If  $V$  is an idempotent variety that is not congruence permutable, then there are subuniverses  $U$  and  $W$  of  $\mathbf{F} := \mathbf{F}_V\{x, y\}$  (the 2-generated free algebra) satisfying*

1.  $x \in U \cap W$
2.  $y \in U^c \cap W^c$
3.  $(U \times F) \cup (F \times W) \leq \mathbf{F}^2$

## COLLAPSE FOR CIBS

Kearnes and Tschantz, “Automorphism groups of squares and of free algebras,” 2007.

### LEMMA

*If  $V$  is an idempotent variety that is not congruence permutable, then there are subuniverses  $U$  and  $W$  of  $\mathbf{F} := \mathbf{F}_V\{x, y\}$  (the 2-generated free algebra) satisfying*

1.  $x \in U \cap W$
2.  $y \in U^c \cap W^c$
3.  $(U \times F) \cup (F \times W) \leq \mathbf{F}^2$

For CIB's,  $U$  or  $W$  will be an ideal.

This implies a CTB and a semilattice.



## REMAINING QUESTION FOR CIBs

### CONCLUSION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \notin \mathbf{V}(\mathbf{A})$ . Then  $\text{CSP}(\mathbf{A})$  is tractable.

## REMAINING QUESTION FOR CIBs

### CONCLUSION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \notin V(\mathbf{A})$ . Then  $\text{CSP}(\mathbf{A})$  is tractable.

### OPEN QUESTION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \in V(\mathbf{A})$ . Is  $\text{CSP}(\mathbf{A})$  tractable?

Recall, for every  $\mathbf{A}$ ,

## REMAINING QUESTION FOR CIBs

### CONCLUSION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \notin V(\mathbf{A})$ . Then  $\text{CSP}(\mathbf{A})$  is tractable.

### OPEN QUESTION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \in V(\mathbf{A})$ . Is  $\text{CSP}(\mathbf{A})$  tractable?

Recall, for every  $\mathbf{A}$ ,

- if  $\mathbf{S}_2 \in V(\mathbf{A})$ , then  $V(\mathbf{A})$  is not CM;

## REMAINING QUESTION FOR CIBs

### CONCLUSION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \notin V(\mathbf{A})$ . Then  $\text{CSP}(\mathbf{A})$  is tractable.

### OPEN QUESTION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \in V(\mathbf{A})$ . Is  $\text{CSP}(\mathbf{A})$  tractable?

Recall, for every  $\mathbf{A}$ ,

- if  $\mathbf{S}_2 \in V(\mathbf{A})$ , then  $V(\mathbf{A})$  is not CM;
- if  $V(\mathbf{A})$  is  $\text{SD}_\wedge$ , then  $\text{CSP}(\mathbf{A})$  is tractable (in fact, always has a solution).

## REMAINING QUESTION FOR CIBs

### CONCLUSION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \notin V(\mathbf{A})$ . Then  $\text{CSP}(\mathbf{A})$  is tractable.

### OPEN QUESTION

Let  $\mathbf{A}$  be a CIB and  $\mathbf{S}_2 \in V(\mathbf{A})$ . Is  $\text{CSP}(\mathbf{A})$  tractable?

Recall, for every  $\mathbf{A}$ ,

- if  $\mathbf{S}_2 \in V(\mathbf{A})$ , then  $V(\mathbf{A})$  is not CM;
- if  $V(\mathbf{A})$  is  $\text{SD}_\wedge$ , then  $\text{CSP}(\mathbf{A})$  is tractable (in fact, always has a solution).

### REVISED QUESTION

Let  $\mathbf{A}$  be a CIB with  $\mathbf{S}_2$  in  $V(\mathbf{A})$ , not  $\text{SD}_\wedge$ . Is  $\text{CSP}(\mathbf{A})$  tractable?

## EXAMPLES

$\cdot$	0	1	2	3
0	0	0	0	1
1	0	1	3	2
2	0	3	2	1
3	1	2	1	3

$*$	0	1	2	3
0	0	0	1	1
1	0	1	3	2
2	1	3	2	1
3	1	2	1	3

$\circ$	0	1	2	3
0	0	0	2	1
1	0	1	3	2
2	2	3	2	1
3	1	2	1	3

Maroti's idea:

Bergman's idea: replace basic binary operation with a term from  $\text{Clo}(\mathbf{A})$ , say  $t(x, y) = (x * y) * x$ .

If  $\langle A, t \rangle$  tractable, then so is  $\langle A, * \rangle$