

CONGRUENCE LATTICES OF FINITE ALGEBRAS

William DeMeo

University of Hawai'i at Mānoa

Ph.D. Dissertation Defense

Committee

Ralph Freese (Advisor & Chair)


Peter Jipsen, Bill Lampe, J.B. Nation

Nick Kaiser (University Rep.)

April 16, 2012

CONGRUENCE DECOMPOSITIONS


CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

CONGRUENCE DECOMPOSITIONS

CONGRUENCE RELATIONS REVEAL STRUCTURAL CHARACTERISTICS.



The shape/structure of **Con A** tells us whether and how the algebra can be decomposed.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice L , does there exist a *finite* algebra A such that $\text{Con } A \cong L$?

DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice \mathbf{L} , does there exist a *finite* algebra \mathbf{A} such that $\mathbf{Con A} \cong \mathbf{L}$?

DEFINITION

We call a finite lattice *representable* if it is isomorphic to the congruence lattice of a finite algebra.

THE FINITE LATTICE REPRESENTATION PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

THEOREM (GRÄTZER-SCHMIDT, 1963)

Every algebraic lattice is isomorphic to the congruence lattice of an algebra.

What if the algebra is finite?

Problem: Given a finite lattice \mathbf{L} , does there exist a *finite* algebra \mathbf{A} such that $\mathbf{Con A} \cong \mathbf{L}$?

DEFINITION

We call a finite lattice **representable** if it is isomorphic to the congruence lattice of a finite algebra.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

• Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.

• Does equality hold in each case?

• $\mathcal{L}_0 = \mathcal{L}_1$

- (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?

- (Whitman, 1946) Yes, but proof requires infinite partition lattices.

- (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.

• **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?

• (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \supsetneq \mathcal{L}_3 \supsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

• Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.

• Does equality hold in each case?

• $\mathcal{L}_0 = \mathcal{L}_1$

• (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?

• (Whitman, 1946) Yes, but proof requires infinite partition lattices.

• (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.

• **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?

• (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \supsetneq \mathcal{L}_3 \supsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

• Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.

• Does equality hold in each case?

• $\mathcal{L}_0 = \mathcal{L}_1$

• (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?

• (Whitman, 1946) Yes, but proof requires infinite partition lattices.

• (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.

• **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?

• (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \supsetneq \mathcal{L}_3 \supsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does not say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \supsetneq \mathcal{L}_3 \supsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does not say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \supsetneq \mathcal{L}_3 \supsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \subsetneq \mathcal{L}_3 \subsetneq \mathcal{L}_4$.

SOME IMPORTANT CLASSES OF FINITE LATTICES

- \mathcal{L}_0 = all finite lattices
- \mathcal{L}_1 = lattices isomorphic to sublattices of finite partition lattices
- \mathcal{L}_2 = ...strong congruence lattices of finite partial algebras
- \mathcal{L}_3 = ...congruence lattices of finite algebras
- \mathcal{L}_4 = ...intervals in subgroup lattices of finite groups
- \mathcal{L}_5 = ...subgroup lattices of finite groups

- Clearly $\mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_2 \supseteq \mathcal{L}_3 \supseteq \mathcal{L}_4 \supseteq \mathcal{L}_5$.
- Does equality hold in each case?
- $\mathcal{L}_0 = \mathcal{L}_1$
 - (Birkhoff, 1935) Is every lattice \cong a sublattice of a partition lattice?
 - (Whitman, 1946) Yes, but proof requires infinite partition lattices.
 - (Pudlák and Tůma, 1980) $\mathcal{L}_0 = \mathcal{L}_1$.
- **Main problem:** Is $\mathcal{L}_0 = \mathcal{L}_3$ true?
- (Pálffy and Pudlák, 1980) $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.
This does **not** say $\mathcal{L}_3 = \mathcal{L}_4$. It's possible that $\mathcal{L}_0 \subsetneq \mathcal{L}_3 \subsetneq \mathcal{L}_4$.

RECAP

THEOREM (PUDLÁK AND TŮMA, 1980)

Every finite lattice can be embedded in $\text{Eq}(X)$ with X finite.

In other words, $\mathcal{L}_0 = \mathcal{L}_1$.

THEOREM (PÁLFY AND PUDLÁK, 1980)

The following statements are equivalent:

- (I) *Every finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (II) *Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

In other words, $\mathcal{L}_0 = \mathcal{L}_3$ if and only if $\mathcal{L}_0 = \mathcal{L}_4$.

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 1 (USE CLOSURE PROPERTIES)

The class \mathcal{L}_3 is closed under the following operations:

- lattice duals (Kurzweil and Netter, 1986)
- interval sublattices (follows from Kurzweil-Netter)
- direct products (Tůma, 1986)
- ordinal sums (McKenzie, 1984; Snow, 2000)
- parallel sums (Snow, 2000)
- certain sublattices of lattices in \mathcal{L}_3 (Snow, 2000)
(namely, those obtained as a union of a filter and ideal)

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 2 (USE A GALOIS CORRESPONDENCE)

- Fix $\theta \subseteq X \times X$, $f : X^n \rightarrow X$.

Say that f **respects** θ and write $f(\theta) \subseteq \theta$ provided

$$(x_i, y_i) \in \theta \Rightarrow (f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \theta.$$

- For $L \subseteq \text{Eq}(X)$ define

$$\lambda(L) = \{f \in X^X \mid (\forall \theta \in L) f(\theta) \subseteq \theta\},$$

the set of unary maps on X which respect all relations in L .

- For $F \subseteq X^X$ define

$$\rho(F) = \{\theta \in \text{Eq}(X) \mid (\forall f \in F) f(\theta) \subseteq \theta\},$$

the set of equivalence relations on X respected by all $f \in F$.

- Then $L \subseteq \rho\lambda(L)$ and $\rho\lambda$ is a *closure operator* on $\text{Sub}[\text{Eq}(X)]$.
(idempotent, extensive, order preserving)
- If a lattice $L \leq \text{Eq}(X)$ is *closed*, i.e. $\rho\lambda(L) = L$, then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 3 (SUBGROUP LATTICE INTERVAL)

Find L as an interval in a subgroup lattice of a finite group.

If $H \leq G$ are finite groups, then the interval above H in $Sub(G)$,

$$[H, G] := \{K \mid H \leq K \leq G\},$$

is isomorphic to $\text{Con} \langle G/H, G \rangle$.

HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.


Suppose $L_0 \cong \text{Con} \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.

Consider $L = \alpha^\uparrow \cup \beta^\downarrow$.

Then there exists a set $F' \subset A^A$ such that

$$L \cong \text{Con} \langle A, F \cup F' \rangle.$$

$$L \leq L_0$$



HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.


Suppose $L_0 \cong \text{Con} \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.

Consider $L = \alpha^\uparrow \cup \beta^\downarrow$.

Then there exists a set $F' \subset A^A$ such that

$$L \cong \text{Con} \langle A, F \cup F' \rangle.$$

$$L \leq L_0$$



HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

METHOD 4 (FILTER+IDEAL)

Find L as the union of a filter and ideal in a representable lattice.


Suppose $L_0 \cong \text{Con} \langle A, F \rangle$, $\alpha, \beta \in L_0 \setminus \{0, 1\}$.

Consider $L = \alpha^\uparrow \cup \beta^\downarrow$.


Then there exists a set $F' \subset A^A$ such that

$$L \cong \text{Con} \langle A, F \cup F' \rangle.$$

$$L \leq L_0$$



LATTICES WITH AT MOST 6 ELEMENTS ARE REPRESENTABLE.




Watatani (1996) *J. Funct. Anal.*


Aschbacher (2008) *JAMS*

Theorem: *Every lattice with at most 6 elements is an interval in the subgroup lattice of a finite group.*


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19}




L_{20}




L_{17}




L_{13}



L_{11}




L_9




L_7


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19}




L_{20}




L_{17}




L_{13}



L_{11}




L_9



L_7

FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$

- The group $G = (A_4 \times A_4) \rtimes C_2$ has a subgroup $H \cong S_3$ such that $[H, G] \cong L_{17}$.
- ...so the dual L_{16} is also representable.




SmallGroup(960,11358)

$$|G : H| = 80$$

- The group $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$ has a subgroup $H \cong A_4$ such that $[H, G] \cong L_{13}$.

FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$

- The group $G = (A_4 \times A_4) \rtimes C_2$ has a subgroup $H \cong S_3$ such that $[H, G] \cong L_{17}$.
- ...so the dual L_{16} is also representable.




SmallGroup(960,11358)

$$|G : H| = 80$$

- The group $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$ has a subgroup $H \cong A_4$ such that $[H, G] \cong L_{13}$.

FINDING REPRESENTATIONS...


...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$

- The group $G = (A_4 \times A_4) \rtimes C_2$ has a subgroup $H \cong S_3$ such that $[H, G] \cong L_{17}$.
- ...so the dual L_{16} is also representable.



SmallGroup(960,11358)


$$|G : H| = 80$$

- The group $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$ has a subgroup $H \cong A_4$ such that $[H, G] \cong L_{13}$.

FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$ is a congruence lattice (of A_4 acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$ is a congruence lattice (of A_4 acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$ is a congruence lattice (of A_4 acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

FINDING REPRESENTATIONS...



...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)



- Let $G = (A_4 \times A_4) \rtimes C_2$.
- G has a subgroup $H \cong C_6$ with $[H, G] \cong N_5$.
- Let $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$.
- $\text{Sub}(G)$ is a congruence lattice, so if there exists a subgroup $K \succ 1$, below β and not below γ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$




- $\text{Sub}(A_4)$ is a congruence lattice (of A_4 acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓




L_9 ✓




L_7


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓



L_9 ✓




L_7

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.

- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.

- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α "doubled" in $\text{Con } \mathbf{A}$.

STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α “doubled” in $\text{Con } \mathbf{A}$.

STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.

Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α “doubled” in $\text{Con } \mathbf{A}$.

STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.


Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α “doubled” in $\text{Con } \mathbf{A}$.



STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$


$$B_2 = \{11, 12, 13, 14, 15\}.$$

STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.


Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α “doubled” in $\text{Con } \mathbf{A}$.



STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$


$$B_2 = \{11, 12, 13, 14, 15\}.$$

STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.

STEP 1 Take a permutational algebra $\mathbf{B} = \langle B, F \rangle$ with congruence lattice $\text{Con } \mathbf{B} \cong M_4$.


Example:



- Let $B = \{0, 1, \dots, 5\}$ index the elements of S_3 and consider the right regular action of S_3 on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$ and $g_1 = (0, 1, 2)(3, 4, 5)$ generate this action group, the image of $S_3 \hookrightarrow S_6$.
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$ with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

Goal: expand \mathbf{B} to an algebra \mathbf{A} that has α “doubled” in $\text{Con } \mathbf{A}$.



STEP 2 Since $\alpha = \text{Cg}^B(0, 2)$, we let $A = B_0 \cup B_1 \cup B_2$ where


$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

STEP 3 Define unary operations $e_0, e_1, e_2, s, g_0 e_0$, and $g_1 e_0$.

CONSTRUCTION OF AN ALGEBRA \mathbf{A} WITH $\text{Con } \mathbf{A} \cong L_9$.




$\text{Con} \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con} \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$


$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$.




$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con } \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

$$\alpha = \alpha^* \cap B^2 = \hat{\alpha} \cap B^2, \quad \beta = \beta^* \cap B^2, \quad \dots$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

B_0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \twoheadrightarrow B_0$$

$$e_1: A \twoheadrightarrow B_1$$

$$e_2: A \twoheadrightarrow B_2$$

$$s: A \twoheadrightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \left\{ \begin{array}{cccccc} & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ 11 & 12 & 2 & 13 & 14 & 15 \end{array} \right\}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

	13	11
	14	12
0	15	2
3	4	5

B_0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \twoheadrightarrow B_0$$

$$e_1: A \twoheadrightarrow B_1$$

$$e_2: A \twoheadrightarrow B_2$$

$$s: A \twoheadrightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15\}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

B_0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

B_1			13	11	
10	9	8	14	12	
7	6	0	15	2	
			3	4	5
			B_0		

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

B_1

10	9	8		
7	6	0	1	2
		3	4	5

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

	B_1	2	5
10	9	8	4
7	6	0	3

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

B_1

10	9	8
7	6	0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



Con $\langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

WHY DOES IT WORK?


$$\text{Con } \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con } \langle A, F_A \rangle$$

Why don't the β classes of B_1 and B_2 mix?

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$ to have non-trivial inverse image $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$.
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



RESIDUATION LEMMA

- Define $\widehat{} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each $\beta \in \text{Con } \mathbf{B}$, let $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$. That is,

$$\beta^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to $\text{Con } \mathbf{B}$.

LEMMA

- (I) $\beta^* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is a residuated mapping with residual $|_B$.
- (II) $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$ is a residuated mapping with residual $\widehat{}$.
- (III) For all $\alpha \in \text{Con } \mathbf{A}$, $\beta \in \text{Con } \mathbf{B}$,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular, $\beta^*|_B = \beta = \widehat{\beta}|_B$.

RESIDUATION LEMMA

- Define $\widehat{} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each $\beta \in \text{Con } \mathbf{B}$, let $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$. That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to $\text{Con } \mathbf{B}$.

LEMMA

- (I) $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is a residuated mapping with residual $|_B$.
- (II) $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$ is a residuated mapping with residual $\widehat{}$.
- (III) For all $\alpha \in \text{Con } \mathbf{A}$, $\beta \in \text{Con } \mathbf{B}$,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular, $\beta^*|_B = \beta = \widehat{\beta}|_B$.

RESIDUATION LEMMA

- Define $\widehat{} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each $\beta \in \text{Con } \mathbf{B}$, let $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$. That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to $\text{Con } \mathbf{B}$.

LEMMA


- (I) $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$ is a residuated mapping with residual $|_B$.
- (II) $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$ is a residuated mapping with residual $\widehat{}$.
- (III) For all $\alpha \in \text{Con } \mathbf{A}$, $\beta \in \text{Con } \mathbf{B}$,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular, $\beta^|_B = \beta = \widehat{\beta}|_B$.*

THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$.

- If $\beta \in \mathbf{Con B}$ is a coatom of $\mathbf{Con B}$ with m congruence classes then the interval $[\beta^*, \hat{\beta}]$ in $\mathbf{Con A}$ is 2^{m-1} .




More generally...

- Suppose $\beta \in \mathbf{Con B}$ has transversal $b_{\beta(1)}, \dots, b_{\beta(m)}$.
- Denote by T_r the set of intersection points in the r -th block of β :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$.

- If $\beta \in \mathbf{Con B}$ is a coatom of $\mathbf{Con B}$ with m congruence classes then the interval $[\beta^*, \hat{\beta}]$ in $\mathbf{Con A}$ is 2^{m-1} .




More generally...

- Suppose $\beta \in \mathbf{Con B}$ has transversal $b_{\beta(1)}, \dots, b_{\beta(m)}$.
- Denote by T_r the set of intersection points in the r -th block of β :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$.

- If $\beta \in \mathbf{Con B}$ is a coatom of $\mathbf{Con B}$ with m congruence classes then the interval $[\beta^*, \hat{\beta}]$ in $\mathbf{Con A}$ is 2^{m-1} .



More generally...

- Suppose $\beta \in \mathbf{Con B}$ has transversal $b_{\beta(1)}, \dots, b_{\beta(m)}$.
- Denote by T_r the set of intersection points in the r -th block of β :


$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

$$\text{Then } [\beta^*, \hat{\beta}] = \{\theta \in \mathbf{Eq(A)} : \beta^* \subseteq \theta \subseteq \hat{\beta}\} \cong \prod_{r=1}^m (\mathbf{Eq} | T_r)^{m-1}.$$


SLIGHTLY MORE GENERAL EXAMPLES...

Returning to our original example, the base algebra **B** is the right regular S_3 -set, and the nontrivial relations in $\text{Con } \mathbf{B}$ are


$$\alpha = |0, 1, 2|3, 4, 5| \quad \beta = |0, 3|1, 4|2, 5| \quad \gamma = |0, 4|1, 5|2, 3| \quad \delta = |0, 5|1, 3|2, 4|$$




$$T = \{0, 1\}$$




$$T = \{0, 1, 2\}$$



$$T = \{0, 2, 3\}$$



$$T = \{0, 1, 2, 3\}$$



$$T = \{0, 2, 3, 5\}$$

$$L \cong 2^2 \times 2^2$$

LIMITATIONS

Two limitations of the foregoing construction:

- 1 The sizes $|T_r|$ of the partition lattice factors in

$$[\beta^*, \hat{\beta}] \cong \prod_{r=1}^m (\text{Eq}|T_r|)^{m-1}$$

are limited by the size of the blocks of β .

- 2 If β is not principal, $[\theta^*, \hat{\theta}]$ may be non-trivial for some $\theta \not\leq \beta$.

A GENERALIZATION

THEOREM

Let $\mathbf{B} = \langle B, F \rangle$ be a finite algebra. Suppose


$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has m blocks and fix $N < \infty$.

There exists an overalgebra $\langle A, F_A \rangle$ such that the interval $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$ is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that $\theta^* = \widehat{\theta}$ for all $\theta \not\geq \beta$ in $\text{Con } \mathbf{A}$.



A GENERALIZATION

THEOREM

Let $\mathbf{B} = \langle B, F \rangle$ be a finite algebra. Suppose


$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has m blocks and fix $N < \infty$.


There exists an overalgebra $\langle A, F_A \rangle$ such that the interval $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$ is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$


Moreover, we can arrange it so that $\theta^* = \widehat{\theta}$ for all $\theta \not\geq \beta$ in $\text{Con } \mathbf{A}$.




SEVEN ELEMENT LATTICES: SUMMARY




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓




L_9 ✓




L_7


SEVEN ELEMENT LATTICES: SUMMARY




L_{19} ✓




L_{20} ✓




L_{17} ✓




L_{13} ✓



L_{11} ✓



L_9 ✓



L_7

HAS ANYONE SEEN THIS LATTICE?

mathoverflow

Questions

Tags

Users

Badges

Unanswered


Ask Question

Given a lattice L with n elements, are there finite groups $H < G$ such that $L \cong$ the lattice of subgroups between H and G ?

13



If there is no restriction on n , this is a famous [open problem](#). I'm wondering if any recent work has been done for small $n > 6$. I believe the question is answered (positively) for $n = 6$ by Watatani (1996) [MR1409040](#) and Aschbacher (2008) [MR2393428](#). I also believe we can answer it for $n = 7$, with one possible exception. The exceptional case is shown below.



So my two questions are these:

1) Does anyone know of recent work on this special case of the problem (specifically for $n = 7$ or $n = 8$)?

2) Has anyone found a finite group G with a subgroup H such that the interval

$$[H, G] = \{K : H \leq K \leq G\}$$

is the lattice shown above?

tagged

[finite-groups](#) × 277

[open-problem](#) × 195

[lattices](#) × 129

[universal-algebra](#) × 53

[congruences](#) × 6

asked

3 months ago

viewed


401 times

Tip: You can search for questions with arbitrary boolean combinations of tags (like [this](#)). See tip 12 for details on how. [See more tips and tricks.](#)

MathJax trouble? [\(Re\)process math with jsMath.](#)


INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$.
What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$. What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.




THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, $L_7 \cong [H, G]$.

- (I) G is a primitive permutation group.
- (II) If $N \triangleleft G$, then $C_G(N) = 1$.
- (III) G contains no non-trivial abelian normal subgroup.
- (IV) G is not solvable.
- (V) G is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, all proper subgroups in the interval $[H, G]$ are core-free.

INTERVAL SUBLATTICE ENFORCEABLE PROPERTIES

- L_7 cannot be obtained using the overalgebra construction.
- A minimal representation of L_7 must come from a transitive G -set.
- Suppose $L_7 \cong [H, G]$ for some finite groups $H < G$. What can we say about the group G ?
- If we prove G must have certain properties, then FLRP has a positive answer iff every finite lattice is an interval in the subgroup lattice of a group *satisfying all of these properties*.



THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, $L_7 \cong [H, G]$.

- (I) G is a primitive permutation group.
- (II) If $N \triangleleft G$, then $C_G(N) = 1$.
- (III) G contains no non-trivial abelian normal subgroup.
- (IV) G is not solvable.
- (V) G is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, all proper subgroups in the interval $[H, G]$ are core-free.

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:


$$UH = \langle U, H \rangle \iff UH = HU.$$

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$




SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee H = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:

$$UH = \langle U, H \rangle \iff UH = HU.$$




$$[U_0, U] := \{V \mid U_0 \leq V \leq U\}$$

SUBGROUP LATTICE BASICS

Let U and H be subgroups of a finite group.

- By UH we mean the set $\{uh \mid u \in U, h \in H\}$.
- $U \vee V = \langle U, H \rangle$ means the group generated by U and H .
- $UH \subseteq \langle U, H \rangle$ and equality holds iff U and H permute:


$$UH = \langle U, H \rangle \iff UH = HU.$$



INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the *H-permuting subgroups*.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the *H-invariant subgroups*: $h V h^{-1} = V$ for all $h \in H$.

INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the *H-permuting subgroups*.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the *H-invariant subgroups*: $h V h^{-1} = V$ for all $h \in H$.

INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $h V h^{-1} = V$ for all $h \in H$.

INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $h V h^{-1} = V$ for all $h \in H$.

INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define

$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $h V h^{-1} = V$ for all $h \in H$.


LEMMA

- $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$
- If $U \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$.
- If $H \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H = [U_0, U]$.

INTERVAL ISOMORPHISMS

- Assume $H \trianglelefteq \langle U, H \rangle$.

Then $UH = \langle U, H \rangle$ and $[U_0, U] \cong [H, UH]$.



- Now drop assumption $H \trianglelefteq \langle U, H \rangle$, assume $UH = \langle U, H \rangle$, and define

$$[U_0, U]^H := \{V \in [U_0, U] \mid VH = HV\}.$$

the ***H-permuting subgroups***.

- If $U \trianglelefteq UH$, define


$$[U_0, U]_H := \{V \in [U_0, U] \mid H \leq N_{UH}(V)\},$$

the ***H-invariant subgroups***: $h V h^{-1} = V$ for all $h \in H$.

LEMMA

- $[H, UH] \cong [U_0, U]^H \leq [U_0, U]$ and $[U, UH] \cong [U_0, H]^U \leq [U_0, H]$
- If $U \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H \leq [U_0, U]$.
- If $H \trianglelefteq UH$, then $[U_0, U]_H = [U_0, U]^H = [U_0, U]$.

THE EXCEPTIONAL SEVEN ELEMENT LATTICE




THEOREM

Suppose $H < G$, $\text{core}_G(H) = 1$, and $L_7 \cong [H, G]$. Then

- (I) G is a primitive permutation group.
- (II) If $N \triangleleft G$, then $C_G(N) = 1$.
- (III) G contains no non-trivial abelian normal subgroup.
- (IV) G is not solvable.
- (V) G is subdirectly irreducible.
- (VI) With the possible exception of at most one maximal subgroup, M_1 or M_2 , all proper subgroups in the interval $[H, G]$ are core-free.

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .


Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .


Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .


Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .


Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!

IDEA OF THE PROOF



Claim: J_1 and J_2 are core-free subgroups of G .

Proof:

- If $N \triangleleft G$ then NH permutes with each subgroup containing H .
- If $1 \neq N \leq J_1$, then $NH = J_1$, so J_1 and K permute.
- Since $J_1 K = G$ and $J_1 \cap K = H$, our lemma yields

$$[J_1, G] \cong [H, K]^{J_1} = \{X \in [H, K] \mid J_1 X = X J_1\}.$$

Impossible!

ACHBACHER-O'NAN-SCOTT THEOREM

Let G be a primitive permutation group of degree d , and let $N := \text{Soc}(G) \cong T^m$ with $m \geq 1$. Then one of the following holds.

① N is regular and

- **Affine type** T is cyclic of order p , so $|N| = p^m$. Then $d = p^m$ and G is permutation isomorphic to a subgroup of the affine general linear group $\text{AGL}(m, p)$.
- **Twisted wreath product type** $m \geq 6$, the group T is nonabelian and G is a group of *twisted wreath product type*, with $d = |T|^m$.

② N is non-regular, non-abelian, and

- **Almost simple** $m = 1$ and $T \leq G \leq \text{Aut}(T)$.
- **Product action** $m \geq 2$ and G is permutation isomorphic to a subgroup of the product action wreath product $P \wr S_{m/l}$ of degree $d = nm/l$. The group P is primitive of type 2.(a) or 2.(c), P has degree n and $\text{Soc}(P) \cong T^l$, where $l \geq 1$ divides m .
- **Diagonal type** $m \geq 2$ and $T^m \leq G \leq T^m \cdot (\text{Out}(T) \times S_m)$, with the diagonal action. The degree $d = |T|^{m-1}$.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways.
- Using these and other methods we have found representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways.
- Using these and other methods we have found representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We developed techniques for putting restrictions on groups that have certain lattices as intervals in their subgroup lattices.

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways.
- Using these and other methods we have found representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We developed techniques for putting restrictions on groups that have certain lattices as intervals in their subgroup lattices.
- By studying the lattice L_7 , we can restrict the class of groups which could possibly represent this lattice. (We believe a slightly modified version of this lattice will restrict to the class of almost simple groups.)

CONCLUSIONS

- We have new methods for building new finite algebras out of old so that the congruence lattice changes in predictable ways.
- Using these and other methods we have found representations for all finite lattices with at most 7 elements with one exception, thus identifying the unique smallest lattice for which there's no known representation.
- We developed techniques for putting restrictions on groups that have certain lattices as intervals in their subgroup lattices.
- By studying the lattice L_7 , we can restrict the class of groups which could possibly represent this lattice. (We believe a slightly modified version of this lattice will restrict to the class of almost simple groups.)
- Future work: Restrict to almost simple groups and then solve the problem using the CFSG Theorem.

