

EXPANSIONS OF FINITE ALGEBRAS  
AND THEIR CONGRUENCE LATTICES

William DeMeo

joint work with

Ralph Freese, Peter Jipsen, Bill Lampe, J.B. Nation

University of Hawai'i at Mānoa

AMS Western Sectional Meeting  
March 4, 2012

## THE PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

**THEOREM (GRÄTZER-SCHMIDT, 1963)**

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

What if the algebra is finite?

**Problem:** Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{Con A} \cong \mathbf{L}$ ?

## THE PROBLEM

There is essentially no restriction on the shape of a congruence lattice of an arbitrary algebra.

### THEOREM (GRÄTZER-SCHMIDT, 1963)

*Every algebraic lattice is isomorphic to the congruence lattice of an algebra.*

What if the algebra is finite?

**Problem:** Given a finite lattice  $\mathbf{L}$ , does there exist a *finite* algebra  $\mathbf{A}$  such that  $\mathbf{Con A} \cong \mathbf{L}$ ?

### DEFINITION

We call a finite lattice **representable** if it is (isomorphic to) the congruence lattice of a finite algebra.

THEOREM (PÁLFY AND PUDLÁK, 1980)

*The following statements are equivalent:*

- (I) *Every finite lattice is isomorphic to the congruence lattice of a finite algebra.*
- (II) *Every finite lattice is isomorphic to an interval in the subgroup lattice of a finite group.*

# HOW TO FIND A REPRESENTATION OF A FINITE LATTICE

## METHOD 0

Write the given lattice in terms of other representable lattices.

- If  $L$  is representable, so is
  - the dual of  $L$
  - any interval sublattice of  $L$
- If  $L_1$  and  $L_2$  are representable, so is
  - the direct product of  $L_1$  and  $L_2$
  - the ordinal sum of  $L_1$  and  $L_2$
  - the parallel sum of  $L_1$  and  $L_2$

# HOW TO FIND A CONCRETE REPRESENTATION OF A FINITE LATTICE

## METHOD 1 (CLOSURE)

Find a “closed” representation of  $L$  in  $\text{Eq}(X)$ .

For  $L \leq \text{Eq}(X)$  define

$$\lambda(L) = \{f \in X^X : (\forall \theta \in L) f(\theta) \subseteq \theta\}$$

For  $F \subseteq X^X$  define

$$\rho(F) = \{\theta \in \text{Eq}(X) : (\forall f \in F) f(\theta) \subseteq \theta\}$$

For every  $L \leq \text{Eq}(X)$  we have  $L \subseteq \rho\lambda(L)$ .

The map  $\rho\lambda$  is a *closure operator* on  $\text{Sub}[\text{Eq}(X)]$ .

(idempotent, extensive, order preserving)

If a lattice  $L \leq \text{Eq}(X)$  is *closed*, i.e.  $\rho\lambda(L) = L$ , then

$$L = \text{Con} \langle X, \lambda(L) \rangle$$

# HOW TO FIND A CONCRETE REPRESENTATION OF A FINITE LATTICE

## METHOD 2 (SUBGROUP LATTICE INTERVAL)

Find  $L$  as an interval in a subgroup lattice of a finite group.

If  $H < G$  are finite groups, then the interval above  $H$  in  $Sub(G)$ ,

$$[H, G] := \{K : H \leq K \leq G\},$$

is isomorphic to  $\text{Con} \langle G/H, G \rangle$ .


## METHOD 3 (FILTER+IDEAL)

Find  $L$  as the union of a filter and ideal in a representable lattice.

## METHOD 3

### LEMMA

Suppose  $L_0 \cong \text{Con} \langle A, F \rangle$ , and  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ .






## METHOD 3

### LEMMA

Suppose  $L_0 \cong \text{Con} \langle A, F \rangle$ , and  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ . Consider  $L = \alpha^\uparrow \cup \beta^\downarrow$ .

$$L \leqslant L_0$$




## METHOD 3

### LEMMA

Suppose  $L_0 \cong \text{Con} \langle A, F \rangle$ , and  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ . Consider  $L = \alpha^\uparrow \cup \beta^\downarrow$ .

There exists a set  $F' \subset A^A$  such that  $L \cong \text{Con} \langle A, F \cup F' \rangle$ .

$$L \leq L_0$$



## METHOD 3

### LEMMA

Suppose  $L_0 \cong \text{Con} \langle A, F \rangle$ , and  $\alpha, \beta \in L_0 \setminus \{0, 1\}$ . Consider  $L = \alpha^\uparrow \cup \beta^\downarrow$ .

There exists a set  $F' \subset A^A$  such that  $L \cong \text{Con} \langle A, F \cup F' \rangle$ .

Proof:

Fix  $\theta \in L_0 \setminus L$ . Then  $\alpha \not\leq \theta \not\leq \beta$ , so

- $\exists (a, b) \in \alpha \setminus \theta$ ,
- $\exists (u, v) \in \theta \setminus \beta$ .

Define  $f_\theta : A \rightarrow A$  by


$$f_\theta(x) = \begin{cases} a & x \in u/\beta, \\ b & x \notin u/\beta. \end{cases}$$

Then


- $(f_\theta(u), f_\theta(v)) = (a, b) \notin \theta$ , so  $f_\theta(\theta) \not\leq \theta$ ,
- $\ker f_\theta \geq \beta$ , so  $f_\theta(\gamma) \subseteq \gamma$  for all  $\gamma \leq \beta$ ,
- $f_\theta(A) \subseteq \{a, b\}$ , so  $f_\theta(\gamma) \subseteq \gamma$  for all  $\gamma \geq \alpha$ .

Let  $F' = \{f_\theta : \theta \in L_0 \setminus L\}$ .

$L \leq L_0$



# LATTICES WITH AT MOST 6 ELEMENTS ARE REPRESENTABLE.




Watatani (1996) *J. Funct. Anal.*

Aschbacher (2008) *JAMS*

**Theorem:** Every lattice with at most 6 elements is an interval in the subgroup lattice of a finite group.

# LATTICES WITH AT MOST 6 ELEMENTS ARE REPRESENTABLE.




Watatani (1996) *J. Funct. Anal.*


Aschbacher (2008) *JAMS*

**Theorem:** *Every lattice with at most 6 elements is an interval in the subgroup lattice of a finite group.*


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$




$L_{20}$




$L_{17}$




$L_{13}$



$L_{11}$




$L_9$




$L_{10}$


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$




$L_{20}$




$L_{17}$




$L_{13}$



$L_{11}$




$L_9$



$L_{10}$


# FINDING REPRESENTATIONS...

...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$



SmallGroup(960,11358)

$$|G : H| = 80$$

- The group  $G = (A_4 \times A_4) \rtimes C_2$  has a subgroup  $H \cong S_3$  such that  $[H, G] \cong L_{17}$ .

- ...so the dual  $L_{16}$  is also representable.

$L_{16}$  can be embedded above diagonal of the direct power of a simple group,

$$L_{16} \hookrightarrow [\Gamma, S^{48}] \cong \text{Eq}(48)^{\text{dual}}.$$


Add the group operations  $G$  which closed  $L_{17}$ , and  $L_{16}$  appears as an upper interval in  $S^{48} \rtimes G$ .

- The group  $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$  has a subgroup  $H \cong A_4$  such that  $[H, G] \cong L_{13}$ .




# FINDING REPRESENTATIONS...

...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$



SmallGroup(960,11358)

$$|G : H| = 80$$

- The group  $G = (A_4 \times A_4) \rtimes C_2$  has a subgroup  $H \cong S_3$  such that  $[H, G] \cong L_{17}$ .
- ...so the dual  $L_{16}$  is also representable.

$L_{16}$  can be embedded above diagonal of the direct power of a simple group,


$$L_{16} \hookrightarrow [\Gamma, S^{48}] \cong \text{Eq}(48)^{\text{dual}}.$$

Add the group operations  $G$  which closed  $L_{17}$ , and  $L_{16}$  appears as an upper interval in  $S^{48} \rtimes G$ .

- The group  $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$  has a subgroup  $H \cong A_4$  such that  $[H, G] \cong L_{13}$ .


# FINDING REPRESENTATIONS...

...AS INTERVALS IN SUBGROUP LATTICES



SmallGroup(288,1025)

$$|G : H| = 48$$



SmallGroup(960,11358)

$$|G : H| = 80$$

- The group  $G = (A_4 \times A_4) \rtimes C_2$  has a subgroup  $H \cong S_3$  such that  $[H, G] \cong L_{17}$ .
- ...so the dual  $L_{16}$  is also representable.

$L_{16}$  can be embedded above diagonal of the direct power of a simple group,

$$L_{16} \hookrightarrow [\Gamma, S^{48}] \cong \text{Eq}(48)^{\text{dual}}.$$


Add the group operations  $G$  which closed  $L_{17}$ , and  $L_{16}$  appears as an upper interval in  $S^{48} \rtimes G$ .

- The group  $G = (C_2 \times C_2 \times C_2 \times C_2) \rtimes A_5$  has a subgroup  $H \cong A_4$  such that  $[H, G] \cong L_{13}$ .

# FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$  is a congruence lattice (of  $A_4$  acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

# FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$  is a congruence lattice (of  $A_4$  acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

# FINDING REPRESENTATIONS...


...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288, 1025)



- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$



- $\text{Sub}(A_4)$  is a congruence lattice (of  $A_4$  acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.

# FINDING REPRESENTATIONS...



...USING SUBGROUP LATTICE INTERVALS AND THE FILTER+IDEAL LEMMA.

SmallGroup(288,1025)



- Let  $G = (A_4 \times A_4) \rtimes C_2$ .
- $G$  has a subgroup  $H \cong C_6$  with  $[H, G] \cong N_5$ .
- Let  $[H, G] = \{H, \alpha, \beta, \gamma, G\} \cong N_5$ .
- $\text{Sub}(G)$  is a congruence lattice, so if there exists a subgroup  $K \succ 1$ , below  $\beta$  and not below  $\gamma$ , then

$$L_{11} \cong K^\downarrow \cup H^\uparrow.$$




- $\text{Sub}(A_4)$  is a congruence lattice (of  $A_4$  acting regularly on itself).
- Therefore,


$$L_{17} \cong V_4^\downarrow \cup P^\uparrow$$

is a congruence lattice.


# ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓




$L_9$  ✓




$L_{10}$


ARE ALL LATTICES WITH AT MOST 7 ELEMENTS REPRESENTABLE?




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓



$L_9$  ✓




$L_{10}$



# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .

Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.

- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .

- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  "doubled" in  $\text{Con } \mathbf{A}$ .

**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .

Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .

**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .

Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$


**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .

**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$




**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$


$$B_2 = \{11, 12, 13, 14, 15\}.$$

**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where

$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$


$$B_2 = \{11, 12, 13, 14, 15\}.$$

**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .

**STEP 1** Take a permutational algebra  $\mathbf{B} = \langle B, F \rangle$  with congruence lattice  $\text{Con } \mathbf{B} \cong M_4$ .


Example:



- Let  $B = \{0, 1, \dots, 5\}$  index the elements of  $S_3$  and consider the right regular action of  $S_3$  on itself.
- $g_0 = (0, 4)(1, 3)(2, 5)$  and  $g_1 = (0, 1, 2)(3, 4, 5)$  generate this action group, the image of  $S_3 \hookrightarrow S_6$ .
- $\text{Con } \langle B, \{g_0, g_1\} \rangle \cong M_4$  with congruences

$$\alpha = |012|345|, \beta = |03|14|25|, \gamma = |04|15|23|, \delta = |05|13|24|.$$

**Goal:** expand  $\mathbf{B}$  to an algebra  $\mathbf{A}$  that has  $\alpha$  “doubled” in  $\text{Con } \mathbf{A}$ .



**STEP 2** Since  $\alpha = \text{Cg}^B(0, 2)$ , we let  $A = B_0 \cup B_1 \cup B_2$  where


$$B_0 = \{0, 1, 2, 3, 4, 5\} = B$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 14, 15\}.$$

**STEP 3** Define unary operations  $e_0, e_1, e_2, s, g_0 e_0$ , and  $g_1 e_0$ .

# CONSTRUCTION OF AN ALGEBRA $\mathbf{A}$ WITH $\text{Con } \mathbf{A} \cong L_9$ .




$\text{Con} \langle B, \{g_0, g_1\} \rangle$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$\text{Con} \langle A, F_A \rangle$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$


$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

# CONSTRUCTION OF AN ALGEBRA **A** WITH $\text{Con } \mathbf{A} \cong L_9$ .




$\text{Con } \langle B, \{g_0, g_1\} \rangle$

$$\alpha = [0, 1, 2 | 3, 4, 5]$$

$$\beta = [0, 3 | 1, 4 | 2, 5]$$

$$\gamma = [0, 4 | 1, 5 | 2, 3]$$

$$\delta = [0, 5 | 1, 3 | 2, 4]$$



$\text{Con } \langle A, F_A \rangle$

$$\hat{\alpha} = [0, 1, 2, 6, 7, 11, 12 | 3, 4, 5 | 8, 9, 10, 13, 14, 15]$$

$$\alpha^* = [0, 1, 2, 6, 7, 11, 12 | 3, 4, 5 | 8, 9, 10 | 13, 14, 15]$$

$$\beta^* = [0, 3, 8 | 1, 4 | 2, 5, 15 | 6, 9 | 7, 10 | 11, 13 | 12, 14]$$

$$\gamma^* = [0, 4, 9 | 1, 5 | 2, 3, 13 | 6, 10 | 7, 8 | 11, 14 | 12, 15]$$

$$\delta^* = [0, 5, 10 | 1, 3 | 2, 4, 14 | 6, 8 | 7, 9, 11, 15 | 12, 13]$$

$$\alpha = \alpha^* \cap B^2 = \hat{\alpha} \cap B^2, \quad \beta = \beta^* \cap B^2, \quad \dots$$



## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

$B_0$

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$

- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations


$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$



$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$



$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

	13	11
	14	12
0	15	2
3	4	5

$B_0$

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array} \}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ \begin{array}{cccccc} 0 & 6 & 7 & 8 & 9 & 10 \end{array} \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \end{array} \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ \begin{array}{cccccc} 11 & 12 & 2 & 13 & 14 & 15 \end{array} \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

0	1	2
3	4	5

$B_0$

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$\begin{array}{cccccc} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \end{array}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$\begin{array}{cccccc} \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \end{array}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

$B_1$			13	11	
10	9	8	14	12	
7	6	0	15	2	
			3	4	5
			$B_0$		

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

# WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

$B_1$					
10	9	8			
7	6	0	1	2	
		3	4	5	

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

	$B_1$	2	5
10	9	8	4
7	6	0	3

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$



## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?

$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$

$B_1$

10	9	8
7	6	0

- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



- $A = B_0 \cup B_1 \cup B_2$
- Unary operations

$$e_0: A \rightarrow B_0$$

$$e_1: A \rightarrow B_1$$

$$e_2: A \rightarrow B_2$$

$$s: A \rightarrow B_0$$

$$B_1 = \{ 0 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \}$$

$$B_0 = \{ 0 \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \}$$

$$B_2 = \{ 11 \quad 12 \quad 2 \quad 13 \quad 14 \quad 15 \}$$

$$ge_0: A \xrightarrow{e_0} B_0 \xrightarrow{g} B_0$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$



## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con} \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con} \langle A, F_A \rangle$$

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## WHY DOES IT WORK?


$$\text{Con } \langle B, \{g_0, g_1\} \rangle$$

$$\alpha = |0, 1, 2|3, 4, 5|$$

$$\beta = |0, 3|1, 4|2, 5|$$

$$\gamma = |0, 4|1, 5|2, 3|$$

$$\delta = |0, 5|1, 3|2, 4|$$



$$\text{Con } \langle A, F_A \rangle$$

*Why don't the  $\beta$  classes of  $B_1$  and  $B_2$  mix?*

$$\hat{\alpha} = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10, 13, 14, 15|$$

$$\alpha^* = |0, 1, 2, 6, 7, 11, 12|3, 4, 5|8, 9, 10|13, 14, 15|$$

$$\beta^* = |0, 3, 8|1, 4|2, 5, 15|6, 9|7, 10|11, 13|12, 14|$$

$$\gamma^* = |0, 4, 9|1, 5|2, 3, 13|6, 10|7, 8|11, 14|12, 15|$$

$$\delta^* = |0, 5, 10|1, 3|2, 4, 14|6, 8|7, 9, 11, 15|12, 13|$$

## VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...



- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \widehat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$







## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## VARIATIONS ON THE SAME EXAMPLE...




- Suppose we want  $\beta = \text{Cg}^{\mathbf{B}}(0, 3) = |0, 3|2, 5|1, 4|$  to have non-trivial inverse image  $\beta|_B^{-1} = [\beta^*, \hat{\beta}]$ .
- Select elements 0 and 3 as intersection points:

$$A = B_0 \cup B_1 \cup B_2 \quad \text{where}$$

$$B_0 = \{0, 1, 2, 3, 4, 5\}$$

$$B_1 = \{0, 6, 7, 8, 9, 10\}$$

$$B_2 = \{11, 12, 13, 3, 14, 15\}.$$



## RESIDUATION LEMMA

- Define  $\widehat{\phantom{x}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to  $\text{Con } \mathbf{B}$ .

### LEMMA

- (I)  $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is a residuated mapping with residual  $|_B$ .
- (II)  $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is a residuated mapping with residual  $\widehat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \widehat{\beta}|_B$ .

## RESIDUATION LEMMA

- Define  $\widehat{\phantom{x}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to  $\text{Con } \mathbf{B}$ .

### LEMMA

- (I)  $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is a residuated mapping with residual  $|_B$ .
- (II)  $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is a residuated mapping with residual  $\widehat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \widehat{\beta}|_B$ .



## RESIDUATION LEMMA

- Define  $\widehat{\phantom{x}} : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  by

$$\widehat{\beta} = \{(x, y) \in A^2 : (ef(x), ef(y)) \in \beta \text{ for all } f \in \text{Pol}_1(\mathbf{A})\}.$$

- For each  $\beta \in \text{Con } \mathbf{B}$ , let  $\beta^* = \text{Cg}^{\mathbf{A}}(\beta)$ . That is,

$$* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$$

is the congruence generation operator restricted to  $\text{Con } \mathbf{B}$ .

### LEMMA


- (I)  $* : \text{Con } \mathbf{B} \rightarrow \text{Con } \mathbf{A}$  is a residuated mapping with residual  $|_B$ .
- (II)  $|_B : \text{Con } \mathbf{A} \rightarrow \text{Con } \mathbf{B}$  is a residuated mapping with residual  $\widehat{\phantom{x}}$ .
- (III) For all  $\alpha \in \text{Con } \mathbf{A}$ ,  $\beta \in \text{Con } \mathbf{B}$ ,

$$\beta = \alpha|_B \quad \Leftrightarrow \quad \beta^* \leq \alpha \leq \widehat{\beta}.$$

In particular,  $\beta^*|_B = \beta = \widehat{\beta}|_B$ .

## THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .




*More generally...*

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

## THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .




*More generally...*

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :

$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

## THE STRUCTURE OF THE INTERVAL $[\beta^*, \hat{\beta}] \leq \mathbf{Con A}$ .

- If  $\beta \in \mathbf{Con B}$  is a coatom of  $\mathbf{Con B}$  with  $m$  congruence classes then the interval  $[\beta^*, \hat{\beta}]$  in  $\mathbf{Con A}$  is  $2^{m-1}$ .



More generally...

- Suppose  $\beta \in \mathbf{Con B}$  has transversal  $b_{\beta(1)}, \dots, b_{\beta(m)}$ .
- Denote by  $T_r$  the set of intersection points in the  $r$ -th block of  $\beta$ :


$$T_r = \bigcup_{k=1}^K B_k \cap b_{\beta(r)}/\beta.$$

$$\text{Then } [\beta^*, \hat{\beta}] = \{\theta \in \mathbf{Eq(A)} : \beta^* \subseteq \theta \subseteq \hat{\beta}\} \cong \prod_{r=1}^m (\mathbf{Eq} | T_r|)^{m-1}.$$


## SLIGHTLY MORE GENERAL EXAMPLES...

Returning to our original example, the base algebra **B** is the right regular  $S_3$ -set, and the nontrivial relations in  $\text{Con } \mathbf{B}$  are


$$\alpha = |0, 1, 2|3, 4, 5| \quad \beta = |0, 3|1, 4|2, 5| \quad \gamma = |0, 4|1, 5|2, 3| \quad \delta = |0, 5|1, 3|2, 4|$$




$$T = \{0, 1\}$$




$$T = \{0, 1, 2\}$$



$$T = \{0, 2, 3\}$$



$$T = \{0, 1, 2, 3\}$$



$$T = \{0, 2, 3, 5\}$$

$$L \cong 2^2 \times 2^2$$

## LIMITATIONS

Two limitations of the foregoing construction:

- 1 The sizes  $|T_r|$  of the partition lattice factors in

$$[\beta^*, \widehat{\beta}] \cong \prod_{r=1}^m (\text{Eq}|T_r|)^{m-1}$$

are limited by the size of the blocks of  $\beta$ .

- 2 If  $\beta$  is not principal,  $[\theta^*, \widehat{\theta}]$  may be non-trivial for some  $\theta \not\leq \beta$ .

# A GENERALIZATION

## THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose


$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is

$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .



# A GENERALIZATION

## THEOREM

Let  $\mathbf{B} = \langle B, F \rangle$  be a finite algebra. Suppose


$$\beta = \text{Cg}^{\mathbf{B}}((a_1, b_1), \dots, (a_{K-1}, b_{K-1}))$$

has  $m$  blocks and fix  $N < \infty$ .

There exists an overalgebra  $\langle A, F_A \rangle$  such that the interval  $\beta|_B^{-1} \leq \text{Con } \mathbf{A}$  is


$$[\beta^*, \widehat{\beta}] \cong (\text{Eq}(N))^{m-1}.$$

Moreover, we can arrange it so that  $\theta^* = \widehat{\theta}$  for all  $\theta \not\geq \beta$  in  $\text{Con } \mathbf{A}$ .







## CONCLUDING REMARKS




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓




$L_9$  ✓




$L_{10}$


## CONCLUDING REMARKS




$L_{19}$  ✓




$L_{20}$  ✓




$L_{17}$  ✓




$L_{13}$  ✓



$L_{11}$  ✓



$L_9$  ✓



$L_{10}$

# HAS ANYONE SEEN THIS LATTICE?

[login](#) | [faq](#) | [how to ask](#) | [meta](#)

mathoverflow

Questions

Tags

Users

Badges


Unanswered

Ask Question

Given a lattice  $L$  with  $n$  elements, are there finite groups  $H < G$  such that  $L \cong$  the lattice of subgroups between  $H$  and  $G$ ?

13

If there is no restriction on  $n$ , this is a famous [open problem](#). I'm wondering if any recent work has been done for small  $n > 6$ . I believe the question is answered (positively) for  $n = 6$  by Watatani (1996) [MR1409040](#) and Aschbacher (2008) [MR2393428](#). I also believe we can answer it for  $n = 7$ , with one possible exception. The exceptional case is shown below.



So my two questions are these:

1) Does anyone know of recent work on this special case of the problem (specifically for  $n = 7$  or  $n = 8$ )?

2) Has anyone found a finite group  $G$  with a subgroup  $H$  such that the interval

$$[H, G] = \{K : H \leq K \leq G\}$$

is the lattice shown above?

Welcome to  
MathOverflow

A place for mathematicians  
to ask and answer  
questions.

tagged

[finite-groups](#) × 262

[open-problem](#) × 190

[lattices](#) × 123

[universal-algebra](#) × 51

[congruences](#) × 5


asked

1 month ago

viewed

358 times


## CONCLUDING REMARKS



$L_{10}$

- $L_{10}$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_{10}$  must come from a transitive  $G$ -set.
- If  $[H, G] \cong L_{10}$  with  $H$  core-free in  $G$  then
  - $G$  is a non-solvable primitive permutation group.
  - If  $N$  is a minimal normal subgroup of  $G$ , then  $N$  is nonabelian.


## CONCLUDING REMARKS



$L_{10}$


- $L_{10}$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_{10}$  must come from a transitive  $G$ -set.
- If  $[H, G] \cong L_{10}$  with  $H$  core-free in  $G$  then
  - $G$  is a non-solvable primitive permutation group.
  - If  $N$  is a minimal normal subgroup of  $G$ , then  $N$  is nonabelian.

## CONCLUDING REMARKS



- $L_{10}$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_{10}$  must come from a transitive  $G$ -set.
- If  $[H, G] \cong L_{10}$  with  $H$  core-free in  $G$  then
  - $G$  is a non-solvable primitive permutation group.
  - If  $N$  is a minimal normal subgroup of  $G$ , then  $N$  is nonabelian.

## CONCLUDING REMARKS



$L_{10}$

- $L_{10}$  cannot be obtained using the overalgebra construction.
- A minimal representation of  $L_{10}$  must come from a transitive  $G$ -set.
- If  $[H, G] \cong L_{10}$  with  $H$  core-free in  $G$  then
  - $G$  is a non-solvable primitive permutation group.
  - If  $N$  is a minimal normal subgroup of  $G$ , then  $N$  is nonabelian.

