

# HW 11

DUE Friday, April 27, 9am

## List of Exercises

**Section 6.1:** 16, (27), 28, (30)

**Section 6.2:** 10, 12, (13), 16, 24, (33)

**Section 6.3:** 6, 8, 12, (13), 16

---

### Section 6.1

**Exercises: 16, (27), 28, (30)**

**6.1.16.** Determine whether  $\mathbf{u} = \begin{bmatrix} 12 \\ 3 \\ -5 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} 2 \\ -3 \\ 3 \end{bmatrix}$  are orthogonal vectors.

**6.1.27.** (recommended)

Suppose a vector  $\mathbf{y}$  is orthogonal to vectors  $\mathbf{u}$  and  $\mathbf{v}$ . Show that  $\mathbf{y}$  is orthogonal to the vector  $\mathbf{u} + \mathbf{v}$ .

**6.1.28.** Suppose  $\mathbf{y}$  is orthogonal to  $\mathbf{u}$  and  $\mathbf{v}$ . Show that  $\mathbf{y}$  is orthogonal to every  $\mathbf{w}$  in  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$ . [Hint: An arbitrary  $\mathbf{w}$  in  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$  has the form  $\mathbf{w} = c_1\mathbf{u} + c_2\mathbf{v}$ ; show that  $\mathbf{y}$  is orthogonal to every such a vector.]

**6.1.30.** (recommended)

Let  $W$  be a subspace of  $\mathbb{R}^n$ , and let  $W^\perp$  be the set of all vectors orthogonal to  $W$ . Show that  $W^\perp$  is a subspace of  $\mathbb{R}^n$  using the following steps.

**a.** Take  $\mathbf{z}$  in  $W^\perp$ , and let  $\mathbf{u}$  represent any element of  $W$ . Then  $\mathbf{z} \cdot \mathbf{u} = 0$ . Take any scalar  $c$  and show that  $c\mathbf{z}$  is orthogonal to  $\mathbf{u}$ . (Since  $\mathbf{u}$  was an arbitrary element of  $W$ , this will show that  $c\mathbf{z}$  is in  $W^\perp$ .)

**b.** Take  $\mathbf{z}_1$  and  $\mathbf{z}_2$  in  $W^\perp$ , and let  $\mathbf{u}$  be any element of  $W$ . Show that  $\mathbf{z}_1 + \mathbf{z}_2$  is orthogonal to  $\mathbf{u}$ . What can you conclude about  $\mathbf{z}_1 + \mathbf{z}_2$ ? Why?

**c.** Finish the proof that  $W^\perp$  is a subspace of  $\mathbb{R}^n$ .

---

### Section 6.2

**Exercises: 10, 12, (13), 16, 24, (33)**

**6.2.10.** Let  $\mathbf{u}_1 = \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}$ ,  $\mathbf{u}_3 = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}$ , and  $\mathbf{x} = \begin{bmatrix} 5 \\ -3 \\ 1 \end{bmatrix}$ . Show that  $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  is an orthogonal basis for  $\mathbb{R}^3$ . Then express  $\mathbf{x}$  as a linear combination of the  $\mathbf{u}$ 's.

**6.2.12.** Compute the orthogonal projection of  $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$  onto the line through  $\begin{bmatrix} -1 \\ 3 \end{bmatrix}$ .

**6.2.13.** (recommended)

Let  $\mathbf{y} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$  and  $\mathbf{u} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$ . Write  $\mathbf{y}$  as the sum of two orthogonal vectors, one in  $\text{Span}\{\mathbf{u}\}$  and the other orthogonal to  $\mathbf{u}$ .

**6.2.16.** Let  $\mathbf{y} = \begin{bmatrix} -3 \\ 9 \end{bmatrix}$  and  $\mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ . Compute the distance from  $\mathbf{y}$  to the line passing through  $\mathbf{u}$  and the origin.

**6.2.24.** Mark each statement True or False. Justify each answer. All vectors are assumed to belong to  $\mathbb{R}^n$ .

- a.** Not every orthogonal set in  $\mathbb{R}^n$  is linearly independent.
- b.** If a set  $S = \{\mathbf{u}_1, \dots, \mathbf{u}_p\}$  has the property that  $\mathbf{u}_i \cdot \mathbf{u}_j = 0$  whenever  $i \neq j$ , then  $S$  is an orthonormal set.
- c.** If the columns of an  $m \times n$  matrix  $A$  are orthonormal, then the linear mapping  $\mathbf{x} \mapsto A\mathbf{x}$  preserves lengths.
- d.** The orthogonal projection of  $\mathbf{y}$  onto  $\mathbf{v}$  is the same as the orthogonal projection of  $\mathbf{y}$  onto  $c\mathbf{v}$  whenever  $c \neq 0$ .
- e.** An orthogonal matrix is invertible.

**6.2.33.** (recommended)

Suppose  $\mathbf{u}$  is a nonzero vector in  $\mathbb{R}^n$ , and let  $L = \text{Span}\{\mathbf{u}\}$ . Show that the mapping  $\mathbf{x} \mapsto \text{proj}_L \mathbf{x}$  is a linear transformation.

**Section 6.3****Exercises: 6, 8, 12, (13), 16**

**6.3.6.** Let  $\mathbf{y} = \begin{bmatrix} 6 \\ 4 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_1 = \begin{bmatrix} -4 \\ -1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ . Verify that  $\{\mathbf{u}_1, \mathbf{u}_2\}$  is an orthogonal set, and then find the orthogonal projection of  $\mathbf{y}$  onto  $\text{Span}\{\mathbf{u}_1, \mathbf{u}_2\}$ .

**6.3.8.** Let  $\mathbf{y} = \begin{bmatrix} -1 \\ 4 \\ 3 \end{bmatrix}$ ,  $\mathbf{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1 \\ 3 \\ -2 \end{bmatrix}$ . Let  $W$  be the subspace spanned by  $\mathbf{u}_1$  and  $\mathbf{u}_2$ , and write  $\mathbf{y}$  as the sum of a vector in  $W$  and a vector orthogonal to  $W$ .

**6.3.12.** Let  $\mathbf{y} = \begin{bmatrix} 3 \\ -1 \\ 1 \\ 13 \end{bmatrix}$ ,  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -2 \\ -1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -4 \\ 1 \\ 0 \\ 3 \end{bmatrix}$ . Find the closest point to  $\mathbf{y}$  in the subspace  $W$  spanned by  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .

**6.3.13.** (recommended)

Let  $\mathbf{z} = \begin{bmatrix} 3 \\ -7 \\ 2 \\ 3 \end{bmatrix}$ ,  $\mathbf{v}_1 = \begin{bmatrix} 2 \\ -1 \\ -3 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ -1 \end{bmatrix}$ . Find the best approximation to  $\mathbf{z}$  by a vector of the form

$$c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2.$$

**6.3.16.** Let  $\mathbf{y}$ ,  $\mathbf{v}_1$ ,  $\mathbf{v}_2$  be as in Exercise 12. Find the distance from  $\mathbf{y}$  to the subspace  $W$  spanned by  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .