MATH 325: Honours Ordinary Differential Equations

William Homier[1]

[1] *McGill University Physics, 3600 Rue University, Montréal, QC H3A 2T8, Canada*

January 6th, 2026

Abstract

Contents

1 Introduction **1**

2 Prerequisite Knowledge **1**
2.1 Real Analysis . . . 1
2.1.1 Norms . . . 1
2.1.2 Sequences and Convergence . . . 1
2.1.3 Uniform Convergence . . . 2
2.1.4 The Mean Value Inequality . . . 2

3 Intro, Classification, Theorem of Existence & Uniqueness **2**
3.1 Intro . . . 2
3.2 Classification . . . 3
3.2.1 The Order . . . 3
3.2.2 Scalar vs. System . . . 4
3.2.3 Linearity . . . 5
3.2.4 Autonomy . . . 5
3.3 Solutions and Initial Value Problems . . . 6
3.4 Existence and Uniqueness Theorem . . . 6
3.4.1 Lipschitz continuity . . . 6
3.4.2 Local Lipschitz continuity . . . 7
3.4.3 Integral Form of the IVP . . . 8
3.4.4 Picard operator . . . 8
3.4.5 Statement of the Theorem . . . 9
3.4.6 Picard Iteration and Existence . . . 10
3.4.7 Uniqueness . . . 10

4 First-Order Scalar Equation **11**
4.1 First Order Linear Equations . . . 11
4.1.1 Integrating Factor Method . . . 11
4.1.2 Existence and Uniqueness for First Order Linear Equations 12
4.2 Separable Equations . . . 12
4.2.1 General Method . . . 13
4.3 Exact Equations . . . 13
4.3.1 Potential Functions and Integral Curves . . . 13
4.3.2 Definition and Exactness Criterion . . . 14
4.3.3 Integrating Factors for Non-Exact Equations . . . 15
4.4 Applications . . . 16

5 Systems of Linear Equations **16**
5.1 From higher order scalar ODEs to linear systems . . . 17
5.2 Theory of First Order Linear Systems . . . 17
5.3 Structure of the Solution Space . . . 18
5.4 Linear homogeneous systems with constant coefficients . . . 21
5.4.1 Complex eigenvalues and real-valued solutions . . . 27

5.4.2 Nondiagonalizable Matrices 29
5.5 Exponential of a Matrix 32
5.6 Building the Solution Set With e^{At} 34
5.6.1 Eigenvectors 34
5.6.2 Generalized Eigenvectors 34
5.7 Non-homogeneous systems 37
5.7.1 Variation of Parameters 37
5.8 Mass-Spring Systems 38
5.9 RLC Circuits 40

6 Stability, Phase Portraits and Orbits **41**
6.1 Orbits and Phase Portraits 41
6.2 Equilibrium Solutions 41
6.3 Stability 42
6.3.1 Definitions 42
6.3.2 Examples 42
6.4 Phase Portraits of Planar Linear Systems 43

7 Laplace Transform **44**

8 Appendix **44**
8.1 Picard Iterations Convergence 44

9 Solutions **45**

10 Useful Links **48**

1 Introduction

Jean-Philippe Lessard (Burnside 1119). Tutorials every wednesday from 9am to 10am, ENGTR 0070, with Eunpyo Bang. Office hours thursday. No textbooks. 25% assignments (2 written assignments 15%, and 5 webworks 10%). 25% Midterm (February 16 - inclass). 50% Final. Since its honours you will deal with analysis. The Midterm is from 4:05 pm to 5:25 pm in MAASS 112, which will include chapters 1, 2, 3 and will be 5 questions long.

2 Prerequisite Knowledge

2.1 Real Analysis

2.1.1 Norms

Definition 1 (Norm)**.** *A norm $\|\cdot\|$ on $\mathbb{R}^n$ is a way of measuring the "size" or "length" of a vector $x \in \mathbb{R}^n$. It must satisfy:*

- $\|x\| \geq 0$, *and* $\|x\| = 0$ *only if* $x = 0$,
- $\|\alpha x\| = |\alpha| \, \|x\|$ *for any scalar* α,
- $\|x + y\| \leq \|x\| + \|y\|$ *(triangle inequality).*

Example. *The three most common norms on $\mathbb{R}^n$ are:*

- ***Euclidean norm:*** $\|x\|_2 = \sqrt{x_1^2 + \cdots + x_n^2}$ *(standard geometric length),*
- ***Supremum norm:*** $\|x\|_\infty = \max\{|x_1|, \ldots, |x_n|\}$ *(size of the largest component),*
- ℓ^1 ***norm:*** $\|x\|_1 = |x_1| + \cdots + |x_n|$ *(sum of all components, also called the taxicab norm).*

Remark 1. *In this course, any norm can be used unless stated otherwise. The choice of norm does not affect whether a solution exists or is unique.*

2.1.2 Sequences and Convergence

Definition 2 (Cauchy Sequence)**.** *A sequence of vectors $\{x_k\} \subset \mathbb{R}^n$ is called a Cauchy sequence if its terms eventually get arbitrarily close to one another: for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that*

$$\|x_m - x_n\| < \varepsilon \quad \text{for all } m, n \geq N.$$

Theorem 1. *Every Cauchy sequence in $\mathbb{R}^n$ converges to a point in $\mathbb{R}^n$.*

Remark 2. *This is the key fact used in Picard iteration: we show that the sequence of approximate solutions $\{y_k\}$ is Cauchy, and therefore it must converge to an actual solution.*

2.1.3 Uniform Convergence

Definition 3 (Uniform Convergence)**.** *A sequence of functions $f_k : J \to \mathbb{R}^n$ converges* uniformly *to $f : J \to \mathbb{R}^n$ if*

$$\lim_{k\to\infty} \sup_{t\in J} \|f_k(t) - f(t)\| = 0.$$

In plain terms: the approximation error $\|f_k(t) - f(t)\|$ becomes small simultaneously *for all $t \in J$, not just pointwise.*

Theorem 2. *If $f_k \to f$ uniformly on J and each f_k is continuous, then f is also continuous. Moreover,*

$$\lim_{k\to\infty} \int_J f_k(t)\,dt = \int_J f(t)\,dt.$$

Remark 3. *The second statement says that for uniformly convergent sequences, you can swap the limit and the integral. This is used in the Picard iteration proof to pass the limit inside the integral and confirm that the limit function actually solves the IVP.*

2.1.4 The Mean Value Inequality

Theorem 3 (Mean Value Inequality)**.** *Let $U \subseteq \mathbb{R}^n$ be open and $f : U \to \mathbb{R}^n$ be C^1. For any compact set $K \subset U$ and all $x, y \in K$,*

$$\|f(y) - f(x)\| \le \left(\sup_{z\in K} \|Df(z)\|\right) \|y - x\|,$$

where Df denotes the Jacobian matrix of f.

Remark 4. *This generalizes the familiar scalar fact $|f(b) - f(a)| \le \sup |f'| \cdot |b - a|$ to higher dimensions. Its main consequence for this course is that any C^1 function is locally Lipschitz: on a compact set K, the sup of $\|Df\|$ is finite, providing a Lipschitz constant. This is why the hypothesis "f is C^1" is often used as a sufficient condition for the Existence and Uniqueness Theorem.*

3 Intro, Classification, Theorem of Existence & Uniqueness

3.1 Intro

January 06, 2026.

Definition 4 (Differential Equation)**.** *A differential equation (DE) is a relation that involves an unknown function and some of its derivatives.*

Definition 5 (Ordinary Differential Equation)**.** *An ordinary differential equation (ODE) is a differential equation whose unknown function depends on one independent variable only.*

Definition 6 (Partial Differential Equation)**.** *A partial differential equation (PDE) is a differential equation whose unknown function depends on more than one independent variable.* ***Will not be taught in this course.***

Example (Motion under gravity and linear drag)**.** *A ball of mass m falls subject to gravity and linear air resistance. Let $v(t)$ be its velocity (positive downward). The net force gives*

$$m\frac{dv}{dt} = mg - \lambda v,$$

where $\lambda \geq 0$ is the drag coefficient. Writing $v = y'$, this becomes the second-order ODE

$$my'' + \lambda y' = mg.$$

Example (The Malthusian Growth Model)**.** *Let $N(t)$ be a population size. Assuming the rate of change is proportional to the current population,*

$$\frac{dN}{dt} = rN,$$

where r is the growth rate ($r > 0$) or decay rate ($r < 0$). Separating variables and applying the initial condition $N(0) = N_0$ gives the unique solution

$$N(t) = N_0 e^{rt}.$$

Example (The Logistic Equation)**.** *If the growth rate depends on the population itself, with carrying capacity K, the simplest choice $R(N) = r(1 - N/K)$ gives*

$$\frac{dN}{dt} = rN\left(1 - \frac{N}{K}\right),$$

called the logistic equation. It models growth when $N < K$ and decay when $N > K$.

Example (Heat Equation)**.** *The Heat Equation is an example of a PDE. Let $u = u(x,t)$, $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$. This PDE denotes the temperature of a body at time t and at position x.*

January 08, 2026.

3.2 Classification

3.2.1 The Order

Definition 7. *The order of an ODE is the order of the highest derivative that appears in the equation.*

Example. *$N' = rN$ is first order. $y'' + 2y' = e^t$ is second order.*

3.2.2 Scalar vs. System

Definition 8 (Scalar ODE)**.** *A scalar ODE has one unknown function.*

Example (Scalar ODE)**.**

$$y'' + y = 0$$

Definition 9 (System of ODEs)**.** *A system of ODEs has several unknown functions.*

Example (System of ODEs)**.**

$$\begin{cases} y_1' = y_2 \\ y_2' = -y_1 \end{cases}$$

Definition 10 (System of first order ODEs)**.** *A first order system has the form*

$$y'(t) = f(y(t), t),$$

where $y(t) \in \mathbb{R}^n$ *and* $f : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R}^n$*. In components,*

$$y_i'(t) = f_i(y_1, \ldots, y_n, t), \quad i = 1, \ldots, n.$$

Any n*th-order ODE*

$$y^{(n)} = G(t, y, y', \ldots, y^{(n-1)})$$

can be written as a first order system by setting

$$y_1 = y,\ y_2 = y',\ \ldots,\ y_n = y^{(n-1)},$$

which gives

$$\begin{cases} y_1' = y_2, \\ y_2' = y_3, \\ \vdots \\ y_{n-1}' = y_n, \\ y_n' = G(t, y_1, \ldots, y_n). \end{cases}$$

Example (System of First Order ODEs)**.** *Consider the second-order ODE*

$$y'' + 2y' + y = e^t.$$

Define new variables

$$y_1 = y, \qquad y_2 = y'.$$

Then the system becomes

$$\begin{pmatrix} y_1' \\ y_2' \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & -2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} + \begin{pmatrix} 0 \\ e^t \end{pmatrix}.$$

Let

$$A = \begin{pmatrix} 0 & 1 \\ -1 & -2 \end{pmatrix}, \quad r(t) = \begin{pmatrix} 0 \\ e^t \end{pmatrix}.$$

Problem 1 (System of First Order ODEs)**.** *Rewrite this third order ODE as a first order system:*

$$y''' + 4y' - y = 0$$

3.2.3 Linearity

Definition 11 (Linearity)**.** *An ODE is linear if the unknown function and its derivatives appear only to the first power, are not multiplied together, and are not composed with nonlinear functions. They may be multiplied by functions of the independent variable.*

Example (Linearity)**.** *Consider the following ODEs:*

- *Linear:* $y' + 3y = 0$
- *Linear:* $y'' - 2xy' + y = cos(x)$
- *Non-linear:* $y' + y^2 = 0$
- *Non-linear:* $y'' + sin(y) = 0$

Example (Lorenz system)**.**

$$\begin{aligned} y_1' &= \sigma(y_2 - y_1), \\ y_2' &= \rho y_1 - y_2 - y_1 y_3, \\ y_3' &= y_1 y_2 - \beta y_3, \end{aligned}$$

where σ, ρ, β are parameters. This is a nonlinear first order system in $\mathbb{R}^3$.

3.2.4 Autonomy

Definition 12 (Autonomy)**.** *The nth order ODE $F(t, y, y', \ldots, y^{(n)}) = 0$ is autonomous if F does not depend explicitly on t, that is, it can be written as $F(y, y', \ldots, y^{(n)}) = 0$. Otherwise, it is non-autonomous.*

Example (Autonomy)**.** *Consider the following ODEs:*

- *Non-autonomous:* $y'' + 2y' + y - e^t = 0$
- *Autonomous:* $N'(t) = rN(t)$
- *Non-autonomous:* $y'(t) = ty(t)$

Example. *A classic, simple example of an autonomous first-order system is the linear growth/decay model:*

$$\frac{dy}{dt} = ky$$

Here, $f(y, t) = ky$, which depends only on y (where k is a constant), making it autonomous.

Remark 5. *The Lorenz system (3.2.3) is an example of an autonomous system.*

3.3 Solutions and Initial Value Problems

Definition 13 (Solution)**.** *A solution of $y'(t) = f(y(t), t)$ on an interval J is a differentiable function $y : J \to D$ such that*

$$y'(t) = f(y(t), t) \quad \forall\, t \in J.$$

Example (Explicit Solution)**.** *Consider the following ODE $y' + y = 1$. We can verify that $y(t) = e^{-t} + 1$, and therefore $y'(t) = -e^{-t}$, is a solution on $\mathbb{R}$. Indeed,*

$$y' + y = -e^{-t} + (e^{-t} + 1) = 1.$$

Example (Implicit Solution)**.** *Consider the ODE $y\frac{dy}{dx} = x$. This is a nonautonomous, nonlinear, first-order scalar ODE. Separating $y\,dy = x\,dx$ and integrating gives the implicit solution $x^2 - y^2 = C, \quad C \in \mathbb{R}$.*

January 13, 2026

Definition 14 (Initial Value Problem (IVP))**.** *An initial value problem consists of an ODE together with a condition $y(t_0) = y_0$, where the value of the solution is prescribed at an initial time t_0.*

3.4 Existence and Uniqueness Theorem

3.4.1 Lipschitz continuity

Definition 15 (Lipschitz Continuity)**.** *A function $f : D \to \mathbb{R}^n$ is Lipschitz continuous on $D \subseteq \mathbb{R}^n$ if there exists $L \geq 0$ such that*

$$z||f(y_1) - f(y_2)|| \leq L||y_1 - y_2|| \quad \forall y_1, y_2 \in D, z \in \mathbb{R}.$$

The smallest such L is called the Lipschitz constant, denoted $\mathrm{Lip}(f)$.

Remark 6 (How to find a Lipschitz constant)**.** *If f is differentiable on $D \subset \mathbb{R}$ and $|f'(y)| \leq M$ for all $y \in D$, then f is Lipschitz with* $\mathrm{Lip}(f) = M$.

Example. *Let $f(y) = 4y - 5$, $D = \mathbb{R}$, and $|| \cdot || = | \cdot |$. Then*

$$|f(y_1) - f(y_2)| = |4y_1 - 4y_2| = 4|y_1 - y_2|.$$

So f is Lipschitz on $\mathbb{R}$ with $Lip(f) = 4$.

Example. *Let*

$$f(y) = \frac{1}{y-1}, \quad f'(y) = -\frac{1}{(y-1)^2}, \quad D = (1, +\infty).$$

This function is not Lipschitz continuous on D since as $y \to 1^+$, the derivative $f'(y)$ approaches $-\infty$, and so the ratio $\frac{|f(y_2)-f(y_1)|}{|y_2-y_1|}$ approaches ∞ for any $y_1, y_2 \in D$. Therefore, there is no Lipschitz constant for f on D.
To make this function Lipschitz continuous, we fix $\delta > 1$ and define $D_\delta =$

$(\delta, +\infty)$. *For* $y_1, y_2 \in D_\delta$, *by the Mean Value Theorem, there exists* $z \in (y_1, y_2)$ *such that*

$$f(y_2) - f(y_1) = f'(z)(y_2 - y_1).$$

So

$$|f(y_2) - f(y_1)| \leq \frac{1}{(z-1)^2}|y_2 - y_1| \leq \frac{1}{(\delta-1)^2}|y_2 - y_1|.$$

Thus f *is Lipschitz on* D_δ *with*

$$Lip(f) = \frac{1}{(\delta-1)^2}.$$

Example. *Suppose* f *is differentiable and satisfies*

$$-4 \leq f'(y) \leq 9 \quad \textit{for all } y \in \mathbb{R}.$$

Then

$$|f'(y)| \leq 9 \quad \forall y,$$

so f *is Lipschitz on* $\mathbb{R}$ *with Lipschitz constant* $L = 9$.

3.4.2 Local Lipschitz continuity

Definition 16 (Locally Lipschitz)**.** *Let* $D \subseteq \mathbb{R}^n$ *be open. A function* $f : D \to \mathbb{R}^n$ *is* locally Lipschitz *if for every compact set* $K \subset D$ *there exists* $L > 0$ *such that*

$$\|f(y_1) - f(y_2)\| \leq L\|y_1 - y_2\| \quad \forall\, y_1, y_2 \in K.$$

Remark 7. *Lipschitz* $\Rightarrow$ *locally Lipschitz, but not conversely. Any* C^1 *function is locally Lipschitz (by the Mean Value Inequality from Section* (2.1.4)*).*

Example (Locally Lipschitz but not Lipschitz)**.**

$$f(y) = y^2 \in \mathbb{R}.$$

Since $f'(y) = 2y$ *is unbounded on* $\mathbb{R}$, *no single constant works on the whole domain, so* f *is not Lipschitz. However, on any bounded set* $K = [-M, M]$, $|f'(y)| \leq 2M$, *so* f *is locally Lipschitz.*

Example (Lipschitz (hence locally Lipschitz))**.**

$$f(y) = \sin y.$$

Since $|f'(y)| = |\cos y| \leq 1$, $\forall y$, f *is Lipschitz on* $\mathbb{R}$, *with* $Lip(f) = 1$.

Example (Continuous but not locally Lipschitz)**.**

$$f(y) = \sqrt{|y|}.$$

The derivative is unbounded near $y = 0$, *so no Lipschitz constant exists even locally. Hence* f *is not locally Lipschitz.*

January 13, 2026

3.4.3 Integral Form of the IVP

Lemma 4. *A function y solves the IVP if and only if*

$$y(t) = y_0 + \int_{t_0}^{t} f(y(s), s)ds.$$

Proof. If $y' = f(y,t)$ and $y(t_0) = y_0$, then by the Fundamental Theorem of Calculus,

$$y(t) = y_0 + \int_{t_0}^{t} f(y(s), s)ds.$$

Conversely, differentiating the right-hand side gives

$$y'(t) = f(y(t), t), \quad y(t_0) = y_0.$$

□

3.4.4 Picard operator

The integral form motivates defining the following operator. Since $(y_0, t_0) \in D \times (a,b)$ and the set is open, choose $\alpha, \delta > 0$ such that

$$D_{\alpha,\delta} = \{(y,t) : \|y - y_0\| \leq \alpha,\ |t - t_0| \leq \delta\} \subset D \times (a,b),$$

and set

$$M_{\alpha,\delta} = \sup_{(y,t)\in D_{\alpha,\delta}} \|f(y,t)\| < +\infty, \qquad \varepsilon = \min\left(\delta, \frac{\alpha}{M_{\alpha,\delta}}\right), \qquad J = (t_0 - \varepsilon, t_0 + \varepsilon).$$

Definition 17 (Picard Operator)**.** *For any $y : J \to \mathbb{R}^n$ with $y(t_0) = y_0$ and $(y(t), t) \in D_{\alpha,\delta}$ for all $t \in J$, define*

$$T(y)(t) = y_0 + \int_{t_0}^{t} f(y(s), s)\, ds.$$

A function y solves the IVP if and only if $T(y) = y$, i.e. y is a fixed point *of T.*

Lemma 5 (Invariance)**.** *If $y(t_0) = y_0$ and $(y(t), t) \in D_{\alpha,\delta}$ $\forall t \in J$, then $T(y)(t_0) = y_0$ and $(T(y)(t), t) \in D_{\alpha,\delta}$ for all $t \in J$.*[1]

Proof. Clearly $T(y)(t_0) = y_0$. For $t \in J$,

$$\|T(y)(t) - y_0\| \leq \int_{t_0}^{t} \|f(y(s), s)\| ds \leq M_{\alpha,\delta}|t - t_0| \leq M_{\alpha,\delta}\epsilon \leq \alpha.$$

Hence $(T(y)(t), t) \in D_{\alpha,\delta}$. □

[1] Basically, If a function stays inside the box, then applying T keeps it inside the box.

Example (Picard operator)**.** *Consider the IVP $y' = y$, $y(0) = 1$. It's integral form is $y(t) = 1 + \int_0^t y(s)ds$. Start with the simplest function that satisfies the initial condition: $y_0(t) = 1$. Now iterate:*

$$y_1 = T(y_0) = 1 + \int_0^t 1ds = 1 + t,$$

$$y_2 = T(y_1) = 1 + \int_0^t (1+s)ds = 1 + t + \frac{t^2}{2},$$

$$y_3 = T(y_2) = 1 + t + \frac{t^2}{2} + \frac{t^3}{6}.$$

Each step adds to the next power of t. We can express this relation as

$$y(t) = \sum_{k=0}^{\infty} \frac{t^k}{k!} = e^t.$$

Indeed, $T(y)(t) = 1 + \int_0^t e^s ds = e^t$, so $T(y) = y$. Therefore, T is an operator that maps a function to a new one, and only the true solution is a fixed point of T.

3.4.5 Statement of the Theorem

Theorem 6 (Existence and Uniqueness)**.** *Let $D \subseteq \mathbb{R}^n$ be open and (a, b) an open interval containing t_0. Suppose $f : D \times (a, b) \to \mathbb{R}^n$ [2] is continuous and locally Lipschitz in y. If $y_0 \in D$, then the IVP*

$$\begin{cases} y' = f(y, t), \\ y(t_0) = y_0 \end{cases}$$

has a unique solution on some open interval J containing t_0.

January 20, 2026

Remark 8 (Choosing ε)**.** *To apply the Existence and Uniqueness Theorem, we choose ε so that the solution remains inside the domain $D \times (a, b)$. Consider the IVP*

$$\begin{cases} y' = f(y, t), \\ y(t_0) = y_0. \end{cases}$$

The theorem guarantees that there exists a unique solution

$$y : (t_0 - \varepsilon, t_0 + \varepsilon) \to \mathbb{R}^n.$$

Choose $\alpha > 0$ and $\delta > 0$ such that

$$D_{\alpha,\delta} := \{(y, t) : \|y - y_0\| \le \alpha,\ |t - t_0| \le \delta\} \subset D \times (a, b).$$

[2] $D \times (a, b)$ denotes the set of all pairs (y, t) with $y \in D \subset \mathbb{R}^n$ and $t \in (a, b)$, i.e., all allowed state-time inputs of f.

Since f is continuous, define

$$M_{\alpha,\delta} = \sup_{(y,t)\in D_{\alpha,\delta}} \|f(y,t)\| < \infty.$$

Then one may take

$$\varepsilon = \min\left(\delta, \frac{\alpha}{M_{\alpha,\delta}}\right).$$

Problem 2 (Existence and Uniqueness Theorem)**.** *Consider the initial value problem*

$$\begin{cases} y' = t^2 + y^2, \\ y(0) = 0, \end{cases}$$

Determine the largest value of ϵ for which the Theorem of Existence and Uniqueness ensures the existence of a solution $y : (-\epsilon, \epsilon) \to \mathbb{R}$.

Problem 3 (LLC)**.** *Consider the initial value problem*

$$\begin{cases} y' = 3y^{2/3}, \\ y(0) = 0. \end{cases}$$

Investigate existence and uniqueness.

January 15, 2026

3.4.6 Picard Iteration and Existence

Define $y_0(t) \equiv y_0$ (constant). For $k \geq 1$ set

$$y_k(t) = T(y_{k-1})(t) = y_0 + \int_{t_0}^{t} f(y_{k-1}(s), s)\, ds.$$

By the invariance lemma, each y_k stays inside $D_{\alpha,\delta}$.

The sequence $\{y_k\}$ is Cauchy in $\mathbb{R}^n$ (see Appendix 8.1) and therefore converges uniformly to a function $y : J \to \mathbb{R}^n$ which satisfies the integral equation, hence solves the IVP.

3.4.7 Uniqueness

Proof of Uniqueness. Assume $y(t)$ and $z(t)$ are two solutions of the IVP with the same initial condition. By the Lipschitz property of f,

$$\|y(t) - z(t)\| = \| \int_{t_0}^{t} f(y(s), s) - f(z(s), s) ds\| \leq \int_{t_0}^{t} L\|y(s) - z(s)\| ds,$$

for some Lipschitz constant $L > 0$. Define

$$g(t) = \int_{t_0}^{t} \|y(s) - z(s)\| ds,$$

so $g'(t) \leq Lg(t)$ and $g(t_0) = 0$. Grönwall's idea[3] tells us this forces $g(t) = 0$ for all t, so $\|y(t) - z(t)\| = 0$. Hence, $y(t) = z(t)$, meaning two solutions with the same initial condition cannot diverge, this proves the solution is unique. $\square$

4 First-Order Scalar Equation

January 22, 2026

A first-order scalar ODE involves a single unknown function $y(t)$ and its first derivative:

$$y' = f(y, t).$$

We study three main classes: linear, separable, and exact equations.

4.1 First Order Linear Equations

Definition 18 (First Order Linear Equation)**.** *A first-order linear equation has the form*

$$a_0(t)y' + a_1(t)y = g(t), \quad a_0(t) \neq 0.$$

Dividing by $a_0(t)$ gives the standard form

$$y' + P(t)y = Q(t).$$

4.1.1 Integrating Factor Method

To solve $y' + P(t)y = Q(t)$, multiply both sides by a function $\mu(t)$ chosen so that the left-hand side becomes an exact derivative:

$$\mu y' + \mu P(t)y = \frac{d}{dt}(\mu y).$$

This requires $\mu'(t) = \mu(t)P(t)$, which gives

$$\mu(t) = e^{\int P(t)\,dt}.$$

The equation then becomes

$$\frac{d}{dt}(\mu y) = \mu Q(t),$$

and integrating both sides yields

$$y(t) = \frac{1}{\mu(t)}\left(\int \mu(t)Q(t)\,dt + C\right), \quad C \in \mathbb{R}.$$

Example. *Solve $y' - 2y = 3e^t$ with $y(0) = 1$.*
Here $P(t) = -2$, $Q(t) = 3e^t$, so $\mu(t) = e^{-2t}$. Multiplying through:

$$\frac{d}{dt}(e^{-2t}y) = 3e^{-t}.$$

[3] Multiplying by $e^{-L(t-t_0)}$ gives $\frac{d}{dt}(e^{-L(t-t_0)}g(t)) \leq 0$, which shows $e^{-L(t-t_0)}g(t)$ is decreasing and thus $g(t) = 0$

Integrating:

$$e^{-2t}y = -3e^{-t} + C \implies y(t) = -3e^{t} + Ce^{2t}.$$

Applying $y(0) = 1$ *gives* $C = 4$*, so*

$$y(t) = -3e^{t} + 4e^{2t}.$$

Remark 9 (Quick Solution Method)**.** *A quick method to solve a first-order linear equation (not presented by the professor) is*

$$y(t) = \frac{\int \mu(t)Q(t)}{\mu(t)}.$$

4.1.2 Existence and Uniqueness for First Order Linear Equations

Theorem 7 (Existence & Uniqueness)**.** *If* P *and* Q *are continuous on an open interval containing* t_0*, then the IVP*

$$y' + P(t)y = Q(t), \quad y(t_0) = y_0$$

has a unique solution on that entire interval.

Remark 10. *Unlike the general E&U theorem from Chapter (3.4.5), this result gives a global solution on the full interval where* P *and* Q *are continuous, not just locally.*

Example. *For the IVP*

$$\begin{cases} y' + \dfrac{1}{t-1}y = \dfrac{1}{\cos t}, \\ y(0) = 1, \end{cases}$$

$P(t)$ *is discontinuous at* $t = 1$ *and* $Q(t)$ *is discontinuous at* $t = \pm\frac{\pi}{2}$*. The largest interval containing* $t_0 = 0$ *on which both are continuous is*

$$J_{\max} = \left(-\frac{\pi}{2}, 1\right).$$

4.2 Separable Equations

January 27, 2026

Definition 19. *The ODE* $\frac{dy}{dx} = f(x, y)$ *is* separable *if* $f(x, y) = f_1(x)f_2(y)$ *for some functions* f_1 *and* f_2*.*

Example. $\dfrac{dy}{dx} = -\dfrac{x}{y}$ *is separable with* $f_1(x) = x$*,* $f_2(y) = -\dfrac{1}{y}$*.*
$\dfrac{dy}{dx} = x^2 + y^2$ *is not separable.*

4.2.1 General Method

Writing the separable ODE as

$$M(x) + N(y)\frac{dy}{dx} = 0, \tag{1}$$

where $M(x) = -f_1(x)$ and $N(y) = \frac{1}{f_2(y)}$, and letting H_1, H_2 be antiderivatives of M, N respectively, the chain rule gives

$$\frac{d}{dx}(H_1(x) + H_2(y(x))) = 0,$$

so the implicit general solution is

$$H_1(x) + H_2(y) = C, \quad C \in \mathbb{R},$$

or equivalently in integral form,

$$\int \frac{dy}{f_2(y)} = \int f_1(x)\,dx + C.$$

Example (Logistic Equation)**.** *The logistic equation*

$$\frac{dN}{dt} = rN\left(1 - \frac{N}{K}\right)$$

is separable. Using partial fractions,

$$\int \frac{dN}{N} + \int \frac{dN}{N-K} = \int r\,dt \implies \ln\left|\frac{N}{N-K}\right| = rt + C.$$

Solving with $N(0) = N_0$ gives

$$N(t) = \frac{KN_0}{N_0 + (K - N_0)e^{-rt}}.$$

As $t \to \infty$, $N(t) \to K$ (the carrying capacity), provided $N_0 \neq 0$.

4.3 Exact Equations

January 29, 2026

4.3.1 Potential Functions and Integral Curves

Definition 20 (Potential Function)**.** *A function $\Psi(x, y)$ is a potential function for the equation (1) if*

$$\frac{\partial \Psi}{\partial x} = M \quad \text{and} \quad \frac{\partial \Psi}{\partial y} = N.$$

The general solution is then given implicitly by the level curves

$$\Psi(x, y) = C.$$

Definition 21 (Integral Curve)**.** *An integral curve is the level curve of Ψ passing through a specific initial condition (x_0, y_0):*

$$\Gamma_{(x_0,y_0)} = \{(x, y) \mid \Psi(x, y) = \Psi(x_0, y_0)\}.$$

Remark 11. *For a separable equation, the general solution $\Psi(x, y) = C$ describes all possible solution curves, and picking a particular C gives the integral curve passing through a given point.*

Example. *For $\frac{dy}{dx} = -\frac{x}{y}$, $y \neq 0$, separating gives $y\,dy + x\,dx = 0$, so the potential function is*

$$\Psi(x, y) = \frac{x^2}{2} + \frac{y^2}{2}.$$

The general solution is the family of circles $x^2+y^2 = C$. For $y(0) = 2$, $\Psi(0, 2) = 2$, so the unique integral curve is

$$y(x) = \sqrt{4 - x^2}, \quad J_{\max} = (-2, 2).$$

4.3.2 Definition and Exactness Criterion

Definition 22 (Exact Equation)**.** *The equation (1) is exact on a domain D if there exists a potential function Ψ such that $\Psi_x = M$ and $\Psi_y = N$. Along any solution $y(x)$,*

$$\frac{d}{dx}\Psi(x, y(x)) = 0,$$

so Ψ is constant along solutions.

Theorem 8. *Let $D \subset \mathbb{R}^2$ be open and simply connected[4]. Suppose M, N, M_y, N_x are continuous on D. Then (1) is exact on D if and only if*

$$M_y(x, y) = N_x(x, y) \quad \forall\,(x, y) \in D.$$

Proof. ($\Rightarrow$) Suppose it is exact, then $M = \Psi_x$ and $N = \Psi_y$, so $M_y = \Psi_{xy} = \Psi_{yx} = N_x$ by equality of mixed partials.

($\Leftarrow$) If $M_y = N_x$ on a simply connected domain and partials are continuous, one can construct Ψ with $\Psi_x = M$ and $\Psi_y = N$ explicitly by integration.[5] $\square$

Example. *Solve*

$$y\cos x + 2xe^y + (\sin x + x^2e^y - 1)\,\frac{dy}{dx} = 0.$$

Identify $M = y\cos x + 2xe^y$ and $N = \sin x + x^2e^y - 1$. Check:

$$M_y = \cos x + 2xe^y = N_x. \quad \checkmark$$

[4] Meaning there are no holes in the domain.

[5] The construction is shown for rectangles in the notes; the general case follows similarly.

So the equation is exact. Integrate $\Psi_x = M$ in x:

$$\Psi = y \sin x + x^2 e^y + c(y).$$

Differentiate in y and match with N:

$$\Psi_y = \sin x + x^2 e^y + c'(y) = \sin x + x^2 e^y - 1 \implies c'(y) = -1 \implies c(y) = -y.$$

The general solution is

$$\Psi(x, y) = y \sin x + x^2 e^y - y = C.$$

Remark 12. *Every separable equation is exact on a simply connected domain. Writing $M(x)\,dx + N(y)\,dy = 0$ gives $M_y = 0 = N_x$ automatically, so exactness always holds. The potential function is simply $\Psi(x, y) = H_1(x) + H_2(y)$.*

February 3, 2026

4.3.3 Integrating Factors for Non-Exact Equations

If $M_y \neq N_x$, we seek a function $\mu(x, y)$ such that $\mu M\,dx + \mu N\,dy = 0$ is exact. The condition $(\mu M)_y = (\mu N)_x$ expands to

$$\mu_y M + \mu M_y = \mu_x N + \mu N_x.$$

Case: $\mu = \mu(x)$ only. Then $\mu_y = 0$ and the condition reduces to

$$\frac{d\mu}{dx} = \mu\,\frac{M_y - N_x}{N}.$$

If $\frac{M_y - N_x}{N}$ depends only on x, this is a separable ODE for $\mu(x)$ that can be solved directly.

Example. *Solve*

$$(3xy + y^2)\,dx + (x^2 + xy)\,dy = 0.$$

Here $M_y = 3x + 2y$ and $N_x = 2x + y$, so $M_y \neq N_x$. Compute:

$$\frac{M_y - N_x}{N} = \frac{x + y}{x(x + y)} = \frac{1}{x},$$

which depends only on x. Solving $\frac{d\mu}{dx} = \frac{\mu}{x}$ gives $\mu(x) = x$. Multiplying through by x:

$$(3x^2 y + xy^2)\,dx + (x^3 + x^2 y)\,dy = 0.$$

Now $M_y = 3x^2 + 2xy = N_x$. ✓ Integrating:

$$\Psi = \int (3x^2 y + xy^2)\,dx = x^3 y + \frac{x^2 y^2}{2} + c(y).$$

Matching $\Psi_y = N$ gives $c'(y) = 0$, so the general solution is

$$x^3 y + \frac{x^2 y^2}{2} = C.$$

4.4 Applications

Example (Falling Object with Air Resistance)**.** *An object of mass m falls under gravity with damping $\gamma > 0$. Setting $v(0) = 0$, the equation*

$$v' + \frac{\gamma}{m} v = g$$

has integrating factor $\mu = e^{\frac{\gamma}{m}t}$, giving

$$v(t) = \frac{gm}{\gamma}\left(1 - e^{-\frac{\gamma}{m}t}\right).$$

As $t \to \infty$, the velocity approaches the terminal value $v_\infty = \frac{gm}{\gamma}$.

Example (Mixing Problem)**.** *A tank holds 100 L of brine with y_0 g of salt. Pure salt solution enters at 50 g/L at rate R L/s, and the well-mixed solution exits at the same rate. The salt amount $y(t)$ satisfies*

$$y' + \frac{R}{100} y = 50R, \quad y(0) = y_0.$$

Using integrating factor $\mu = e^{\frac{R}{100}t}$,

$$y(t) = 5000 + (y_0 - 5000)e^{-\frac{R}{100}t}.$$

Regardless of y_0, the salt amount converges to 5000 g as $t \to \infty$.

5 Systems of Linear Equations

February 3, 2026

Many problems involve several unknown functions that evolve together and influence one another. Instead of a single differential equation, we therefore study systems of first-order linear ODEs. Writing the unknown functions as a vector allows us to treat the entire system using matrix algebra and apply the tools of linear algebra (matrices, eigenvalues, and fundamental matrices).

We consider systems of the form

$$y'(t) = A(t)y(t) + r(t), \quad t \in Q, \tag{2}$$

where $y(t)$ is the unknown vector, $A(t)$ is a matrix of coefficients, $r(t)$ is a forcing term, and $Q \subseteq \mathbb{R}$. Explicitly,

$$y(t) = \begin{pmatrix} y_1(t) \\ \vdots \\ y_n(t) \end{pmatrix} \in \mathbb{R}^n, \quad A(t) \in M_n(\mathbb{R}), \quad r(t) = \begin{pmatrix} r_1(t) \\ \vdots \\ r_n(t) \end{pmatrix},$$

5.1 From higher order scalar ODEs to linear systems

Example. *Convert the third order equation*

$$y''' - 2y'' + 4y' - 5y = \sin t + t^3$$

into a first order system. Let

$$y_1 = y, \quad y_2 = y', \quad y_3 = y''.$$

Then

$$\begin{pmatrix} y_1' \\ y_2' \\ y_3' \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 5 & -4 & 2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \sin t + t^3 \end{pmatrix}.$$

Thus the equation is written in the form $y' = Ay + r$, with $J_{\max} = \mathbb{R}$.

5.2 Theory of First Order Linear Systems

Theorem 9 (Existence and uniqueness)**.** *If every entry of $A(t)$ and $r(t)$ is continuous on an interval Q, then for any*

$$t_0 \in Q, \quad y_0 \in \mathbb{R}^n,$$

the IVP

$$\begin{cases} y'(t) = A(t)y(t) + r(t), \\ y(t_0) = y_0 \end{cases}$$

has a unique solution on all of Q. Hence $J_{\max} = Q$.

Definition 23. *If $r(t) = 0$, the system is homogeneous. Otherwise it is non-homogeneous.*

For a homogeneous system

$$y'(t) = A(t)y(t), \tag{3}$$

the set of all solutions forms a vector space: it contains the zero function, and any linear combination of solutions is also a solution (26). To describe this space concretely, we select n linearly independent solutions; arranging them as the columns of a matrix gives a fundamental matrix, which allows every solution to be expressed as a linear combination of these columns.

The non-homogeneous system $y' = A(t)y + r(t)$ can be solved using a fundamental matrix $\Phi(t)$ of the homogeneous system:

$$y(t) = \Phi(t)c + \Phi(t) \int_{t_0}^{t} \Phi(s)^{-1} r(s) ds,$$

where $c \in \mathbb{R}^n$ is determined by the initial condition $y(t_0) = y_0$.

5.3 Structure of the Solution Space

Definition 24 (Linear Independence)**.** *Functions $y_1, \ldots, y_n$ in the solution set S*[6] *are linearly independent if*

$$c_1 y_1(t) + \cdots + c_n y_n(t) = 0 \quad \forall t \in Q \implies c_1 = \cdots = c_n = 0.$$

Theorem 10 (Solution Set)**.** *The solution set S is an n-dimensional vector subspace of $C^1(Q)$.*

Proof. To prove that S is a vector space of dimension n, we show the following:
Step 1: Vector space. If $y_1, y_2 \in S$ and $\alpha_1, \alpha_2 \in \mathbb{R}$,

$$(\alpha_1 y_1 + \alpha_2 y_2)' = \alpha_1 y_1' + \alpha_2 y_2' = A(t)(\alpha_1 y_1 + \alpha_2 y_2),$$

so any linear combination is in S, and the zero function is also in S.

Step 2: Construct a basis. Let $e_1, \ldots, e_n$ be the standard basis of $\mathbb{R}^n$, and let y_i be the unique solution of

$$\begin{cases} y' = A(t)y, \\ y(t_0) = e_i \end{cases}$$

for $i = 1, \ldots, n$. Define $B = \{y_1, \ldots, y_n\}$. By uniqueness, any solution $z \in S$ can be written as a linear combination of the y_i, so B spans S. Evaluating a linear combination at t_0 shows the y_i are independent. Hence, B is a basis and $\dim S = n$. □

February 5, 2026

Definition 25 (Matrix Solution)**.** *A matrix solution $y(t)$ of the equation*

$$y' = A(t)y \tag{4}$$

is an $n \times n$ matrix function whose columns are solutions to (4).

Definition 26 (Principle of Superposition)**.** *The principle of superposition states that if $y_1(t), \ldots, y_n(t)$ are solutions of $y' = A(t)y$, then any linear combination of these solutions is also a solution.*

By the POS (principle of superposition), for any fixed vector

$$C = \begin{pmatrix} C_1 \\ C_2 \\ \vdots \\ C_n \end{pmatrix} \in \mathbb{R}^n, \quad y(t)c = c_1 y_1(t) + c_2 y_2(t) + \cdots + c_n y_n(t) \in S.$$

Proposition 1. *Let $y(t)$ be a matrix solution of (4). Either*

$$\det(y(t)) \neq 0 \quad \forall t \in Q$$

or

$$\det(y(t)) = 0.$$

[6] $y_1, \ldots, y_n$ are solutions of $y' = A(t)y$

Proof. Assume there exists $t_0 \in Q$ such that $\det(y(t_0)) = 0$. Then $y(t_0)$ is not invertible. Hence, by linear algebra, there exists a nonzero vector $v \in \mathbb{R}^n$ such that

$$y(t_0)v = 0.$$

Define

$$y_1(t) := y(t)v.$$

Since each column of $y(t)$ is a solution of $y' = A(t)y$, any linear combination of the columns is also a solution. Therefore $y_1(t)$ is a solution, i.e. $y_1 \in S$. Also define the zero solution

$$y_2(t) := 0,$$ [7]

which satisfies $y_2' = A(t)y_2$, so $y_2 \in S$. Now,

$$y_1(t_0) = y(t_0)v = 0 = y_2(t_0).$$

Thus y_1 and y_2 are two solutions with the same initial condition. By the existence and uniqueness theorem,

$$y_1(t) = y_2(t) \quad \text{for all } t \in Q.$$

Hence

$$y(t)v = 0 \quad \text{for all } t.$$

Therefore the columns of $y(t)$ are linearly dependent for every t, which implies

$$\det(y(t)) = 0 \quad \text{for all } t \in Q.$$

□

Definition 27 (Wronskian)**.**

$$w(t) = w(y_1, \ldots, y_n)(t) = \det(y(t))$$

is called the Wronskian of $y_1, \ldots, y_n$.

Example (Wronskian)**.** *Let*

$$y_1(t) = \begin{pmatrix} cos(2t) \\ -sin(2t) \end{pmatrix}, \quad y_2(t) = \begin{pmatrix} -2sin(2t) \\ -2cos(2t) \end{pmatrix},$$

Compute the Wronskian to determine whether the functions $y_1(t)$ *and* $y_2(t)$ *are linearly independent.*

$$W(t) = \det \begin{pmatrix} cos(2t) & -2sin(2t) \\ -sin(2t) & -2cos(2t) \end{pmatrix} = -2.$$

Therefore $y_1(t)$ *and* $y_2(t)$ *are linearly independent since* $W(t) \neq 0$.

Definition 28 (Fundamental Matrix Solution)**.** *A matrix solution with nonvanishing*[8] *Wronskian is called a fundamental matrix solution.*

Proposition 2 (Recovering the coefficient matrix)**.** *Let $\Phi(t)$ be a fundamental matrix solution of the homogeneous system (4) on an interval Q. Then $\Phi(t)$ is invertible for all $t \in Q$, and*

$$A(t) = \Phi'(t)\,\Phi(t)^{-1}.$$

In particular, if $A(t) = A$ is constant, then the right-hand side is independent of t.

Proof. Since $\Phi(t)$ is a matrix solution, its columns satisfy (4), hence

$$\Phi'(t) = A(t)\Phi(t).$$

Because $\Phi(t)$ is a fundamental matrix, $\det \Phi(t) \neq 0$ for all $t \in Q$, so $\Phi(t)$ is invertible. Multiplying both sides on the right by $\Phi(t)^{-1}$ gives

$$A(t) = \Phi'(t)\Phi(t)^{-1}.$$

□

Example. *Suppose the general solution of a linear system is given by*

$$y(t) = \begin{pmatrix} e^{2t} & 0 \\ 0 & e^{-t} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}.$$

Then a fundamental matrix is

$$\Phi(t) = \begin{pmatrix} e^{2t} & 0 \\ 0 & e^{-t} \end{pmatrix}.$$

Differentiating,

$$\Phi'(t) = \begin{pmatrix} 2e^{2t} & 0 \\ 0 & -e^{-t} \end{pmatrix},$$

and since

$$\Phi(t)^{-1} = \begin{pmatrix} e^{-2t} & 0 \\ 0 & e^{t} \end{pmatrix},$$

we obtain

$$A = \Phi'(t)\Phi(t)^{-1} = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}.$$

Theorem 11 (Liouville's formula)**.** *Let $y(t)$ be a matrix solution to (4) satisfying the initial condition $y(t_0) = y_0 \in M_n(\mathbb{R})$. Then*

$$\det(y(t)) = \det(y_0) e^{\int_{t_0}^{t} \operatorname{tr}(A(s))\, ds}.$$

Here,

$$\operatorname{tr}(M) = \textit{sum of the diagonal entries of } M \in M_n(\mathbb{R}).$$

[8] Nonvanishing in this context means not equal to zero.

Remark 13. *For the homogeneous system (4), the goal is to find n linearly independent solutions*

$$y_1, y_2, \dots, y_n$$

such that $W(t_0) \neq 0$.

Since $y_1, \dots, y_n$ are linearly independent, they form a basis for the solution space. Therefore every solution has the form

$$y(t) = c_1 y_1(t) + c_2 y_2(t) + \cdots + c_n y_n(t), \qquad c_1, \dots, c_n \in \mathbb{R}.$$

Given an initial condition

$$y(t_0) = y_0 \in \mathbb{R}^n,$$

$$y(t_0) = c_1 y_1(t_0) + \cdots + c_n y_n(t_0) = \begin{pmatrix} | & | & | \\ y_1(t_0) & y_2(t_0) & y_n(t_0) \\ | & | & | \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix} = y_0 = \begin{pmatrix} y_0' \\ y_0'' \\ \vdots \\ y_0^{(n)} \end{pmatrix}.$$

The unique $(c_1, \dots, c_n)$ is given by

$$\begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} | & | \\ y_1(t_0) & y_n(t_0) \\ | & | \end{pmatrix}^{-1} \begin{pmatrix} y_0' \\ y_0'' \\ \vdots \\ y_0^{(n)} \end{pmatrix}.$$

5.4 Linear homogeneous systems with constant coefficients

Consider (4), where $A \in M_n(\mathbb{R})$ with constant entries. We look for solutions of the form

$$y(t) = e^{\lambda t} u \tag{5}$$

where u is a constant vector.[9] Differentiating (5) gives

$$y'(t) = \frac{d}{dt}(e^{\lambda t} u) = e^{\lambda t}(\lambda u),$$

and substituting (5) into the differential equation (4) gives

$$y'(t) = Ay = A(e^{\lambda t} u) = e^{\lambda t}(Au).$$

Hence if we combine both equations, we get

$$e^{\lambda t}(\lambda u) = e^{\lambda t}(Au) \implies Au = \lambda u. \tag{6}$$

In other words, λ is an eigenvalue of A and u is a corresponding eigenvector.

[9] Recall the analogous scalar case $a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_{n-1} y' + a_n y = 0$, where we look for solutions of the form e^{rt}.

Definition 29 (Kernel)**.** *Given a matrix $A \in M_n(\mathbb{C})$, define*

$$\ker A = \{u \mid Au = 0\},$$

called the kernel (or null space) of A.

Equation (6) is equivalent to

$$Au - \lambda u = 0 \iff (A - \lambda I)u = 0,$$

where I is the identity matrix in $M_n(\mathbb{R})$.

Definition 30 (Characteristic Polynomial)**.** *Let $A \in M_n(\mathbb{R})$. The characteristic polynomial of A, denoted $p(\lambda)$, is defined to be*

$$p(\lambda) = \det(A - \lambda I)$$

Definition 31 (Eigenvalue)**.** *An eigenvalue of A is a root of $p(\lambda)$.*

Definition 32 (Spectrum)**.** *The set*

$$\sigma(A) = \{\lambda \mid p(\lambda) = 0\}$$

is called the spectrum of A.

Example. *Consider*

$$y' = Ay, \qquad y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \qquad A = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}.$$

To solve the system, we first find the eigenvalues and eigenvectors of A. Define the characteristic polynomial to be

$$p(\lambda) = \det(A - \lambda I) = \begin{vmatrix} 1-\lambda & 3 \\ 3 & 1-\lambda \end{vmatrix} = (1-\lambda)^2 - 9.$$

Solving $(1-\lambda)^2 - 9 = 0$, gives us two eigenvalues $\lambda_1 = -2$ and $\lambda_2 = 4$. Hence the spectrum of A is

$$\sigma(A) = \{-2, 4\}.$$

To find the eigenvectors, we need to solve $(A - \lambda I)u = 0$.

For $\lambda_1 = -2$:

$$(A - \lambda I)u_1 = 0 \implies \begin{pmatrix} 3 & 3 \\ 3 & 3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \implies b = -a.$$

You can pick any values of a, and b is the negative of a; choose $a = 1$:

$$u_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

For $\lambda_2 = 4$, solving $(A - 4I)u_2 = 0$ similarly yields

$$u_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Each eigenpair (λ, u) produces a solution of the system of the form $y(t) = e^{\lambda t}u$. Hence

$$y_1(t) = c_1 e^{-2t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \in S, \qquad y_2(t) = c_2 e^{4t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \in S,$$

where c_1 and c_2 are arbitrary constants that will be determined when applying the initial condition to form the general solution.

To ensure that these solutions form a complete basis for the solution space, we check their linear independence using the Wronskian. The wronskian of y_1 and y_2 is

$$w(t) = \det \begin{pmatrix} e^{-2t} & e^{4t} \\ -e^{-2t} & e^{4t} \end{pmatrix}.$$

Evaluating at $t = 0$,

$$w(0) = \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} = 2 \neq 0,$$

so y_1 and y_2 are linearly independent. Therefore they form a basis of the solution space,

$$S = \langle \{y_1, y_2\} \rangle,$$

and the general solution is

$$y(t) = c_1 e^{-2t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + c_2 e^{4t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} e^{-2t} & e^{4t} \\ -e^{-2t} & e^{4t} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

If we set an initial condition such as $y(0) = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$, then we can solve for c_1 and c_2:

$$y(0) = \begin{pmatrix} 1 \\ 2 \end{pmatrix} = c_1 \begin{pmatrix} -1 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

$$\implies \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1/2 \\ 3/2 \end{pmatrix}.$$

Therefore, the general solution of the system with the initial condition is

$$y(t) = \frac{1}{2} e^{-2t} \begin{pmatrix} -1 \\ 1 \end{pmatrix} + \frac{3}{2} e^{4t} \begin{pmatrix} -1 \\ 1 \end{pmatrix},$$

which is the unique solution satisfying $y(0) = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

February 10, 2026

Example (3 by 3 system)**.** *Consider the system*

$$y' = Ay, \quad A - \lambda I = \begin{pmatrix} 3-\lambda & 1 & -2 \\ -1 & 5-\lambda & -2 \\ -3 & 3 & -2-\lambda \end{pmatrix}.$$

Step 1: Find eigenvalues.
The characteristic polynomial is

$$p(\lambda) = \det(A - \lambda I) = -(\lambda+2)(\lambda-4)^2 = 0,$$

giving eigenvalues $\lambda_1 = -2$ *and* $\lambda_2 = \lambda_3 = 4$.
Step 2: Find eigenvectors.
For $\lambda_1 = -2$*, solve*

$$(A + 2I)u_1 = 0.$$

A solution is $u_1 = \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}$*, yielding* $y_1(t) = e^{-2t}u_1$.
For $\lambda = 4$*, solve*

$$(A - 4I)u = 0.$$

If we do Gaussian elimination (row reduction), we find that $a = b - 2c$*, so* b *and* c *are free variables. Thus*

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} = b \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + c \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}.$$

Since every solution is a linear combination of these two vectors, they span the solution space (the kernel). A basis for the kernel is

$$\ker(A - 4I) = \mathit{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

This gives the eigenvectors

$$u_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad u_3 = \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix},$$

giving

$$y_2(t) = e^{4t}u_2, \quad y_3(t) = e^{4t}u_3.$$

Step 3: Verify linear independence.
The Wronskian at $t = 0$ *is*

$$W(0) = \det \begin{pmatrix} 1 & 1 & -2 \\ 1 & 1 & 0 \\ 3 & 0 & 1 \end{pmatrix} = 6 \neq 0,$$

so the solutions are linearly independent.
Step 4: General solution.
Therefore, the general solution is

$$y(t) = c_1 e^{-2t} \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix} + c_2 e^{4t} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + c_3 e^{4t} \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}.$$

Definition 33 (Diagonalizable matrices)**.** *A matrix $A \in M_n(\mathbb{R})$ is diagonalizable if there exists an invertible matrix P such that $P^{-1}AP$ is a diagonal matrix.*

Theorem 12 (Diagonalization)**.** *Let $A \in M_n(\mathbb{R})$ and assume that all eigenvalues of A are real and distinct, then A is diagonalizable.*

Proof. Denote by $\lambda_1, \ldots, \lambda_n$ the real, distinct eigenvalues of A. For each eigenvalue λ_i, let $u_i \in \mathbb{R}^n$ be a corresponding eigenvector, so

$$Au_i = \lambda_i u_i.$$

Since the eigenvalues are distinct, a standard result from linear algebra states that the eigenvectors $u_1, \ldots, u_n$ are linearly independent. Therefore they form a basis of $\mathbb{R}^n$. Let P be the matrix

$$P = \begin{bmatrix} u_1 & u_2 & \ldots & u_n \end{bmatrix},$$

whose columns are these eigenvectors. Because the columns are linearly independent, P is invertible. Now observe how A acts on the columns of P. For the ith standard basis vector e_i,

$$APe_i = Au_i = \lambda_i u_i.$$

Thus multiplying by A simply scales the ith column of P by λ_i. This means that multiplying P on the right by the diagonal matrix

$$D = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$$

has exactly the same effect, since PD also scales the ith column by λ_i. Hence

$$AP = PD.$$

Finally, multiplying on the left by P^{-1} gives

$$P^{-1}AP = D,$$

so A is similar to the diagonal matrix D, and therefore diagonalizable. □

Theorem 13 (Diagonalization 2)**.** *A is diagonalizable if and only if it has a basis of eigenvectors.*

Example. *Let*

$$A = \begin{pmatrix} 3 & -1 & -2 \\ -1 & 5 & -2 \\ -3 & 3 & -2 \end{pmatrix}$$

Find the basis of eigenvectors of A.
Step 1: Find the eigenvalues.
Compute the characteristic polynomial:

$$p(\lambda) = \det(A - \lambda I) = -(\lambda + 2)(\lambda - 4)^2.$$

Hence the eigenvalues are $\lambda = -2$ and $\lambda = 4$ (with multiplicity 2).
Step 2: Find eigenvectors.
For $\lambda = -2$, solve $(A + 2I)u = 0$. Row reduction gives one free variable, so the eigenspace is one-dimensional. A basis vector is

$$u_1 = \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}.$$

For $\lambda = 4$, solve $(A - 4I)u = 0$. Row reduction gives two free variables, so the eigenspace is two-dimensional. A convenient basis is

$$u_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad u_3 = \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}.$$

Step 3: Form a basis.
Thus the eigenvectors

$$\{u_1, u_2, u_3\} = \left\{ \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\}$$

are linearly independent and form a basis of $\mathbb{R}^3$. Since A has a basis of eigenvectors, it is diagonalizable.

Example. *Let*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

Step 1: Find the eigenvalues.
Since A is upper triangular, the eigenvalues are the diagonal entries. Thus

$$\lambda_1 = \lambda_2 = 1.$$

Step 2: Find the eigenvectors.
Solve

$$(A - I)u = 0.$$

Then

$$A - I = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Let $u = (a, b)^T$. *The system gives* $b = 0$, *while* a *is free. Hence every eigenvector has the form*

$$u = a \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Step 3: Describe the eigenspace.
Therefore the eigenspace is

$$\text{span}\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\},$$

which is one-dimensional. Although the eigenvalue has multiplicity 2, *we only get one independent eigenvector. Hence there is not enough to form a basis of* $\mathbb{R}^2$, *and* A *is not diagonalizable.*

Theorem 14 (General solution via diagonalization)**.** *Let* $A \in M_n(\mathbb{R})$ *and consider the system* (4). *If* A *is diagonalizable with eigenvalues* $\lambda_1, \ldots, \lambda_n$ *and corresponding eigenvectors* $u_1, \ldots, u_n$ *(possibly complex), then the general solution is*

$$y(t) = \sum_{i=1}^{n} c_i e^{\lambda_i t} u_i,$$

where $c_1, \ldots, c_n$ *are constants.*

5.4.1 Complex eigenvalues and real-valued solutions

Consider the system (4). If A has real entries and a complex eigenpair (λ, u) with $\lambda = \alpha + i\beta$ and $u = p + iq$ (where $p, q \in \mathbb{R}^n$), then a complex solution is

$$y(t) = e^{(\alpha + i\beta)t}(p + iq). \tag{7}$$

Definition 34 (Euler's Formula)**.** *Let* $x \in \mathbb{R}$. *Then the Euler's formula is*

$$e^{ix} = \cos(x) + i\sin(x). \tag{8}$$

This identity follows from the complex exponential power series

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}, \quad z \in \mathbb{C}.$$

Using Euler's formula (8), we expand (7) as

$$y(t) = e^{(\alpha + i\beta)t}(p + iq) = e^{\alpha t} e^{i\beta t}(p + iq) = e^{\alpha t}(\cos(\beta t) + i\sin(\beta t))(p + iq).$$

Multiplying and separating real and imaginary parts gives

$$y(t) = e^{\alpha t}(\cos(\beta t)p - \sin(\beta t)q) + i\, e^{\alpha t}(\sin(\beta t)p + \cos(\beta t)q).$$

Taking real and imaginary parts therefore yields two real solutions:

$$y_1(t) = e^{\alpha t}\big(p\cos(\beta t) - q\sin(\beta t)\big), \qquad y_2(t) = e^{\alpha t}\big(p\sin(\beta t) + q\cos(\beta t)\big). \tag{9}$$

Proposition 3 (Complex conjugate eigenpairs)**.** *Let $A \in M_n(\mathbb{R})$. If (λ, u) is an eigenpair with $\lambda \in \mathbb{C}$, then $(\overline{\lambda}, \overline{u})$ is also an eigenpair.*

Proof. From $Au = \lambda u$, take complex conjugates:

$$\overline{Au} = \overline{\lambda u}.$$

Since A is real, $\overline{A} = A$, so

$$A\overline{u} = \overline{\lambda}\,\overline{u}.$$

□

Example. *Let*

$$A = \begin{pmatrix} 2 & -1 \\ 1 & 2 \end{pmatrix}$$

Step 1: Find the eigenvalues.
Define the characteristic polynomial to be

$$p(\lambda) = \begin{vmatrix} 2-\lambda & -1 \\ 1 & 2-\lambda \end{vmatrix} = (\lambda - 2)^2 + 1 = 0$$

Giving us two possible eigenvalues $\lambda_1 = 2 + i$ and $\lambda_2 = 2 - i$.
Step 2: Find the eigenvectors.
Solve $(A - \lambda_1 I)u = 0$. Then

$$A - (2+i)I = \begin{pmatrix} -i & -1 \\ 1 & -i \end{pmatrix}.$$

Let $u = (a, b)^T$. The system gives $b = ia$, so taking $a = 1$ we obtain

$$u_1 = \begin{pmatrix} 1 \\ i \end{pmatrix}.$$

By the proposition (3), the second eigenvector is

$$u_2 = \overline{u_1} = \begin{pmatrix} 1 \\ -i \end{pmatrix}.$$

Problem 4 (Complex eigenvalues and real-valued solutions)**.** *Consider the linear system*

$$\vec{y}' = \begin{pmatrix} -3 & -2 \\ 5 & 3 \end{pmatrix} \vec{y}$$

(a) Find the eigenvalues and eigenvectors for the coefficient matrix.

(b) Find the real-valued solution to the initial value problem.

$$\begin{cases} y_1' = -3y_1 - 2y_2, & y_1(0) = 7 \\ y_2' = 5y_1 + 3y_2, & y_2(0) = -15 \end{cases}$$

Use t as the independent variable in your answers.

February 12, 2026

Proposition 4. *Assume that $\lambda = \alpha + i\beta \in \sigma(A)$ is an eigenvalue associated to the eigenvector $u = p + iq$, where $\beta \neq 0$. Then the functions (9) are real-valued linearly independent solutions of (4).*

Reviewed pages 14-17 from Chapter 3 prof notes

5.4.2 Nondiagonalizable Matrices

Remark 14. *Assume that $A \in M_n(\mathbb{R})$ does not admit a basis of eigenvectors. Then there are not enough solutions of the form $y(t) = e^{\lambda t}u$ to build the general solution.*

Definition 35 (Multiplicity of Eigenvalues)**.** *Let $\lambda_0 \in \sigma(A)$.*

1. *The algebraic multiplicity of λ_0 is the largest integer $m \geq 1$ such that $(\lambda - \lambda_0)^m$ divides $p(\lambda)$.*
2. *The geometric multiplicity of λ_0 is the number $\dim ker(A - \lambda_0 I)$. In other words, it is the number of linearly independent eigenvectors associated to λ_0.*

The defect of λ_0 is given by $d_{\lambda_0} = m - \dim ker(A - \lambda_0 I) \geq 0$. We call an eigenvalue defective if $d_{\lambda_0} > 0$.

Example. *Let*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad p(\lambda) = (\lambda - 1)^2.$$

Hence the spectrum is $\sigma(A) = \{1\}$, so the algebraic multiplicity is $m = 2$. To find the geometric multiplicity, compute the kernel of $A - I$:

$$A - I = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Solve

$$(A - I)x = 0, \quad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

Then

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

which gives $x_2 = 0$ and x_1 free. Thus

$$\ker(A - I) = \left\{ \begin{pmatrix} x_1 \\ 0 \end{pmatrix} : x_1 \in \mathbb{R} \right\} = \operatorname{span}\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}.$$

This space is one-dimensional, so $\dim(\ker(A - I)) = 1$. Therefore the geometric multiplicity is $1 < 2$, and the defect is $d_{\lambda_0} = 2 - 1 = 1 > 0$. Hence $\lambda_0 = 1$ is defective.

Example. *Let*

$$A = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & -2 & -1 & -1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \quad p(\lambda) = \lambda^3(\lambda + 1).$$

Hence $\sigma(A) = \{0, -1\}$*, and the algebraic multiplicity of* $\lambda_0 = 0$ *is* $m = 3$*. To compute the geometric multiplicity, solve*

$$Ax = 0, \quad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}.$$

Row reduction gives one free parameter, yielding

$$\ker(A) = \operatorname{span}\left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

Thus $\dim(\ker A) = 1$*. Therefore the defect of* $\lambda_0 = 0$ *is* $d_{\lambda_0} = 3 - 1 = 2 > 0$*, so* λ_0 *is defective with defect* 2*.*

Example. *Let*

$$A = \begin{pmatrix} 3 & 1 & -2 \\ -1 & 5 & -2 \\ -3 & 3 & -2 \end{pmatrix}, \quad p(\lambda) = (\lambda + 2)(\lambda - 4)^2.$$

Hence $\sigma(A) = \{-2, 4\}$*, and the algebraic multiplicity of* $\lambda_0 = 4$ *is* $m = 2$*. To compute the geometric multiplicity, solve*

$$(A - 4I)x = 0, \quad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

Row reduction gives two free parameters, yielding

$$\ker(A - 4I) = \operatorname{span}\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Thus $\dim(\ker(A - 4I)) = 2$*. Therefore the defect of* $\lambda_0 = 4$ *is* $d_{\lambda_0} = 2 - 2 = 0$*, so* λ_0 *is not defective.*

Definition 36 (Generalized Eigenvector)**.** *Let* $\lambda \in \sigma(A)$ *with algebraic multiplicity* m*. A generalized eigenvector* u *associated to* λ *satisfies*

$$(A - \lambda I)^m u = 0.$$

Theorem 15 (Existence of a Generalized Eigenvector Basis)**.** *Let* $\lambda \in \sigma(A)$ *have algebraic multiplicity* m*. Then*

$$\dim(\ker(A - \lambda I)^m) = m.$$

As a consequence, every matrix $A \in M_n(\mathbb{R})$ *has a basis consisting of generalized eigenvectors.*

Denote

$$k_j = \ker(A - \lambda I)^j, \quad \lambda \in \sigma(A).$$

Then

$$k_0 = \{0\}, \quad k_1 = \{\text{eigenvectors associated to } \lambda\},$$

and $\dim(k_0) = 0 < \dim(k_1) < \dim(k_m) = m$.

Lemma 16. *For all j,*

$$k_j \subseteq k_{j+1}.$$

Indeed, if $u \in k_j$, then $(A - \lambda I)^j u = 0$ implies $(A - \lambda I)^{j+1} u = 0$. Hence

$$k_0 \subsetneq k_1 \subsetneq k_2 \subsetneq \cdots \subsetneq k_j = k_{j+1} = \cdots = k_m.$$

Lemma 17. *If for some $j \in \{1, \ldots, m\}$ we have $k_j = k_{j+1}$, then*

$$k_j = k_{j+1} = k_{j+2} = \cdots = k_m.$$

Proof. Assume $k_j = k_{j+1}$. Since $k_{j+1} \subseteq k_{j+2}$, let $w \in k_{j+2}$. Set $u = (A - \lambda I)w \in k_{j+1} = k_j$. Then

$$(A - \lambda I)^{j+1} u = (A - \lambda I)^{j+2} w = 0 \implies (A - \lambda I)^j u = 0 \implies (A - \lambda I)^{j+1} w = 0.$$

Thus $w \in k_{j+1}$, so $k_{j+1} = k_{j+2}$. Iterating this gives the result. □

Algorithm 1 (Computing Generalized Eigenvectors)**.** *Assume we have a basis of k_i with $i \geq 1$ and $\dim(k_i) < m$. To find new vectors in $k_{i+1} \setminus k_i$, look for $u \in k_{i+1}$ such that*

$$(A - \lambda I)u = w \in k_i,$$

for all vectors w in the basis of k_i.

Example. *Let*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad k_0 = \{0\}, \quad k_1 = \operatorname{span}\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}, \quad k_2 = ?$$

To find $u \in k_2 \setminus k_1$, solve $(A - I)u = w \in k_1$:

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

This gives $b = 1$ and a free. Pick $a = 0$:

$$\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \in k_2 \setminus k_1.$$

Check:

$$(A - I)^2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = (A - I) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

February 17, 2026

5.5 Exponential of a Matrix

Definition 37 (Matrix Exponential)**.** *For $A \in M_n(\mathbb{R})$, the exponential of A is defined by the power series*[10]

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!} = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \cdots .$$

Definition 38 (Induced Matrix Norm)**.** *Let $\|\cdot\|$ be a norm on $\mathbb{R}^n$. The induced (operator) matrix norm on $M_n(\mathbb{R})$ is defined by*

$$\|A\| = \sup_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \sup_{\|x\|=1} \|Ax\|.$$

Proposition 5 (Basic Properties)**.** *For all $A, B \in M_n(\mathbb{R})$ and all $x \in \mathbb{R}^n$:*

$$\|Ax\| \leq \|A\| \cdot \|x\|,$$

$$|||AB||| \leq |||A||| \cdot |||B|||.$$ [11]

Proof. By definition,

$$\|A\| = \sup_{\|x\|=1} \|Ax\|.$$

For any nonzero x,

$$\|Ax\| = \left\| A\left(\frac{x}{\|x\|}\right) \right\| \|x\| \leq \|A\| \cdot \|x\|.$$

For submultiplicativity, let $\|x\| = 1$. Then

$$\|ABx\| \leq \|A\| c \dot{\|} Bx\| \leq \|A\| \cdot \|B\|.$$

Taking the supremum over all unit vectors x gives

$$\|AB\| \leq \|A\| \cdot \|B\|.$$

□

Proposition 6 (Convergence)**.** *For every $A \in M_n(\mathbb{R})$, the series defining e^A converges absolutely.*

Proof. Using the triangle inequality[12],

$$|||\sum_{k=0}^{N} \frac{A^k}{k!}||| \leq \sum_{k=0}^{N} \frac{|||A|||^k}{k!}.$$

Since $\sum_{k=0}^{\infty} \frac{\|A\|^k}{k!} = e^{\|A\|}$ converges, the matrix series converges. □

[10] This definition is motivated by the scalar identity $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$.
[11] triple bar is norm for a matrix while double bar is the norm of a vector.
[12] $||A + B|| \leq ||A|| + ||B||$

Proposition 7 (Basic Properties)**.** *For all $A \in M_n(\mathbb{R})$ and $t, s \in \mathbb{R}$,*

- $e^{A \cdot 0} = I$,
- $e^{A(t+s)} = e^{At}e^{As}$,
- $e^{-At} = (e^{At})^{-1}$,
- $\frac{d}{dt}e^{At} = Ae^{At} = e^{At}A$.

If $AB = BA$, then

$$e^{A+B} = e^A e^B.$$

Proof of the Derivative Formula.

$$\frac{d}{dt}e^{At} = \frac{d}{dt}\sum_{k=0}^{\infty}\frac{A^k t^k}{k!} = \sum_{k=1}^{\infty}\frac{A^k t^{k-1}}{(k-1)!} = A\sum_{k=0}^{\infty}\frac{A^k t^k}{k!} = Ae^{At}.$$

□

Theorem 18 (Fundamental Matrix Solution)**.** *The matrix function e^{At} is a fundamental matrix solution of (4). Hence the general solution is*

$$y(t) = e^{At}C, \quad C \in \mathbb{R}^n.$$

The unique solution of the initial value problem

$$\begin{cases} y' = Ay, \\ y(0) = y_0 \end{cases}$$

is

$$y(t) = e^{At}y_0.$$

Proof. Since $\frac{d}{dt}e^{At} = Ae^{At}$, each column of e^{At} satisfies the system. Moreover,

$$e^{A \cdot 0} = I,$$

so $\det(e^{A0}) = 1 \neq 0$, and e^{At} is invertible for all t. □

February 19, 2026

Let $y(t)$ be any fundamental matrix solution of $y' = Ay$. Then there exists an invertible matrix $C \in M_n(\mathbb{R})$ such that

$$e^{At} = y(t)C.$$

If we denote the columns of e^{At} by $w_1(t), \ldots, w_n(t)$, each column can be expressed as a linear combination of the columns of $y(t)$:

$$w_j(t) = y(t)C^{(j)}, \quad C^{(j)} \in \mathbb{R}^n.$$

Evaluating at $t = 0$, we get

$$e^{A \cdot 0} = I = y(0)C \implies C = y(0)^{-1}.$$

Hence, in general,

$$e^{At} = y(t)y(0)^{-1}.$$

Example. *Let*

$$A = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}, \quad \lambda_1 = -2,\, \lambda_2 = 4, \quad u_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix},\, u_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

A fundamental matrix solution is

$$y(t) = \begin{pmatrix} -e^{-2t} & e^{4t} \\ e^{-2t} & e^{4t} \end{pmatrix}, \quad y(0) = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Compute the inverse:

$$y(0)^{-1} = \frac{1}{2}\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Thus, the matrix exponential is

$$e^{At} = y(t)y(0)^{-1} = \begin{pmatrix} \frac{1}{2}e^{-2t} + \frac{1}{2}e^{4t} & -\frac{1}{2}e^{-2t} + \frac{1}{2}e^{4t} \\ -\frac{1}{2}e^{-2t} + \frac{1}{2}e^{4t} & \frac{1}{2}e^{-2t} + \frac{1}{2}e^{4t} \end{pmatrix}.$$

Verifying at $t = 0$:

$$e^{A\cdot 0} = I.$$

Hence, if $y_0 \in \mathbb{R}^n$ is the initial condition, the solution of (4) with $y(0) = y_0$ is

$$y(t) = e^{At}y_0.$$

5.6 Building the Solution Set With e^{At}

Given any vector $v \in \mathbb{R}^n$, the function $e^{At}v$ is a solution of (4).

5.6.1 Eigenvectors

Assume u is an eigenvector associated to $\lambda \in \sigma(A)$. Then

$$e^{At}u = \sum_{k=0}^{\infty} \frac{A^k t^k}{k!}u = \sum_{k=0}^{\infty} \frac{\lambda^k t^k}{k!}u = e^{\lambda t}u.$$

Thus eigenvectors generate solutions of the form $e^{\lambda t}u$.

5.6.2 Generalized Eigenvectors

Suppose u is a generalized eigenvector associated to $\lambda \in \sigma(A)$. Then there exists $j \in \mathbb{N}$ such that

$$(A - \lambda I)^{j+1}u = 0.$$

Write

$$A = (A - \lambda I) + \lambda I.$$

Since $(A - \lambda I)$ commutes with λI,

$$e^{At} = e^{((A-\lambda I)+\lambda I)t} = e^{\lambda t}e^{(A-\lambda I)t}.$$

Hence

$$e^{At}u = e^{\lambda t}\sum_{k=0}^{\infty}\frac{(A-\lambda I)^k t^k}{k!}u.$$

Because $(A-\lambda I)^k u = 0$ for all $k > j$, the sum is finite:

$$e^{At}u = e^{\lambda t}\sum_{k=0}^{j}\frac{(A-\lambda I)^k t^k}{k!}u.$$

Theorem 19 (Generalized Eigenvector Basis)**.** *Every matrix $A \in M_n(\mathbb{R})$ admits n linearly independent generalized eigenvectors $u_1, \ldots, u_n$.*

Proposition 8. *The general solution of (4) is*

$$y(t) = \sum_{i=1}^{n} c_i e^{At} u_i, \quad c_i \in \mathbb{R},$$

where u_i are generalized eigenvectors of A.

Example. *Consider*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \sigma(A) = \{1, 1\}.$$

Define the chain of generalized eigenspaces

$$k_0 \subsetneq k_1 \subsetneq k_2 \subsetneq \cdots \subsetneq k_j \subsetneq k_{j+1},$$

where

$$k_m = \ker(A-\lambda I)^m.$$

Here,

$$k_1 = \left\langle \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\rangle \subsetneq k_2 = \left\langle \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\rangle.$$

Since

$$\dim(\ker(A-\lambda I)^2) = 2,$$

we choose

$$u_2 \in k_2 = \ker(A-\lambda I)^2 \quad \textit{with } u_2 \notin k_1.$$

The eigenvector is

$$u_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

and u_2 satisfies

$$(A-I)u_2 = u_1.$$

Take

$$u_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Then

$$y_1(t) = e^t u_1,$$

and

$$y_2(t) = e^{At} u_2 = e^t(u_2 + tu_1) = e^t \begin{pmatrix} t \\ 1 \end{pmatrix}.$$

Thus the general solution is

$$y(t) = c_1 e^t \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 e^t \begin{pmatrix} t \\ 1 \end{pmatrix}.$$

Example. *Consider*

$$A = \begin{pmatrix} 2 & 4 \\ -1 & -2 \end{pmatrix}.$$

The characteristic polynomial is

$$p(\lambda) = (2-\lambda)(-2-\lambda) + 4 = \lambda^2,$$

so $\lambda = 0$ *with multiplicity* 2 *or* $\sigma(A) = \{0, 0\}$.

$$k_1 = \ker(A) = \left\langle \begin{pmatrix} -2 \\ 1 \end{pmatrix} \right\rangle.$$

To find $u_2 \in k_2 \setminus k_1$, *solve*

$$A \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}.$$

This gives $a = -2b - 1$. *Choosing* $b = 0$,

$$u_2 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}.$$

Then

$$y_1(t) = u_1 = \begin{pmatrix} -2 \\ 1 \end{pmatrix},$$

$$y_2(t) = e^{At} u_2 = u_2 + tu_1 = \begin{pmatrix} -1-2t \\ t \end{pmatrix}.$$

Hence

$$y(t) = c_1 \begin{pmatrix} -2 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} -1-2t \\ t \end{pmatrix}.$$

A fundamental matrix is

$$Y(t) = \begin{pmatrix} -2 & -1-2t \\ 1 & t \end{pmatrix}.$$

The Wronskian is

$$W(t) = \det Y(t) = 1 \neq 0.$$

5.7 Non-homogeneous systems

Consider

$$y'(t) = Ay(t) + r(t), \tag{10}$$

where $A \in M_n(\mathbb{R})$ and $r : I \to \mathbb{R}^n$ is continuous. Let $Y(t)$ be a fundamental matrix solution of the homogeneous system (4). Suppose $y_1(t)$ and $y_2(t)$ are two solutions of (10). Define

$$y(t) = y_2(t) - y_1(t).$$

Then

$$y'(t) = y_2'(t) - y_1'(t) = (Ay_2 + r) - (Ay_1 + r) = A(y_2 - y_1) = Ay(t).$$

Thus $y(t)$ solves the homogeneous system (4). Hence there exists $C \in \mathbb{R}^n$ such that

$$y_2(t) - y_1(t) = Y(t)C.$$

Therefore,

$$y_2(t) = Y(t)C + y_1(t).$$

Theorem 20 (General Solution of Non-homogeneous Systems)**.** *Let $y_p(t)$ be any particular solution of (10). Then the general solution is*

$$y(t) = Y(t)C + y_p(t), \quad C \in \mathbb{R}^n.$$

5.7.1 Variation of Parameters

To compute a particular solution, look for

$$y_p(t) = Y(t)C_p(t),$$

where $C_p(t)$ is an unknown vector function. Differentiate:

$$y_p'(t) = Y'(t)C_p(t) + Y(t)C_p'(t).$$

Since $Y'(t) = AY(t)$,

$$y_p'(t) = AY(t)C_p(t) + Y(t)C_p'(t) = Ay_p(t) + Y(t)C_p'(t).$$

For y_p to satisfy (10), we require

$$y_p'(t) = Ay_p(t) + r(t).$$

Thus

$$Ay_p(t) + Y(t)C_p'(t) = Ay_p(t) + r(t),$$

which implies

$$Y(t)C_p'(t) = r(t).$$

Since $Y(t)$ is invertible,

$$C_p'(t) = Y^{-1}(t)r(t).$$

Integrating,

$$C_p(t) = \int Y^{-1}(t)r(t)\,dt.$$

Therefore,

$$y_p(t) = Y(t)\int Y^{-1}(t)r(t)\,dt.$$

Hence the general solution of (10) is

$$y(t) = Y(t)C + Y(t)\int Y^{-1}(t)r(t)\,dt = Y(t)\left[C + \int Y^{-1}(t)r(t)\,dt\right], \quad C \in \mathbb{R}^n.$$

5.8 Mass-Spring Systems

February 24, 2026

Consider a mass attached to a vertical spring.



At equilibrium, gravity and the spring force balance:

$$mg + kL = 0.$$

Definition 39 (Hooke's Law)**.** *The restoring force exerted by a spring is proportional to its displacement from equilibrium:*

$$F_s = -kx,$$

where $k > 0$ is the spring constant.

Let $u(t)$ denote the displacement from equilibrium, with $u > 0$ downward. Additional forces include damping $F_d = -\gamma u'(t)$ and an external force $F(t)$. Applying Newton's second law gives

$$mu''(t) = mg - k(L + u(t)) - \gamma u'(t) + F(t).$$

Using the equilibrium relation, this simplifies to

$$mu''(t) + \gamma u'(t) + ku(t) = F(t). \tag{11}$$

To express this as a first-order system, set

$$y_1 = u, \qquad y_2 = u', \qquad y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}.$$

Then

$$y' = \begin{pmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{\gamma}{m} \end{pmatrix} y + \begin{pmatrix} 0 \\ \frac{1}{m}F(t) \end{pmatrix}.$$

We first analyze the homogeneous system $y' = Ay$, where

$$A = \begin{pmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{\gamma}{m} \end{pmatrix}.$$

The characteristic polynomial is

$$p(\lambda) = \det(A - \lambda I) = \lambda^2 + \frac{\gamma}{m}\lambda + \frac{k}{m}.$$

Thus,

$$\lambda_\pm = \frac{1}{2m}\left(-\gamma \pm \sqrt{\gamma^2 - 4km}\right).$$

Let $\Delta = \gamma^2 - 4km$.

- $\Delta > 0$ (Overdamped)

  Two distinct real negative eigenvalues. The solution is

  $$u(t) = c_1 e^{\lambda_1 t} + c_2 e^{\lambda_2 t},$$

  which decays to zero without oscillation.

- $\Delta = 0$ (Critically damped).

  A repeated eigenvalue $\lambda = -\frac{\gamma}{2m}$. The solution is

  $$u(t) = (c_1 + c_2 t)e^{\lambda t},$$

  which decays to zero as quickly as possible without oscillating.

- $\Delta < 0$ (Underdamped).

  Complex conjugate eigenvalues

  $$\lambda = -\frac{\gamma}{2m} \pm i\omega, \qquad \omega = \frac{1}{2m}\sqrt{4km - \gamma^2}.$$

  The solution becomes

  $$u(t) = e^{-\frac{\gamma}{2m}t}\left(c_1 \cos(\omega t) + c_2 \sin(\omega t)\right),$$

  which represents exponentially decaying oscillations.

5.9 RLC Circuits

Consider a series RLC circuit.



Let $Q(t)$ denote the charge on the capacitor at time t. The current is

$$I(t) = Q'(t).$$

Definition 40 (Kirchhoff's Voltage Law)**.** *The sum of the voltage drops across the elements of a closed circuit equals the applied voltage $E(t)$.*

Voltage drops across each component:

- Resistor: $RI(t)$
- Capacitor: $\frac{Q(t)}{C}$
- Inductor: $LI'(t)$

Applying Kirchhoff's law,

$$RI + \frac{Q}{C} + LI' = E(t).$$

Substituting $I = Q'$, we obtain

$$LQ''(t) + RQ'(t) + \frac{1}{C}Q(t) = E(t). \tag{12}$$

This has the same mathematical form as the mass-spring system, with the correspondence

$$m \leftrightarrow L, \quad \gamma \leftrightarrow R, \quad k \leftrightarrow \frac{1}{C}.$$

6 Stability, Phase Portraits and Orbits

March 10, 2026

Let $y'(t) = f(y(t))$, f locally Lipschitz, and $y_0 \in \mathbb{R}^n$. Denote by

$$\Gamma_{y_0} = \{y(t) \mid y'(t) = f(y(t)),\ y(0) = y_0,\ t \in J_{\max}\},$$

the *orbit* of y_0, where $J_{\max}$ is the maximal interval of existence. The orbit is a curve in $\mathbb{R}^n$.

6.1 Orbits and Phase Portraits

Definition 41 (Phase Portrait)**.** *The phase portrait is the collection of all orbits of the system $y' = f(y)$.*

Example. *Let $y' = y(1-y)$. Since this is a separable ODE, the solution is*

$$y(t) = \frac{y_0}{y_0 + (1-y_0)e^{-t}}.$$

The orbits are:

- $\Gamma_0 = \{0\}$,
- $\Gamma_1 = \{1\}$,
- $\Gamma_{y_0} = (0,1)$ *for* $y_0 \in (0,1)$,
- $\Gamma_{y_0} = (1,+\infty)$ *for* $y_0 > 1$,
- $\Gamma_{y_0} = (-\infty,0)$ *for* $y_0 < 0$.

The values 0 *and* 1 *are the equilibrium solutions.*

6.2 Equilibrium Solutions

Definition 42 (Equilibrium Solution)**.** *A vector $\tilde{y} \in \mathbb{R}^n$ is called an* equilibrium solution *(or steady state) of $y' = f(y)$ if $f(\tilde{y}) = 0$. Since $\tilde{y}' = 0 = f(\tilde{y})$, the solution satisfying $y(0) = \tilde{y}$ is constant: $\tilde{y}(t) = \tilde{y}$ for all $t \in \mathbb{R}$, and its orbit is the single point $\Gamma_{\tilde{y}} = \{\tilde{y}\}$.*

Example. *For the linear system $y' = Ay$, equilibrium solutions satisfy $Au = 0$, i.e. they lie in* $\ker(A)$.

- *If A is invertible, then* $\ker(A) = \{0\}$, *so $\tilde{y} = 0$ is the only equilibrium.*
- *If A is not invertible,* $\ker(A)$ *contains more equilibria.*

Example. *Let* $A = \begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix}$. *Solving $Au = 0$ gives $a = b$, so*

$$\ker(A) = \left\langle \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\rangle = \left\{ c \begin{pmatrix} 1 \\ 1 \end{pmatrix} \,\middle|\, c \in \mathbb{R} \right\}.$$

Every point on this line is an equilibrium solution.

Example (Nonlinear Pendulum)**.** *From Newton's second law,*

$$\theta'' + \gamma\theta' + \frac{g}{l}\sin(\theta) = 0.$$

Setting $y_1 = \theta$, $y_2 = \theta'$, the system becomes

$$\begin{pmatrix} y_1' \\ y_2' \end{pmatrix} = \begin{pmatrix} y_2 \\ -\frac{g}{l}\sin(y_1) - \gamma y_2 \end{pmatrix} =: f(y).$$

Equilibria satisfy $f(y) = 0$: we need $y_2 = 0$ and $\sin(y_1) = 0$, giving

$$y_1 = n\pi, \quad n \in \mathbb{Z}.$$

6.3 Stability

6.3.1 Definitions

Definition 43 (Stable Equilibrium)**.** *An equilibrium $\tilde{y}$ of $y' = f(y)$ is* stable *if for every $\epsilon > 0$ there exists $\delta = \delta(\epsilon) > 0$ such that every solution with $\|y(0) - \tilde{y}\| \leq \delta$ satisfies*

$$\|y(t) - \tilde{y}\| \leq \epsilon \quad \forall\, t \geq 0.$$

Intuitively: solutions that start close to $\tilde{y}$ remain close for all future time.

Definition 44 (Unstable Equilibrium)**.** *An equilibrium $\tilde{y}$ is* unstable *if it is not stable: there exists $\epsilon > 0$ such that for every $\delta > 0$, there is a solution $y(t)$ with $\|y(0) - \tilde{y}\| \leq \delta$ and a time $T > 0$ for which $\|y(T) - \tilde{y}\| > \epsilon$.*

Definition 45 (Asymptotic Stability)**.** *An equilibrium $\tilde{y}$ is* asymptotically stable *if it is stable and there exists $\alpha > 0$ such that*

$$\|y(0) - \tilde{y}\| \leq \alpha \implies \lim_{t \to +\infty} \|y(t) - \tilde{y}\| = 0.$$

6.3.2 Examples

March 12, 2026

Example (Stable equilibrium)**.** *Let $y' = -2y$. The unique equilibrium is $\tilde{y} = 0$, and the solution is $y(t) = e^{-2t}y(0)$. For $t \geq 0$,*

$$|y(t) - 0| = e^{-2t}|y(0)| \leq |y(0)|.$$

Given $\epsilon > 0$, choose $\delta = \epsilon$. Then $|y(0)| \leq \delta$ implies $|y(t)| \leq \epsilon$ for all $t \geq 0$, so $\tilde{y} = 0$ is stable.

Example (Logistic equation)**.** *Let $y' = y(1 - y)$. The equilibria are $\tilde{y} = 0$ and $\tilde{y} = 1$.*

$\tilde{y} = 0$ ***is unstable.*** *Take $\epsilon = \frac{1}{2}$ and any $0 < \delta < 1$. Choose $y(0) = \delta$; the solution is*

$$y(t) = \frac{\delta}{\delta + (1 - \delta)e^{-t}}.$$

There exists $T = T(\delta)$ such that $y(T) \geq \frac{1}{2}$, so $|y(T) - 0| \geq \epsilon$. Hence $\tilde{y} = 0$ is unstable.

$\tilde{y} = 1$ ***is asymptotically stable.*** *For any $y_0 > 0$,*

$$|y(t) - 1| = \left| \frac{(y_0 - 1)e^{-t}}{y_0 + (1 - y_0)e^{-t}} \right| = \frac{|y_0 - 1|}{y_0 e^t + (1 - y_0)} \leq |y_0 - 1|,$$

since $y_0(e^t - 1) + 1 \geq 1$ for $t \geq 0$. Choosing $\delta = \epsilon$ gives stability. Moreover, as $t \to +\infty$ the numerator $|y_0 - 1|e^{-t} \to 0$, so $y(t) \to 1$ and $\tilde{y} = 1$ is asymptotically stable.

Example (Linear system)**.** *Let $y' = Ay$ with*

$$A = \begin{pmatrix} -\frac{3}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{3}{2} \end{pmatrix}.$$

Since $\det(A) \neq 0$, the only equilibrium is $\tilde{y} = 0$.

A fundamental matrix solution is

$$Y(t) = \begin{pmatrix} -e^{-2t} & e^{-t} \\ e^{-2t} & e^{-t} \end{pmatrix},$$

giving

$$e^{At} = Y(t)Y(0)^{-1} = \frac{1}{2} \begin{pmatrix} e^{-2t} + e^{-t} & -e^{-2t} + e^{-t} \\ -e^{-2t} + e^{-t} & e^{-2t} + e^{-t} \end{pmatrix}.$$

Using the induced sup-norm $\|x\|_\infty = \max\{|x_1|, |x_2|\}$ and the corresponding matrix norm

$$\|M\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^{n} |M_{ij}|,$$

one checks that $\|e^{At}\|_\infty \leq 2e^{-t}$. Therefore

$$\|y(t)\|_\infty = \|e^{At} y(0)\|_\infty \leq 2e^{-t} \|y(0)\|_\infty.$$

Stability: *given $\epsilon > 0$, choose $\delta = \frac{\epsilon}{2}$. Then $\|y(0)\|_\infty \leq \delta$ implies $\|y(t)\|_\infty \leq 2e^{-t}\delta \leq \epsilon$ for all $t \geq 0$.*

Asymptotic stability: *for any $\alpha > 0$, if $\|y(0)\|_\infty \leq \alpha$ then $\|y(t)\|_\infty \leq 2\alpha e^{-t} \to 0$ as $t \to +\infty$. Hence $\tilde{y} = 0$ is asymptotically stable.*

6.4 Phase Portraits of Planar Linear Systems

Let (4) hold with $A \in M_2(\mathbb{R})$ and $\det(A) \neq 0$.

1. A is diagonalizable with real eigenvalues. The general solution is

$$y(t) = c_1 e^{\lambda_1 t} u_1 + c_2 e^{\lambda_2 t} u_2.$$

7 Laplace Transform

8 Appendix

8.1 Picard Iterations Convergence

Proof of Picard Iterations Convergence. Pick $t \in [t_0, t_0 + \epsilon]$ (the proof is similar for $t \in [t_0 - \epsilon, t_0]$). The goal is to show that $\{y_k(t)\}_{k=0}^{\infty}$ is a Cauchy sequence[13] in $\mathbb{R}^n$.[14] We prove by induction that

$$(**) \quad ||y_m(t) - y_{m-1}(t)|| \leq L^{m-1} M_{\alpha,\delta} \frac{(t-t_0)^m}{m!}, \ \forall m \geq 1.$$

Base case $m = 1$:

$$||y_1(t) - y_0(t)|| = \left\| \int_{t_0}^{t} f(y_0, s) ds \right\| \leq \int_{t_0}^{t} ||f(y_0, s)|| ds \leq M_{\alpha,\delta} |t - t_0| \leq \alpha.$$

Induction step:

$$||y_{m+1}(t) - y_m(t)|| \leq \int_{t_0}^{t} ||f(y_m(s), s) - f(y_{m-1}(s), s)|| ds.$$

Since $D_{\alpha,\delta}$ is compact and f is Lipschitz on $D_{\alpha,\delta}$, there exists L such that

$$||f(x,t) - f(y,t)|| \leq L||x - y||.$$

Thus,

$$\leq L \int_{t_0}^{t} ||y_m(s) - y_{m-1}(s)|| ds.$$

Using (**),

$$\leq L^m M_{\alpha,\delta} \frac{1}{(m-1)!} \int_{t_0}^{t} (s - t_0)^{m-1} ds = L^m M_{\alpha,\delta} \frac{(t-t_0)^m}{m!}.$$

Hence (**) holds. In particular, for all $\rho \geq 1$,

$$||y_\rho(t) - y_{\rho-1}(t)|| \leq M_{\alpha,\delta} \frac{(L(t-t_0))^\rho}{(\rho)!} < \frac{M_{\alpha,\delta}}{L} \frac{(L\epsilon)^\rho}{\rho!}.$$

Let $m, p \geq 1$:

$$||y_{m+p}(t) - y_{m+1}(t)|| \leq \sum_{k=1}^{p-1} ||y_{m+k+1}(t) - y_{m+k}(t)||.$$

[13] Cauchy sequence is a sequence that has a limit in a metric space $\mathbb{R}^n$.

[14] Evaluated on compacted cylinder, understanding epsilon, the L Lipschitz constant coming from somewhere, not on the analysis background such as the Banach fixed point theorem.

So,

$$< \frac{M_{\alpha,\delta}}{L} \sum_{j=m+2}^{m+p} \frac{(L\epsilon)^j}{j!}.$$

Since $e^{L\epsilon} = \sum_{j=0}^{\infty} \frac{(L\epsilon)^j}{j!}$ converges,

$$\to_{m,p\to+\infty} 0.$$

Thus $\{y_k(t)\}$ is Cauchy and converges to $y(t)$. Taking limits in the iteration,

$$y(t) = y_0 + \int_{t_0}^{t} f(y(s), s)ds.$$

□

9 Solutions

Solution 1 (System of First Order ODEs)**.** *Define new variables:*

$$y_1 = y, \quad y_2 = y', \quad y_3 = y''.$$

Now we can rewrite the third order ODE into a first order system:

$$\begin{cases} y_1' = y_2 \\ y_2' = y_3 \\ y_3' = -4y_2 + y_1 \end{cases}$$

Solution 2 (Existence and Uniqueness Theorem)**.** *Let $f(y,t) = t^2 + y^2$. This function is continuous and has continuous partial derivatives $\forall (t,y) \in \mathbb{R}^2$, ensuring a unique local solution. To find the guaranteed interval of existence ϵ, we define a rectangular domain D centered at the origin:*

$$D_{\alpha,\delta} = \{(y,t) : \|y\| \leq \alpha, |t| \leq \delta\}.$$

Now we need to bound the function. We find the maximum value M of $|f(t,y)|$ on this domain:

$$M = \sup_{(t,y)\in D_{\alpha,\delta}} |t^2 + y^2| = \delta^2 + \alpha^2.$$

Now we determine the Existence Interval. According to the theorem, the solution is guaranteed to exist for $|t| \leq \epsilon$, where:

$$\epsilon = \min\left(\delta, \frac{\alpha}{M}\right) = \min\left(\delta, \frac{\alpha}{\delta^2 + \alpha^2}\right).$$

To find the largest possible value for ϵ, we first maximize $h(\alpha) = \frac{\alpha}{\delta^2+\alpha^2}$ with respect to α for a fixed δ:

$$h'(\alpha) = \frac{(\delta^2 + \alpha^2)(1) - \alpha(2\alpha)}{(\delta^2 + \alpha^2)^2} = \frac{\delta^2 - \alpha^2}{(\delta^2 + \alpha^2)^2}.$$

Setting $h'(\alpha) = 0$ gives $\alpha = \delta$.Substituting $\alpha = \delta$ back into our expression for ϵ:

$$\epsilon = \min\left(\delta, \frac{\delta}{\delta^2 + \delta^2}\right) = \min\left(\delta, \frac{1}{2\delta}\right).$$

The maximum value of this minimum occurs when the two terms are equal:

$$\delta = \frac{1}{2\delta} \implies \delta^2 = \frac{1}{2} \implies \delta = \frac{1}{\sqrt{2}}.$$

The largest guaranteed interval of existence ϵ using this rectangular domain is:

$$\epsilon = \frac{1}{\sqrt{2}} \approx 0.707.$$

Solution 3 (LLC)**.** *We have*

$$f(y) = 3y^{2/3}, \qquad f'(y) = 2y^{-1/3}.$$

This derivative is not bounded near $y = 0$, so f is not locally Lipschitz at $y = 0$. The function

$$y_1(t) = 0$$

is a solution. In Chapter 2, solving the separable equation (assuming $y \neq 0$) gives another solution

$$y_2(t) = \begin{cases} t^3, & t \geq 0, \\ 0, & t < 0. \end{cases}$$

Indeed,

$$y_2'(t) = 3t^2 = 3(y_2(t))^{2/3}.$$

Thus, the solution is not unique.

Solution 4 (Complex eigenvalues and real-valued solutions)**.** *We are given the system*

$$\vec{y'} = A\vec{y}, \qquad A = \begin{pmatrix} -3 & -2 \\ 5 & 3 \end{pmatrix}.$$

(a) Eigenvalues and eigenvectors. *The characteristic polynomial is*

$$p(\lambda) = \det(A - \lambda I) = \begin{vmatrix} -3-\lambda & -2 \\ 5 & 3-\lambda \end{vmatrix} = (-3-\lambda)(3-\lambda) + 10 = \lambda^2 + 1.$$

Setting $p(\lambda) = 0$ gives us the eigenvalues $\lambda_1 = i$ and $\lambda_2 = -i$.

For $\lambda_1 = i$, we solve $(A - iI)u = 0$:

$$\begin{pmatrix} -3-i & -2 \\ 5 & 3-i \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 0.$$

From the first row, $(-3-i)a - 2b = 0$ implies $a = \frac{2}{-3-i}b$. Choosing $b = 1$, we obtain our first solution

$$u_1 = \begin{pmatrix} \frac{2}{-3-i} \\ 1 \end{pmatrix}.$$

For $\lambda_2 = -i$, we similarly solve $(A + iI)u = 0$:

$$\begin{pmatrix} -3+i & -2 \\ 5 & 3+i \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 0,$$

which gives $a = \frac{2}{-3+i} b$. Again choosing $b = 1$, we obtain our second solution

$$u_2 = \begin{pmatrix} \dfrac{2}{-3+i} \\ 1 \end{pmatrix}.$$

Hence,

$$\boxed{\lambda_1 = i, \quad \vec{v}_1 = \begin{pmatrix} \dfrac{2}{-3-i} \\ 1 \end{pmatrix}, \qquad \lambda_2 = -i, \quad \vec{v}_2 = \begin{pmatrix} \dfrac{2}{-3+i} \\ 1 \end{pmatrix}.}$$

(b) Real-valued solution of the initial value problem.
From the eigenpair (λ_1, u_1)[15], a complex-valued solution is

$$\vec{y}(t) = e^{it} u_1.$$

Using Euler's formula (8), we write

$$\vec{y}(t) = (\cos t + i \sin t) \begin{pmatrix} \dfrac{2}{-3-i} \\ 1 \end{pmatrix} = (\cos t + i \sin t) \begin{pmatrix} -\dfrac{3}{5} + \dfrac{1}{5} i \\ 1 \end{pmatrix}.$$

Separating real and imaginary parts yields two real-valued solutions:

$$\vec{y}_1(t) = \begin{pmatrix} -3\cos t - \sin t \\ 5 \cos t \end{pmatrix}, \qquad \vec{y}_2(t) = \begin{pmatrix} \cos t - 3 \sin t \\ 5 \sin t \end{pmatrix}.$$

Therefore, the general real solution is

$$\vec{y}(t) = c_1 \vec{y}_1(t) + c_2 \vec{y}_2(t).$$

Applying the initial conditions $y_(0) = \begin{pmatrix} 7 \\ -15 \end{pmatrix}$, we compute

$$\vec{y}_1(0) = \begin{pmatrix} -3 \\ 5 \end{pmatrix}, \qquad \vec{y}_2(0) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Thus,

$$\begin{pmatrix} 7 \\ -15 \end{pmatrix} = c_1 \begin{pmatrix} -3 \\ 5 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

[15] Since the matrix has real coefficients, the second eigenpair (λ_2, u_2) is simply the complex conjugate of (λ_1, u_1), so it produces the same two real solutions after taking real and imaginary parts. Therefore, working with one eigenpair is sufficient.

Solving gives

$$c_1 = -3, \qquad c_2 = -2.$$

Therefore, the real-valued solution of the initial value problem is

$$\vec{y}(t) = c_1\vec{y}_1(t) + c_2\vec{y}_2(t) = -3\begin{pmatrix} -3\cos t - \sin t \\ 5\cos t \end{pmatrix} - 2\begin{pmatrix} -3\sin t + \cos t \\ 5\sin t \end{pmatrix}.$$

You can simplify each component if desired:

$$\vec{y}(t) = \begin{pmatrix} 9\cos t + 3\sin t + 6\sin t - 2\cos t \\ -15\cos t - 10\sin t \end{pmatrix} = \begin{pmatrix} 7\cos t + 9\sin t \\ -15\cos t - 10\sin t \end{pmatrix}.$$

10 Useful Links