

PHYS 350: Honours Electricity and Magnetism

W.R.YCH.

August 31st, 2026

Abstract

NOT FINISHED

Contents

1	Introduction	1
2	Vector Analysis	1
2.1	Scalars, Vectors, and Fields	1
2.2	Divergence, Gradient, and Curl	1
2.3	Integrals	3
2.4	The Three Fundamental Theorems	3
2.5	Second Derivatives	4
2.6	Index Notation	5
2.7	Proving the Identities	6
2.8	Irrotational and Solenoidal Fields	7
3	Electrostatics	7
3.1	Gauss' Law and the Inverse Square Law	7
4	Potentials	9
5	Electric Fields in Matter	9
6	Magnetostatics	9
7	Magnetic Fields in Matter	9
8	Electrodynamics	9
A	Appendix	9
B	Useful Links	9

1 Introduction

Textbook used is David J. Griffiths - Introduction to Electrodynamics Fifth Edition. Taught by Debanjan Sarkar & Adrian C. Liu. Mondays, Wednesdays and Fridays from 13:35 to 14:25 in rutherford 112. Course grading is divided as follows:

- 35% problem sets (must be handwritten scans).
- 25% midterm.
- 40% final exam.

2 Vector Analysis

2.1 Scalars, Vectors, and Fields

Definition 1 (Scalar). *A scalar is a quantity with just a magnitude*

Definition 2 (Vector). *Vector is a quantity with a magnitude and a direction. A vector can be written in the form of*

$$\vec{v} = v_x \hat{x} + v_y \hat{y} + v_z \hat{z}.$$

Definition 3 (Cross Product). *The cross product is a vector perpendicular to both, set by the right hand rule, and it vanishes for parallel vectors, and can be mathematically written as*

$$\vec{v} \times \vec{w} = |\vec{v}| |\vec{w}| \sin \theta \hat{n} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ v_x & v_y & v_z \\ w_x & w_y & w_z \end{vmatrix} = \hat{x}(v_y w_z - w_y v_z) - \hat{y}(v_x w_z - w_x v_z) + \hat{z}(v_x w_y - w_x v_y).$$

Definition 4 (Dot Product). *The dot product measures how much of $\vec{v}$ points along $\vec{w}$, so it is zero for perpendicular vectors, and can be mathematically written as*

$$\vec{v} \cdot \vec{w} = v_x w_x + v_y w_y + v_z w_z = |\vec{v}| |\vec{w}| \cos \theta.$$

Remark 1. *Note $\vec{v} \times \vec{w} = -\vec{w} \times \vec{v}$, and $|\vec{v}| = \sqrt{\vec{v} \cdot \vec{v}}$.*

Definition 5 (Field). *A field is something with a value at every point in space. A scalar field assigns a number to each point, like temperature $T(x, y, z)$ or the height of a hill. A vector field assigns a vector to each point, like fluid velocity or $\vec{E}$.*

2.2 Divergence, Gradient, and Curl

Definition 6 (Del operator). *The del operator*

$$\vec{\nabla} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right),$$

means nothing until it is fed a field.

$$\vec{\nabla} f = \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z} \quad (\text{gradient: scalar} \rightarrow \text{vector}), \quad (1)$$

$$\vec{\nabla} \cdot \vec{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z} \quad (\text{divergence: vector} \rightarrow \text{scalar}), \quad (2)$$

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix} \quad (\text{curl: vector} \rightarrow \text{vector}). \quad (3)$$

For the divergence we differentiate each component along its own direction. For the curl we mix directions, which is why terms like $\partial F_y / \partial x$ show up. The direction of the curl comes from the right hand rule: curl your fingers the way the paddle wheel turns and your thumb points along $\vec{\nabla} \times \vec{F}$.

The gradient points in the steepest uphill direction, and its length grows where the hill is steeper. It is always perpendicular to the contours of constant f .

Remark 2 (Reading divergence and curl off a picture). *Divergence measures sources and sinks. Arrows pointing away from a point mean fluid is being created there (positive divergence); arrows converging on a point mean fluid is being destroyed (negative divergence). Think of a small box: non-zero divergence means the inflow and outflow through its faces do not balance.*

Curl is tested with a small paddle wheel. If the flow makes it spin, the curl is non-zero at that point. A uniform flow does not spin it, but a shear flow, where the arrows are all parallel yet longer on one side, does: one blade is pushed harder than the other, giving a net rotation. Straight field lines therefore do not imply zero curl.

Example. *Judging fields by eye:*

- $y\hat{x} - x\hat{y}$ circulates clockwise about the origin; $-y\hat{x} + x\hat{y}$ circulates counterclockwise. Both have non-zero curl.
- $y\hat{x}$ and $x\hat{y}$ have straight arrows whose length changes across the flow. These are shear fields: zero divergence, non-zero curl.
- Radial arrows growing with distance from a point: $\vec{\nabla} \cdot \vec{F} \neq 0$, $\vec{\nabla} \times \vec{F} = 0$.
- Hyperbolic flow, in along one axis and out along the other: what enters a small box leaves it, and nothing rotates, so both vanish.
- A spiral converging on a point: both are non-zero.

2.3 Integrals

An integral is an instruction to move along a region, evaluate something at each step, and add it to a running total. The multivariable versions are:

$$\int_C \vec{F} \cdot d\vec{l} \quad \text{line integral, the work done if } \vec{F} \text{ is a force,} \quad (4)$$

$$\int_S \vec{F} \cdot d\vec{a} = \int_S \vec{F} \cdot \hat{n} da \quad \text{flux, how much of the field pokes through the surface,} \quad (5)$$

$$\int_V f d^3\vec{r} = \int_V f dx dy dz \quad \text{volume integral, the mass if } f \text{ is a density.} \quad (6)$$

We write $\oint$ when the curve is a closed loop or the surface is closed.

Example. Take $\vec{F} = y\hat{x}$ along a counterclockwise quarter circle of radius R . On the circle $d\vec{l} = R d\phi(-\sin\phi\hat{x} + \cos\phi\hat{y})$ and $y = R\sin\phi$, so

$$\int_C \vec{F} \cdot d\vec{l} = -R^2 \int_0^{\pi/2} \sin^2\phi d\phi = -\frac{R^2\pi}{4}.$$

It is negative because $\vec{F}$ points along $+\hat{x}$ while we travel in the $-\hat{x}$ direction over most of the arc.

2.4 The Three Fundamental Theorems

In single variable calculus, $\int_a^b (\partial f / \partial x) dx = f(b) - f(a)$. The same pattern holds in higher dimensions: integrating a derivative over a region reduces to evaluating the original object on the boundary of that region. The three theorems differ only in the dimension of the region.

Theorem 1 (Gradient theorem). *The boundary of a curve is two points:*

$$\int_a^b \vec{\nabla} h \cdot d\vec{l} = h(b) - h(a).$$

To find my altitude change climbing a hill, I can compare before and after, or add up all the little changes along the way.

Theorem 2 (Stokes' theorem). *The boundary of a surface is a curve:*

$$\int_S (\vec{\nabla} \times \vec{F}) \cdot d\vec{a} = \oint_C \vec{F} \cdot d\vec{l}.$$

To get the total swirl, I can work out the rotation at the boundary, or add up all the little whirlpools inside. Neighbouring whirlpools cancel along their shared edges, leaving only the boundary.

Theorem 3 (Gauss' divergence theorem). *The boundary of a volume is a surface:*

$$\int_V (\vec{\nabla} \cdot \vec{F}) d^3\vec{r} = \oint_S \vec{F} \cdot d\vec{a}.$$

To find the total flux leaking out through the surface, I can add up all the sources inside. Flux between neighbouring cells cancels, leaving only the outer surface.

Remark 3. *Maxwell's equations come in a differential form written with $\vec{\nabla} \cdot$ and $\vec{\nabla} \times$, and an equivalent integral form written with $\oint \vec{F} \cdot d\vec{a}$ and $\oint \vec{F} \cdot d\vec{l}$. These two theorems are what let us pass between them. Gauss' law and Ampère's law are the first examples.*

Definition 7 (Current density). *The current I through a wire is not a local quantity, since a wire has finite thickness. The local version is the current density*

$$\vec{J} = \frac{d\vec{I}}{da_{\perp}},$$

where $da_{\perp}$ is the differential cross section perpendicular to the flow. The divergence theorem applied to $\vec{J}$ is how the conservation of charge gets written down.

2.5 Second Derivatives

Since we have three kinds of first derivative, we can mix and match to build second derivatives. Taking the divergence of a gradient gives

$$\vec{\nabla} \cdot (\vec{\nabla} f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \equiv \nabla^2 f,$$

where ∇^2 is the Laplacian. Mathematicians often write Δ , but in physics we use ∇^2 . Acting on a scalar field it returns a scalar field. It can also act on a vector field, component by component, returning a vector field.

Example (Sign of the Laplacian). *Let f have a local maximum at some point. The gradient always points uphill, so from a bird's eye view $\vec{\nabla} f$ points inward toward the maximum from every side. That field has negative divergence there, so $\nabla^2 f < 0$. The one step version: each $\partial^2 f / \partial x_i^2$ is negative at a maximum and the Laplacian is their sum. By the same argument $\nabla^2 f > 0$ at a local minimum.*

At a point where all first derivatives vanish the Laplacian can still have any sign. At a saddle point the second derivatives have opposite signs in different directions and can cancel, giving zero. So a flat point tells you nothing about $\nabla^2 f$ on its own.

The full list of second derivative identities:

$$\vec{\nabla} \cdot (\vec{\nabla} f) = \nabla^2 f \quad (\text{generally non-zero}), \quad (7)$$

$$\vec{\nabla} \times (\vec{\nabla} f) = 0 \quad (\text{curl of a gradient}), \quad (8)$$

$$\vec{\nabla} \cdot (\vec{\nabla} \times \vec{F}) = 0 \quad (\text{divergence of a curl}), \quad (9)$$

$$\vec{\nabla} \times (\vec{\nabla} \times \vec{F}) = \vec{\nabla}(\vec{\nabla} \cdot \vec{F}) - \nabla^2 \vec{F} \quad (\text{the BAC-CAB of vector calculus}). \quad (10)$$

These can be proved by writing out every partial derivative, which is tedious, or with index notation.

2.6 Index Notation

Write the i th component of $\vec{v}$ as v_i , so $v_1 = v_x$, $v_2 = v_y$, $v_3 = v_z$. A matrix carries two indices, A_{ij} , where i says which row and j says which column. The transpose swaps them, $(A^T)_{ij} = A_{ji}$. A symmetric matrix has $A_{ij} = A_{ji}$, and an antisymmetric one has $B_{ij} = -B_{ji}$.

Definition 8 (Kronecker delta). *The identity matrix is written*

$$\delta_{ij} = \begin{cases} +1 & i = j \\ 0 & i \neq j \end{cases}.$$

Matrix multiplication $\vec{w} = A\vec{v}$ becomes $w_i = \sum_j A_{ij}v_j$. Multiplying by the identity gives $\sum_j \delta_{ij}v_j = v_i$, since the delta kills every term except $j = i$. This is the main trick: a delta summed against something sets one index equal to the other.

The dot product is $\vec{v} \cdot \vec{w} = \sum_i v_i w_i$, with no free index because the result is a scalar. The cross product needs a second symbol:

$$(\vec{v} \times \vec{w})_i = \sum_{j,k} \varepsilon_{ijk} v_j w_k.$$

Definition 9 (Levi-Civita symbol). ε_{ijk} is defined by

- $\varepsilon_{123} = +1$, and cyclic permutations keep the sign: $\varepsilon_{123} = \varepsilon_{231} = \varepsilon_{312}$.
- Swapping any two indices flips the sign, so $\varepsilon_{213} = \varepsilon_{132} = \varepsilon_{321} = -1$.
- If any two indices are equal, $\varepsilon_{ijk} = 0$.

A quick way to remember the signs: place 1, 2, 3 around a circle. Reading them clockwise from any starting point gives +1, counterclockwise gives -1.

Example. For $i = 1$, only the terms with $(j, k) = (2, 3)$ and $(3, 2)$ survive:

$$(\vec{v} \times \vec{w})_1 = \varepsilon_{123} v_2 w_3 + \varepsilon_{132} v_3 w_2 = v_2 w_3 - v_3 w_2.$$

Likewise $(\vec{v} \times \vec{w})_2 = v_3 w_1 - v_1 w_3$ and $(\vec{v} \times \vec{w})_3 = v_1 w_2 - v_2 w_1$, which matches the determinant.

The identity that does all the work is

$$\sum_i \varepsilon_{ijk} \varepsilon_{imn} = \delta_{jm} \delta_{kn} - \delta_{jn} \delta_{km}, \quad (11)$$

which can be checked component by component.

Writing $\partial_i \equiv \partial/\partial x_i$, so $\partial_1 = \partial/\partial x$, $\partial_2 = \partial/\partial y$, $\partial_3 = \partial/\partial z$, the derivatives become

$$\vec{\nabla} \cdot \vec{F} = \sum_i \partial_i F_i, \quad (\vec{\nabla} \times \vec{F})_i = \sum_{j,k} \varepsilon_{ijk} \partial_j F_k.$$

The same bookkeeping as for the cross product gives the curl components,

$$(\vec{\nabla} \times \vec{F})_1 = \partial_2 F_3 - \partial_3 F_2, \quad (\vec{\nabla} \times \vec{F})_2 = \partial_3 F_1 - \partial_1 F_3, \quad (\vec{\nabla} \times \vec{F})_3 = \partial_1 F_2 - \partial_2 F_1.$$

2.7 Proving the Identities

Set $\vec{G} \equiv \vec{\nabla} \times (\vec{\nabla} \times \vec{F})$. Then

$$G_i = \sum_{j,k} \varepsilon_{ijk} \partial_j \left(\sum_{m,n} \varepsilon_{kmn} \partial_m F_n \right) \quad (12)$$

$$= \sum_{j,m,n} \left(\sum_k \varepsilon_{kij} \varepsilon_{kmn} \right) \partial_j \partial_m F_n \quad (13)$$

$$= \sum_{j,m,n} (\delta_{im} \delta_{jn} - \delta_{in} \delta_{jm}) \partial_j \partial_m F_n. \quad (14)$$

In the second line we used the cyclic property $\varepsilon_{ijk} = \varepsilon_{kij}$ to move the summed index to the front so the identity applies. The first term gives

$$\sum_j \partial_j \partial_i F_j = \partial_i \left(\sum_j \partial_j F_j \right) = \left(\vec{\nabla} (\vec{\nabla} \cdot \vec{F}) \right)_i,$$

and the second gives $-\sum_j \partial_j \partial_j F_i = -(\nabla^2 \vec{F})_i$. Together they reproduce the identity claimed above.

For the curl of a gradient,

$$\left(\vec{\nabla} \times (\vec{\nabla} f) \right)_i = \sum_{j,k} \varepsilon_{ijk} \partial_j \partial_k f. \quad (*)$$

Now j and k are internal to the sum, so their names do not matter. Swap the labels:

$$= \sum_{j,k} \varepsilon_{ikj} \partial_k \partial_j f = - \sum_{j,k} \varepsilon_{ijk} \partial_k \partial_j f = - \sum_{j,k} \varepsilon_{ijk} \partial_j \partial_k f. \quad (**)$$

The second step uses the antisymmetry of ε and the third uses the fact that partial derivatives commute. Comparing $(*)$ and $(**)$, the expression equals minus itself, and the only such number is zero. So $\vec{\nabla} \times (\vec{\nabla} f) = 0$.

2.8 Irrotational and Solenoidal Fields

The two vanishing identities let us classify fields.

- Since $\vec{\nabla} \times (\vec{\nabla} \psi) = 0$, writing $\vec{F} = \vec{\nabla} \psi$ guarantees $\vec{\nabla} \times \vec{F} = 0$. Such a field is **irrotational**.
- Since $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{A}) = 0$, writing $\vec{F} = \vec{\nabla} \times \vec{A}$ guarantees $\vec{\nabla} \cdot \vec{F} = 0$. Such a field is **solenoidal**.

Any vector field can be written as $\vec{\nabla} \psi + \vec{\nabla} \times \vec{A}$.

Theorem 4 (Irrotational fields). *The following four statements are equivalent:*

- (i) $\vec{\nabla} \times \vec{F} = 0$ everywhere.
- (ii) $\int_a^b \vec{F} \cdot d\vec{l}$ is independent of path.
- (iii) $\oint_C \vec{F} \cdot d\vec{l} = 0$ for any closed loop.
- (iv) $\vec{F} = \vec{\nabla} \psi$ for some scalar ψ .

Theorem 5 (Solenoidal fields). *The following four statements are equivalent:*

- (i) $\vec{\nabla} \cdot \vec{F} = 0$ everywhere.
- (ii) $\int_S \vec{F} \cdot d\vec{a}$ is independent of the surface, for surfaces sharing a boundary.
- (iii) $\oint_S \vec{F} \cdot d\vec{a} = 0$ for any closed surface.
- (iv) $\vec{F} = \vec{\nabla} \times \vec{A}$ for some vector field $\vec{A}$.

Remark 4. In electrostatics we will find $\vec{\nabla} \times \vec{E} = 0$. That is the irrotational case, and it is what licenses us to write $\vec{E} = -\vec{\nabla} V$ and introduce the electric potential. In magnetostatics we will find $\vec{\nabla} \cdot \vec{B} = 0$, which is the solenoidal case and licenses $\vec{B} = \vec{\nabla} \times \vec{A}$, the vector potential.

3 Electrostatics

3.1 Gauss' Law and the Inverse Square Law

Coulomb's law for two point charges,

$$F = k \frac{q_1 q_2}{r^2},$$

has the same $1/r^2$ shape as Newton's gravity, $F = Gm_1 m_2 / d^2$. The reason is geometric rather than a coincidence, and Gauss' law makes that explicit.

Law 1 (Gauss’ law). *The flux of $\vec{E}$ through any closed surface is set by the charge inside it:*

$$\oint_S \vec{E} \cdot d\vec{a} = \frac{Q_{enc}}{\epsilon_0}.$$

Surround a point charge with a sphere of radius r . The flux is fixed by Q , but the surface area over which it is spread grows like r^2 , so $E \propto 1/r^2$. The exponent is really a count of dimensions.

Example (Coulomb’s law in d dimensions). *In a universe with four spatial dimensions (x, y, z, w) , volumes are four-dimensional and the surfaces enclosing them are three-dimensional, with “area” scaling like r^3 . Gauss’ law then gives $E \propto 1/r^3$. In general, in d spatial dimensions a point charge has $E \propto 1/r^{d-1}$. The inverse square law is what three dimensions look like.*

Remark 5 (Compactified dimensions). *Suppose an extra dimension exists but is curled up, the way a flat sheet with coordinates (x, y) can be rolled into a cylinder whose second coordinate w is periodic with a small radius. At distances much smaller than that radius, field lines spread out into all the dimensions and the field falls off faster than $1/r^2$. At distances much larger than it, there is only room to spread out in the large dimensions and the usual $1/r^2$ is recovered. So a measured deviation from the inverse square law at short range would be evidence for extra dimensions.*

This has been tested for gravity by the Eöt-Wash torsion balance experiments (2004, 2007, 2020), which checked the $1/r^2$ law down to separations of $52\,\mu\text{m}$. Newtonian gravity fit the data, ruling out gravitational strength Yukawa interactions with ranges above $38.6\,\mu\text{m}$ at 95% confidence.

Remark 6 (Field lines). *Field lines start on positive charges and end on negative ones, and their density measures the field strength. This is the same statement as Gauss’ law: the number of lines through a closed surface counts the enclosed charge. Two like charges have lines that bend away from each other. An infinite sheet of charge has lines that cannot spread out at all, so its field is uniform and does not fall off with distance.*

- 4 Potentials
- 5 Electric Fields in Matter
- 6 Magnetostatics
- 7 Magnetic Fields in Matter
- 8 Electrodynamics
- A Appendix
- B Useful Links