

MATH 475: Honours Partial Differential Equations

W.R.YCH.

August 31st, 2026

Abstract

Contents

1	Introduction	1
1.1	Ways to Write a PDE	1
2	Potential Theory	1
2.1	Newton's Problem	1
2.2	Toolkit	2
2.3	Radial Chain Rule	2
2.4	The Laplacian	2
2.5	Harmonic Radial Functions	3
A	Appendix	4
B	Solutions	4
C	Useful Links	4

1 Introduction

Taught by Adam Oberman. Mondays and Wednesdays from 14:35 to 15:55 in Steward Biology Building N2/2. Course grading is divided as follows:

- 10% 5-6 Assignments.
- 10% Attendance and participation.
- 10% 3-5 quizzes.
- 20% midterm.
- 50% final exam.

1.1 Ways to Write a PDE

August 31,
2026

Many of the classical equations are built from a single operator, the Laplacian

$$\Delta u = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2}. \quad (1)$$

Equation	Form	Type
Heat	$u_t = \Delta u, \quad t \geq 0$	parabolic
Poisson	$\Delta u = f$	elliptic (stationary)
Laplace	$\Delta u = 0$	elliptic (stationary)
Wave	$u_{tt} = \Delta u$	hyperbolic

Not every PDE fits this pattern. Nonlinear examples include Burgers' equation

$$u_t + uu_x = 0, \quad (2)$$

and the Navier-Stokes equations.

2 Potential Theory

2.1 Newton's Problem

Place a point mass M at the origin and a test mass m at $x \in \mathbb{R}^3 \setminus \{0\}$. The gravitational force on m has magnitude $GmM/\|x\|^2$ and points back toward the origin:

$$\vec{F}(x) = -\frac{GmM}{\|x\|^2} \cdot \frac{x}{\|x\|} = -\frac{GmM}{\|x\|^3} x. \quad (3)$$

2.2 Toolkit

Lemma 1 (Derivative of distance is direction). *Let $r(x) = \|x\|$ with $x = (x_1, x_2, x_3) \in \mathbb{R}^3$. Then for $x \neq 0$,*

$$\frac{\partial r}{\partial x_i} = \frac{x_i}{r}, \quad \nabla r(x) = \frac{x}{r} = \hat{x}. \quad (4)$$

Proof. Start from $r^2 = x_1^2 + x_2^2 + x_3^2$ and differentiate both sides with respect to x_i :

$$2r \frac{\partial r}{\partial x_i} = 2x_i \implies \frac{\partial r}{\partial x_i} = \frac{x_i}{r}. \quad (5)$$

This requires $r \neq 0$, so the identity fails at the origin. $\square$

Remark 1. ∇r is the unit vector pointing radially outward, so $\|\nabla r\| = 1$ everywhere it is defined.

2.3 Radial Chain Rule

Proposition 1. *Let $f : (0, \infty) \rightarrow \mathbb{R}$ be differentiable and set $g(x) = f(r(x)) = f(\|x\|)$. Then for $x \neq 0$,*

$$\nabla g(x) = f'(r) \nabla r(x) = f'(r) \frac{x}{r}. \quad (6)$$

Proof. Componentwise, by the chain rule,

$$\frac{\partial g}{\partial x_i} = f'(r(x)) \frac{\partial r}{\partial x_i} = f'(r) \frac{x_i}{r}. \quad (7)$$

$\square$

So the gradient of any radial function points radially, and f' alone controls the magnitude.

Example (Gradient refresher, non-radial). *For $g(x) = x_1 x_2 + x_3^2$,*

$$\nabla g(x) = \left(\frac{\partial g}{\partial x_1}, \frac{\partial g}{\partial x_2}, \frac{\partial g}{\partial x_3} \right) = (x_2, x_1, 2x_3). \quad (8)$$

Remark 2. *The chain rule result above, $\nabla g(x) = f'(r(x)) \nabla r(x) = f'(r) \frac{x}{r}$, is the tool we use whenever a function depends only on distance from the origin.*

2.4 The Laplacian

Everything below works in $\mathbb{R}^d$ for any $d \geq 1$. Write $x = (x_1, \dots, x_d)$.

First form: sum of second partials

Definition 1 (Laplacian).

$$\Delta = \sum_{i=1}^d \frac{\partial^2}{\partial x_i^2}, \quad \Delta f(x) = \sum_{i=1}^d \frac{\partial^2 f}{\partial x_i^2}(x). \quad (9)$$

In $d = 3$ this reads

$$\Delta f(x) = \frac{\partial^2 f}{\partial x_1^2}(x) + \frac{\partial^2 f}{\partial x_2^2}(x) + \frac{\partial^2 f}{\partial x_3^2}(x). \quad (10)$$

Example. Let $f(x_1, x_2, x_3) = x_1^2 - x_2^2$. Then

$$\frac{\partial^2 f}{\partial x_1^2} = 2, \quad \frac{\partial^2 f}{\partial x_2^2} = -2, \quad \frac{\partial^2 f}{\partial x_3^2} = 0, \quad (11)$$

so $\Delta f = 2 - 2 + 0 = 0$. A function with $\Delta f = 0$ is called harmonic.

Second form: divergence of the gradient

Definition 2 (Divergence). For a vector field $\vec{v}(x) = (v^1(x), \dots, v^d(x))$, where the superscripts label components and are not powers,

$$\nabla \cdot \vec{v} = \operatorname{div} \vec{v} = \sum_{i=1}^d \frac{\partial v^i}{\partial x_i}(x). \quad (12)$$

Applying this to the gradient field $\vec{v} = \nabla f$, whose i -th component is $\partial f / \partial x_i$, gives

$$\nabla \cdot (\nabla f) = \sum_{i=1}^d \frac{\partial}{\partial x_i} \left(\frac{\partial f}{\partial x_i} \right) = \sum_{i=1}^d \frac{\partial^2 f}{\partial x_i^2} = \Delta f. \quad (13)$$

Proposition 2.

$$\Delta = \nabla \cdot \nabla = \operatorname{div} \circ \operatorname{grad}. \quad (14)$$

Remark 3. This second form is the useful one for radial functions: combining it with the radial chain rule turns Δ into an ODE in r .

2.5 Harmonic Radial Functions

Proposition 3 (Radial Laplacian). For $g(x) = f(r)$, $r = \|x\|$ in $\mathbb{R}^d$, $x \neq 0$:

$$\Delta g = f''(r) + \frac{d-1}{r} f'(r). \quad (15)$$

Solve $\Delta f = 0$ in $d = 3$ with the ansatz $f(r) = r^a$:

$$\Delta f = a(a-1)r^{a-2} + \frac{2}{r} ar^{a-1} = a(a+1)r^{a-2}, \quad (16)$$

which vanishes for $r > 0$ iff $a = 0$ or $a = -1$. So

$$g(x) = A + \frac{B}{\|x\|}. \quad (17)$$

Remark 4. $1/\|x\|$ is the Newtonian potential: $\nabla(1/r) = -x/\|x\|^3$, the inverse-square field. Harmonic away from the origin only.

A Appendix

B Solutions

C Useful Links