PHYS 350: Honours Electricity and Magnetism

W.R.YCH.

August 31st, 2026

Abstract

Contents

1 Introduction **1**

2 Vector Analysis **1**
2.1 Vector Algebra 1
2.2 Differential Calculus 1
2.3 Integral Calculus 2
2.3.1 The Fundamental Theorems 3
2.4 Spherical Coordinates 4
2.5 Index Notation 5
2.6 Second Derivatives Identities 6
2.7 The Dirac Delta Function 6
2.8 The Theory of Vector Fields 7

3 Electrostatics **7**
3.1 Electric Field 8
3.1.1 Continuous Charge Distributions 9
3.2 Divergence and Curl of Electrostatic Fields 10
3.2.1 Gauss's Law 10
3.2.2 Applications of Gauss's Law 10
3.2.3 The Curl of $\vec{E}$ 12
3.3 Electric Potential 13
3.3.1 Poisson's Equation and Laplace's Equation 14
3.3.2 The Potential of a Localized Charge Distribution 14
3.3.3 Boundary Conditions 14
3.4 Work and Energy in Electrostatics 15
3.4.1 The Energy of a Point Charge Distribution 15
3.4.2 The Energy of a Continuous Charge Distribution 16
3.4.3 Comments on Electrostatic Energy 16
3.5 Conductors 16
3.5.1 Induced Charges 17
3.5.2 Surface Charge and the Force on a Conductor 18

4 Potentials **19**
4.1 Laplace's Equation 19
4.2 The Method of Images 21
4.3 Separation of Variables: Cartesian 22
4.4 Separation of Variables: Spherical 23
4.5 Multipole Expansion 24

5 Electric Fields in Matter **27**

6 Magnetostatics **27**
6.1 Current Density 27

7 Magnetic Fields in Matter **27**

8 Electrodynamics **27**

A Extra Material **28**

B Proofs **28**

C Useful Links **30**

1 Introduction

Textbook used is David J. Griffiths - Introduction to Electrodynamics Fifth Edition. Course grading is divided as follows:

- 35% problem sets (must be handwritten scans).
- 25% midterm.
- 40% final exam.

2 Vector Analysis

August 31 - September 4, 2026

2.1 Vector Algebra

Definition 1 (Scalar)**.** *A scalar is a quantity with just a magnitude.*

Definition 2 (Vector)**.** *A vector is a quantity with a magnitude and a direction, i.e.* $\vec{v} = v_x\hat{x} + v_y\hat{y} + v_z\hat{z}$.

Definition 3 (Cross Product)**.** *The cross product is a vector perpendicular to both, set by the right hand rule, and it vanishes for parallel vectors:*

$$\vec{v}\times\vec{w} = |\vec{v}||\vec{w}|\sin\theta\,\hat{n} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ v_x & v_y & v_z \\ w_x & w_y & w_z \end{vmatrix} = \hat{x}(v_yw_z - w_yv_z) - \hat{y}(v_xw_z - w_xv_z) + \hat{z}(v_xw_y - w_xv_y). \tag{1}$$

Definition 4 (Dot Product)**.** *The dot product measures how much of* $\vec{v}$ *points along* $\vec{w}$*, so it is zero for perpendicular vectors:*

$$\vec{v}\cdot\vec{w} = v_xw_x + v_yw_y + v_zw_z = |\vec{v}||\vec{w}|\cos\theta. \tag{2}$$

Remark 1. *Note* $\vec{v}\times\vec{w} = -\vec{w}\times\vec{v}$*, and* $|\vec{v}| = \sqrt{\vec{v}\cdot\vec{v}}$.

Definition 5 (Field)**.** *A field is something with a value at every point in space. A scalar field assigns a number to each point. A vector field assigns a vector to each point.*

2.2 Differential Calculus

Definition 6 (Del operator)**.** *The del operator*

$$\vec{\nabla} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}\right).$$

Definition 7 (Gradient)**.** *The gradient turns a scalar into a vector, pointing in the direction of steepest increase of* f*:*

$$\vec{\nabla}f = \frac{\partial f}{\partial x}\hat{x} + \frac{\partial f}{\partial y}\hat{y} + \frac{\partial f}{\partial z}\hat{z}. \tag{3}$$

Remark 2. *The gradient points in the steepest uphill direction, and its length grows where the hill is steeper. It is always perpendicular to the contours of constant f.*

Definition 8 (Divergence)**.** *Divergence turns a vector into a scalar, measuring sources and sinks. For a small box, non-zero divergence means the inflow and outflow through its faces do not balance: arrows spreading from a point (positive) mean fluid is created there, arrows converging (negative) mean it is destroyed.*

$$\vec{\nabla} \cdot \vec{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}. \tag{4}$$

Definition 9 (Curl)**.** *Curl turns a vector into a vector, measuring local rotation. Picture a small paddle wheel: if the flow spins it, the curl is non-zero there. Uniform flow does not spin it, but shear flow (parallel arrows, longer on one side) does, so straight field lines do not imply zero curl.*

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix} = \left(\frac{\partial}{\partial y} F_z - \frac{\partial}{\partial z} F_y, -(\frac{\partial}{\partial x} F_z - \frac{\partial}{\partial z} F_x), \frac{\partial}{\partial x} F_y - \frac{\partial}{\partial y} F_x \right). \tag{5}$$

Remark 3. *The direction of the curl follows the right-hand rule: curl your fingers along the paddle wheel's rotation and your thumb points along $\vec{\nabla} \times \vec{F}$.*

Remark 4. *Divergence and curl are themselves fields: they can vanish in some places and not others, and a general field has both.*

2.3 Integral Calculus

Definition 10 (Line Integral)**.** *A line integral sums $\vec{F}$ along a curve, picking up only the component tangent to the path:*

$$\int_C \vec{F} \cdot d\vec{l}.$$

Example. *Take $\vec{F} = y\hat{x}$ along a counterclockwise quarter circle of radius R.*




Figure 1: The field $\vec{F} = y\hat{x}$ (gray) and the quarter circle C. At a typical point, $\vec{F}$ points along $+\hat{x}$ while $d\vec{l}$ has a $-\hat{x}$ component, so $\vec{F} \cdot d\vec{l} < 0$.

On the circle $d\vec{l} = \frac{d\vec{r}}{d\phi} d\phi = R(-\sin\phi\,\hat{x} + \cos\phi\,\hat{y})d\phi$ *and* $y = R\sin\phi$, *so*

$$\int_C \vec{F} \cdot d\vec{l} = \int_0^{\pi/2} \begin{pmatrix} R\sin\phi \\ 0 \end{pmatrix} \cdot R \begin{pmatrix} -\sin\phi \\ \cos\phi \end{pmatrix} d\phi = -R^2 \int_0^{\pi/2} \sin^2\phi\, d\phi = -\frac{R^2\pi}{4}.$$

It is negative because $\vec{F}$ *points along* $+\hat{x}$ *while we travel in the* $-\hat{x}$ *direction over most of the arc.*

Definition 11 (Flux)**.** *Flux measures how much of the field pokes through a surface:*

$$\int_S \vec{F} \cdot d\vec{a} = \int_S \vec{F} \cdot \hat{n}\, da.$$

Remark 5. *Flux is the total amount of a vector field flowing through a surface. Divergence is the flux per unit volume at a specific point.*

Definition 12 (Volume Integral)**.** *A volume integral sums a scalar over a region:*

$$\int_V f\, d^3\vec{r} = \iiint_V f\, dx\, dy\, dz.$$

Remark 6. *We write* $\oint$ *when the curve is a closed loop or the surface is closed.*

2.3.1 The Fundamental Theorems

Theorem 1 (The Fundamental Theorem of Calculus)**.** *The boundary of an interval is two points:*

$$\int_a^b f'(x)\, dx = f(b) - f(a).$$

Theorem 2 (The Fundamental Theorem for Gradients)**.**

$$\int_a^b \vec{\nabla} T \cdot d\vec{l} = T(b) - T(a).$$

The boundary of a curve is its two endpoints. To find the Eiffel Tower's height, you can sum the rise of every step (left side) or subtract altimeter readings at the top and bottom (right side).

Theorem 3 (The Fundamental Theorem for Curls)**.** *Another name for this theorem is Stokes' theorem.*

$$\iint_S (\vec{\nabla} \times \vec{F}) \cdot d\vec{S} = \oint_C \vec{F} \cdot d\vec{l}. \tag{6}$$

Geometric meaning: to get the total swirl, I can work out the rotation at the boundary, or add up all the little whirlpools inside. Neighbouring whirlpools cancel along their shared edges, leaving only the boundary.

Theorem 4 (The Fundamental Theorem for Divergence)**.** *It has 2 important names: Gauss's theorem (or the divergence theorem) and Green's theorem. Respectively:*

$$\oiint_S (\vec{F} \cdot \hat{n}) dS = \iiint_V (\vec{\nabla} \cdot \vec{F}) dV,$$

$$\oint_C P dx + Q dy = \iint_R (\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}) dx dy.$$

2.4 Spherical Coordinates

Definition 13 (Spherical coordinates)**.** *r is the distance from the origin, $\theta \in [0, \pi]$ the angle down from the $+z$ axis, and $\phi \in [0, 2\pi)$ the angle around it. The unit vectors $\hat{r}, \hat{\theta}, \hat{\phi}$ change direction from point to point, so they cannot be pulled out of an integral; convert to $\hat{x}, \hat{y}, \hat{z}$ first. The elements are*

$$d\vec{l} = dr\,\hat{r} + r\,d\theta\,\hat{\theta} + r\sin\theta\,d\phi\,\hat{\phi}, \qquad d\tau = r^2 \sin\theta\,dr\,d\theta\,d\phi, \qquad d\vec{a}_{sphere} = R^2 \sin\theta\,d\theta\,d\phi\,\hat{r}.$$

Proposition 1 (Derivatives in spherical coordinates)**.**

$$\vec{\nabla} f = \frac{\partial f}{\partial r}\hat{r} + \frac{1}{r}\frac{\partial f}{\partial \theta}\hat{\theta} + \frac{1}{r\sin\theta}\frac{\partial f}{\partial \phi}\hat{\phi},$$

$$\vec{\nabla} \cdot \vec{F} = \frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2 F_r\right) + \frac{1}{r\sin\theta}\frac{\partial}{\partial \theta}\left(\sin\theta\, F_\theta\right) + \frac{1}{r\sin\theta}\frac{\partial F_\phi}{\partial \phi},$$

$$\vec{\nabla} \times \vec{F} = \frac{1}{r\sin\theta}\left[\frac{\partial}{\partial \theta}(\sin\theta\, F_\phi) - \frac{\partial F_\theta}{\partial \phi}\right]\hat{r} + \frac{1}{r}\left[\frac{1}{\sin\theta}\frac{\partial F_r}{\partial \phi} - \frac{\partial}{\partial r}(r F_\phi)\right]\hat{\theta} + \frac{1}{r}\left[\frac{\partial}{\partial r}(r F_\theta) - \frac{\partial F_r}{\partial \theta}\right]\hat{\phi},$$

$$\nabla^2 f = \frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2 \frac{\partial f}{\partial r}\right) + \frac{1}{r^2\sin\theta}\frac{\partial}{\partial \theta}\left(\sin\theta \frac{\partial f}{\partial \theta}\right) + \frac{1}{r^2\sin^2\theta}\frac{\partial^2 f}{\partial \phi^2}.$$

Remark 7. *For a purely radial field $\vec{F} = F_r(r)\hat{r}$, only the first term of the divergence survives, and the curl vanishes. So every central field $f(r)\hat{r}$ is curl-free.*

2.5 Index Notation

Remark 8. *A matrix carries two indices, A_{ij}, where i says which row and j says which column.*

Definition 14 (Transpose)**.** *The transpose swaps the indices, $(A^T)_{ij} = A_{ji}$.*

Definition 15 (Symmetry)**.** *A symmetric matrix has $A_{ij} = A_{ji}$, and an anti-symmetric one has $B_{ij} = -B_{ji}$.*

Definition 16 (Kronecker delta)**.** *The Kronecker delta equals $+1$ when its condition is fulfilled and 0 otherwise, so it could be the identity matrix in index notation:*

$$\delta_{ij} = \begin{cases} +1 & i = j \\ 0 & i \neq j \end{cases}.$$

Example.

$$\sum_j \delta_{ij} v_j = v_i.$$

Proposition 2 (Matrix Multiplication)**.**

$$\vec{w} = A\vec{v} \implies w_i = \sum_j A_{ij} v_j.$$

Proposition 3 (Dot Product)**.**

$$\vec{v} \cdot \vec{w} = \sum_i v_i w_i; \vec{\nabla} \cdot \vec{F} = \sum_i \partial_i F_i.$$

Proposition 4 (Cross Product)**.**

$$(\vec{v} \times \vec{w})_i = \sum_{j,k} \varepsilon_{ijk} v_j w_k; (\vec{\nabla} \times \vec{F})_i = \sum_{j,k} \varepsilon_{ijk} \partial_j F_k.$$

Definition 17 (Levi-Civita symbol)**.** *ε_{ijk} is defined by*

- $\varepsilon_{123} = \varepsilon_{231} = \varepsilon_{312} = +1$.
- $\varepsilon_{321} = \varepsilon_{213} = \varepsilon_{132} = -1$.
- *If any two indices are equal, $\varepsilon_{ijk} = 0$.*

Example. *For $i = 1$, only the terms with $(j, k) = (2, 3)$ and $(3, 2)$ survive:*

$$(\vec{v} \times \vec{w})_1 = \varepsilon_{123} v_2 w_3 + \varepsilon_{132} v_3 w_2 = v_2 w_3 - v_3 w_2.$$

Proposition 5 (Epsilon-delta identity)**.** *Contracting two Levi-Civita symbols on a shared index gives*

$$\sum_i \varepsilon_{ijk} \varepsilon_{imn} = \delta_{jm}\delta_{kn} - \delta_{jn}\delta_{km}.$$

2.6 Second Derivatives Identities

Definition 18 (Laplacian)**.** *Taking the divergence of a gradient gives*

$$\vec{\nabla} \cdot (\vec{\nabla} f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \equiv \nabla^2 f, \tag{7}$$

where ∇^2 *is the Laplacian*[1]*.*

Remark 9. ∇^2 *also acts on vector fields, component by component:* $(\nabla^2 \vec{F})_i = \nabla^2 F_i$*.*

The full list of second derivative identities:

Name	**Identity**
Divergence of a gradient	$\vec{\nabla} \cdot (\vec{\nabla} f) = \nabla^2 f$
Curl of a gradient	$\vec{\nabla} \times (\vec{\nabla} f) = 0$
Divergence of a curl	$\vec{\nabla} \cdot (\vec{\nabla} \times \vec{F}) = 0$
Curl of a curl	$\vec{\nabla} \times (\vec{\nabla} \times \vec{F}) = \vec{\nabla}(\vec{\nabla} \cdot \vec{F}) - \nabla^2 \vec{F}$

Table 1: Second derivative identities.

The curl of a curl identity and the divergence of a gradient identity are proved in Appendix **??**.

2.7 The Dirac Delta Function

Definition 19 (Dirac delta)**.** $\delta(x)$ *is zero everywhere except at* $x = 0$*, where it is an infinitely tall spike of unit area:*

$$\int_{-\infty}^{\infty} \delta(x)\, dx = 1.$$

Proposition 6 (Properties)**.** *(i) Sifting:* $\int_a^b f(x)\, \delta(x - x_0)\, dx = f(x_0)$ *if* $a < x_0 < b$*, and* 0 *if* x_0 *is outside* $[a, b]$*.*

(ii) Scaling: $\delta(kx) = \frac{1}{|k|}\delta(x)$*.*

(iii) In 3D: $\int_V f(\vec{r})\, \delta^3(\vec{r} - \vec{r}_0)\, d\tau = f(\vec{r}_0)$ *if* $\vec{r}_0 \in V$*.*

Example. $\int_0^3 x^3 \delta(x + 1)\, dx = 0$ *since* $x = -1$ *is outside the range, while*

$$\int_{-1}^{1} 9x^2\, \delta(3x + 1)\, dx = \frac{1}{3} \int_{-1}^{1} 9x^2\, \delta\big(x + \tfrac{1}{3}\big)\, dx = \frac{1}{3} \cdot 9 \cdot \frac{1}{9} = \frac{1}{3}.$$

[1] Mathematicians often write Δ, but in physics we use ∇^2.

Theorem 5 (Divergence of $\hat{r}/r^2$)**.** *By the spherical divergence,* $\vec{\nabla}\cdot\left(\hat{r}/r^2\right) = \frac{1}{r^2}\partial_r(1) = 0$ *for* $r \neq 0$*, yet its flux through any sphere is* $\oint \frac{1}{R^2} R^2 \sin\theta \, d\theta \, d\phi = 4\pi$*. By the divergence theorem, everything sits at the origin:*

$$\vec{\nabla}\cdot\left(\frac{\hat{r}}{r^2}\right) = 4\pi\,\delta^3(\vec{r}).$$

So for a point charge, $\vec{\nabla}\cdot\vec{E} = \frac{q}{\epsilon_0}\delta^3(\vec{r}) = \rho/\epsilon_0$*, as Gauss' law requires.*

2.8 The Theory of Vector Fields

Remark 10. *The condition at infinity is needed: a field can be both curl-free and divergence-free without being zero, e.g.* $\vec{F} = yz\,\hat{x} + zx\,\hat{y} + xy\,\hat{z} = \vec{\nabla}(xyz)$*. Adding it to any field changes neither* $\vec{\nabla}\cdot\vec{F}$ *nor* $\vec{\nabla}\times\vec{F}$*.*

Theorem 6 (Irrotational fields)**.** *Let* $\vec{F}$ *be a* C^1 *vector field on a simply connected domain (e.g. all of* $\mathbb{R}^3$*). The following are equivalent:*

(i) $\vec{\nabla}\times\vec{F} = 0$ *everywhere.*

(ii) $\int_a^b \vec{F}\cdot d\vec{l}$ *is independent of path, for any given end points.*

(iii) $\oint_C \vec{F}\cdot d\vec{l} = 0$ *for every closed loop* C*.*

(iv) $\vec{F} = -\vec{\nabla}\psi$ *for some scalar field* ψ*.*

Theorem 7 (Solenoidal fields)**.** *Let* $\vec{F}$ *be a* C^1 *vector field on a domain with no enclosed voids (e.g. all of* $\mathbb{R}^3$*). The following are equivalent:*

(i) $\vec{\nabla}\cdot\vec{F} = 0$ *everywhere.*

(ii) $\int_S \vec{F}\cdot d\vec{a}$ *is independent of surface, for any given boundary line.*

(iii) $\oint_S \vec{F}\cdot d\vec{a} = 0$ *for any closed surface* S*.*

(iv) $\vec{F} = \vec{\nabla}\times\vec{A}$ *for some vector field* $\vec{A}$*.*

Remark 11. *In electrostatics we will find* $\vec{\nabla}\times\vec{E} = 0$*, which licenses* $\vec{E} = -\vec{\nabla}V$ *and the electric potential. In magnetostatics we will find* $\vec{\nabla}\cdot\vec{B} = 0$*, which licenses* $\vec{B} = \vec{\nabla}\times\vec{A}$*, the vector potential.*

Theorem 8 (Helmholtz)**.** *If* $\vec{\nabla}\cdot\vec{F}$ *and* $\vec{\nabla}\times\vec{F}$ *are specified everywhere and* $\vec{F}\to 0$ *at infinity, then* $\vec{F}$ *is uniquely determined. It can be written as* $\vec{F} = \vec{\nabla}\psi + \vec{\nabla}\times\vec{A}$*, an irrotational part plus a solenoidal part.*

3 Electrostatics

September 11-16 & 21-25, 2026

Definition 20 (Electrostatics)**.** *Electrostatics is the study of stationary electric charges.*

3.1 Electric Field

Goal of electrodynamics: given source charges with known positions, find the force on a test charge Q.

Proposition 7 (Principle of Superposition)**.** *The force between two charges is unaffected by other charges, so*

$$\vec{F} = \sum_i \vec{F}_i.$$

Law 1 (Coulomb's law)**.** *The force on a test charge Q due to a point charge q is*

$$\vec{F} = \frac{1}{4\pi\epsilon_0}\frac{qQ}{\mathfrak{r}^2}\,\hat{\mathfrak{r}}\,[N]\,, \tag{8}$$

where $\vec{\mathfrak{r}} = \vec{r} - \vec{r'}$ is the separation vector from the location of q ($\vec{r'}$) to Q ($\vec{r}$).

Remark 12. *Coulomb's law has the same shape as Newton's gravity. The reason is geometric rather than a coincidence, and Gauss' law makes that explicit: in d spatial dimensions the flux spreads over a $(d-1)$-dimensional sphere, so $F \propto 1/\mathfrak{r}^{d-1}$, and the 4π is the area of the unit sphere in $d = 3$.*

Definition 21 (Permittivity of free space)**.** *The permittivity of free space measures how easily an electric field can form in a classical vacuum:*

$$\epsilon_0 = 8.85 \times 10^{-12}\ \left[\frac{C^2}{N \cdot m^2}\right].$$

By the principle of superposition, the total force on a test charge Q due to several point charges is

$$\vec{F} = \frac{Q}{4\pi\epsilon_0}\sum_{i=1}^{n}\frac{q_i}{\mathfrak{r}_i^2}\,\hat{\mathfrak{r}}_i.$$

Which leads us to the definition of the electric field

$$\vec{F} = Q\vec{E}.$$

Definition 22 (Electric field)**.** *The electric field is the force per unit test charge.*

$$\vec{E}(\vec{r}) = \frac{1}{4\pi\epsilon_0}\sum_{i=1}^{n}\frac{q_i}{\mathfrak{r}_i^2}\,\hat{\mathfrak{r}}_i. \tag{9}$$

$\vec{\mathfrak{r}}_i \equiv \vec{r} - \vec{r}_i$, where $\vec{r}$ is the location of the field point, where we want to know what force a charge would feel, and $\vec{r}_i$ is the location of the source charge q_i.

3.1.1 Continuous Charge Distributions

Definition 23 (Continuous distributions)**.** *For a charge along a line, over a surface, or through a volume, replace q_i by $dq = \lambda\, dl' = \sigma\, da' = \rho\, d\tau'$ and the sum becomes an integral:*

$$\vec{E}(\vec{r}) = \frac{1}{4\pi\epsilon_0}\int \frac{dq}{\mathfrak{r}^2}\,\hat{\mathfrak{r}}, \qquad \vec{\mathfrak{r}} = \vec{r} - \vec{r}', \tag{10}$$

where $\vec{r}$ is the field point and $\vec{r}'$ the source point.

Example (Line of charge)**.** *A straight segment of length $2L$, centred at the origin on the x axis, carries a uniform line charge density λ. Find $\vec{E}$ at the point $P = (0, 0, z)$, a height z above its midpoint.*




Figure 2: Setup for the finite line of charge. The source point $\vec{r}' = x'\hat{x}$ runs from $-L$ to L; the field point is P at $\vec{r} = z\hat{z}$; the separation vector is $\vec{\mathfrak{r}} = \vec{r} - \vec{r}'$.

With $\vec{r} = z\hat{z}$ and $\vec{r}' = x'\hat{x}$,

$$\vec{\mathfrak{r}} = z\hat{z} - x'\hat{x}, \qquad \mathfrak{r} = \sqrt{z^2 + x'^2},$$

so

$$\vec{E} = \frac{\lambda}{4\pi\epsilon_0}\int_{-L}^{L} \frac{z\hat{z} - x'\hat{x}}{(z^2 + x'^2)^{3/2}}\,dx' = \frac{\lambda}{4\pi\epsilon_0}\left[\hat{z}\int_{-L}^{L}\frac{z}{(z^2 + x'^2)^{3/2}}dx' - \hat{x}\int_{-L}^{L}\frac{x'}{(z^2 + x'^2)^{3/2}}dx'\right].$$

The second integral vanishes by symmetry: its integrand is odd in x', so the pieces at x' and $-x'$ cancel. The first is done with the substitution $x' = z\tan\theta$, giving

$$\vec{E} = \frac{\lambda}{4\pi\epsilon_0}\,\frac{1}{z}\left[\frac{x'}{\sqrt{x'^2 + z^2}}\right]_{-L}^{L}\hat{z} = \frac{\lambda}{4\pi\epsilon_0}\,\frac{2L}{z\sqrt{L^2 + z^2}}\,\hat{z} = \frac{\lambda}{2\pi\epsilon_0 z}\cdot\frac{L}{\sqrt{L^2 + z^2}}\,\hat{z}.$$

This has two limits:

- *Close to the line ($z \ll L$), it looks like an infinite line:* $\vec{E} \to \dfrac{\lambda}{2\pi\epsilon_0 z}\hat{z}$.
- *Far from the line ($z \gg L$), it looks like a point charge:* $\vec{E} \to \dfrac{Q}{4\pi\epsilon_0 z^2}\hat{z}$ *with* $Q = 2\lambda L$.

3.2 Divergence and Curl of Electrostatic Fields

3.2.1 Gauss's Law

Law 2 (Gauss' law)**.** *The flux of $\vec{E}$ through any closed surface is set by the charge inside it:*

$$\oiint_S \vec{E} \cdot d\vec{a} = \frac{Q_{enc}}{\epsilon_0}. \tag{11}$$

Here Q_{enc} is the total charge of the object inside the gaussian surface S.

Proposition 8 (Differential form)**.** *Apply the divergence theorem (Theorem 4) to the left side and write $Q_{enc} = \iiint_V \rho\, d\tau$:*

$$\iiint_V (\vec{\nabla} \cdot \vec{E})\, d\tau = \frac{1}{\epsilon_0} \iiint_V \rho\, d\tau,$$

where ρ is the charge density. Since this holds for every volume V, the integrands match:

$$\vec{\nabla} \cdot \vec{E} = \frac{\rho}{\epsilon_0}. \tag{12}$$

Definition 24 (Gaussian Surface)**.** *The Gaussian surface just means an imaginary surface (nothing is physically there).*

Remark 13. *If there is no charge inside the Gaussian surface, the net flux is zero: every field line that enters on one side leaves on another.*

3.2.2 Applications of Gauss's Law

Gauss's law always holds but is only useful with symmetry: we need $\vec{E}$ parallel to $d\vec{a}$ and $|\vec{E}|$ constant over the Gaussian surface to pull $|\vec{E}|$ out of the integral. Only three symmetries work: spherical, cylindrical, and planar.

Spherical Symmetry.

Example (Uniform solid sphere)**.** *Let a uniform solid sphere of radius R and total charge q. Imagine a gaussian sphere of radius r, so*

$$\vec{E} \oiint d\vec{a} = \frac{Q_{enc}}{\epsilon_0},$$

let $Q_{enc} = q\,(r^3/R^3)$ *for* $r < R$ *and* $Q_{enc} = q$ *for* $r > R$. *The flux is* $\oiint d\vec{a} = 4\pi r^2$, *so*

$$\vec{E} = \begin{cases} \dfrac{q\,r}{4\pi\epsilon_0 R^3}\hat{r} & r < R \\ \frac{q}{4\pi\epsilon_0 R^2}\hat{r} & r = R \\ \dfrac{q}{4\pi\epsilon_0 r^2}\hat{r} & r > R \end{cases}.$$

Example. *A thin shell instead gives*

$$\vec{E} = \begin{cases} 0 & r < R \\ \frac{\sigma}{\epsilon_0}\hat{r} & r = R \\ \dfrac{q}{4\pi\epsilon_0 r^2}\hat{r} & r > R \end{cases}.$$

Remark 14. *A uniformly charged shell that pulses in and out,* $R(t) = R_0 + A\sin\omega t$, *keeps the constant field* $\frac{Q}{4\pi\epsilon_0 r^2}\hat{r}$ *outside: accelerating charges, but nothing escapes to infinity, so no radiation.*

Cylindrical Symmetry.

Example (Infinite line of charge)**.** *For a line with charge* λ *per unit length, take the Gaussian surface to be a coaxial cylinder of radius* z *(the distance from the line to the field point, not the radius of the line) and length* d. $\vec{E}$ *points radially outward, so* $\vec{E}\cdot d\vec{a} = 0$ *on the end caps and only the curved side contributes:*

$$E\,(2\pi z d) = \frac{\lambda d}{\epsilon_0} \implies E = \frac{\lambda}{2\pi\epsilon_0 z}.$$

Example (Finite versus infinite line)**.** *Near its midpoint, a finite line has a weaker field than an infinite one: the same enclosed charge, but some flux leaks through the caps.*

Problem 1 (Midterm 2025, Q2)**.** *An infinite non-conducting tube along the* x *axis has* $\rho(s) = A\,s/a$ *for* $s < a$ *and* 0 *elsewhere, with* s *the distance from the axis. Find* $\vec{E}$ *everywhere.*

Planar Symmetry.

Example (Infinite plane of charge)**.** *For a plane with surface charge density* σ, *take the Gaussian surface to be a pillbox straddling the plane, with caps of area* A. $\vec{E}$ *points away from the plane on both sides, so there is no flux through the side wall and* EA *through each cap:*

$$2EA = \frac{\sigma A}{\epsilon_0} \implies E = \frac{\sigma}{2\epsilon_0}.$$




Figure 3: Side view of the pillbox straddling the plane.

Example (Two planes)**.** *Superpose two sheets a distance d apart, each giving $\sigma/2\epsilon_0$, pointing away from a positive sheet and toward a negative one.*

- *$+\sigma$ and $-\sigma$ (parallel plates):*

$$E = \begin{cases} \frac{\sigma}{\epsilon_0} & \text{between the plates,} \\ 0 & \text{outside the plates.} \end{cases}$$

- *$+\sigma$ and $+\sigma$:*

$$E = \begin{cases} 0 & \text{between the plates,} \\ \frac{\sigma}{\epsilon_0} & \text{outside the plates.} \end{cases}$$

Remark 15 (Fall-off)**.** *The fall-off is how fast E decreases with the distance r from the source: $E \propto 1/r^2$ for a point charge, $1/r$ for an infinite line, and r^0 (uniform) for an infinite plane. To see why, picture a narrow cone opening from the field point toward the source. Its cross section at distance r grows like r^2, which alone gives the $1/r^2$ of a point charge, but for extended sources the charge inside the cone grows too: it intercepts a length $\propto r$ of a line, giving $E \propto r/r^2 = 1/r$, and an area $\propto r^2$ of a sheet, giving $E \propto r^2/r^2$, a constant.*

Remark 16. *For two equal positive charges, no field lines cross the midplane: the field there is parallel to the plane and vanishes at the midpoint. Gauss's law still holds, but without symmetry it cannot solve for $\vec{E}$.*

3.2.3 The Curl of $\vec{E}$

Theorem 9 (Electrostatic fields are curl-free)**.** *For any electrostatic field,*

$$\oint \vec{E} \cdot d\vec{l} = 0, \qquad \vec{\nabla} \times \vec{E} = 0. \tag{13}$$

Proof. Let there be a point charge at the origin. The field is radial, $\vec{E} = \frac{q}{4\pi\epsilon_0 r^2}\hat{r}$, so the line integral from some point a to some other point b in three dimensions

$\int_a^b \vec{E} \cdot d\vec{l}$, in spherical coordinates $d\vec{l} = dr\,\hat{r} + r\,d\theta\,\hat{\theta} + r\sin\theta\,d\phi\,\hat{\phi}$, we get

$$\vec{E} \cdot d\vec{l} = \frac{q}{4\pi\epsilon_0 r^2} dr.$$

Therefore,

$$\int_a^b \vec{E} \cdot d\vec{l} = \frac{q}{4\pi\epsilon_0} \int_{r_a}^{r_b} \frac{dr}{r^2} = \frac{q}{4\pi\epsilon_0} \left(\frac{1}{r_a} - \frac{1}{r_b} \right).$$

The integral around a closed path is evidently zero (for then $r_a = r_b$):

$$\oint \vec{E} \cdot d\vec{l} = 0,$$

and hence, by Stokes' theorem (6), $\vec{\nabla} \times \vec{E} = 0$. □

3.3 Electric Potential

Definition 25 (Electric potential)**.** *Since $\vec{\nabla} \times \vec{E} = 0$, the irrotational fields theorem 6 guarantees a scalar V with*

$$\vec{E} = -\vec{\nabla} V. \tag{14}$$

The minus sign is convention: positive charges are pushed toward lower potential energy.

Proposition 9 (Potential from the field)**.**

$$V(\vec{r}) = -\int_{ref}^{\vec{r}} \vec{E} \cdot d\vec{l} \left[\frac{J}{C} = volts\ (v) \right]. \tag{15}$$

The path does not matter since the integral is path-independent[2]*. By convention the reference is at infinity, unless the charge distribution extends to infinity.*

Proposition 10 (Potential of a Charge Distribution Extending to Infinity)**.** *If the charge distribution extends to infinity, the reference point for the potential is typically taken at a finite distance where the potential vanishes, for example a point on the plane.*

Definition 26 (Equipotential)**.** *An equipotential is a region, typically a surface, on which V is constant.*

Example (Point charge)**.** *Along a radial path,*

$$V(r) = -\frac{q}{4\pi\epsilon_0} \int_\infty^r \frac{dr'}{r'^2} = \frac{q}{4\pi\epsilon_0 r}.$$

[2] Changing the reference point adds a constant to V, which leaves $\vec{E} = -\vec{\nabla} V$ unchanged (gauge freedom). Statement (ii) of the irrotational fields theorem.

Example (Uniform solid sphere)**.** *Using the field of Example 3.2.2, outside* $V = \frac{q}{4\pi\epsilon_0 r}$*. Inside, integrate in from infinity through both regions:*

$$V(r) = -\int_{\infty}^{R} \frac{q\,dr'}{4\pi\epsilon_0 r'^2} - \int_{R}^{r} \frac{q\,r'\,dr'}{4\pi\epsilon_0 R^3} = \frac{q}{8\pi\epsilon_0 R}\left(3 - \frac{r^2}{R^2}\right).$$

Problem 2 (Midterm drills)**.** *Find* V *(i) at the centre of a ring of radius* a *and charge* Q*; (ii) at* $x = 2a$ *for a line of charge* Q *from* $x = -a$ *to* $x = a$*.*

3.3.1 Poisson's Equation and Laplace's Equation

Proposition 11 (Poisson's equation)**.** *Combining* $\vec{\nabla}\cdot\vec{E} = \rho/\epsilon_0$ *with* $\vec{E} = -\vec{\nabla}V$*,*

$$\nabla^2 V = -\frac{\rho}{\epsilon_0}. \tag{16}$$

Corollary 9.1 (Laplace's equation)**.** *In regions where there is no charge, so* $\rho = 0$*, Poisson's equation reduces to Laplace's equation:*

$$\nabla^2 V = 0. \tag{17}$$

3.3.2 The Potential of a Localized Charge Distribution

Proposition 12 (Potential of a distribution)**.** *By superposition* (7)*,*

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0}\int \frac{\rho(\vec{r}')}{\mathfrak{r}}\,d\tau', \tag{18}$$

where $\rho(\vec{r}')d\tau' = dq$*. Unlike the integral for* $\vec{E}$*, there is no* $\hat{\mathfrak{r}}$*, so computing* V *first and then* $\vec{E} = -\vec{\nabla}V$ *is often easier.*

3.3.3 Boundary Conditions

A surface charge σ makes $\vec{E}$ jump as you cross it. To fix this we use Gauss's law.




Figure 4: Side views of a charged surface.

Proposition 13 (Normal component)**.** *Take a pillbox of cap area A straddling the surface and shrink its thickness $\varepsilon \to 0$, so the side wall carries no flux. Only the caps count, and the charge inside is σA:*

$$E^{\perp}_{above}A - E^{\perp}_{below}A = \frac{\sigma A}{\epsilon_0} \quad \Longrightarrow \quad E^{\perp}_{above} - E^{\perp}_{below} = \frac{\sigma}{\epsilon_0}.$$

Proposition 14 (Tangential component)**.** *Take a thin rectangular loop of length l straddling the surface, again with $\varepsilon \to 0$ so the short sides contribute nothing. Since $\oint \vec{E} \cdot d\vec{l} = 0$,*

$$E^{\parallel}_{above}\, l - E^{\parallel}_{below}\, l = 0 \quad \Longrightarrow \quad \vec{E}^{\parallel}_{above} = \vec{E}^{\parallel}_{below}.$$

Theorem 10 (Boundary condition on $\vec{E}$)**.** *With $\hat{n}$ the unit normal pointing from below to above, the two results combine into*

$$\vec{E}_{above} - \vec{E}_{below} = \frac{\sigma}{\epsilon_0}\hat{n}. \tag{19}$$

The normal component jumps by σ/ϵ_0; the tangential component is continuous.

Remark 17. *Check with the infinite plane: $\vec{E} = +\frac{\sigma}{2\epsilon_0}\hat{n}$ above and $-\frac{\sigma}{2\epsilon_0}\hat{n}$ below, a jump of exactly σ/ϵ_0.*

Theorem 11 (Boundary conditions on V)**.** *The potential is continuous across any boundary, since*

$$V_{above} - V_{below} = -\int_{\vec{a}}^{\vec{b}} \vec{E} \cdot d\vec{l} \overset{\text{path length}\to 0}{\Longrightarrow} V_{above} = V_{below}.$$

However, the gradient of V is not continuous: substituting $\vec{E} = -\vec{\nabla}V$ into Theorem 10 gives

$$\vec{\nabla}V_{above} - \vec{\nabla}V_{below} = -\frac{\sigma}{\epsilon_0}\hat{n}. \tag{20}$$

3.4 Work and Energy in Electrostatics

Proposition 15 (Potential of a test charge)**.** *Bringing Q in from infinity against the field takes*

$$W = -Q\int_{\infty}^{\vec{r}} \vec{E} \cdot d\vec{l} = QV(\vec{r}). \tag{21}$$

3.4.1 The Energy of a Point Charge Distribution

Proposition 16 (Energy of N point charges)**.** *The first charge costs no work, since no field exists yet, and each later one feels the potential $V(\vec{r}_i)$ of those already placed, giving one term per pair:*

$$W = \frac{1}{2}\sum_{i=1}^{n} q_i \left(\sum_{j\neq i}^{n} \frac{q_j}{4\pi\epsilon_0 |\vec{r}_i - \vec{r}_j|} \right) = \frac{1}{2}\sum_{i=1}^{n} q_i V(\vec{r}_i), \tag{22}$$

where the $\frac{1}{2}$ corrects for counting each pair twice.

3.4.2 The Energy of a Continuous Charge Distribution

Theorem 12 (Energy stored in the field)**.** *For a volume charge distribution, the energy is*

$$W = \frac{1}{2}\int \rho V \, d\tau = \frac{\epsilon_0}{2}\int_{all\ space} E^2 \, d\tau. \tag{23}$$

3.4.3 Comments on Electrostatic Energy

Definition 27 (Energy Density)**.** *From Theorem 12, the energy per unit volume stored in the field is*

$$u = \frac{\epsilon_0}{2}E^2. \tag{24}$$

The proof is in Appendix B.

Problem 3 (Midterm drill)**.** *Two plates of area A, a distance d apart, carry $+\sigma$ and $-\sigma$ ($A \gg d^2$). Find the energy.*

Problem 4 (Problem Set 4, Q4)**.** *Concentric shells of radii $a < b$ carry $+q$ and $-q$. Find the energy.*

3.5 Conductors

Definition 28 (Insulator)**.** *An insulator, such as glass or ruber, each electron is on a short leash, attached to a particular atom.*

Definition 29 (Conductor)**.** *A conductor is a material with charges that are free to move (in a metal, the electrons).*

Proposition 17. *In an external field, a conductor's free charges move to its surface until their field $\vec{E}_{int}$ cancels $\vec{E}_{ext}$ inside. In practice, it's the negative charges (the electrons) that do the moving, leaving a positive charge on the far side.*




Figure 5: A conductor in an external field.

Proposition 18 (Properties of a conductor in equilibrium)**.**

(i) $\vec{E} = 0$ inside, otherwise the free charges would still be moving.

(ii) $\rho = 0$ *inside, since* $\rho = \epsilon_0 \vec{\nabla} \cdot \vec{E} = \epsilon_0 \cdot 0 = 0$.

(iii) Any net charge therefore sits on the surface.

(iv) A conductor is an equipotential: for any two points within (or at the surface of) a conductor, $V(\vec{b}) - V(\vec{a}) = -\int_{\vec{a}}^{\vec{b}} \vec{E} \cdot d\vec{l} = 0$.


(v) Just outside, $\vec{E}$ *is perpendicular to the surface. A tangential part would push the surface charges along the surface, and they would not be in equilibrium.*

Remark 18. *Grounding a conductor connects it to* $V(\infty) = 0$*, so its whole body, not just its surface, sits at* $V = 0$*. Positive charge flows to the ground if the conductor would otherwise be at* $V > 0$*, or from the ground if it would be at* $V < 0$*, until* $V = 0$.

3.5.1 Induced Charges

Remark 19 (Induced Charges)**.** *If a conductor has a cavity with a charge* q *inside,* q *induces* $-q$ *on the cavity wall and* $+q$ *on the outer surface (for a neutral conductor), and the outside cannot tell how the charge inside is arranged. In particular, for a sphere the outer charge spreads uniformly even if the cavity is off centre, so the field outside is that of a point charge at the centre of the sphere, not of the cavity.*

Remark 20 (Induced Charges with External Field)**.** *Same setup as the previous remark, but now add an external field: the charge on the cavity wall is still* $-q$ *and the outer surface still carries* $+q$*, but the external field induces a nonuniform distribution on the outer surface. The field inside the cavity is unchanged, since it is due only to* q *and the wall charge.*



Remark 21 (Faraday Cage)**.** *If a conductor has an empty cavity, the field inside the cavity is zero: its wall is an equipotential, and a solution of Laplace's equation with constant boundary values is constant (no maxima or minima, see the Laplace's equation section).*

3.5.2 Surface Charge and the Force on a Conductor

Proposition 19 (Field at the surface of a conductor)**.** *Theorem 10 with $\vec{E}_{below} = 0$ (inside the metal) gives, just outside,*

$$\vec{E} = \frac{\sigma}{\epsilon_0}\hat{n}.$$

Remark 22.

Example (Neutral conductor near a charged rod)**.** *A neutral metal block sits near an insulating rod with uniform positive charge. Negative charge is induced on the side facing the rod and an equal positive charge on the far side. The negative charges are closer, and Coulomb's force falls off with distance, so the attraction beats the repulsion: the neutral conductor is pulled toward the rod.*

Example (Conducting versus insulating disk)**.** *For a disk with uniform σ (an insulator), the disk is not an equipotential and the field lines are not perpendicular to it. That arrangement cannot last on a conductor: the field along the surface pushes charge outward, i.e. "downhill" in potential, and it piles up at the rim. So a charged conducting disk has a higher σ near its edge than near its centre, its surface is an equipotential, and its field lines leave it at right angles.*

Remark 23. *In general, charge on a conductor concentrates where the surface is most sharply curved (edges, corners, points). Since $E = \sigma/\epsilon_0$ there, the field is strongest at sharp points, which is how lightning rods work.*

Example (Concentric conducting shells)**.** *A metal sphere of radius R with charge q sits inside a thick neutral metal shell of inner radius a and outer radius b.*

(i) A Gaussian sphere inside the shell's metal ($a < r < b$) has $\vec{E} = 0$ on it, so it encloses zero charge: the inner surface carries $-q$. The shell is neutral, so the outer surface carries $+q$. By symmetry each is uniform:

$$\sigma_R = \frac{q}{4\pi R^2}, \qquad \sigma_a = -\frac{q}{4\pi a^2}, \qquad \sigma_b = \frac{q}{4\pi b^2}.$$

(ii) The field is $\frac{q}{4\pi\epsilon_0 r^2}\hat{r}$ for $r > b$ and $R < r < a$, and zero in both metals. Integrating in from infinity,

$$V(0) = -\int_{\infty}^{b} \frac{q\,dr}{4\pi\epsilon_0 r^2} - \int_{a}^{R} \frac{q\,dr}{4\pi\epsilon_0 r^2} = \frac{q}{4\pi\epsilon_0}\left(\frac{1}{b} + \frac{1}{R} - \frac{1}{a}\right).$$

(iii) Grounding the outer surface sets $V(b) = 0 = V(\infty)$, so there can be no field outside and $\sigma_b = 0$. The Gauss argument in (i) is unchanged, so σ_R and σ_a stay the same, and the first integral in (ii) disappears: $V(0) = \frac{q}{4\pi\epsilon_0}\left(\frac{1}{R} - \frac{1}{a}\right)$.

(iv) The outer surface now has zero charge; the $+q$ flowed away to the ground, leaving the shell with net charge $-q$. If the outer surface kept a charge Q', the total charge inside a sphere just outside b would be Q', giving $V(b) = \frac{Q'}{4\pi\epsilon_0 b} \neq 0$.

Problem 5 (Problem Set 4, Q3)**.** *A neutral conducting sphere of radius R has two spherical cavities, of radii a and b, with point charges q_a and q_b at their centres. (i) Find σ_a, σ_b, σ_R; (ii) the field outside; (iii) the field in each cavity; (iv) the force on q_a and q_b; (v) what changes if a charge q_c is brought near the outside?*

Problem 6 (Midterm 2025, Q3)**.** *A thick conducting shell (inner radius a, outer radius b, empty cavity) has net charge $-Q$. (a) Where is the charge? (b) Sketch $E_r(r)$. (c) Sketch $V(r)$ with $V(\infty) = 0$. (d) A charge $+q$ with $q > Q$ is placed outside: sketch the negative charge. (e) Find the total charge on the inner surface and (f) on the outer surface.*

4 Potentials

September 18-25, 2026

4.1 Laplace's Equation


We want V because $\vec{E} = -\vec{\nabla}V$. The integral $V = \frac{1}{4\pi\epsilon_0}\int \frac{\rho}{\mathfrak{r}}\, d\tau'$ is exact but often impossible to do, and the multipole expansion is only an approximation. Often we do not even know ρ everywhere: we know the potentials of some conductors instead, and the charge on them is whatever it has to be. In the empty space between the charges, Poisson's equation simplifies.

Law 3 (Laplace's equation)**.** *In a region with no charge,*

$$\nabla^2 V = \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0. \tag{25}$$

Remark 24. *Point charges, charged objects and conductors are where $\nabla^2 V = -\rho/\epsilon_0$ holds with $\rho \neq 0$. Everywhere else $\nabla^2 V = 0$, and the charges only enter as boundary conditions on that empty region.*

Remark 25 (The Laplacian compares a point with its surroundings)**.** *In one dimension, $V'' = 0$ gives a straight line, $V = mx + b$, and then $V(x) = \frac{1}{2}[V(x+a) + V(x-a)]$: every point is the average of its neighbours. In general $\nabla^2 V$ measures how far a point sits from the average of its surroundings, and since $\nabla^2 V = -\rho/\epsilon_0$, that tells you the sign of the charge there.*



below the average
$\nabla^2 V > 0$, so $\rho < 0$

equal to the average
$\nabla^2 V = 0$, no charge

above the average
$\nabla^2 V < 0$, so $\rho > 0$

Figure 6: The middle point compared with the average of its two neighbours (open circle).

Theorem 13 (Mean value property)**.** *If $\nabla^2 V = 0$, the value of V at any point equals its average over any sphere of radius R centred on that point (containing no charge):*

$$V(\vec{r}_0) = \frac{1}{4\pi R^2} \oint_{sphere} V \, da. \tag{26}$$

Corollary 13.1 (No local maxima or minima)**.** *A solution of Laplace's equation has no local maxima or minima inside the region; extreme values occur only on the boundary. A maximum would be larger than every point around it, so it could not equal their average.*

Theorem 14 (Earnshaw's theorem)**.** *A charged particle cannot be held in stable equilibrium by electrostatic forces alone.*

Proof. Stable equilibrium for a charge q needs its potential energy qV to have a local minimum at that point. But the other charges' V obeys Laplace's equation there, so it has no local minimum. Concretely, if $\frac{\partial^2 V}{\partial x^2} > 0$ (stable along x), then $\frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} < 0$, so the charge escapes sideways. For example, a charge between two equal charges is stable along the line joining them but slips out perpendicular to it. □

Theorem 15 (Uniqueness theorem)**.** *The solution of Laplace's equation in a volume is uniquely determined if V is specified on the whole boundary of that volume. Similarly, if the region contains conductors, $\vec{E}$ is uniquely determined once the charge density ρ and the total charge on each conductor are given.*

Remark 26. *The significance: any function that satisfies Laplace's equation and the boundary conditions is* the *answer, no matter how it was found. Guessing is allowed, which is what the method of images and separation of variables exploit.*

Proposition 20 (Method of relaxation)**.** *On a grid, the mean value property says each cell equals the average of its neighbours. In two dimensions,*

$$V_{i,j} = \frac{1}{4}\left(V_{i+1,j} + V_{i-1,j} + V_{i,j+1} + V_{i,j-1}\right).$$

This gives a numerical method:

1. *Initialize the grid with a guess for V.*
2. *Sweep through the grid, replacing each value by the average of its nearest neighbours, keeping the boundary values fixed.*
3. *Repeat until the values stop changing.*

4.2 The Method of Images

Remark 27 (A solution in search of a problem)**.** *Maxwell pointed out that the inverse problem is easier: given a potential, finding conductors that fit it is simpler than finding the potential for given conductors. Any equipotential surface of a known potential can be replaced by a conductor held at that potential, and by uniqueness the field outside it does not change.*

Example (Point charge above a grounded plane)**.** *A charge q sits at height d above an infinite grounded conducting plane at $z = 0$. For $z > 0$ we need $\nabla^2 V = 0$ (apart from at q), $V = 0$ on the plane, and $V \to 0$ far away. Forget the plane and put an image charge $-q$ at $z = -d$ instead. Every point of the plane is equidistant from the two charges, so this pair gives $V = 0$ there, and it has the same charge in the region $z > 0$. By uniqueness it is the answer for $z \geq 0$:*

$$V(x, y, z) = \frac{1}{4\pi\epsilon_0}\left[\frac{q}{\sqrt{x^2 + y^2 + (z-d)^2}} - \frac{q}{\sqrt{x^2 + y^2 + (z+d)^2}}\right].$$




Figure 7: The real problem (left) and the image problem (right) have the same charge and the same boundary values in $z > 0$, so they have the same potential there.

Remark 28. *The image charge is not real; it is bookkeeping. Physically, the field comes from the negative surface charge induced on the plane, and the image reproduces its effect in the region we care about. The formula says nothing about $z < 0$, where $V = 0$ inside the conductor. The image must sit outside the region being solved, otherwise it would change ρ there.*

Remark 29. *The induced charge on the plane totals $-q$, and q is attracted to the plane with the force of its image, but the energy is only half of the image pair's, since there is no field in $z < 0$.*

Example (Two charges above a grounded plane)**.** *Charges $+Q$ at height d and $+3Q$ at height $2d$ sit above a grounded plane. Each charge on its own is handled by one image, so by superposition we need two: $-Q$ at $-d$ and $-3Q$ at $-2d$. The method works whenever the problem is linear (superposition) and the boundary values come out right (uniqueness).*

4.3 Separation of Variables: Cartesian

Proposition 21 (Separation)**.** *Look for solutions $V(x,y,z) = X(x)Y(y)Z(z)$. Substituting into $\nabla^2 V = 0$ and dividing by V,*

$$\frac{X''}{X} + \frac{Y''}{Y} + \frac{Z''}{Z} = 0.$$

Each term depends on a different variable, which can be varied alone, so each must be a constant: $X'' = c_x X$, $Y'' = c_y Y$, $Z'' = c_z Z$ with $c_x + c_y + c_z = 0$. One PDE becomes three ODEs.

Remark 30. *Most solutions are not separable (e.g. $1/r$), but Laplace's equation is linear, so sums of separable solutions can build general ones.*

Proposition 22 (Three kinds of solution)**.** *For each direction,*

$$c = +k^2 :\ X = Ae^{kx} + Be^{-kx}, \qquad c = -k^2 :\ X = A\sin kx + B\cos kx, \qquad c = 0 :\ X = A + Bx.$$

Since the constants sum to zero, they cannot all have the same sign unless all vanish.

Remark 31 (Which direction gets which sign)**.** *A sinusoid crosses zero infinitely often; an exponential (or* cosh, sinh*) at most once. So a direction in which V must vanish at two places gets the sinusoid, and the other gets the exponential. Symmetry picks the parity: even $\Rightarrow$* cos, cosh*; odd $\Rightarrow$* sin, sinh*.* cosh *never vanishes;* sinh *vanishes only at* 0*.*

Example (Square pipe)**.** *A long pipe along z with square cross section: sides $x = \pm a/2$ grounded, top and bottom $y = \pm a/2$ held at V_0. Nothing depends on z, so $c_z = 0$ and $c_x = -c_y$. V vanishes at two values of x, so X is sinusoidal and Y hyperbolic; the problem is even in both x and y, so*

$$V = C\cos kx\,\cosh ky.$$

$V(\pm a/2, y) = 0$ quantises k: $\cos(ka/2) = 0 \Rightarrow k = (2n+1)\pi/a$. Superposing,

$$V(x,y) = \sum_{n=0}^{\infty} C_n \cos\frac{(2n+1)\pi x}{a}\cosh\frac{(2n+1)\pi y}{a}.$$

Setting $y = a/2$, multiplying by $\cos\frac{(2m+1)\pi x}{a}$ and integrating over x (Fourier's trick, using orthogonality $\int_{-a/2}^{a/2}\cos\cos\,dx = \frac{a}{2}\delta_{mn}$) picks out one term:

$$V(x,y) = \frac{4V_0}{\pi}\sum_{n=0}^{\infty}\frac{(-1)^n}{2n+1}\cos\frac{(2n+1)\pi x}{a}\,\frac{\cosh\left[(2n+1)\pi y/a\right]}{\cosh\left[(2n+1)\pi/2\right]}.$$

Check: rotating by $90°$ swaps hot and grounded faces, so $V(0,0) = V_0/2$ exactly; the series gives $0.5074\,V_0$ with one term. The centre is a saddle, as $\nabla^2 V = 0$ requires. Near the corners the series overshoots (Gibbs phenomenon), because the boundary data jumps there.

Remark 32 (Summary)**.** *1. Separate: one PDE becomes ODEs with constants summing to zero.*

2. *Choose signs: two or more zeros in a direction $\Rightarrow$ sinusoid; otherwise exponential/hyperbolic.*
3. *Use symmetry to keep only even or odd functions.*
4. *Impose the $V = 0$ boundaries: these quantise k.*
5. *Superpose and use Fourier's trick for the coefficients.*
6. *Uniqueness guarantees the result is the answer.*

4.4 Separation of Variables: Spherical

Proposition 23 (Axially symmetric solutions)**.** *If nothing depends on ϕ, separating $V(r,\theta) = R(r)\Theta(\theta)$ in the spherical Laplacian gives radial solutions r^l and $r^{-(l+1)}$ and angular solutions the Legendre polynomials $P_l(\cos\theta)$, so*

$$V(r,\theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos\theta), \tag{27}$$

with $P_0 = 1$, $P_1 = x$, $P_2 = (3x^2-1)/2$, and orthogonality $\int_0^\pi P_m P_n \sin\theta\, d\theta = \frac{2}{2m+1}\delta_{mn}$.

Proposition 24 (Sphere with given surface potential $V_0(\theta)$)**.**

- *Inside, V must be finite at $r = 0$, so $B_l = 0$.*
- *Outside, $V \to 0$ at infinity, so $A_l = 0$.*
- *Write $V_0(\theta) = \sum_l C_l P_l(\cos\theta)$ (by inspection if it is a simple polynomial in $\cos\theta$, else by orthogonality). Continuity at R then gives*

$$V_{in} = \sum_l C_l \left(\frac{r}{R}\right)^l P_l(\cos\theta), \qquad V_{out} = \sum_l C_l \left(\frac{R}{r}\right)^{l+1} P_l(\cos\theta).$$

- *The surface charge comes from the jump in slope, $\sigma = -\epsilon_0 \left(\frac{\partial V_{out}}{\partial r} - \frac{\partial V_{in}}{\partial r} \right)_{r=R}$, giving*

$$\sigma(\theta) = \frac{\epsilon_0}{R} \sum_l (2l+1) C_l P_l(\cos\theta).$$

Remark 33. *Rewrite powers of $\cos\theta$ in Legendre polynomials, e.g. $\cos^2\theta = \frac{2}{3}P_2 + \frac{1}{3}P_0$. Each l outside is a multipole: $l = 0$ gives $V \propto 1/r$ with total charge $Q = 4\pi\epsilon_0 R C_0$.*

Problem 7 (Midterm 2025, Q5)**.** *A spherical shell of radius R has $V(R,\theta) = V_0(1+\cos\theta)$, no charge elsewhere, and $V(\infty) = 0$. Find V outside, V inside, and $\sigma(\theta)$.*

Solution 1. *$V_0(1+\cos\theta) = V_0P_0 + V_0P_1$, so $C_0 = C_1 = V_0$ and all others vanish:*

$$V_{out} = V_0\left(\frac{R}{r} + \frac{R^2}{r^2}\cos\theta\right), \qquad V_{in} = V_0\left(1 + \frac{r}{R}\cos\theta\right),$$

$$\sigma(\theta) = \frac{\epsilon_0 V_0}{R}\left(1 + 3\cos\theta\right).$$

Check: $V_{in} = V_0 + V_0 z/R$ is a uniform field inside, and V_{out} is a monopole plus a dipole.

4.5 Multipole Expansion

Example (Two charges on the z axis)**.** *Place q_1 at $z = d/2$ and q_2 at $z = -d/2$, and measure θ from the z axis.*




Figure 8: Two charges on the z axis and a far field point P at distance r and angle θ.

Then

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0}\left(\frac{q_1}{\mathfrak{r}_1} + \frac{q_2}{\mathfrak{r}_2}\right).$$

To get $\mathfrak{r}_1$, write $\vec{\mathfrak{r}}_1 = \vec{r} - \frac{d}{2}\hat{z}$ and square it, using $\vec{r}\cdot\hat{z} = r\cos\theta$ (this is just the law of cosines):

$$\mathfrak{r}_1^2 = r^2 + \left(\frac{d}{2}\right)^2 - 2r\frac{d}{2}\cos\theta = r^2 + \left(\frac{d}{2}\right)^2 - rd\cos\theta.$$

For q_2 the source sits at $-\frac{d}{2}\hat{z}$, so the cross term flips sign. Both cases in one line, top sign for q_1 and bottom sign for q_2:

$$\mathfrak{r}_{1,2}^2 = r^2 + \left(\frac{d}{2}\right)^2 \mp rd\cos\theta.$$

Writing $1/\mathfrak{r}_{1,2} = \frac{1}{r}(1+\epsilon_\mp)^{-1/2}$ with $\epsilon_\mp = \frac{d}{2r}\left(\frac{d}{2r} \mp 2\cos\theta\right)$ and expanding for $d \ll r$, $(1+\epsilon)^{-1/2} = 1 - \frac{1}{2}\epsilon + \frac{3}{8}\epsilon^2 - \ldots$, gives

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0}\left[\underbrace{\frac{q_1+q_2}{r}}_{monopole} + \underbrace{\frac{(q_1-q_2)d\cos\theta}{2r^2}}_{dipole} + \underbrace{\frac{(q_1+q_2)d^2}{4}\frac{3\cos^2\theta-1}{2r^3}}_{quadrupole} + \ldots\right].$$

Each term gains a power of $1/r$, so $E \sim 1/r^2,\ 1/r^3,\ 1/r^4, \ldots$

Remark 34. *Setting $q_1 = -q_2$ kills the monopole term, leaving $V \sim 1/r^2$ and $E \sim 1/r^3$ at leading order.*

Definition 30 (Dipole Moment)**.** *For charges $+q$ and $-q$ separated by $\vec{d}$ (pointing from $-q$ to $+q$), the dipole moment is*

$$\vec{p} = q\vec{d}, \qquad [\vec{p}] = \mathrm{C}\cdot\mathrm{m}. \tag{28}$$

For any distribution,

$$\vec{p} = \sum_i q_i\vec{r}_i = \int \vec{r}'\,\rho(\vec{r}')\,d\tau', \tag{29}$$

which reduces to $q\vec{d}$ for the pair. It measures how far apart the positive and negative charge sit, and sets the dipole term of the potential,

$$V_{dip} = \frac{1}{4\pi\epsilon_0}\frac{\vec{p}\cdot\hat{r}}{r^2}. \tag{30}$$

Proposition 25 (Dipole field)**.** *For $\vec{p} = p\hat{z}$, $V_{dip} = \frac{p\cos\theta}{4\pi\epsilon_0 r^2}$, and $\vec{E} = -\vec{\nabla}V$ in spherical coordinates gives*

$$\vec{E}_{dip}(r,\theta) = \frac{p}{4\pi\epsilon_0 r^3}\left(2\cos\theta\,\hat{r} + \sin\theta\,\hat{\theta}\right). \tag{31}$$

Definition 31 (Ideal dipole)**.** *Take $q_1 = q$ and $q_2 = -q$. The monopole and quadrupole terms vanish, and the dipole term is $\frac{1}{4\pi\epsilon_0}\frac{qd\cos\theta}{r^2}$. Every term in the expansion carries a factor $q\,d^{\,n}$, which we can write as $(qd)\,d^{\,n-1}$. Now shrink $d \to 0$ while raising $q \to \infty$ so that the product qd stays fixed:*

- *The dipole term ($n = 1$) contains only qd, so it is unchanged.*
- *Every later term carries at least one extra factor of d, so it goes to zero.*

(q must grow, otherwise the dipole term would vanish too.) What is left is a point object with dipole moment $\vec{p} = q\vec{d}$ and exactly $V = \frac{1}{4\pi\epsilon_0}\frac{\vec{p}\cdot\hat{r}}{r^2}$.

Remark 35. *The names count charges in powers of 2: monopole (1), dipole (2), quadrupole (4), octupole (8), and so on. Each is built from two opposite copies of the previous one, placed close together so the previous order cancels. For example, two opposite dipoles give a quadrupole with $V \sim 1/r^3$.*

Theorem 16 (Multipole expansion)**.** *For r large compared with the size of the distribution,*

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0}\left(\frac{Q}{r} + \frac{\vec{p}\cdot\hat{r}}{r^2} + \frac{\sum_{i,j}\hat{r}_i Q_{ij}\hat{r}_j}{r^3} + \dots\right), \tag{32}$$

with

$$Q = \int \rho(\vec{r}')\,d\tau', \qquad \vec{p} = \int \vec{r}'\rho(\vec{r}')\,d\tau', \qquad Q_{ij} = \int \frac{1}{2}\left(3r'_i r'_j - r'^2\delta_{ij}\right)\rho(\vec{r}')\,d\tau'.$$

Griffiths writes the same series with Legendre polynomials; see Appendix 40.

Remark 36. *Each term is smaller than the previous one by a factor of roughly d/r, where d is the size of the charge distribution. Far away ($r \gg d$) the first nonzero term dominates and the rest are small corrections. Close up ($r \sim d$) the terms no longer shrink and the expansion is useless. So to find the leading behaviour: compute Q; if it vanishes, compute $\vec{p}$; if that vanishes too, the leading term is the quadrupole.*

Remark 37 (Origin dependence)**.** *Shifting the origin by $\vec{a}$ changes $\vec{p} \to \vec{p} - Q\vec{a}$, so $\vec{p}$ is well defined only when $Q = 0$ (a neutral molecule, not an ion). In general only the first non-vanishing moment is origin-independent.*

Problem 8 (Midterm drill)**.** *Charges $3q$ at $a\hat{z}$, q at $-a\hat{z}$, and $-2q$ at $\pm a\hat{y}$. (i) Sketch how to find the energy. (ii) Find V to dipole order.*

Solution 2. *(i) Sum $\frac{q_i q_j}{4\pi\epsilon_0 \mathfrak{r}_{ij}}$ over the 6 pairs: the two z charges are $2a$ apart, as are the two y charges, and each z charge is $\sqrt{2}a$ from each y charge:*

$$U = \frac{1}{4\pi\epsilon_0 a}\left[\frac{3q^2}{2} + \frac{4q^2}{2} + \frac{2(3q)(-2q) + 2(q)(-2q)}{\sqrt{2}}\right] = \frac{q^2}{4\pi\epsilon_0 a}\left(\frac{7}{2} - 8\sqrt{2}\right).$$

(ii) $Q = 3q + q - 4q = 0$, and $\vec{p} = 3qa\hat{z} - qa\hat{z} - 2qa\hat{y} + 2qa\hat{y} = 2qa\hat{z}$, so

$$V \approx \frac{1}{4\pi\epsilon_0}\frac{2qa\cos\theta}{r^2}.$$

Problem 9 (Midterm 2025, Q4)**.** *Charges $+q$ and $-q$, a distance a apart, sit a height d above a conductor filling $z < 0$. (a) Find $\vec{E}$ and V for $z < 0$. (b) Describe how to find the exact V for $z > 0$. (c) Find how V and $|\vec{E}|$ fall off for $r \gg d$.*

Solution 3. *(a)* $\vec{E} = 0$ *inside the conductor, so* V *is constant, independent of* z *(*$V = 0$*, matching the plane at infinity).*

(b) Method of images: place $-q$ *at depth* d *directly below* $+q$*, and* $+q$ *directly below* $-q$*. The four charges give* $V = 0$ *on* $z = 0$*, and the images lie outside* $z > 0$*, so by uniqueness*

$$V = \frac{1}{4\pi\epsilon_0} \sum_{i=1}^{4} \frac{q_i}{|\vec{r} - \vec{r}_i|} \quad \text{for } z > 0.$$

(c) Total charge is 0*. The real pair has* $\vec{p} = qa\hat{x}$ *and the image pair* $-qa\hat{x}$*, so the dipole moment also cancels. The leading term is the quadrupole:* $V \sim 1/r^3$ *and* $|\vec{E}| \sim 1/r^4$*.*

5 Electric Fields in Matter

6 Magnetostatics

6.1 Current Density

Definition 32 (Current density)**.** *The current* I *through a wire is not a local quantity, since a wire has finite thickness. The local version is the current density*

$$\vec{J} = \frac{d\vec{I}}{da_\perp},$$

where $da_\perp$ *is the differential cross section perpendicular to the flow.*

Remark 38. *Applying the divergence theorem (Theorem 4) to* $\vec{J}$ *is how conservation of charge gets written down: the charge inside* V *can only decrease by flowing out through* S*,*

$$\frac{d}{dt} \iiint_V \rho \, dV = -\oiint_S \vec{J} \cdot d\vec{A} = -\iiint_V \vec{\nabla} \cdot \vec{J} \, dV \quad \Longrightarrow \quad \frac{\partial \rho}{\partial t} = -\vec{\nabla} \cdot \vec{J}.$$

7 Magnetic Fields in Matter

8 Electrodynamics

A Extra Material

Remark 39 (Compactified dimensions)**.** *Suppose an extra dimension exists but is curled up, the way a flat sheet with coordinates (x, y) can be rolled into a cylinder whose second coordinate w is periodic with a small radius. At distances much smaller than that radius, field lines spread out into all the dimensions and the field falls off faster than $1/r^2$. At distances much larger than it, there is only room to spread out in the large dimensions and the usual $1/r^2$ is recovered. So a measured deviation from the inverse square law at short range would be evidence for extra dimensions.*

This has been tested for gravity by the Eöt-Wash torsion balance experiments (2004, 2007, 2020), which checked the $1/r^2$ law down to separations of $52\,\mu$m. Newtonian gravity fit the data, ruling out gravitational strength Yukawa interactions with ranges above $38.6\,\mu$m at 95% confidence.

Remark 40. *Griffiths' form of the multipole expansion, with α the angle between $\vec{r}$ and $\vec{r}'$ and P_n the Legendre polynomials:*

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0}\sum_{n=0}^{\infty}\frac{1}{r^{n+1}}\int (r')^n P_n(\cos\alpha)\rho(\vec{r}')\,d\tau'.$$

$P_0 = 1$ gives the monopole, $P_1 = \cos\alpha$ the dipole (since $r'\cos\alpha = \vec{r}'\cdot\hat{r}$), and $P_2 = \frac{1}{2}(3\cos^2\alpha - 1)$ the quadrupole.

Remark 41 (Self-energy)**.** *The energy of a charge configuration can be written as a pair sum or as a field integral,*

$$W_{pair} = \frac{1}{2}\frac{1}{4\pi\epsilon_0}\sum_{i\neq j}\frac{q_i q_j}{\mathfrak{r}_{ij}}, \qquad W_{field} = \frac{\epsilon_0}{2}\int E^2\,d\tau,$$

but the two do not always agree. The pair sum only counts the work of bringing different charges together ($i \neq j$); it ignores the work needed to build each point charge itself. The field integral counts everything, including that self-energy. So W_{field} is always positive (since $E^2 \geq 0$), while W_{pair} can be negative, e.g. $+q$ and $-q$ at distance d give $W_{pair} = -\frac{q^2}{4\pi\epsilon_0 d}$. For a true point charge the self-energy is infinite: near it $E^2 \sim 1/r^4$, and $\int \frac{1}{r^4}\,r^2\,dr$ blows up at $r = 0$. So use the pair sum for point charges and the field integral for continuous distributions.

B Proofs

Proof. The curl of a curl. Set $\vec{G} \equiv \vec{\nabla}\times(\vec{\nabla}\times\vec{F})$. Then,

$$G_i = \sum_{j,k}\varepsilon_{ijk}\partial_j\sum_{m,n}\varepsilon_{kmn}\partial_m F_n = \sum_{j,k,m,n}\varepsilon_{ijk}\varepsilon_{kmn}\partial_j\partial_m F_n.$$

Using the cyclic permutation $\varepsilon_{ijk} = \varepsilon_{kij}$, so that the summed index k sits first in both epsilons, the epsilon-delta identity 5 gives

$$\sum_k \varepsilon_{kij}\varepsilon_{kmn} = \delta_{im}\delta_{jn} - \delta_{in}\delta_{jm}.$$

Reading off the proposition with the summed index k removed:

$$G_i = \sum_{j,m,n} (\delta_{im}\delta_{jn} - \delta_{in}\delta_{jm})\, \partial_j \partial_m F_n.$$

Each Kronecker delta collapses one sum by setting its indices equal. In the first term δ_{im} forces $m = i$ and δ_{jn} forces $n = j$; in the second, δ_{in} forces $n = i$ and δ_{jm} forces $m = j$:

$$G_i = \sum_j \partial_j \partial_i F_j - \sum_j \partial_j \partial_j F_i \overset{3}{=} \partial_i \sum_j \partial_j F_j - \sum_j \partial_j \partial_j F_i = \left(\vec{\nabla}(\vec{\nabla}\cdot\vec{F})\right)_i - (\nabla^2 \vec{F})_i.$$

$$\therefore \vec{\nabla} \times (\vec{\nabla} \times \vec{F}) = \vec{\nabla}(\vec{\nabla}\cdot\vec{F}) - \nabla^2 \vec{F}.$$

□

Proof. Curl of a gradient. Let f be twice continuously differentiable. With $(\vec{\nabla} f)_k = \partial_k f$,

$$\left(\vec{\nabla} \times (\vec{\nabla} f)\right)_i = \sum_{j,k} \varepsilon_{ijk} \partial_j \partial_k f.$$

The indices j and k are summed over, so they can be renamed $j \leftrightarrow k$:

$$\left(\vec{\nabla} \times (\vec{\nabla} f)\right)_i = \sum_{j,k} \varepsilon_{ikj} \partial_k \partial_j f.$$

Now use antisymmetry, $\varepsilon_{ikj} = -\varepsilon_{ijk}$, and the fact that partial derivatives commute, $\partial_k \partial_j f = \partial_j \partial_k f$:

$$\left(\vec{\nabla} \times (\vec{\nabla} f)\right)_i = -\sum_{j,k} \varepsilon_{ijk} \partial_j \partial_k f = -\left(\vec{\nabla} \times (\vec{\nabla} f)\right)_i.$$

A number equal to its own negative is zero, so every component vanishes and $\vec{\nabla} \times (\vec{\nabla} f) = \vec{0}$. □

Proof. Using the energy of N point charges, $U = \frac{1}{2}\sum_i q_i V(\vec{r}_i)$. For a continuous distribution this becomes $U = \frac{1}{2}\int \rho V \, d\tau$. Substituting $\rho = -\epsilon_0 \nabla^2 V$ from Poisson's equation,

$$U = -\frac{\epsilon_0}{2} \int V \nabla^2 V \, d\tau.$$

To rewrite $V\nabla^2 V$, use the product rule $\partial_i(fg) = (\partial_i f)g + f(\partial_i g)$ with $f = V$ and $g = \partial_i V$, summed over i:

$$\underbrace{\sum_i \partial_i(V\partial_i V)}_{\vec{\nabla}\cdot(V\vec{\nabla}V)} = \underbrace{\sum_i (\partial_i V)^2}_{|\vec{\nabla}V|^2} + \underbrace{V\sum_i \partial_i\partial_i V}_{V\nabla^2 V} \quad \Longrightarrow \quad V\nabla^2 V = \vec{\nabla}\cdot(V\vec{\nabla}V) - |\vec{\nabla}V|^2.$$

[3] This equality uses $\partial_j\partial_i = \partial_i\partial_j$, valid for $\vec{F}$ twice continuously differentiable.

Substituting, the overall minus sign flips both terms:

$$U = -\frac{\epsilon_0}{2}\int \left[\vec{\nabla}\cdot(V\vec{\nabla}V) - |\vec{\nabla}V|^2\right] d\tau = \frac{\epsilon_0}{2}\int |\vec{\nabla}V|^2\, d\tau - \frac{\epsilon_0}{2}\int \vec{\nabla}\cdot(V\vec{\nabla}V)\, d\tau.$$

The second integral vanishes. To see this, integrate over a huge sphere of radius R instead of all space, and let $R \to \infty$ at the end. The divergence theorem turns the integral into a flux through the sphere's surface, $\oint V\vec{\nabla}V \cdot d\vec{a}$. If the charge sits in a finite region, then far away $V \sim 1/R$ and $|\vec{\nabla}V| \sim 1/R^2$, so the integrand is $\sim 1/R^3$, while the area is $4\pi R^2$. The flux is therefore $\sim 1/R$, which goes to zero. With $\vec{E} = -\vec{\nabla}V$, $U = \frac{\epsilon_0}{2}\int E^2\, d\tau$. □

C Useful Links

- Laplace relaxation demo:
  https://claude.ai/artifact/J2wPx8V3ajec8vDLsARnKX
- Electrostatics simulation:
  https://icphysweb.z13.web.core.windows.net/simulation.html