

LIPID PANEL QUICK REFERENCE

Reading Your Lipid Results

Test	Desirable	Borderline	High / Low Risk	What It Means
Total Cholesterol	<200	200–239	≥240 mg/dL	Overall screening value — always interpret with LDL & HDL
LDL-Cholesterol	<100	100–129	≥130 mg/dL (very high risk: <55–70)	Primary "bad" cholesterol — main target of diet and medication
HDL-Cholesterol	≥60	40–59 M / 50–59 F	<40 M / <50 F	Protective "good" cholesterol — higher is better
Triglycerides (TG)	<150	150–199	≥200 mg/dL (≥500 = pancreatitis risk)	Reflects sugar, white rice, alcohol, and processed food intake
Non-HDL-C (TC–HDL)	<130	130–159	≥160 mg/dL	Includes all atherogenic particles — often more predictive than LDL alone
TG / HDL Ratio	<3.5	3.5–5.0	>5.0	High ratio = insulin resistance + small dense LDL (most dangerous type)

DIETARY FAT TYPES AND THEIR EFFECTS

How Different Fats Move Your Lipid Panel

Fat Type	LDL-C	HDL-C	Triglycerides	Common Philippine Sources
Saturated Fat (SFA)	↑↑	—	—	Coconut oil (VCO, Minola), lard, butter, gata (coconut cream), chicharon, fatty pork (liempo, lechon), full-fat cheese
Trans Fat (TFA) — worst	↑↑	↓↓	—	Partially hydrogenated margarine, some imported biscuits & cookies, commercial fried food — avoid entirely
Monounsaturated (MUFA)	↓	—	—	Olive oil, canola oil, avocado, pili nuts, cashews, almonds
Omega-3 PUFA (EPA/DHA)	—	↑ slightly	↓↓	Sardinas (lata), galunggong, bangus, salmon — aim for 2 servings/week
Soluble Fiber	↓	—	—	Oatmeal, monggo, okra, kamote, psyllium husk — traps bile acids in the gut, 5–10 g/day lowers LDL ~5–11 mg/dL
Sugar & Refined Carbs	—	↓	↑↑	White rice (large portions), softdrinks, sweetened juices, instant noodles, white bread, pastries — biggest TG driver in Filipino diet
Alcohol	—	—	↑↑	Beer, gin, lambanog, wine — even moderate use raises TG significantly; stopping alcohol often drops TG 30–50% within weeks

HOW TO READ THE HEAT MAP (PAGES 2–3)

Legend

ROW COLOR = Overall Heart Rating

Avoid — high cardiovascular risk **Limit** — okay occasionally, small portion
 Good — eat freely, heart-healthy

HEAT CELLS = Nutrient Impact per Serving

H = High (harmful) **M** = Moderate **L** = Low (safe)

Columns: **SF** = Saturated Fat | **SUG** = Sugar / Refined Carb (TG driver) | **CHOL** = Dietary Cholesterol

 Coconut Oil (VCO, Minola) — the most common hidden risk in Filipino cooking

Coconut oil and gata (coconut cream) are 92% saturated fat — the highest of any cooking fat. A single serving of kare-kare or laing can contain 15–25 g of saturated fat. Replace with **canola oil or olive oil** for everyday cooking. Occasional use in traditional recipes is acceptable — but not as a daily staple.

Educational use only. This handout does not replace individualized medical advice. Consult your physician for your personal LDL target based on your cardiovascular risk profile. References: ACC/AHA 2025 Lipid Guidelines · AHA 2021 Dietary Fats Advisory · KDIGO 2024 Lipids in CKD.

williamriveromd.com
cholesterol-diet-guide.html

Food Item (Filipino Name)	Typical Serving	SF	SUG	CHOL	Rating	Key Note
A · LUTONG-BAHAY — PORK & BEEF DISHES						
Lechon baboy (whole roasted pig)	1 slice 100g	H	L	H	AVOID	Skin = pure saturated fat; organ stuffing raises dietary cholesterol further
Crispy pata (deep-fried pork knuckle)	1 piece 150g	H	L	H	AVOID	Pork skin + knuckle cartilage = very high saturated fat and dietary cholesterol
Chicharon bulaklak (pork mesentery, fried)	50 g	H	L	H	AVOID	Organ membrane deep-fried in lard; among the highest sat-fat snacks
Dinuguan (pork blood stew)	1 cup	H	L	H	AVOID	Pork blood + innards in pork fat broth; very high in cholesterol and sat fat
Laing (taro leaves in coconut cream)	1 cup	H	L	L	AVOID	Coconut cream = 15–25 g saturated fat per serving; use reduced amount of light gata
Bicol express (pork + gata + chili)	1 cup	H	L	M	AVOID	Double fat load: pork belly + thick coconut cream sauce
Kare-kare + bagoong (peanut stew)	1 bowl	H	M	M	AVOID	Peanut sauce + oxtail fat; bagoong adds very high sodium on top
Longganisa Vigan / sweet (cured fatty pork)	2 links 80g	H	M	M	AVOID	High sat fat + high sodium + sugar; nitrite preservatives add extra risk
Paksiw na lechon / lechon sauce	1 cup	H	M	H	AVOID	Leftover lechon fat recooked in liver-rich sauce; double cholesterol hit
Adobo (pork belly, with skin)	1 cup	H	L	M	LIMIT	Use pork loin instead of belly; remove skin; drain fat before serving
Bulalo (bone marrow beef soup)	1 bowl	H	L	H	LIMIT	Bone marrow is very high cholesterol; eat meat, discard broth and marrow
Sinigang na baboy (pork)	1 bowl	M	L	M	LIMIT	Use lean pork (kasim), skip broth (purines leach in); sinigang na isda is a better swap
Menudo / Afritada / Caldereta (pork)	1 cup	M	M	M	LIMIT	Use lean pork; liver spread in caldereta adds dietary cholesterol — reduce amount
Nilaga (pork) — meat + broth	1 bowl	M	L	M	LIMIT	Eat the lean meat; skip broth and potato; substitute beef brisket with chicken
Tocino (sweet-cured pork belly)	2 slices 80g	H	M	M	LIMIT	High sat fat + added sugar; prefer homemade lean pork versions; occasional only
Arroz caldo (pork or chicken rice porridge)	1 bowl	L	H	L	LIMIT	Low fat but high refined carbs (TG risk); use brown rice; top with tokwa instead of pork
Tinola (chicken + malunggay/sayote)	1 bowl	L	L	L	GOOD	Remove skin; eat chicken + vegetables; limit broth (sodium); excellent everyday meal
Adobo sa puti (chicken, no skin, no liver)	1 cup	L	L	L	GOOD	Lean chicken breast adobo; vinegar-based cooking preserves flavor without extra fat
Nilagang manok (boiled chicken)	1 bowl	L	L	L	GOOD	Excellent — remove skin; eat meat and vegetables; discard broth to reduce sodium
B · ISDA AT PAGKAIN SA DAGAT — FISH & SEAFOOD						
Crispy fried bangus (whole, with skin)	1 small	M	L	M	AVOID	Frying in coconut oil or lard adds large sat fat load; eat grilled or steamed instead
Sugpo / hipon (shrimp)	8 pcs medium	L	L	H	LIMIT	Low sat fat but high dietary cholesterol (~150mg/serving); limit to 1–2x/week
Pusit (squid, grilled or adobo)	1 cup	L	L	M	LIMIT	Moderate dietary cholesterol; avoid deep-fried tempura version

Talaba (oysters, raw or grilled)	6 pieces	L	L	M	LIMIT	Rich in zinc and omega-3; moderate cholesterol content; avoid fried versions
Sinigang na bangus / tilapia / labahita	1 bowl	L	L	L	GOOD	Best Filipino soup — fish + vegetables; limit broth intake for sodium
Inihaw na bangus (grilled milkfish, no skin)	1 medium	L	L	L	GOOD	Excellent omega-3 source; remove skin before eating to cut sat fat further
Inihaw na galunggong (round scad)	2–3 pieces	L	L	L	GOOD	Best budget omega-3 fish in PH markets; grill or steam; avoid frying
Sardinas sa kamatis (canned sardines, tomato)	1 can	L	L	L	GOOD	Affordable EPA/DHA omega-3; choose in tomato sauce or water, NOT oil
Tilapia (steamed or grilled)	1 medium	L	L	L	GOOD	Lean white fish; mild omega-3; affordable; cook without coconut oil
Tuyo (dried herring, grilled not fried)	2 pieces	L	L	L	GOOD	Good omega-3 but very high sodium; limit to 1–2x/week; do NOT fry in extra oil
C · GULAY AT LEGUMES — VEGETABLES & LEGUMES (EAT FREELY)						
Monggo guisado (mung bean stew)	1 cup	L	L	L	GOOD	Best soluble fiber staple; LDL-lowering; use canola oil, not lard, for sautéing
Pinakbet (no bagoong, canola oil)	1 cup	L	L	L	GOOD	Okra, ampalaya, eggplant — all fiber-rich and LDL-lowering; skip the bagoong
Ginisang kangkong (sautéed water spinach)	1 cup	L	L	L	GOOD	Use canola oil + garlic; add tokwa for protein; affordable heart-healthy staple
Ampalaya con huevo (bitter melon + egg)	1 cup	L	L	M	GOOD	Ampalaya lowers LDL and blood sugar; egg adds moderate dietary cholesterol — limit to 1 egg
Lumpiang sariwa (fresh spring rolls)	2 rolls	L	L	L	GOOD	Excellent — vegetables + tokwa wrapped; avoid the fried lumpia version
D · STREET FOOD — PAGKAIN SA KALYE						
Chicharon (pork rinds, bagged)	1 bag 40g	H	L	H	AVOID	12–15 g sat fat per small bag; most popular but among the worst snacks for LDL
Tokneneng / kwek-kwek (battered fried eggs)	3 pieces	M	L	H	AVOID	Egg yolk cholesterol + deep-fried batter in recycled street food oil
Balut (duck embryo)	1 piece	M	L	H	AVOID	High dietary cholesterol (~600mg) from developing embryo yolk; limit strictly if LDL elevated
Isaw (grilled pig/chicken intestines)	5 sticks	H	L	H	AVOID	Organ meat high in sat fat and cholesterol; grilling doesn't reduce fat content
Betamax (coagulated pork/chicken blood)	3 cubes	M	L	H	AVOID	Blood products high in dietary cholesterol; frequently grilled in lard basting
Fishball (commercial, deep-fried)	5 pieces	M	M	M	LIMIT	Processed fish product with fillers; fried in frequently reused oil; sweet sauce adds sugar
Kikiam (Chinese sausage roll, fried)	3 pieces	M	L	M	LIMIT	Processed meat product; moderate sat fat; avoid dipping in bagoong-based sauce
Banana cue (fried saba + brown sugar)	3 pieces	L	H	L	LIMIT	Low fat but high sugar → raises TG; try boiled saba instead
Turon (fried banana + jackfruit roll)	2 pieces	M	H	L	LIMIT	Sugar + lard frying wrapper; occasional treat; baked turon is a better option
Pork siomai (steamed)	4 pieces	M	L	M	LIMIT	Steamed is better than fried; moderate fat from ground pork; limit calamansi-soy dip
Boiled mais (corn on cob, no butter)	1 piece	L	L	L	GOOD	Whole grain fiber; filling snack; do NOT add butter or margarine
Boiled kamote (sweet potato, no sugar)	1 medium	L	L	L	GOOD	Soluble fiber + low glycemic index; replaces white rice snack; excellent LDL-lowering

E · KAKANIN AT MATAMIS — FILIPINO SWEETS & DESSERTS

Leche flan	1 slice 100g	H	H	H	AVOID	Triple threat: egg yolks (cholesterol) + condensed milk (sat fat + sugar) — all three cells red
Biko / sinukmani (glutinous rice + gata)	1 piece	H	H	L	AVOID	Coconut cream + refined glutinous rice + sugar = high sat fat AND high TG
Maja blanca (coconut milk pudding)	1 square	H	H	L	AVOID	Thick coconut cream + condensed milk + cornstarch; high sat fat and sugar
Polvoron (lard shortbread)	2 pieces	H	H	M	AVOID	Traditionally made with lard + condensed milk; some brands use vegetable shortening (check label)
Ensaymada (butter + cream cheese)	1 large	H	M	M	AVOID	Butter filling + cream cheese topping = high sat fat; check for partially hydrogenated margarine
Halo-halo (full serving)	1 large glass	H	H	M	AVOID	Coconut cream + condensed milk + ube ice cream + multiple sugary toppings
Yema (egg yolk + condensed milk candy)	4 pieces	H	H	H	AVOID	Pure egg yolk + condensed milk = concentrated cholesterol and saturated fat
Bibingka (plain, no extra butter)	1 medium	M	M	M	LIMIT	Rice flour + egg + coconut milk; moderate in all; avoid extra butter topping
Champorado (chocolate rice porridge)	1 bowl	L	H	L	LIMIT	High refined carb + sugar → raises TG; skip condensed milk topping; use low-fat milk
Puto (plain, steamed rice cake)	3 pieces	L	M	L	LIMIT	Lower fat than most kakanin; refined rice starch raises TG if eaten in large amounts
Fresh papaya, melon, watermelon	1 cup cubed	L	L	L	GOOD	High fiber; natural sweetness without TG spike; best dessert choice after any meal
Boiled saba banana (no sugar)	2 pieces	L	L	L	GOOD	Resistant starch + fiber; healthy snack; do NOT fry or add sugar

SF = Saturated Fat · SUG = Sugar/Refined Carb (TG driver) · CHOL = Dietary Cholesterol · H=High · M=Moderate · L=Low | Educational use only. Consult your physician for individual dietary advice.

williamriveromd.com

Food Item (Brand / Chain)	Typical Serving	SF	SUG	CHOL	Rating	Key Note
F · FAST FOOD — JOLLIBEE · MANG INASAL · KFC · MCDONALD'S · CHOWKING						
Jollibee Chickenjoy (with skin + gravy)	1 piece	H	M	H	AVOID	Skin + batter = phosphate additives absorbed at 100%; gravy adds sodium + starch thickeners
KFC Original Recipe (with skin)	1 piece	H	L	H	AVOID	11-herb batter seals high sat fat in; skin contains most of the phosphate additives
Jolly Hotdog Classic (with cheese)	1 sandwich	H	M	M	AVOID	Hotdog = high sat fat + sodium phosphate preservatives; cheese adds more sat fat
Jollibee Jolly Spaghetti	1 regular	M	H	M	AVOID	Sweet banana ketchup sauce = high sugar → TG↑↑; hotdog slices add sat fat
McD McSpicy / McCrispy (with skin)	1 sandwich	H	M	H	AVOID	High sat fat from breading + palm oil frying; remove breading and skin to reduce impact
Burger King Whopper / McD Big Mac	1 burger	H	M	H	AVOID	Double beef patty + cheese + mayo = ~20g sat fat per burger; cheese and mayo worst contributors
Chowking Lauriat (fried chicken + pork)	1 set	H	M	H	AVOID	Fried protein + sweet sauce + white rice combo = high sat fat + high TG
Mang Inasal Liempo (grilled pork belly)	1 slab	H	L	M	AVOID	Even grilled, pork belly fat remains; avoid the marinade-basted skin portions
Jollibee Chickenjoy (skin removed, no gravy)	1 piece	M	L	M	LIMIT	Removing skin cuts sat fat by ~40%; no gravy reduces sodium and starch thickeners significantly
Mang Inasal Paa (thigh, with skin)	1 piece	M	L	M	LIMIT	Grilled is far better than fried; remove skin; use vinegar dip, not the chicken oil
McDo Filet-O-Fish	1 sandwich	M	M	M	LIMIT	Better than beef/chicken options; tartar sauce adds sat fat; ask for less sauce
Chowking Congee / Arroz caldo	1 bowl	L	H	L	LIMIT	Low fat but refined carb-heavy → TG↑; filling option post-dialysis; avoid extra toppings
Jollibee Palabok Fiesta (small)	1 small	M	H	M	LIMIT	Shrimp sauce + pork guisado + egg = moderate across all; skip the chicharong topping
Mang Inasal Pecho (breast, no skin, vinegar)	1 piece	L	L	L	GOOD	Best fast food order in PH; grilled chicken breast + vinegar dip; skip chicken oil and unlimited rice
Chowking Tofu Sisig	1 serving	L	L	L	GOOD	Plant protein; low sat fat and cholesterol; watch sodium in sauce
McDo side corn / side salad (no dressing)	1 side	L	L	L	GOOD	Best side dish choices; request no dressing or use lemon wedge instead
G · GROCERY & PACKAGED FOODS — LATA · NAKA-PACK · NABIBILI SA SUPERMARKET						
Spam Original (Hormel)	2 slices 57g	H	L	M	AVOID	10g sat fat per serving + very high sodium; among the most used processed meats in PH
Purefoods / CDO Hotdog (regular)	2 pieces	H	L	M	AVOID	High sat fat + sodium phosphate additives (100% absorbed); check label for "phosphate"
Argentina / CDO Corned Beef	½ can	H	L	M	AVOID	High sat fat + very high sodium; carrageenan filler in lower-grade brands
Ligo / CDO Vienna Sausage	1 can	H	L	M	AVOID	Processed pork + high sodium + phosphate preservatives; no nutritional benefit
Lard / mantika ng baboy (pork fat)	1 tbsp	H	L	L	AVOID	Still widely used for sautéing in Filipino kitchens; replace entirely with canola oil

Coconut cream / gata (Kara, Aroy-D, San Miguel)	½ cup	H	L	L	AVOID	~20g sat fat per ½ cup; most common hidden cholesterol-raiser in Filipino home cooking
Condensed milk (Alaska, Magnolia, Nestlé)	2 tbsp	M	H	M	AVOID	Very high sugar + sat fat from whole milk; used in coffee, kakanin, and leche flan
Lucky Me Instant Pancit Canton / cup noodles	1 pack	M	H	L	AVOID	Palm-oil fried noodles + very high sodium + MSG; regular use raises TG and LDL
Butter (Anchor, Gold Ribbon, Lady's Choice)	1 tbsp	H	L	L	AVOID	7g sat fat per tbsp; replace with canola oil for cooking; use sparingly on bread
Eden / Quickmelt / Kraft Cheese	1 slice	H	L	M	AVOID	Processed cheese = high sat fat + sodium; Filipino processed cheese brands highest in sat fat
Whole egg (native or commercial)	1 large	L	L	H	LIMIT	1 egg/day acceptable for most; hyper-responders (25% of people) should limit to 4/week
Peanut butter (Lady's Choice, Skippy)	2 tbsp	M	M	L	LIMIT	Moderate sat fat from palm oil filler; choose natural/no-sugar-added versions when available
Whole fresh milk (Anchor, Alaska, Bear Brand)	1 glass 250mL	M	M	M	LIMIT	Replace with low-fat (1%) or skim milk; or switch to Vitasoy soy milk
Canned tuna in oil (Century, Mega)	1 can	L	L	L	LIMIT	Tuna itself is healthy; oil packing adds unnecessary fat — choose water-packed version instead
Sardines in tomato sauce (555, Ligo, San Marino)	1 can	L	L	L	GOOD	Affordable omega-3; LDL and TG lowering; eat 3–4x/week; NOT in oil or cream sauce
Canned tuna in water (Century, Mega)	1 can	L	L	L	GOOD	Lean protein + omega-3; versatile; watch sodium content in flavored variants
Quaker Rolled Oats (plain)	½ cup dry	L	L	L	GOOD	β-glucan fiber lowers LDL 5–8%; best breakfast; do NOT add condensed milk or sugar
Canola oil (Baguio Gold, Mazola Canola)	1 tbsp	L	L	L	GOOD	Best all-purpose cooking oil in PH: high MUFA, low SFA; replaces lard and coconut oil
Tokwa / firm tofu (fresh or packed)	½ block	L	L	L	GOOD	Plant protein; replaces pork in adobo, menudo, scrambles; soy protein modestly lowers LDL
Brown rice (various brands)	1 cup cooked	L	M	L	GOOD	More fiber + lower glycemic index than white rice; replace at least half of daily white rice intake
Pili nuts (Bicolano native nut)	30 g handful	L	L	L	GOOD	Native Philippine nut high in MUFA + PUFA; cardioprotective; best snack alternative to chicharon
H · COOKING OILS & FATS — MANTIKA AT TABA						
Coconut oil (VCO, Minola, Natura)	1 tbsp	H	L	L	AVOID	92% saturated fat — the highest of any oil; consistent LDL-raiser in clinical trials
Lard / baboy fat (mantika)	1 tbsp	H	L	L	AVOID	Still standard in many Filipino kitchens and carinderia; replace with canola oil
Palm oil / palm shortening	1 tbsp	H	L	L	AVOID	50% sat fat; used in commercial frying, instant noodles, and many biscuit brands
Partially hydrogenated margarine (trans fat)	1 tbsp	H	L	L	AVOID	Raises LDL AND lowers HDL simultaneously; worst fat for heart; check label for "hydrogenated"
Regular vegetable oil blend (generic)	1 tbsp	L	L	L	LIMIT	Often contains palm oil blend; check label; use less than you think you need
Canola oil (Baguio Gold, Mazola)	1 tbsp	L	L	L	GOOD	Best for high-heat sautéing and frying; high MUFA, very low SFA; neutral taste

Extra virgin olive oil (various brands)	1 tbsp	L	L	L	GOOD	Best for dressings, dipping, low-heat cooking; polyphenols add anti-inflammatory benefit
I · INUMIN — DRINKS & BEVERAGES						
Softdrinks (Coke, Sprite, Pepsi, Royal)	1 can 330mL	L	H	L	AVOID	35–40g sugar per can → TG↑↑; habitual intake is one of the fastest ways to raise triglycerides
Red Horse / San Miguel Beer	1 bottle 330mL	L	H	L	AVOID	Alcohol directly raises TG; stopping beer often drops TG 30–50% within weeks
Ginebra gin / Tanduay rum	1 shot + mixer	L	H	L	AVOID	Alcohol metabolism in liver generates fat; spirits mixed with juice/cola = compounded TG risk
3-in-1 coffee (Nescafe 3-in-1, Great Taste)	1 sachet	L	H	L	AVOID	Non-dairy creamer (palm oil) + sugar = high TG driver; switch to black brewed coffee
Sago't gulaman / buko pandan (sweetened)	1 large glass	L	H	L	AVOID	Arnibal (brown sugar syrup) + coconut cream = high sugar + sat fat; beloved but very high TG
Powdered juice drinks (Tang, 8 O'Clock, Zesto)	1 glass	L	H	L	AVOID	Almost pure sugar + artificial flavoring; no nutritional value; major TG raiser in Filipino diet
Milo / Ovaltine (hot or iced)	1 cup	L	M	L	LIMIT	Moderate sugar; use less powder than directed; prepare with skim or low-fat milk
Fresh buko (coconut) juice, no sugar added	1 coconut	L	M	L	LIMIT	Natural sugars; refreshing; acceptable if not diabetic; avoid bottled sweetened versions
Fresh fruit juice (no added sugar)	1 glass	L	M	L	LIMIT	Fructose in juice still raises TG; eat whole fruit instead — the fiber slows sugar absorption
Black coffee (barako, brewed, no sugar)	1 cup	L	L	L	GOOD	Unfiltered (French press, barako boiled) may raise LDL slightly; paper-filtered is safer
Salabat (fresh ginger tea, no sugar)	1 cup	L	L	L	GOOD	Anti-inflammatory; traditional Filipino tea; small LDL-lowering effect; ideal post-meal drink
Unsweetened calamansi juice (fresh)	1 glass	L	L	L	GOOD	Vitamin C; very low sugar if no arnibal added; excellent with grilled fish
Plain water / mineral water	Any amount	L	L	L	GOOD	Best beverage; aim for 6–8 glasses/day; add calamansi slices for flavor

LIFESTYLE — BEYOND DIET

Changes That Move Your Lipid Panel

 Aerobic Exercise ↑ HDL 3–9% · ↓ TG 10–20% 150 min/week brisk walking, swimming, or cycling. Shifts LDL particles to larger, less dangerous type.	 Weight Loss (5–10%) ↓ TG 20–30% · ↑ HDL 5–8% Even partial weight loss measurably improves all lipid values — every kilogram matters.	 Reduce White Rice ↓ TG fast · ↓ TG/HDL ratio De novo lipogenesis: your liver converts excess refined carbs to VLDL fat. Smaller portions + brown rice.
 Quit Smoking ↑ HDL within weeks Smoking lowers HDL 4–6 mg/dL. Quitting is the single most impactful cardiovascular intervention.	 Stop Alcohol ↓ TG 30–50% in weeks If TG is elevated, stopping Red Horse and gin is often faster than any dietary change.	 Sleep 7–8 Hours Modest lipid benefit Poor sleep raises LDL and TG independently. Chronic stress raises cortisol → more VLDL secretion.

✔ **Sample Heart-Healthy Filipino Day**

Breakfast	Quaker oatmeal + sliced banana + 1 boiled egg (or tokwa scramble) + black barako coffee (no 3-in-1)
Lunch	½ cup brown rice + monggo guisado with malunggay + inihaw na galunggong + tomato-onion ensalada + calamansi juice
Snack	30g pili nuts or cashews + fresh papaya or melon slices (NOT juice)
Dinner	Sardines in tomato sauce (lata) + steamed kangkong with garlic in canola oil + small brown rice portion + salabat