



MEDICATION DOSES: EPI 1 mg q3–5 min | AMIO 300 mg → 150 mg | LIDO 1–1.5 then 0.5–0.75 mg/kg | ADEN 6 mg → 12 mg (rapid IV push) | ATRO 1 mg q3–5 min (max 3 mg) | Ca gluconate 10% 1–2 g/10–20 mL (or CaCl₂ 1 g) | NaHCO₃ 50 mEq/50 mL

TREATMENT: CPR — Absent : 0 / Present : ✓ (x2-min rounds) | SHOCK — Biphasic : 120–200 J / Monophasic : 360 J | TIME in military format

[illegible]

Recording grid · Event conclusion · Personnel on duty

MEDICATION DOSES: EPI 1 mg q3–5 min | AMIO 300 mg → 150 mg | LIDO 1–1.5 then 0.5–0.75 mg/kg | ADEN 6 mg → 12 mg (rapid IV push) | ATRO 1 mg q3–5 min (max 3 mg) | Ca gluconate 10% 1–2 g/10–20 mL (or CaCl₂ 1 g) | NaHCO₃ 50 mEq/50 mL
TREATMENT: CPR — Absent : 0 / Present : ✓ (×2-min rounds) | SHOCK — Biphasic : 120–200 J / Monophasic : 360 J | TIME in military format

TIME MILITARY	BP	HR	RHYTHM	O ₂ SAT	PULSE ABS 0 / ✓	CPR ✓ /2MIN	SHOCK JOULES	EPI 1MG	AMIO 300/150	LIDO MG/KG	ADEN 6/12	ATRO 1MG	CA GLUC 1–2G	NAHCO ₃ 50MEQ	COMMENTS BVM · INTUBATION · IV FLUIDS · PACING · CARDIOVERSION · LINES · LABS · ETC

CODE BLUE EVENT CONCLUSION

Time event ended: _____

Patient outcome: ☐ ROSC achieved ☐ Expired / efforts terminated

Reason resuscitation ended: ☐ ROSC ☐ No benefit to continue

Family notification: ☐ At bedside ☐ By phone ☐ Unable to reach ☐ No family ☐ Pending

Name of person notified: _____

Relationship to patient: _____

PERSONNEL ON DUTY

_____ PRIMARY NURSE	_____ CHARGE NURSE	_____ RECORDER
_____ CPR STAFF	_____ MEDICATION STAFF	_____ RESPIRATORY THERAPIST
_____ RESIDENT ON DUTY (ROD)	_____ PHYSICIAN	_____ ON-CALL NEPHROLOGIST