



Fiber-Rich Filipino Food Guide

How much fiber is in what you eat — and what to do about it

W. G. M. Rivero, MD, FPCP, DPSN · williamriveromd.com

14–18

g

Avg. Filipino
gets

25 g

Women / day

38

g

Men /
day

Most
Filipinos
get
only
14–
18
g/
day
—
well
below
the
recommended
target.

2 Filipino Foods Ranked by Fiber

Food	Fiber per serving	CKD / Dialysis note
● HIGH — ≥ 7 g per serving		
Black beans (cooked, 1 cup)	15 g	Limit in CKD 4–5/dialysis (K [↑] , P [↑])
Garbanzos / chickpeas (1 cup)	12 g	Limit in CKD 4–5/dialysis (K [↑] , P [↑])
Chia seeds (2 tbsp)	10 g	Kidney-friendly in small doses
Monggo / mung beans (½ cup cooked)	8 g	Limit in CKD 4–5/dialysis (K [↑] , P [↑])
● GOOD — 4–6 g per serving		
Guava (1 cup)	8 g	Limit in CKD 4–5/dialysis (K [↑])
Avocado (1 whole)*	10 g*	Very high K — CKD 4–5/dialysis: ½ max
Saluyot / jute leaves (1 cup cooked)	5 g	Low K/P — generally safe
Oatmeal (1 serving, 40 g dry)	4 g	Kidney-friendly; lowers LDL
Brown rice (1 cup cooked)	3.5 g	Slightly more P than white rice
● MODERATE — 2–4 g per serving		
Kamote / sweet potato (1 medium)	3.8 g	Moderate K — leach before cooking
Malunggay leaves (1 cup)	3.5 g	Low K/P — generally safe
Saba banana (1 medium)	3.5 g	Moderate K — limit in CKD 4–5
Okra (1 cup cooked)	3.2 g	Low K/P — very kidney-friendly
Sitaw / yard-long beans (1 cup)	3 g	Generally safe; low K
Sampaloc / tamarind (100 g pulp)†	3–5 g†	Only if pulp is eaten; high K
Kangkong / water spinach (1 cup)	2.5 g	Low K/P — safe in usual amounts
Pako / fern salad (1 cup)	2.5 g	Low K — generally safe
■ LOW — < 2 g per serving		
Pechay / bok choy (1 cup cooked)	2.5 g	Low K/P — safe
Eggplant / talong (1 cup)	2.5 g	Low K/P — safe
Pineapple (1 cup)	2.3 g	Low K — safe in moderation
White rice (1 cup cooked)	0.6 g	Lowest K/P of all staples

*Whole avocado (full fruit). †Sampaloc pulp only — strained sinigang broth = 0 g fiber.

3 Fiber Supplements

Psyllium husk (C-Lium)

Shirataki rice

Chia seeds

Flaxseed (ground)

Oat bran

3–5 g / serving

Best all-round; very kidney-friendly. Drink extra water.

~1 g / cup

Ultra kidney-friendly; near-zero K & P. Good rice substitute.

10 g / 2 tbsp

Easy to add to lugaw or water. Omega-3 bonus.

5–6 g / 2 tbsp

Omega-3 bonus. Use ground for absorption.

6–8 g / ½ cup

Beta-glucan lowers LDL. Good in porridge.

4 CKD & Dialysis Notes

Important for Kidney Patients

- **Psyllium and shirataki** are the most kidney-friendly options — low potassium, low phosphorus.
- **CKD Stage 4–5 / dialysis:** limit avocado, monggo, and guava due to high potassium and phosphorus — check amounts with your dietitian.
- **Prunes** work well for constipation (7 g fiber + sorbitol effect) but are **HIGH in potassium** — limit to 2–3 pieces in CKD 4–5/dialysis.
- **Sinigang mix powder** contains 0 g fiber. Use fresh sampaloc pulp (eaten, not strained) to get the fiber benefit from tamarind.

5 The Fiber Myth

COMMON MISCONCEPTION

"A big green salad $\approx$ 2–3 g fiber. Half cup monggo $\approx$ 8 g. Fiber $\neq$ vegetables."