Acceptance Study for SciBooNE Charged-Current Coherent Pion Production Technical Note Rough Draft

Jonathan Asaadi*1 and Zachary Williams^{†1}

¹Department of Physics, The University of Texas at Arlington

April 11, 2017

Abstract

Should there be an abstract?

1 Introduction

The goal of this document is to provide a reference for the acceptance study performed for the SciBooNE charged current coherent pion re-analysis as well as provide documentation to the code used in this study (in the event anything needs to be revisited in the future).

The code currently lives in this github repository labeled SciBooNE-MC and the corresponding ROOT files used in the simulation can be downloaded from here (insert dropbox/Google Drive Link here)

The paper is structured such that Section 2 outlines samples used in this study, Section 3 describes the SciBooNE detector as it was simulated in this study, Section 5 gives a high level summary of the results including the event-reduction table as well as the CC-Coh- π acceptance results.

Sections ?? - ?? provide supporting plots which are used to generate the acceptance tables found in Section ??.

The appendix is left to explain how the code is run and the details of the scripts within.

1.1 Goal

The goal of the reanalysis is to examine the acceptance modeling for the SciBooNE results in the presence of modern neutrino generators and updated models in order to understand why SciBooNE did not observe Charged-Current Coherent Pion Production at low neutrino energy. The purpose of this acceptance study is to blah blah blah... (coming back to this later...)

2 Samples

Five different samples were used in this study, three samples for ν -mode and two samples in $\bar{\nu}$ -mode. ¹ Table 1 summarizes these samples. Details on these samples can be found in Appendix

¹All of these samples were generated by Callum Wilkinson (many thanks)

Summary of samples

Mode	NEUT version	Pion-Model	Number of simulated events
$\overline{\nu}$	5.3.6	Rein-Sehgal	1,000,000
$\overline{\nu}$	5.3.6	Berger-Sehgal	1,000,000
$\overline{\nu}$	X.X.X	Rein-Sehgal	100,000
$\bar{\nu}$	5.3.6	Rein-Sehgal	1,000,000
$ar{ u}$	5.3.6	Berger-Sehgal	1,000,000

Table 1: Summary of the samples used to build the acceptance model for this study.

3 Simulation

This section is intended to detail the nuances of this acceptance model, and to detail what assumptions are made in the acceptance modeling to result in accurate classifications of events as Charged-Current Coherent Pion Production.

3.1 The Detector

For the purposes of this acceptance study, the SciBooNE experiment as being composed of two sub-detectors. The first (and the more upstream) of the sub-detectors, is the Scintillator Bar Tracker (SciBar) which was originally conceived and constructed to function as the near detector for the K2K experiment [reference]. The second (and more downstream) of the sub-detectors, is the Muon Range Detector (MRD), which is the detector designed and constructed specifically for SciBooNE for measuring the momentum of muons produced from charged-current neutrino interactions up to 1.2~GeV/c by using the observed range of the trajectory of the muon. These detectors and the corresponding coordinate system we will use throughout this note are shown in Figure 3.1




Figure 1: Representation of the SciBooNE detector and the coordinate frame we use in this study

3.1.1 The Scintillator Bar Tracker (SciBar)

The Scintillator Bar Tracker (SciBar) sub-detector is a scintillator detector (go into more detail?). In this acceptance study, the z-direction starts at 0 at the front face of the SciBar sub-detector. The direction of the beam is in the z-direction, which means the xy-plane is perpindicular to the beam. The dimensions of the sub-detector have the x and y dimensions of the same length of 3.0 m. The

z dimension is 1.7 m. This simulation models the scintillator materials as having a constant energy deposition (dE/dx) value of 2.04 MeV/cm.

3.1.2 The Muon Range Detector (MRD)

The Muon Range Detector (MRD), depicted in Figure 3.1.2 is located 0.55 m downstream of SciBar in the z-direction, and is a composition of 2 sets of 13 alternating slabs of steel-scintillator layers, where the scintillator layers alternate between being horizontally oriented or vertically oriented, in the xy-plane. The steel layers have a z-direction thickness of 5.08 cm and the scintillator layers have a z-direction thickness of 0.6 cm. Combining all the layers of the different alternating materials results in 26 scintillator layers that "sandwich" 25 steel layers inbetween and gives a total z-direction dimension of being 1.37m. The xy-plane is modeled as a square again (as was the case with SciBar, too) with dimensions in the x-direction and the y-direction of 2.6 m. The energy deposition (dE/dx) of the muon for the scintillator layers is again a constant of 2.04 MeV/cm and the energy deposition for the steel layers are a constant with value 11.43 MeV/cm.




Figure 2: Depiction of the Muon Range Detector (MRB) which consists of alternating layers of horizontal scintillator (shown in pink) steel slabs (showin in grey) and vertical scintillator (shown in purple)

4 Event Selection

4.1 Event Classifications

There were three different classifications for events that qualified as CC-Coh Pion Production events, that the muon made it to the MRD, and the muon penetrated at least three layers of steel. These categories will be referenced multiple times throughout the remainder of this paper, which makes pertinent that the reader has an understanding of what each of the three specifically mean for any event that falls under any of these classifications.

4.1.1 Stopped

An event is classified as "Stopped," if the event qualified as being a CC-Coh Pion Production event, the muon of the event reached the MRD, penetrated three layers of steel, and stopped (or embedded) in the MRD without exiting the sides or the back face. Events that are classified as "Stopped" are included in the combined samples of this acceptance study and are called "Good" events. Maybe put in the dimensions of the xy-plane and z-direction that meet this classification for the MRD? This is shown in the figure below.




Figure 3: Depiction of an event that was classified as "Stopped."

4.1.2 Not-Stopped

An event is classified as "Not-Stopped," if the event qualified as being a CC-Coh Pion Production event, the muon of the event reached the MRD, and the muon passed out the back face of the MRD without stopping. Events that are classified as "Not-Stopped" are included in the combined samples of this acceptance study and are also called "Good" events. Maybe put in the dimensions of the xy-plane and z-direction that meet this classification for the MRD? This is shown in the figure below.




Figure 4: Depiction of an event that was classified as "Not-Stopped."

4.1.3 Out-Side

An event is classified as "Out-Side," if the event qualified as being a CC-Coh Pion Production event, the muon of the event reached the MRD, penetrated three layers of steel, and then passed through one of the sides of the MRD (not including the back face) without stopping. Events that are classified as "Out-Side" are not included in the combined samples because there was not enough material traversal for an accurate reconstruction of the particles momentum and energy to be made. Maybe put in the dimensions of the xy-plane and z-direction that meet this classification for the MRD? This is shown in the figure below.




Figure 5: Depiction of an event that was classified as "Out-Side."

4.2 Stepping Through the Detectors Along an Event

I do not know if I am really going to include this here or not...

5 Results

A Appendix: Sample Details

Appendix on samples

A.1 ν -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000 ν interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE_numu_coh_RooTrack.root

found at the following link (put link to sample here).

A.2 ν -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000 ν interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE_numu_coh_RooTrack_NEW.root

found at the following link (put link to sample here).

A.3 ν -Mode Rein-Sehgal NEUTvx.x.x

A sample of $100,000 \nu$ interactions were simulated using the NEUT generator (vx.x.x, believed to be the version used by the SciBooNE collaboration in the original publication) and the corresponding older Rein-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE_numu_coh_OLDNEUT_RooTrack.root

found at the following link (put link to sample here).

A.4 barv-Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000 $\bar{\nu}$ interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE_numubar_coh_RooTrack.root

found at the following link (put link to sample here).

A.5 $\bar{\nu}$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000 $\bar{\nu}$ interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE_numubar_coh_RooTrack_NEW.root

found at the following link (put link to sample here).

A.6 Vertex Distributions

The events were all given a random initial point that was generated with the goal that the vertex distributions of this simulation would closely match the vertex distributions that Hiraide (need to put a reference) showed in his thesis. This was done by... etc.

Put in the code for how we made the vertex distributions of the interactions.




Figure 6: Vertex distributions of the events in the new Rein-Sehgal sample.

A.7 NewNMReinSehgal.C

This file is the macro that corresponds to the "NewNMReinSehgal.h" file, which connects with this file: "SciBooNE_numu_coh_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms). When this macro is run (which can take a while), it also plots a few different histograms. The histograms that are plotted are the ones shown in the figures below with descriptions included with the corresponding figures. The order that the histograms appear in this paper is the same order they will be shown when this macro is run in root.




Figure 7: New Rein-Sehgal X-Y vertex distributions for muons that made it to the MRD.




Figure 8: New Rein-Sehgal Z vertex distributions for the interactions.




Figure 9: New Rein-Sehgal Y vertex distributions for the interactions.




Figure 10: New Rein-Sehgal X vertex distributions for the interactions.




Figure 11: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 12: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".




Figure 13: This histogram depicts the amount of muons that embedded in what layer of steel.




Figure 14: This is a plot right here. Nice of you to notice!




Figure 15: This is a plot right here. Nice of you to notice!

The NewNMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{1}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{2}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either θ_{μ} , or θ_{π} , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal

angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{3}$$

$$\theta_{\pi} = tan^{-1}(\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}/P_{\pi_z}) \tag{4}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are Q^2 and |t|. The Q^2 corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{5}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute Q^2 :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$

$$(6)$$

 Q^2 is reported in units of $(MeV/c)^2$.

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{7}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(8)

|t| is reported in units of $(MeV/c)^2$.

A.8 NewNMBergerSehgal.C

This file is the macro that corresponds to the "NewNMBergerSehgal.h" file, which connects with this file: "SciBooNE_numu_coh_RooTrack_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 16: This is a plot right here. Nice of you to notice!




Figure 17: This is a plot right here. Nice of you to notice!




Figure 18: This is a plot right here. Nice of you to notice!




Figure 19: This is a plot right here. Nice of you to notice!




Figure 20: This is a plot right here. Nice of you to notice!




Figure 21: This is a plot right here. Nice of you to notice!




Figure 22: This is a plot right here. Nice of you to notice!




Figure 23: This is a plot right here. Nice of you to notice!




Figure 24: This is a plot right here. Nice of you to notice!

The NewNMBergerSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{9}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{10}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either θ_{μ} , or θ_{π} , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{11}$$

$$\theta_{\pi} = tan^{-1} \left(\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2} / P_{\pi_z} \right) \tag{12}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are Q^2 and |t|. The Q^2 corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{13}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute Q^2 :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(14)

 Q^2 is reported in units of $(MeV/c)^2$.

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{15}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
 (16)

|t| is reported in units of $(MeV/c)^2$.

A.9 OldNMReinSehgal.C

This file is the macro that corresponds to the "OldNMReinSehgal.h" file, which connects with this file: "SciBooNE_numu_coh_OLDNEUT_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoOBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 25: This is a plot right here. Nice of you to notice!




Figure 26: This is a plot right here. Nice of you to notice!




Figure 27: This is a plot right here. Nice of you to notice!




Figure 28: This is a plot right here. Nice of you to notice!




Figure 29: This is a plot right here. Nice of you to notice!




Figure 30: This is a plot right here. Nice of you to notice!




Figure 31: This is a plot right here. Nice of you to notice!




Figure 32: This is a plot right here. Nice of you to notice!




Figure 33: This is a plot right here. Nice of you to notice!

The OldNMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{17}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{18}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either θ_{μ} , or θ_{π} , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal

angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{19}$$

$$\theta_{\pi} = tan^{-1}(\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}/P_{\pi_z}) \tag{20}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are Q^2 and |t|. The Q^2 corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{21}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute Q^2 :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(22)

 Q^2 is reported in units of $(MeV/c)^2$.

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}|$$
(23)

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(24)

|t| is reported in units of $(MeV/c)^2$.

A.10 NewANMReinSehgal.C

This file is the macro that corresponds to the "NewANMReinSehgal.h" file, which connects with this file: "SciBooNE_numubar_coh_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 34: This is a plot right here. Nice of you to notice!




Figure 35: This is a plot right here. Nice of you to notice!




Figure 36: This is a plot right here. Nice of you to notice!




Figure 37: This is a plot right here. Nice of you to notice!




Figure 38: This is a plot right here. Nice of you to notice!




Figure 39: This is a plot right here. Nice of you to notice!




Figure 40: This is a plot right here. Nice of you to notice!




Figure 41: This is a plot right here. Nice of you to notice!




Figure 42: This is a plot right here. Nice of you to notice!

The NewANMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{25}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{26}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either θ_{μ} , or θ_{π} , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{27}$$

$$\theta_{\pi} = tan^{-1}(\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}/P_{\pi_z}) \tag{28}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are Q^2 and |t|. The Q^2 corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{29}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute Q^2 :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(30)

 Q^2 is reported in units of $(MeV/c)^2$.

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{31}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(32)

$$|t| \text{ is reported in units of } (MeV/c)^2.$$

A.11 NewANMBergerSehgal.C

This file is the macro that corresponds to the "NewANMBergerSehgal.h" file, which connects with this file: "SciBooNE_numubar_coh_RooTrack_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 43: This is a plot right here. Nice of you to notice!




Figure 44: This is a plot right here. Nice of you to notice!




Figure 45: This is a plot right here. Nice of you to notice!




Figure 46: This is a plot right here. Nice of you to notice!




Figure 47: This is a plot right here. Nice of you to notice!




Figure 48: This is a plot right here. Nice of you to notice!




Figure 49: This is a plot right here. Nice of you to notice!




Figure 50: This is a plot right here. Nice of you to notice!




Figure 51: This is a plot right here. Nice of you to notice!

The NewANMBergerSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{33}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{34}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either θ_{μ} , or θ_{π} , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal

angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{35}$$

$$\theta_{\pi} = tan^{-1}(\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}/P_{\pi_z}) \tag{36}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are Q^2 and |t|. The Q^2 corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{37}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute Q^2 :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(38)

 Q^2 is reported in units of $(MeV/c)^2$.

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}|$$
(39)

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(40)

|t| is reported in units of $(MeV/c)^2$.

A.12 NMCombinedPlots.C

I need to come back and insert all of my images here.




Figure 52: This is a plot right here. Nice of you to notice!




Figure 53: This is a plot right here. Nice of you to notice!




Figure 54: This is a plot right here. Nice of you to notice!




Figure 55: This is a plot right here. Nice of you to notice!




Figure 56: This is a plot right here. Nice of you to notice!




Figure 57: This is a plot right here. Nice of you to notice!




Figure 58: This is a plot right here. Nice of you to notice!




Figure 59: This is a plot right here. Nice of you to notice!




Figure 60: This is a plot right here. Nice of you to notice!




Figure 61: This is a plot right here. Nice of you to notice!




Figure 62: This is a plot right here. Nice of you to notice!




Figure 63: This is a plot right here. Nice of you to notice!




Figure 64: This is a plot right here. Nice of you to notice!




Figure 65: This is a plot right here. Nice of you to notice!




Figure 66: This is a plot right here. Nice of you to notice!




Figure 67: This is a plot right here. Nice of you to notice!




Figure 68: This is a plot right here. Nice of you to notice!




Figure 69: This is a plot right here. Nice of you to notice!




Figure 70: This is a plot right here. Nice of you to notice!




Figure 71: This is a plot right here. Nice of you to notice!




Figure 72: This is a plot right here. Nice of you to notice!




Figure 73: This is a plot right here. Nice of you to notice!




Figure 74: This is a plot right here. Nice of you to notice!




Figure 75: This is a plot right here. Nice of you to notice!




Figure 76: This is a plot right here. Nice of you to notice!

A.13 NMPionPlotting.C




Figure 77: This is a plot right here. Nice of you to notice!




Figure 78: This is a plot right here. Nice of you to notice!




Figure 79: This is a plot right here. Nice of you to notice!




Figure 80: This is a plot right here. Nice of you to notice!




Figure 81: This is a plot right here. Nice of you to notice!




Figure 82: This is a plot right here. Nice of you to notice!




Figure 83: This is a plot right here. Nice of you to notice!




Figure 84: This is a plot right here. Nice of you to notice!




Figure 85: This is a plot right here. Nice of you to notice!




Figure 86: This is a plot right here. Nice of you to notice!




Figure 87: This is a plot right here. Nice of you to notice!




Figure 88: This is a plot right here. Nice of you to notice!

A.14 NMFourSquaredPlotting.C




Figure 89: This is a plot right here. Nice of you to notice!




Figure 90: This is a plot right here. Nice of you to notice!




Figure 91: This is a plot right here. Nice of you to notice!




Figure 92: This is a plot right here. Nice of you to notice!




Figure 93: This is a plot right here. Nice of you to notice!




Figure 94: This is a plot right here. Nice of you to notice!

A.15 ANMCombinedPlots.C




Figure 95: This is a plot right here. Nice of you to notice!




Figure 96: This is a plot right here. Nice of you to notice!




Figure 97: This is a plot right here. Nice of you to notice!




Figure 98: This is a plot right here. Nice of you to notice!




Figure 99: This is a plot right here. Nice of you to notice!




Figure 100: This is a plot right here. Nice of you to notice!




Figure 101: This is a plot right here. Nice of you to notice!




Figure 102: This is a plot right here. Nice of you to notice!




Figure 103: This is a plot right here. Nice of you to notice!




Figure 104: This is a plot right here. Nice of you to notice!




Figure 105: This is a plot right here. Nice of you to notice!




Figure 106: This is a plot right here. Nice of you to notice!




Figure 107: This is a plot right here. Nice of you to notice!




Figure 108: This is a plot right here. Nice of you to notice!




Figure 109: This is a plot right here. Nice of you to notice!




Figure 110: This is a plot right here. Nice of you to notice!




Figure 111: This is a plot right here. Nice of you to notice!




Figure 112: This is a plot right here. Nice of you to notice!

A.16 ANMPionPlotting.C




Figure 113: This is a plot right here. Nice of you to notice!




Figure 114: This is a plot right here. Nice of you to notice!




Figure 115: This is a plot right here. Nice of you to notice!




Figure 116: This is a plot right here. Nice of you to notice!




Figure 117: This is a plot right here. Nice of you to notice!




Figure 118: This is a plot right here. Nice of you to notice!




Figure 119: This is a plot right here. Nice of you to notice!




Figure 120: This is a plot right here. Nice of you to notice!




Figure 121: This is a plot right here. Nice of you to notice!




Figure 122: This is a plot right here. Nice of you to notice!




Figure 123: This is a plot right here. Nice of you to notice!




Figure 124: This is a plot right here. Nice of you to notice!

A.17 ANMFourSquaredPlotting.C




Figure 125: This is a plot right here. Nice of you to notice!




Figure 126: This is a plot right here. Nice of you to notice!




Figure 127: This is a plot right here. Nice of you to notice!




Figure 128: This is a plot right here. Nice of you to notice!




Figure 129: This is a plot right here. Nice of you to notice!




Figure 130: This is a plot right here. Nice of you to notice!

B Steps for Running the Code

The instructions on how to run the code and the order the files need to run in so that there are no resulting error messages, or other issues while running the code, are detailed in this section.

- Step 1: This is the first step. (Run the NewNM macros and the NewANM macros and the OldNM macro.)
- Step 2: This is the second step. (Run the combined plotting macros.)
- Step 3: This is the third step. (Run the Pion Plotting macros.)
- Step 4: Etc. (Run the FourSquaredMomentum macros.)

C Closing Remarks and Cautions

These are just a few cautionary suggestions for potential issues that might be encountered while trying to use this code. This will also be where and further closing remarks can be made.

D Acknowledgements

Thank everyone who helped, and thank everyone who gave their inputs into your acceptance study. YOU NEED TO GIVE A HUGE AND SPECIAL THANKS TO DR. ASAADI RIGHT HERE! (He has been suuuuuuuper patient...)

E Appendix

E.1 List of Figures

There will eventually be a huge list of figures here.

E.2 List of Tables

There will eventually be the event reduction tables and 2D histogram tables here.

	_																																
	1950.3111	0.828.29	0.713615	2011407	0,000	0.403846	0,381662	0,333333																			_						
	1300 1350	7830.67	1623833	1691252		0.25074	9	=	2					_				_	_	_	_		_	_	_	_	_				_	_	
	870,1910	813096	6000	9899	_		231623	- 52	_				_						_	_				_	_						_	_	
	STUST 1	2000	_	18390	-		_	0.0714286	~	-		_	_	-			-	-	_	_	-			_	_	-	-	-		-	_	_	-
	730180 18	708 02	_	1000		_	ß		3		-	_	_	0.0				_	_	_		-		_	_	-	-	-	-		_	_	
	2007/20 175	201369 0.7	_	50 6350	_	_	ŭ	0.25	900	8		0	0	-			-	0				-			0	0 1	=			-	-	0	0.0
	ш		ŭ			-	_	36 02	_		-	0		-	-		-					-					=			-	-		
	ㅁ	0.0021569	_	38 0.648718	-	34 0.34058	ŭ	22	0.1875			-	_	-	-				-	_				_	_	-	=				-	_	
	1910/1920	80 8	_	_	O COLOR	2 0.046	_	_	0.1875						-					_				_		-	=						
	130,100	0.50E/IR	_	0.064068	-	-	_	0.191489	_	900		0			- 1												=						
	1300130	0.74383	0.727468	0.657848	10000	0.48575	0.29418	0,435556						- 0	- 1					_				_		-	=				_		
	14501500	0.7342	0,716946	90000	900	0.3840.5	0,355191	932	0,303	0.0768231	1 -	_	_		-				_	_				_	_	-	-					_	
_	3501400 14001450	2000	0.7200	0.08173	00000	0,438053	20713E.0	1611970	0.168014	292110					-											-	-						
ıga	3301400	77406	0,721683	0001465	100000	330000	911/08/	182432	974710	818110	_	_	_	_	_		_	_	_	_	_		_	_	_	_	_			_	_	_	
Sek	13001330	1261181	735724	67453	90100	45.187	30008	32300	_	10645161		_	_	-				Ī	_	_				_	_							_	
tein-Sehgal	20130 1	0 0 0	7987	98319		7000	11483 0	_	-	0.113636 0.	1 2 2 2		_			= 0	-		_	_			_	_	_	-	-			-	_	_	-
7	Η	280,000 02 1,289,74 0.3	7.46NB 0.7	708806 00	TO STATE OF	AGENS O	379548 0.	_	_	182827 0.1	3 23	_	_	-	-				_	_			-	_	_	-	=				-	_	0.0
\geq	Б	Sarah 0.28	Ξ	_	00730	-	_	_	_	12231 018	13	0	0	-		= 0	-	0	0		00	-			0	0 1	=			-	-	0	0.0
ew [1001150 1150	8 8	-				37.3 0.36	-	Ť		_	0		-	-		-					-				-	=			-	-		
Ζ.		74 0.88785	Ť	_	0.000000	36 0.481908	200 95	_	_	75 0216867	_	-	_	-	-			-	_	_				_	_	-	=				-	_	
for	IOOLIGO TGOLLOO	2 0.852174	_	_	0.00000		_	_	_	2 0.168675	0 0			0.0	-		0.0				0.0						=	0 0		0.0			
	н	0.8345.22	0.74026	0.704342	-	0.08278	0.452253	_	0.351297	0.144772	, 0	0	0.136364		-												=						
gram	0.0010036	0.8195.0	0.770838	0.722747	2000	0.30578	0.450833	0,36223E	0.284238	0221135	0.081485	0.0405415		- 0													-				_		
Histog	036006	0.877E04	0,763163	0.713876	000/000	0.52,016	0.438182	0.371585	0.236571	0.2389.6	0.12272	0.084121	0.0253002		-					_				_	_	-	-				_	_	
Ξ	0.000	0.798611	0.784038	0.71437	001/100	0.532462	0,455387	0.378802	0.250735	0.239492	0.106557	0.08837.3	9009000	5700	-											-	-						
7	П	0.796748	DE220	0.70867	92.000	051236	0.465472	0.404237	0312306	025280	1287	0.074184		0.0131579	_	_	_	_	_	_	_		_	_	_	_	_	_		_	_	_	
for	20.80	283	92222	23332	100.00	28225	1.88222	146728	2201025	1025347	881881	500500	3014586	302222				_	_	_			_	_	_		_				_	_	
ole 1	200750	8000	9000		00000	100000	÷	SINC!	23800	18896	135764	0.0845225	00724638	0.0234114	none o				_	_				_	_	_	_					_	
Ľab	F	S15455 0.7	_	72838 0.7	-	_	_	A10802 0.4	Ĭ.	2000000	_	00 988890	ŭ	900000	3	= 0									-	-	-	- 0			_	-	0 0
- ~i	Н	18(25/7) 0.81	_		0000000		_	_	_	1291561 0.26	-		_	201730 0.00	-	= 0	-			_	00	-		_		-	-		-	-	-		
: e	H	20	_	·		_	50000	-	_	0.0	-	_			-	- 0	0.0	_	_	_	00			_	_	-	=	0 0	-	0.0		_	
Ľab	2000	2000000	2	2 0.74278		- 6	3 0.48988	_	-	34 024964	_	798 0.0831667	NI 0.0423498	386 0.0121873	-		-		_	_				_	_	-	=			-		_	
- '	300,550	Distant	0.79541			-	0.456	-	_	0.2063		0.074786	_	Septimore 12	- 1					_				_		-	=						
	2030	0716049	0.030122	2000	0,000,0	0.46743	0379450	0.336716	02000	7210120	0100105	00557781	_	000023857	- 1				_	_				_	_	-	=					_	
	100450	10000	0,582	0.524072	1010111	0.381508	0.32289	0.273638	0.22549	017771	0.0810899	0.0418139	90009000														_						
	20400	570	791677	20105	20000	22862	230065	1228074	16271	012807	004531	0.0165622	_	_	_			_	_	_	_		_	_	_	_	_				_	_	
	310,350	237763	331136	200000	2000	200517	215553	3167825	0.129648	0.0861688			_					Ī	_	_			_	_	Ī						_	Ī	
	20000 3	2005	_	21210			_			0.0461908			_	-			-		_	_			-	_	_	-	-			-		_	0.0
	Ī	2002.0		-	1000	- 12	_	÷	_	200000000	-	_	_	-				_	_	_		-	-	_	_	-	-				-	_	0.0
	30.23		_	_	-		_	900	ã	8.	-	0	0	-	- 1		-					-				0 1	=			-	-		
	150.31	0.08675	O.OPSTG7.	0.0096318	000000	002835	0.00618557					0			-												=						
	20100 100150			- 0					-			0	-	0.0	-		0.0		0	-	0.0			-			-	0 0		0.0			0 0
	(feV/c 50a)				- 0	-	0		-			0	-	-		= 0	0.00		0	-				-		-	=	0 0		0.0	_		
	RS 0.30 N		-	- c		0 0	-		0 1	00		-	-	00	= 1	= c		-	0	0 20	9 1	0 0	0 50	0 84	0 84	56 F6	n Bar	50 0	0 0	0 0	0 20	0 84	0 0
	New NM R.	0.45Dg 45.9De	9.13.5 Deg	135-18 Deg	TO AND THE	7.35 De	31.5.35 Deg	36.40.5 Deg	40.5-45 Deg	45-49.5 Deg	54.38.5 Dec	585-68 Dec	63.67.5 Deg	67.5.72 Deg	(2-00 Lb)	CONTRACTOR	855 m De	90.915 Deg	945.90 Deg	99-1035 Deg	1055108 Deg	1125.117 Dec	117121 5 Deg	121.51261.	126130.5 Deg	13054351	Seriogo (se	1205144 Deg	1485153 Dec	153157.5 Dec	157.5-162 Deg	162,166.5 Deg	1665171 Deg
	ك	_	_	_	_		Ť	÷	_	_			Ť	Ť.	_	_			Ť	Ė	_		_	Ė	Ť	_	_	_	_	_	_	Ť	_

	_																																		
	1980-300	083333	0.0000	112/29/0	125.747	0.454545	0.308825	0.1350	8181810	- 0									_		-						-								
	19001950	1000000	0.755783	212120	0.497696	0.461538	1285714	23806	1282714	_	-					_	_	_	_	_	-			_	_	_	-			-	_			_	
	18301900	0.794872	Green and a	200000	057332	0.496500	0.353636	g.	oleees	-	-						_	-	_	_	-			_	_	_	-			-	_			_	
	1810181	1861111	7979	Series .	9229	0.40113	0.402430	orlesies	0.333333	9							_		_	_				_	_		-			-	_	_	_		
	17304800	_		_		1,482,87			_	9000		_	_		_	_	_	_	-				_	_	_						_	_	_		_
	7004750	-	1897	-	-	0.4most 0	-	_	_	200000						_	_		_				_	_	_						_	_	_		
	600 1700	-	7662	-	_	-	_	Ĕ.		97							_		_					_	_						_	_			
	0.001000	_	1,280	-	_	_	<u> </u>	Ξ.	0.181818		4				-	_	_		_	-			-	_	_						_	_	-		-
	Soulem le	-	777823 0	-	0.38736 0		_	o Seeling o		0.054836 0	200				0.00	_	_	01	_	-				_	_	00				0.0	_	_	-	-	0.0
	S004580 15	-	1000	_	_	Ť	_	_	52 52	5					-	_	0	-	_	- 0		= =		_	-	- 0		= =		0.0	_	_	-	= 0	0.0
	490-1300 13	-	9 6		-	0.538936 0.4	<u> </u>	0.20223	_	0.15/885 0.2					0		0	-	_	-	-			0		0.0	-			-	_	0	0 1		
	1400-1450 148	-	200	_	_	_	_	_	_	100	_		-			_	0	0 1	=	- 0	0 1	= =		_	_	- 0		= =	0 0		_	0	-	= 0	-
gal	3301400 140	_	100	_		÷	-	_	-	1231881	_		-	-	0	-	_	-	=	-				_	-	0.0		-		0.0	_	-	-		
\hat{s}^{eh}	1300-1350 1354	_	000000	-	-	_	_	_	_	019/388 023	-		-	-		-	-	-	=	-	- 1			_	-	00	- 1		0 0		_		-		0.0
er-	2501300 130	-		-	_	_	÷	=	_	-	_	=	-				0	-	=	- 0	-										_	0			-
Berger-Sehgal	1201120 120	_	CaleTi n.Carino	_		Ť	Ť	_	÷	_			_				0	0 1	=	- 0	-			_		0.0	-			0.0	_	-	-	- 0	0.0
NM-B	_	-		-		_	_	<u> </u>	_	_	sel nordered	7	-		-			-	=	- 0	= 1		-	_	-		= 1			-	_		-		
	130 1130 1200	_	1000	-		Ť	<u> </u>	18 0.257/64	-	201236	-		_	-			0	-	=	- 0	-	= =				- 0	-	= =			_	0			
lew		08372				_	-	_	_	9778			-	t)	-	-		-	=	- 0	= 1		-	_	-		= 1				-	-	-		
\leftarrow	60 lecolloo	_	0 10 10 10 10 10 10 10 10 10 10 10 10 10	-	-	_	0.490654	_	_	2000	-			Secolo Se			0	-	=	-	- 1						- 1				_		-		
for	Н	-	100000	-		_	_	_	_	3 0.59150	-	_		67 no7LOss			0		=	-	-						-			0.0	_				
me.	950-1000	0.8485		-		-	-	-	-	033323		-	_	73 nothers	_	_		-	=	- 0	- 1			_	_		- 1				_	-	-		
Listogram	0064006	-		1 7 Charl		05/08/0	_		_	_	1,000	-	-			_	_	-	=	- 0	-			_	_		- 1				-	_	-		
Ŧist	830900	-	7,3466	-	=	-	ŏ	Ĕ.	Ť	903690	-	0.118903			0.03125		_		=	- 0	-			_	_		-				_		-		
	800830	-	0.76225	-	_	0.001238	_	-	<u> </u>		0.161453		S DOTNERS		_		_		=		-			_	_		-				_	_	-		
for 2	20800	0888833	178.00	07824m	791900	0622386	055257	049436	192520	200	2000	II SI II	UNSTEEN	-	-	00123457			=	- 0				_	_		- 1				_	_	-		
le fc	2007	0.843373	900	0.7670	10168910	0.63963	100 (20)	0.470036	0.40834	238602	3 alone	012180	DIESE CO.	0000000	0.0120482	0.00549351			_		-			_	_		- 1					-	-		
abl	002-029	38586	20142	740480	067725	0.61545	2554788	0.492452	5000	33566/	lofin	1 2008	THE REAL PROPERTY.	SCHOOL	onlon67	OUGSTR00	_	_	_	_			_	_	_						_	_	_		
H		1880435	127180	23m22	3687027	362088	_	200		13126/9	37076	Darea	UNABER	000000	0.0195072	_	_	_	_	_			_	_	_						_	_	_		
le 3	30.60	8,		7887	-	165657	-	0.472900	-	=:	18731	_	n/Arche		0014213	_	_		_					_	_	-					_	_			
abl	300500	Т	Series Series		exedit.	JASES C	7250	130444	99929	2000	lores	l locas	NO SECOND	20000	0.001/1087		_		_					_	_						_				
L	Г		70255	100	13	715 08	191	182	919	200			_	_	19	-	_	0 1	_	0.0				_	-	0.0				0.0	_	0	-	- 0	0.0
	450.50	-		100	200	18 053	15	18 1 18 1	22	8	at 0 Herry	-	-	-	_		0	0 1	=	- 0	-					0 0	-			0.0	_	-	-	- 0	
	400450		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-	100	B 0447	36 0.2563	10.3333	25/2/2	8 02248	-	_			_		0	-	=	_	-			_		0.0	- 1			0.0	_	-	-		
	30.40	990	1000	1488	0.498	0.38126	03286	62836	17333	0.1922318	-					_	0	-	=	_	-			_			-				_		-		
	330330	0.422	1355	0.58/088	0.318462	0.321879	0.2338	1585	0.170164	907	98900	Cool or	0.000 d 280						_		-			_			_				_				
	23030	8	225.20	Secure	28145	197461	152830	200	100,790	006/89/2	online 2	milean				_	_	_	_	_	-			_	_		-				_	_	-		
	200230	98.5	9095	14000	1515760	90,000,000	0.085743	0.0815217	10000	2000						-	_							_	_						_				
	130-20 20	6		In SOM	70 100000			0.0055477 0.0	ď	2 3	0.0		0.00	-	0.0	_	0	-	=	-	-			_		0.0	-				_	0	-		
	100130 130	-	3 5	000	8	900	8	8		-	0 0		-	-	0		_	01	=	-	-					0 0	-		0 0	0.0	_	-	0 1	- 0	0
	30 too 10					0	_	-								_	0	-	=		_		_	_	-		_				0	0	-		
	DED MeV c		_	_			_	_	_	_	_	_	_							_	_	_		_	_	_	_	_	_		_	_	_	_	_
	ew NMBS 0	Deg	Page 1	S.Dec	225 Deg	SZTDg 0	315 Dg 0	536Deg	HOSDE 0	25505	Section of	S.Dec	Scale	675 Dec 0	572Dg	2765 Dg 0	d Deg 0	5Deg	200	notice De	2000	Maria Lag	08-1125 Deg 0	1125.117.Dg 0	2.5 Deg 0	12151261bg	a contract	Section of	of the party of	44.148.5 Deg	148.5153.Dig 0	53-157.5 Deg 0	157.5.162.Deg 0	2000	171 175 5 Deg
	New	15 Dec	9135 De	135,18 D	182	2252	27.33	8	9	9 4	9 9	desch.	8	250	6.57	72.76	765.8	81855 Day	2000	8 8	5 1	100	108	125	12	250				1	1485	2	10.5	100	E

	_																														
	903111	1085	9 1	022220	9222	LQ.																									
	ㅁ							0 0		0	0		0	0	0		0 0	0		0	0		0 0	-	0	-	0 1	==		0 1	
	1900-1950	0,533333	0.00	0.00	70	검	0,000																				-				
	8301900	0.882383	SIZIG	9.9	4	525																									
	STUISO 1	_	_		_	_	299			-	_		-	-	_		-	-		-	-		-	-	_	0 1	0 1		-	-	
		-	_	0.47619	_	0.375	_	- 0			_														0		-			-	
	0.810221	70.00000	NESSON.	0.47619	95759	0,33333	0.22222																								
	02110041	27.74	8	9 15		_	0.571429																								
	-										0	0.0				0.0		0	00		0	00	0 0		0	0	-			-	
	16301700	1 0.823529	0.000	17520	0.47384	0,454545	770	0.2857																							
	Salen lenten	0.666667	9888	0.461538	0.44444	384615	0.352941	0.0	0																						
	1000							_		_	_		-	_	_		-	_		-	_		-	-	_	-	-	-	-	-	
	H	88	_	0634146	_		0,533333				0		0								0					0	-		-	-	
	13001330	0.508081	0.68880	0.38889	0.000057	0.381662	0.325	0.38888																							
	150,150	98888	90000	0 190	8000	347836	7900000	//																							
	щ		_	_	-	_	_	_	 12	0	_		0		0		0	0		0	0		0 0		0	0 1	0 1	0.0		0 1	
7	1400-1430	180	9 6	9 8			2	8 1 2	0.333333																						
183	13201400	3825	70000	1652174	413756	0.487179	23866	2 5	q																						
Sel	1300,1350 1		00000	-	_	_	÷	_		_	_	-	_	_	_	-		_	-		_		-	-	_	-	-		-	_	
u-T		-	_		-	-	-		S 153		_		0								0					0	-			0 1	
ĕ.	12301300	1 0.861111	/900000	0.5655	0,52	728715.0	0.348837	0.3886.2	0.168867		0,35555																				
I-F	0.1230	0.75	9062	9 19	_	_	1929271	1285714	0.088883																						
NM-Rein-Sehgal	001130 1130120 12101230	-	_	_	_	-	_	_		0	_			0	0		0	0		0	0		0.0		0	0 1	0 1	0.0		0 1	
7	11301	8 8	0,683544	0.00000	0.501338	0.465116	0.376812	0.272727	10		_																				
ŏ	1001130	0815780	3,76316	2000	7,8817	5210TS	0.6566	26/85/	111111	143857	_		_	_	_			_			_				_	_	_		_	_	
for (0.001100	1.00 STATES		-	_	ŭ	-	8 8	2 5					-	_	-	-	-		-	_		-	-	-	-	-	-	_		
	Е	-		_	_	_	_	100	38	0.1	0	0125 0									0					0	-			-	
Histogram	1000,1050	0.8 0.800067	0.747253	9000	0.52880	0,531034	0.471588	0.317318	870	0.071428	3																				
g	0.0010036	0.766231	0.70187	061646	0,654054	0.1935.0	0.416957	0.3016.25	285714	230.02	35/143	18																			
to,	25 025005		7,46688 0.	0 196194				0.407407 0.	8 62	190		0 0			-	-	-		-	-	-		-		_	-	-	-	-	-	
Η̈́	8		_		-	_	_	# E	8 8	0.02	0	0 0	0		0		0	0		0	0		0 0		0	0	0 1			0 1	
$\overline{}$	83030		0.965079	9 0	0.503.58	0.524887	œ.	0.2047.57	0.23620	0.122807	0.131579	9000															-				
$\overline{2}$	SUSO SOON	0.90 EBB	0.744681	98339	7897.190	7350557	0.4508578	3,479,672	27.24.3	112857	17671	97202	1111110		_						_				_	-	-			_	
-	0.802	S7.5 S7.750	7007	78190	6.0074	SIGN	Hezses	Success	STORES.	26210		0.075/217			_											_					
e J	H	99	_	-	_	_	-			_	_		-	0				0			0	00	- 0		0	0	-			-	
Pable	200730	0.7438	0.725		0.632350	0,55078	0.45631	0.42222	1620	0.23828		0.0487816																			
Ë	9700	r- oq	1387	61869	100000	5406	7,01017	12649	28136	23588	138.78	JUNE 402	0.022222																		
4	9 02	01857145 0.65		-	-	222	<u> </u>	8 1	88	_	_	ne.	0.000000	_	_	-	-	-		-	_		-	-	-	-	-	-	_	_	
le	900							9 8	0.26360	_	_			_			0		0 0	0			0 0		-	0	-			-	
<u> </u>	20.60	- 69	0.72519	OFFILE OF	0.63818	1949210	0.433025	0.37704	0.2300	0.16638	0.110701	0.0569562	0.022706																		
Η	300,530	504762	766657	647482	6,000	230100	25525		21233	171588	0.107266	0.0724638																			
	П	-	-		_	_	-			_	-			0	-	0.0		0	0.0		-	0.0	0.0	0.0	-	0 1	-			-	0.0
	9030	0.00	0.65257	9	0.465600	0.46608	0.3640	0.28296	0.13622			0000000															-				
	02700	799997	8	2000	38797	3000	381239	939	163462	0.160819	0.0826613	0.0897554																			
	H				_	38	9					0.01230022		-	-		-	-		-	_		0.0	-	_	0 1	-			-	
	320,400		0.451613		-	0.236236	_	0.23507			_	Ē.														0	-				
	300330	0.5 0.285714	0.285714	I STORES	0.280488	910	024444	0.155172		100030301	0.023622																				
	250.30	10	0.181818	1184211	0247050	0.1375	1-	0.13333	0134783	9030100																					
			8		_		_	2 6	88	90	_	0.0				0.0			0.0					0.0	_	0	-			-	
	311.23	0.00	0.000023		2 0.04	_	_	700																		-	-				
	130,300		0	11671428	79991100	0.0217391	723018170																								
	100130	Ī		_		_	_			_	_	_			_	_		_					_		_	_			_	_	
	20100	00	_	_	_	_	_	_			_	-	_	-	_	-		_			_				_	-	_		_	_	
	MeV/c St	00	ے د	2 =		0	0	ے د	20	0	0	00		0	0	00		0	00		0		-	20	0	0 1	ا ت	20		0 1	20
	0.30 Me					_	_	- 0			_														_					-	
	6d NM R.S	345 Deg 459 Deg	9.13.5 Deg	S.225 Dec	25.27 Dec	27.31.5 Deg	31.5-36 Deg	Standing of fire	5 495 Deg	195-54 Deg	54-38.5 Deg	385.03 Deg	67.5.72 Dec	72-76.5 Deg	76.5-81 Deg	81.85.5 Deg	90.945 Dec	945.99 Deg	SOURS Deg	08.112.5 Dec	125-117 Deg	17.12.5 Dg	26.130.5 Deg	35 130.5 Deg	395144 Dg	44.14S.5 Deg	485-153 Deg	52 54 50 Dec	62.165.5 Deg	1665-171 Deg	7.1.16.5.Dg
	PiO	0.45Deg 459Deg	913	18.0	22.5	27.33	ii B	9 9	9 9	9	80.00	, j 6 8 8	100	22.78	76.5.1	82 W	8	9151	18	9	1125	171	126.1	ij	1395	7	1485		162.1	98	1755

	_					_					_				_							_			_				_
	lgen-2mm	0.705882	0688776	5715200	9																								
	1900-1950	20,000	0.719988	162230	125		_		-		_	_			_			_	_		_	_			_				
	18301900 1	78378										_										_			_				
	Brodso 18	131962 0.				_	0 1			0.00		_	0.0	0.0	_	0.0			0 1			_	0.0	-	_	0 0	0.0	0.0	_
	1730-1800 18		0.71859 0.7	_			0 1			0.0		_	00	0.0	_				0 1			_			_	0 0			-
						_	0 1			-		_		-	_	00	-	-	0 1	= =	-	-		-	_		0.0		
	160 1700 1700 1750		7 0.70146 92 0.08188			120	-		- 0			_			_	00	-		-						_		0.0		
	-	4 0.733146		_	4 0388	3 0333	2					_				0.0						-			_	0 0	0.0		
	o lemico		0.724876		0.3461	0.3333			- 0							00													
	15504600	80.00	0.716617	182/38	0.402052	0.302303						_			_							_			_				
	1300-1550	0.797.08	0.708791	0.59465	03869	0.28361	0.12857																						
	4501500	250 FEB.	0.71233	0.5973BS	161286	20023	21					_	_		_			_	_		_	_			_	_			
	1400-1450	0.735301		Constan		706720	lecars.	3.			_	_			_			_				_			Ī				_
Rein-Sehgal	1330,1400,1	0.78624 0.79623 0		10880		_		02/2/2/		-	_	_		-	_			_	-	==		_		-	_	-	-		
Set			0.725786 0.7		-	_	_	11111		-		_		-	0	00	-	_	-	= =		-		-	_		0.0	0.0	
in-	IZadan landan			_	_	_	-		- 0	0.0		_		0 0	_	00						0	0.0		-	0 0	0.0	0.0	
-	20 1230	16 0.820552 36 0.816524		97 0.296406 7 0.25406		_	_	2 T	- 6	3 -		_	- 0	0		00			0			-			_	0 0			
ANM	1150120 120120		0.742802 0.700468			_	_	125391						0 0		00			0 1							0 0	0.0		
	113042		0.756123		0.487885	_	_	0.307002				_													_				
New	10501100 11001150	0.0465	0.788636	0.62355	0.5382	0.30346	135912	2 2	0.0833333			_													_				
	10501100	1888 0.78672	0.756151	0.649333	20000	0.42007	0.338079	23043	olskis																				
for	londes	191181	764215 708415	Delicats	1986	12528	34745	25160	0.125 0.155 0.155 0.155 0.155	2000	_	_			_			_	_		_	_		_	_				_
am	0001000	184722	0,777,286	75.000	2232	1420201	3.3861.28	128107	0.116279		_	_		_	_			_	_		_	_		_	_	_			_
Histogram	05006				1000	0.42842	137/221	25938	Distraction of the state of the		25	_			_	-		_	_			_		_	_				_
ist(820900 9		72316			_	_	2400s		005555		_			_			_	-			_		-	_		-		
Ξ	800800		0.778019 0.77291		_	_	-	2000	0175556		0.125	-				00						0	0.0	-	_		0.0	0.0	
$\frac{2}{1}$	70-80	_	7,2247 0.7	200					olsecs oly	2007	2	_				00		_		==		0	0.0	-	_		0.0	0.0	
for ,	_	==												-	-		-	-	-	==					_		0.0		-
lable i	90.50	0.846154		-		_	_	028062				265 (M289855)			_	00	-		-						_		0.0		
Ta!	650-70	0.807.565		0.086785	9.0		-	1000	0.190678			8 00074			_							-			_				
	0.000		0.746473								0.043912	0.014498													_				
je	230600	0.836902	0.78400	0.700024	0.50658	7,080,0	0.38282	22072	0.178556	0.0817391	0.0423729	nolumes																	
E	00000	8722	7,8963	.000F13	2000	42900	Sciani	200	2500	00721161	0.0374384	1000E3564																	
	98.00	7435		538125	-	aple28	33369	212046	0140620		0.0226148 0	_	-	-	_	-		_	-			_	-	-	_	-	-		
	母母	585054 611905 07	20 2	(F) (A)	9 70	23	18 J	88	12kgl 0.1		m	_	00	0 0	_	-			0 1			0			_	0 0	0.0		-
	ģ		88	200	2 22	350	20.0	8 2			=		0 0	0 0	_	00			0 1		-				_	0 0			
	9.00	==	0.6271	90000		0.308		0.1313	2 oleans		_	-				00			-						-				
	30.50	0.339333	0.372408	0.32415	0.237522	1282	0.16816	0.0300.0	0.0053882	-		_			_							-			_				
	20.30	1881451	132861	120064	157462	140581	1107342	10/5/2/2	3011405			_			_	-			-			_			_				
	20020	179487	175821	Listons	0779102	7170000	Sissis	10169103			_	_			_			_				_			Ī				_
	Ľ	000000	nest286	0.047723			0 1	00	0.0			_	0.0		_	0.0			0 1	0.0		_	-	-	_	0.0	0.0	-	_
	100130 13020	88	99	100	8	8	0 1		0.0	0 0			0 0	0 0	_	00			0 1	0.0		0				0 0			
	Solon 100				-					-	-	_			_						-	_							-
	0.50 MeV/c 5	_ 0			_ 0			_ 0		_ 0		_	_ 0	_ 0						2		_				_ 0			
	RS	0 0			0 0		B 1	50 50		0 00		B	E 0	0 00		50 5	ء قرم	Deg	Deg	0 G	Deg	neg 0	Deg d	Dec D	Deg	0 0 0 0 0 0 0 0 0 0	Des	Deg d	Deg
	New ANM	045Dg 459Dg	9-135 Deg 135-18 Deg	18.22.5 Dg	Zals De	31 5.35 De	36-40.5 Dg	65.55 5.45 5.45 5.45 5.45 5.45 5.45 5.45	49.554 Dg	385.63 D.	63675 De	675-72 Dg	72.65 Lig	81855 Dg	85590 D.	Supta Pe	online De	lessies Deg	Ins.1125.D	1712 5 D	121.5-126 De	12613051	1205125De 1751205De	1305-144 De	144.148.5 Deg	1485453	1575 162 Deg	1621665 Deg	17.17551
	_	_	_	_	_	_	_		_	_	-	_	-	_	_	-	-	_	_	_	_	-	_	-	_	_	_	-	-

	_																																	
	1950,310	0.876524	7055507	12321910	0,487179	0,482759	0,333333				-						-											_	_			_		
	1900-1950	0.82529	0.706827	OGGRESS	0.553632	0.455405	0,000	52	_	-					_		_	-	-	_	_		-		_	-	_	_				_		
	18501910	0.774194	62016970	17582010	052250	0.28257	0,53555	0.333553	_						_	-							-			0	_	_	_			_		
	SILIST	0815780	0,000	3627737	13250	0.443623	138961		_						_	_			_	_	_				_	_	_	_				_		
	0.81057	78787	_	_	_	0.445783	_	_							_					_	_				_	_		_				_		
	7004750 13	0.824074 0.	_	_	_	0,445299 0.	_	0.166667	0.0	= 0	-		-		_	0 0			-	_	_	0.0		_	_	_	-	_	0.0	-		_	0.0	
	670,700 17	28/2/13 08	_	-	Ť	÷	_	0.0	-	= 0	0.0				-	-		-	-			-				_	-	=	-		0.0	_	-	
	1600,1620, 146	81756 0.8	-	Ť	Ť	÷	-	135555	8.	= 0		-	-		0	0 0		0.0	-	_	_	0.0	-	-	_	-	-	=	0.0		0.0	0	0.0	
	50,100 10	22530 03		_	=	-	2	_	0.166957 0	3 .	= =	-	-			-			-	_	_	0.0	-	-	_	-	-	=	00		0.0		0.0	
	30,130 130	S23.29 0.82	-	-	÷	÷	_			3 .	= =	-			-	- 0	-		-	_	_	-			_	-	-	=			-	_	-	-
			-	÷	÷	_		_	SI.	c		-			-		= 1		-	-	_	-	-	-	_	0	-	=			-	-	-	-
	14501500	7939067	-	-	Ť	Ť	-	_	0.157866	= 0		-	-		-	- 0	= 1		-	-	-	-		-	-		-	=					-	-
ıgal	301400 14001450	1 0.83	_	-	Ť	_	_		02020			-				- 0	-	= =	-	_	_	- 0			_		-	=					- 0	
Set		1800180	-	_	Ť	÷	_		0.410256		3 -	-					- 1		-						_	_		=						
er-	13001320	0.89005	0.730774	0.689633	0.386435	0.548585	0.4447	0.53648	0.338983	UZIEWO	==	. =					-		. =								_	=						
ANM-Berger-Sehgal	138130	0.840741	0.7530	0.66013	0,581667	0,53848	0.4887	03226	22	7/05/1		. =							. =									=						
Æ.	130c1230	0.867.33	0.7256	912290	0.628604	0,530,06	0.467143	0,451791	031380	O SON ED	10000																	-						
\leq	11301200	983380	0.735.898	8106290	1050190	0.53726	0.457645	0,408455	0.355866	1000	/02/10				_	_							-			0	_	_	_			_	_	
×	11001130	983636	1300	90000	68889	51369	1,4943	988	0.000	900	23071			_	_	_			_	_	_				_	_	_	_			_	_		
New A	10504100	841432	_	SSIDE	2607290	781882		3000	-	2000	_	_		_						_	_				_			_						
Ö.	1 0001000	18863		_	_	_		_		70/00	-	_	_	_	_	-		-		_	_	-		_	_	_		=	-			_	-	
	0001709	33133		_	=	_	_	Ξ.		3 10				-	_	0 0			-	_	_	0.0		-	_	_		_	0.0	-	0.00	_	0.0	
ogram i	30000	SESSE	, 0		_	_	_	=	37268	N COLUMN					_	-		-		_	_				_	_		_			-	_		
listo	0608	38,672	762567		-	95200	_	Ĕ.	3000		-	-	_		_										_	_		_				_		
田	8.0808	813138 0	-	_	_	_	_	_	0.00000	0 0 0 0	-	-	-	-	_	-				_	_				_	_		_			-	_		
$\frac{5}{2}$	8 08057	0 12208	_	_	Ť	Ť	_	_	_	2000		-	_		_	0.088888				0		-					-	=			-	_	-	
for	00.70	200 1922		_	-	ŭ	_	_	<u> </u>	-	12655	_				3 0	-		-	_	_	-	-	-	_	_	-	=				_	-	
$^{\circ}$	F	80 AST80	_	Ť	-	_		_		_		_		_	IO.	0.0000000000000000000000000000000000000		-	-	_	_	0.0	-	-	_	0	-	=		0	0.0	_	0.0	
Tab	02020	-		ŭ	-	Ĕ	_	_	118747 0.42	_	-	-	-	-		88.0	-	= =	-			0.0					-	=	00		0.00	0	0.0	-
6:	0.0910	G 0.8311.84		_	-	-	-	_			-		21	101	-		- 1		-	0	0	- 0			0		_	=			-	-	- 0	-
ble	2000	0.796703	_	÷	Ī	<u> </u>	0.57483	2000	0.000	U.S. BERK		_	_	_	89 0.00.014SC		-		-	-	_	-			_		_	=					-	
Ta	200530	0.86.86	127.127.0	0.72838	0,655512	0,000	0.512134	0,431607	0.333191		010010	0130010		_	0.00573390		-		. =	_	_				_			=						
	93050	710217	0.68130	0.638381	0,57533	0.516755	0.445354	0.30223	0.327250	02/200	0.148818	n Index	0.050,082		0,00182983		-											-						
	00710	1968161	50000	53E281	7220010	746873	78487	7,232,0	28685	7,46,00	0.000	DAING:	1622501	DIFFERE	_	_			_	_	_	_		_	_	_	_	_	_			_	_	
	20000	20005	781622	414129	123521	1882	2303	575008	23525	1/208	010240	902290	=	=	_					_	_			_	_	_		_				_		
	300550	121381	0.353812	0,339000 0	319025	33.257 0	2681	33000	9	0.13/218					-	-		-		_	_	-			_	_		_			-	-		
	20.30	38386		35597 03	_	_	_	=		University U.					-	-	0 1			0	0	-	-		0	0	-	=			-	_	-	
	200200	274470	-	-	_	ŏ			0.044774 0.1		3 2	-			_	-			-			0.0	-				-	=	-		0.0	_	-	-
	t		-	-	-		10 99280	8	88	3 .	0 0		0		-	- 0				0	0	-	-		0		-	=		0 0	-	_	-	
	10.150 150.20	220000	6	000	7200	0.019	8	_	-	= 0		-			-	- 0	- 1		-	_	_	-	-		_		-	=			0.0	-	-	
	30100	00		-	-	_	-	_	-	= 0	-	-				-	-	-	-	-	-	-			-		_	=	-				-	
	0.50 MeV/c 3	9.0		0	0	_	ا د	٥	ە د	_ 0	- 0				_	٥ ك	- 1	_ 0	_	0	0	90			0	0	ا ف	_	ه د		- 0		90	
	(BS 0.3			<u>۔</u>	<u>۔</u> عر	<u>۔</u>	= 1 E	- 2°	= c	= c	e e	9 2	- 2	0	<u>۔</u>	e e	20	2° 1	-	0	D. S.	Deg o	54	o d	Deg	Deg 0	e i	n Sari	0 0 0 0 0 0		0.0	Deg	Deg o	
	New ANM B.S.	0.45 Deg	9-135 Deg	13548 L	18.22.5 De	22527 De	Zi a 5 Deg	31.5-36 Deg	36.40.5 Deg	distributed in	00550 Dec	54.85 De	38,5,63 De	63.67.5 Des	67.5-72 Deg	72.85 De	10000	N 800 Dec	90,915 Dec	945.99 Deg	99 IGS D-g	1035108 Deg	1125.117 Dec	117.121.5 Deg	121.5126	126130.5 Deg	1305135 Deg	Barren Barren	1305144 Deg	1001000	153 157 5 Dec	157.5.162 Deg	1621665 Deg	120 125 5 Dec