# SciBooNE Charged-Current Coherent Pion Production Acceptance Study Technical Note

Jonathan Asaadi<sup>1</sup> and Zachary Williams<sup>1</sup>

<sup>1</sup>Department of Physics, The University of Texas at Arlington

June 5, 2017

# 1 Introduction

This document is intended to serve as a reference for the acceptance study performed for the SciBooNE charged current coherent pion production (CC-Coh  $\pi^{+/-}$ ) re-analysis, as well as provide documentation of the code used in this study (in the event anything needs to be revisited in the future). The code resides in the github repository labeled as and linked here: SciBooNE-MC. The corresponding ROOT files that were used in this acceptance study can be downloaded from here: SciBooNE-MC-ROOTFiles.

The paper is structured such that Section 2 outlines Monte Carlo samples used in this study, Section 3 describes the SciBooNE detector as it was simulated in this study, Section 4 describes the various event samples that were used to both validate and generate the acceptance studies for the CC-Coh  $\pi^{+/-}$  sample. Section 5 gives a high level summary of the results including the event-reduction tables as well as the CC-Coh  $\pi^{+/-}$  acceptance results.

The appendix is left to explain how the code is run and the details of the scripts within. The appendix also details the order in which the macros should be run in, and the important plots that each macro produces that play a role in making the plots shown in Section 5 (the Results section).

## 1.1 Goal of the Re-Analysis

The goal of the re-analysis is to examine the acceptance modeling for the SciBooNE results in the presence of modern neutrino generators and updated models in order to hopefully shed light on why SciBooNE did not observe charged-current coherent pion production at low neutrino energy.

This study is intended to examine the effects of the acceptance modeling for a sample of coherent pion interactions inside the SciBooNE detector and compare what these would have been for various coherent pion production models. We utilize a simple, but robust, simulation of the SciBooNE detector and the NEUT neutrino generator to select and classify these neutrino events.

# 2 Samples

Five different samples were used in this study, three samples were generated in neutrino mode ( $\nu$ -mode) and two samples in antineutrino mode ( $\bar{\nu}$ -mode.)<sup>1</sup> Table 2 summarizes these samples. Details on these samples can be found in the Appendix.

<sup>&</sup>lt;sup>1</sup>All of these samples were generated by Callum Wilkinson (Thanks, Callum!)

#### **Summary of Samples**

Mode	NEUT version	Pion-Model	Number of simulated events
$\overline{\nu}$	5.3.6	Rein-Sehgal	1,000,000
$\overline{\nu}$	5.3.6	Berger-Sehgal	1,000,000
$\overline{\nu}$	5.0.1	Rein-Sehgal	100,000
$\bar{\bar{\nu}}$	5.3.6	Rein-Sehgal	1,000,000
$\bar{\bar{\nu}}$	5.3.6	Berger-Sehgal	1,000,000

Table 2: Summary of the samples used to build the acceptance model for this study.

## 3 Detector Simulation

This section is intended to detail the detector simulation done in this acceptance model, and to describe the assumptions made in order to accomplish accurate classifications of simulated events as charged-current coherent pion production.

#### 3.1 The Detector

For the purposes of this acceptance study, the SciBooNE experiment is composed of two sub-detectors. The first (and the more upstream) of the sub-detectors, is the Scintillator Bar Tracker (SciBar) which was originally conceived and constructed to function as the near detector for the K2K experiment. The second (and more downstream) of the sub-detectors, is the Muon Range Detector (MRD), which is the detector designed and constructed specifically for SciBooNE for measuring the momentum of muons produced from charged-current neutrino interactions up to  $1.2 \ GeV/c$  by using the observed range of the trajectory of the muon. The coordinante system used throughout this study, and illustrated in Figure 3.1, puts the origin in the lower corner of the SciBar detector, has z along the beam direction, y opposite to gravity, and x to beam left.




Figure 3.1: Representation of the SciBooNE detector and the coordinate frame we use in this study

#### 3.1.1 The Scintillator Bar Tracker (SciBar)

The Scintillator Bar Tracker (SciBar) sub-detector is a scintillator detector which was used to identify neutrino interactions within SciBooNE. The dimensions of the SciBar detector used in this simulation are 0 < x < 3.0 m, 0 < y < 3.0 m, and 0 < z < 1.7 m. This simulation models the scintillator materials as having a constant energy deposition per unit length (dE/dx) for both

muons and pions of 2.04 MeV/cm based on previous SciBooNE analyses and on mean values for typical particle momentum listed in the particle data group (PDG).

## 3.1.2 The Muon Range Detector (MRD)

The Muon Range Detector (MRD), depicted in Figure 3.1.2, is located 0.55 m downstream of SciBar in the z-direction, and is a composition of two sets of thirteen alternating slabs of steel-scintillator layers, where the scintillator layers alternate between being horizontally oriented or vertically oriented, in the xy-plane. The steel layers have a z-direction thickness of 5.08 cm and the scintillator layers have a z-direction thickness of 0.6 cm. Combining all the layers of the different alternating materials results in 26 scintillator layers that "sandwich" twenty five steel layers inbetween and gives a total z-direction dimension of being 1.37m. The xy-plane is modeled as a square again (as was the case with SciBar, too) with dimensions in the x-direction and the y-direction of 2.6 m. The energy deposition per unit length (dE/dx) of a muon penetrating the scintillator layers is assumed to be a constant 2.04 MeV/cm while the energy deposition for the muon in the steel layers is assumed to be a greater value of 11.43 MeV/cm. Both values are typical for muons at the energy range produced in SciBooNE and taken from the PDG.




Figure 3.1.2: Depiction of the Muon Range Detector (MRD) which consists of alternating layers of horizontal scintillator (shown in pink) steel slabs (shown in grey) and vertical scintillator (shown in purple)

#### 4 Event Selection

Two main samples are used in this study to generate the acceptance tables. The first is a charged current inclusive (CC-Inclusive) sample which requires a muon was created in the neutrino interaction and this muon intersects the MRD. This sample is described in Section 5.1 and is used to validate the building of the acceptance model by comparing it to previous SciBooNE analyses.

The second sample is the charged current coherent pion (CC-Coh  $\pi^{+/-}$ ) sample which requires a muon and charged pion are created in the neutrino interaction exclusively (e.g. no other final state particles in the event). This sample is described in Section 5.2.

Both of these samples are selected using NEUT MC-truth flags which ensure we are treating pure samples which are classified by the neutrino generator as belonging to the appropriate sample.

Whether or not the event identified by our selection makes it into the final sample used in the acceptance study depends on the behavior of the muon with respect to the MRD. A muon which enters the MRD from a neutrino interaction will either come to stop in the MRD, exit out the back of the MRD (assuming it's momentum is great enough), or exit out the side of the MRD. In the next sections we explain this classification further.

# 4.1 Muon Stops within the MRD ("Stopped")

The requirement to classify a neutrino interaction as a "stopped" event requires the muon from the interaction to have reached the MRD, penetrated at least three layers of steel (giving activity in three layers of scintillator), and to then deposit all of its remaining energy prior to reaching a boundary of the MRD. An illustration of a CC-Coh  $\pi^{+/-}$  event which would be classified as "stopped" is shown in Figure 4.1.




Figure 4.1: Depiction of an event that was classified as "Stopped."

These events allow for complete reconstruction of the muon's momentum based on the number of layers which the muon penetrated and the muons incident angle.

#### 4.2 Muon exits out the back of the MRD ("Out-the-back")

The classification of a neutrino interaction as "out-the-back" requires that the muon from the interaction to have reached the MRD and to have had sufficient energy to have exited out the back face of the MRD without stopping. An illustration of such an event is given in Figure 4.2.




Figure 4.2: Depiction of an event that was classified as "out-the-back".

The exact momentum of muons which pass completely through the MRD could not be made in reconstruction, so these events were classified as having the minimum energy required to penetrate all the steel and scintillator layers of the MRD.

## 4.3 Muon exits out the side of the MRD ("Out-the-side")

The classification of a neutrino interaction as "out-the-side" requires that the muon from the interaction reached the MRD, penetrated at least three layers of steel, and then to have exited out the side of the active volume of the MRD (excluding the very back face). Events which are classified as "out-the-side" are excluded from this study because no accurate reconstruction of the muons momentum can be made when the muon exits out the side of the MRD. An illustration of such an excluded event which exits out the side of the MRD is given in Figure 4.3.




Figure 4.3: Depiction of an event that was classified as "Out-Side."

# 5 Results

The results of this acceptance study can be broken down into two different classification schemes of events. Those that met the conditions to qualify as CC-Inclusive events, and those that met the

conditions of classification as Charged-Current Coherent Pion events. The former is used to validate the acceptance modeling and detector simulation to reasonably reproduce previously published CC-Inclusive studies from SciBooNE, while the latter is used for the reanalysis.

#### 5.1 Charged-Current Inclusive Events

Here we define the charged current inclusive sample (CC-Inclusive) which we use to validate our acceptance model against previously performed simulation studies.

#### 5.1.1 $\nu$ -mode Charged-Current Inclusive Events

Table 5.1.1 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the neutrino mode ( $\nu$ -mode) Monte Carlo.

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal	
Total Sample	1,000,000	1,000,000	100,000	
CC-Inclusive Interaction	725,730	727,278	69,363	
$\mu$ ( $\mu$ + n-other particles in SciBar)				
Muon enters the MRD	263,698	262,608	24,250	
Muon enters the MRD and	231,089	230,054	21,001	
penetrates $\geq 3$ layers of steel				
"Stopped"-Events	177,406	175,799	16,062	
"Out-the-back"-Events	15,389	15,952	1,421	
"Out-the-side"-Events	38,294	38,303	3,518	
Good CC-Inclusive Events	192,795	191,751	17,483	

 $\nu$ -mode CC-Inclusive Event Reduction

Table 5.1.1: Event reduction table for a sample of  $\nu$ -mode CC-Inclusive events simulated in the SciBooNE geometry.

The first quantity that is calculated for the different events is the momentum of both the muon and the pion, which are both found from the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{1}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{2}$$

where  $|\vec{p}_{\mu}|$  represents the magnitude of the momentum for the corresponding particle, and  $P_{\mu_x}$  represents the component of the four momentum for the corresponding particle. The momentum is reported in units of MeV/c.

The next quantity calculated is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1} \left( \frac{\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}}{P_{\mu_z}} \right) \tag{3}$$

$$\theta_{\pi} = tan^{-1} \left( \frac{\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}}{P_{\pi_z}} \right) \tag{4}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of charged-current coherent pion production, the angle should never be larger than 90°.

Figure 5.1.1 shows the momentum and angular ( $\theta$ ) distribution for the sample of  $\nu$ -mode CC-Inclusive events passing all our requirements for all three models considered in this study (NEUT v5.3.6 Rein-Sehgal, NEUT v5.3.6 Berger-Sehgal, NEUT v5.0.1 Rein-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 5.1.1: Muon Momentum (left) and Muon Angle (right) for  $\nu$ -mode CC-Inclusive interactions for all three models included in this study. These samples kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive

Figure 5.1.1 represents the one-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events for this study using all three different models compared to results derived from Hiraide's thesis <sup>2</sup> using the full SciBooNE Monte Carlo simulation. A few reference points are illustrated using dashed lines to guide the readers eye. A few perecent difference is seen, but overall agreement between the two simulations hold.

<sup>&</sup>lt;sup>2</sup> Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide dt.pdf




Figure 5.1.1: One-dimension efficiency plots for the  $\nu$ -mode CC-Inclusive sample.

Figure 5.1.2 shows the two-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events. The left hand side is a reference plot provided by Morgan and the right hand side is for the Rein-Sehgal MC used in this study.

Below are the two dimensional efficiency plots for the CC Inclusive events in  $\nu$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section.




Figure 5.1.2: Two-dimensional efficiency plots for the  $\nu$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 5.1.2: Two-dimensional efficiency plots for the  $\nu$ -mode Berger-Sehgal CC-Inclusive sample.




Figure 5.1.1: Two-dimensional efficiency plots for the  $\nu$ -mode Old Rein-Sehgal CC-Inclusive sample.

## 5.1.2 $\bar{\nu}$ -mode Charged-Current Inclusive Events

Similar to before, Table 5.1.2 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the antineutrino mode ( $\bar{\nu}$ -mode) Monte Carlo.

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal			
Total Sample	1,000,000	1,000,000			
CC-Inclusive Interaction	699,239	704,327			
$\mu$ ( $\mu$ + n-other particles in SciBar)					
Muon enters the MRD	380,362	380,869			
Muon enters the MRD and	336,373	337,979			
penetrates $\geq 3$ layers of steel					
"Stopped"-Events	288,289	288,206			
"Out-the-back"-Events	7,608	7,857			
"Out-the-side"-Events	40,476	41,916			
Good CC-Inclusive Events	295,897	296,063			

 $\bar{\nu}$ -mode CC-Inclusive Event Reduction

Table 5.1.2: Event reduction table for a sample of  $\bar{\nu}$ -mode CC-Inclusive evnets simulated in the SciBooNE geometry.

Figure 5.1.1 shows the momentum and angular distribution for the sample of  $\bar{\nu}$ -mode CC-Inclusive events passing all our requirements for both models considered in this study (NEUT v5.3.6 Rein-Sehgal, and NEUT v5.3.6 Berger-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 5.1.2: Muon Momentum (left) and Muon Angle (right) for  $\bar{\nu}$ -mode CC-Inclusive interactions for all three models included in this study. These samples kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive

Figure 5.1.2 represents the one-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events for this study. No similar reference sample exists to be compared directly against, however we note that the shape and magnitude of the acceptance is nearly unchanged between  $\bar{\nu}$  and  $\nu$ -mode samples (as expected).




Figure 5.1.2: One-dimension efficiency plots for the  $\bar{\nu}$ -mode CC-Inclusive sample. Muon's Momentums is on the right and the Muon's Angles is on the left.

Figure 5.1.2 shows the two-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events for this study compared to results derived from Morgan's reference sample.

Below are the two dimensional efficiency plots for the CC Inclusive events in  $\bar{\nu}$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section.




Figure 5.1.2: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 5.1.2: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode Berger-Sehgal CC-Inclusive sample.

## 5.2 Charged-Current Coherent Pion Production Events

Here we define the Charged-Current Coherent Pion Production sample (CC-Coh  $\pi^{+/-}$ ) which we use to validate our acceptance model against previous simulation studies which were done.

#### 5.2.1 $\nu$ -mode Charged-Current Coherent Pion Events

Table 5.2.1 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi^{+/-}$  events from the neutrino mode ( $\nu$ -mode) Monte Carlo.

#### ν-mode CC-Coherent Pion Event Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal
Total Sample	1,000,000	1,000,000	100,000
CC-Coherent Pion Interaction	12,186	2,576	1,320
$(\mu + \pi + \varnothing \text{ in SciBar})$			
Both muon and pion are	8,535	1,845	884
forward going			
Muon enters the MRD and	7,407	1,592	767
penetrates $\geq 3$ layers of steel			
"Stopped"-Events	6,448	1,350	669
"Out-the-back"-Events	530	150	56
"Out-the-side"-Events	429	92	42
Good Coherent Pion Events	6,978	1,500	725

Table 1: Event reduction table for a sample of  $\nu$ -mode Charged Current Coherent Pion events simulated in the SciBooNE geometry.




Figure 1: Muon Momentum for all of the muons of the events that made it to the MRD and penetrated at least three layers (left) and Muon Angle for the muons of the events that made it to the MRD and penetrated at least three layers (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study. The "Total" classification means that all CC-Coh  $\pi^{+/-}$  events are included in these histograms.




Figure 2: Muon Momentum of both the "stopped" and "not-stopped" samples (left) and Muon Angle of both the "stopped" and "not-stopped" samples (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study. The "Good" classification means that only the stopped and not-stopped CC-Coh  $\pi^{+/-}$  events are included for these histograms.

The last two quantities that are calculated are the two different types of four-momentum transfers

specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{5}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(6)

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{7}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(8)

|t| is reported in units of  $(MeV/c)^2$ .

 $\nu$ -Mode |t| and  $Q^2$  plots are below:




Figure 3: The |t| Momentum Transfer for the "stopped" and "not-stopped" events (left) and  $Q^2$  Momentum Transfer for the "stopped" and "not-stopped" events (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for the three models included in this study.




Figure 5.2.1: These two plots are the one dimensional efficiency plots that make the two dimensional efficiency plots that follow. The left plot is the muon angle efficiency plot and the right is the muon momentum efficiency plot for  $\nu$  mode.

Below are the two dimensional efficiency plots for the CC Coherent Pion events in  $\nu$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section.




Figure 5.2.1: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.




Figure 5.2.1: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Berger-Sehgal CC-Coherent sample.




Figure 5.2.1: Two-dimensional efficiency plot for the old NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.

## 5.2.2 $\bar{\nu}$ -mode Charged-Current Coherent Pion Events

Similar to before, Table 5.2.2 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi^{+/-}$  events from the anti-neutrino mode ( $\bar{\nu}$ -mode) Monte Carlo.

Below are the plots for CC-Coh  $\pi^{+/-}$  Events for  $\bar{\nu}$ -mode. The layout of the rest will be very similar to  $\nu$ -mode, and the equations used previously are the same equations used for the plots below.

## $\bar{\nu}$ -mode CC-Coherent Pion Event Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal
Total Sample	1,000,000	1,000,000
CC-Coherent Pion Interaction	36,669	7,790
$(\mu + \pi + \varnothing \text{ in SciBar})$		
Both muon and pion are	24,675	5,477
forward going		
Muon enters the MRD and	20,445	4,517
penetrates $\geq 3$ layers of steel		
"Stopped"-Events	18,935	4,203
"Out-the-back"-Events	372	82
"Out-the-side"-Events	1,138	232
Good Coherent Pion Events	19,307	4,285

Table 2: Event reduction table for a sample of  $\bar{\nu}$ -mode Charged Current Coherent Pion events simulated in the SciBooNE geometry.




Figure 4: Muon Momentum (left) and Muon Angle (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study.

The structure of the plots in Figure 5.2.2: very closely resembles the plots for the  $\nu$ -mode above, and the rest of the plots in this section have that same characteristic.




Figure 5: Muon Momentum (left) and Muon Angle (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi^{+/-}$  interactions for both the "stopped" and "not-stopped" samples of events.

 $\bar{\nu}$ -mode |t| and  $Q^2$  plots are below. They also have the same overall shape as the plots for  $\nu$ -mode above.




Figure 6: The |t| Momentum Transfer (left) and  $Q^2$  Momentum Transfer (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi^{+/-}$  interactions for both of the models included in this study which are the "stopped" and "not-stopped" events.




Figure 5.2.2: These two plots are the one dimensional efficiency plots that make the two dimensional efficiency plots that follow. The left plot is the muon angle efficiency plot and the right is the muon momentum efficiency plot for  $\bar{\nu}$  mode.

Below are the two dimensional efficiency plots for the CC Coherent Pion events in  $\bar{\nu}$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section.




Figure 5.2.2: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Rein-Sehgal CC-Coherent sample.




Figure 5.2.2: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Berger-Sehgal CC-Coherent sample.

# A Appendix: Sample Details

Appendix on samples

## A.1 $\nu$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$  interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

### A.2 $\nu$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$  interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack\_NEW.root

found at the following link (SciBooNE-MC-ROOTFiles).

## A.3 $\nu$ -Mode Rein-Sehgal NEUTv5.0.1

A sample of  $100,000 \nu$  interactions were simulated using the NEUT generator (v5.0.1, believed to be the version used by the SciBooNE collaboration in the original publication) and the corresponding older Rein-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

# A.4 $\bar{\nu}$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$  interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

# A.5 $\bar{\nu}$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$  interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack\_NEW.root

found at the following link (SciBooNE-MC-ROOTFiles).

#### A.6 Vertex Distributions

The events were all given a random initial point that was generated with the goal that the vertex distributions of this simulation would closely match the vertex distributions that Hiraide <sup>3</sup> showed in his thesis. This was done using the code below and a sample plot for one of the samples is shown below the code.

```
\label{eq:trandom3} \begin{array}{lll} TRandom3 * randX = new & TRandom3(); \\ TRandom3 * randY = new & TRandom3(); \\ TRandom3 * flat = new & TRandom3(); \\ randX->SetSeed(jentry/2); \\ randY->SetSeed(jentry*jentry); \\ flat->SetSeed(jentry*jentry*jentry); \\ \textbf{double} & Xpos = randX->Gaus(1.5,1.3); \\ \textbf{while} & (Xpos<0 \mid \mid Xpos>3.0) & Xpos = randX->Gaus(1.5,1.3); \\ \textbf{double} & Ypos = randY->Gaus(1.5,1.05); \\ \textbf{while} & (Ypos<0 \mid \mid Xpos>3.0) & Ypos = randY->Gaus(1.5,1.05); \\ \textbf{double} & Zpos = flat->Uniform(0,1.7); \\ \end{array}
```




Figure 7: Vertex distributions of the events in the new Rein-Sehgal sample in  $\nu$ -mode.

## A.7 NewNMReinSehgal.C

This file is the macro that corresponds to the "NewNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms). When this macro is run (which can take a while), it also plots a few different histograms. The histograms that are plotted are the ones shown in the figures below with descriptions included with the corresponding figures. The order that the histograms appear in this paper is the same order they will be shown when this macro is run in root.

<sup>&</sup>lt;sup>3</sup>Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide dt.pdf




Figure 8: Vertex distributions of the events in the new Rein-Sehgal sample in  $\nu$ -mode.




Figure 9: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Inclusive events in the new Rein-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Inclusive.




Figure 10: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Coherent Pion events in the new Rein-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Coherent Pion.

The NewNMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{9}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{10}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1} \left( \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2} / P_{\mu_z} \right) \tag{11}$$

$$\theta_{\pi} = tan^{-1} \left( \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2} / P_{\pi_z} \right) \tag{12}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{13}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(14)

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{15}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
 (16)

|t| is reported in units of  $(MeV/c)^2$ .

# A.8 NewNMBergerSehgal.C

This file is the macro that corresponds to the "NewNMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack\_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 11: Vertex distributions of the events in the new Berger-Sehgal sample in  $\nu$ -mode.




Figure 12: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Inclusive events in the new Berger-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Inclusive.




Figure 13: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Coherent Pion events in the new Berger-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Coherent Pion.

## A.9 OldNMReinSehgal.C

This file is the macro that corresponds to the "OldNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoOBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 14: Vertex distributions of the events in the old Rein-Sehgal sample in  $\nu$ -mode.




Figure 15: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Inclusive events in the old Rein-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Inclusive.




Figure 16: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Coherent Pion events in the old Rein-Sehgal sample in  $\nu$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Coherent Pion.

## A.10 NewANMReinSehgal.C

This file is the macro that corresponds to the "NewANMReinSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 17: Vertex distributions of the events in the new Rein-Sehgal sample in  $\bar{\nu}$ -mode.




Figure 18: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Inclusive events in the new Rein-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Inclusive.




Figure 19: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Coherent Pion events in the new Rein-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Coherent Pion.

# A.11 NewANMBergerSehgal.C

This file is the macro that corresponds to the "NewANMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack\_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBSBar.root" inside the

"ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 20: Vertex distributions of the events in the new Berger-Sehgal sample in  $\bar{\nu}$ -mode.




Figure 21: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Inclusive events in the new Berger-Sehgal sample in  $\bar{\nu}$  mode. The left only contain events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Inclusive.




Figure 22: These are the two two dimensional histograms that when the left is divided by the right it returns the two dimensional efficiency histogram for the CC Coherent Pion events in the new Berger-Sehgal sample in  $\bar{\nu}$  mode. The left only contain events that stopped or went out the back and penetrated the front face of the MRD. The right contains all of the events that were classified as CC Coherent Pion.

## A.12 NMCombinedPlots.C

This is the file that performs the main plotting operations for the neutrino mode samples using the muon's information. All of the muon efficiency plots for neutrino mode are made with this file.




Figure 23: These are the  $\nu$  mode one dimensional efficiency plots of the two dimensional CC Inclusive efficiency plots that are below. The left is the angle efficiency plot and the right is the momentum efficiency plot for the muon.




Figure 24: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 25: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Berger-Sehgal CC-Inclusive sample.




Figure 26: Two-dimensional efficiency plot for the old NEUT  $\nu$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 27: These are the two plots for muon angles in  $\nu$  mode that are used to make the one dimensional efficiency plot for the angles in CC Coh Pion events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either stopped or went out the back face. The right plot is the muon angles for all of the events that were classified as CC Coh Pion. The left divided by the right gives the efficiency plot for the muon angles in CC Coh Pion events.




Figure 28: This is the one dimensional muon angles efficiency plot for the CC Coh Pion events from the  $\nu$  mode samples.




Figure 29: These are the two plots for muon momentums in  $\nu$  mode that are used to make the one dimensional efficiency plot for the momentums in CC Coh Pion events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either stopped or went out the back face. The right plot is the muon momentums for all of the events that were classified as CC Coh Pion. The left divided by the right gives the efficiency plot for the muon momentums in CC Coh Pion events.




Figure 30: This is the one dimensional muon momentums efficiency plot for the CC Coh Pion events from the  $\nu$  mode samples.




Figure 31: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.




Figure 32: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Berger-Sehgal CC-Coherent sample.




Figure 33: Two-dimensional efficiency plot for the old NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.

# A.13 NMPionPlotting.C

This is the file that performs the main plotting operations for the neutrino mode samples using the pion's information. All of the pion efficiency plots for neutrino mode are made with this file.




Figure 34: These are the angles of the pions in CC Coh Pion events in the  $\nu$  mode samples. The left plot is the angles for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion angles for all of the events that were classified as CC Coh Pion.




Figure 35: These are the momentums of the pions in CC Coh Pion events in the  $\nu$  mode samples. The left plot is the angles for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion momentums for all of the events that were classified as CC Coh Pion.




Figure 36: These are the energies of the pions in CC Coh Pion events in the  $\nu$  mode samples. The left plot is the energies for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion energies for all of the events that were classified as CC Coh Pion.

# A.14 NMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for neutrino mode.




Figure 37: The |t| Momentum Transfer for the "stopped" and "not-stopped" events (left) and  $Q^2$  Momentum Transfer for the "stopped" and "not-stopped" events (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for the three models included in this study.

#### A.15 ANMCombinedPlots.C

This is the file that performs the main plotting operations for the anti-neutrino mode samples using the muon's information. All of the muon efficiency plots for anti-neutrino mode are made with this file.




Figure 38: These are the  $\bar{\nu}$  mode one dimensional efficiency plots of the two dimensional CC Inclusive efficiency plots that are below. The left is the angle efficiency plot and the right is the momentum efficiency plot for the muon.




Figure 39: Two-dimensional efficiency plots for the  $\bar{\nu}$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 40: Two-dimensional efficiency plots for the  $\bar{\nu}$ -mode Berger-Sehgal CC-Inclusive sample.




Figure 41: These are the two plots for muon angles in  $\bar{\nu}$  mode that are used to make the one dimensional efficiency plot for the angles in CC Coh Pion events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either stopped or went out the back face. The right plot is the muon angles for all of the events that were classified as CC Coh Pion. The left divided by the right gives the efficiency plot for the muon angles in CC Coh Pion events.




Figure 42: This is the one dimensional muon angles efficiency plot for the CC Coh Pion events from the  $\bar{\nu}$  mode samples.




Figure 43: These are the two plots for muon momentums in  $\bar{\nu}$  mode that are used to make the one dimensional efficiency plot for the momentums in CC Coh Pion events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either stopped or went out the back face. The right plot is the muon momentums for all of the events that were classified as CC Coh Pion. The left divided by the right gives the efficiency plot for the muon momentums in CC Coh Pion events.




Figure 44: This is the one dimensional muon momentums efficiency plot for the CC Coh Pion events from the  $\bar{\nu}$  mode samples.




Figure 45: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Rein-Sehgal CC-Coherent sample.




Figure 46: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Berger-Sehgal CC-Coherent sample.

## A.16 ANMPionPlotting.C

This is the file that performs the main plotting operations for the anti-neutrino mode samples using the pion's information. All of the pion efficiency plots for anti-neutrino mode are made with this file.




Figure 47: These are the angles of the pions in CC Coh Pion events in the  $\bar{\nu}$  mode samples. The left plot is the angles for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion angles for all of the events that were classified as CC Coh Pion.




Figure 48: These are the momentums of the pions in CC Coh Pion events in the  $\bar{\nu}$  mode samples. The left plot is the momentums for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion momentums for all of the events that were classified as CC Coh Pion.




Figure 49: These are the energies of the pions in CC Coh Pion events in the  $\bar{\nu}$  mode samples. The left plot is the energies for the pions where the muon made it to the MRD and either stopped or went out the back face of the MRD. The right plot is the pion energies for all of the events that were classified as CC Coh Pion.

## A.17 ANMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for anti-neutrino mode.




Figure 50: The |t| Momentum Transfer (left) and  $Q^2$  Momentum Transfer (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi^{+/-}$  interactions for both of the models included in this study which are the "stopped" and "not-stopped" events.

## B Efficiency Tables

These are the corresponding tables to the 2D Efficiency plots for CC-Inclusive events.

Table 3: Table for 2D Histogram for New NM-Rein-Sehgal

1930-2000	82323	0.7857.4	0.713615	2011/00	0.5625	0.435185	9-829-0	0.380932	033333		_		_	_	_		_		_						_	_	_	_	_	_						_		
1900-1950		_	_	0.630252		_				2							_		_				. =		_													
830-19III	861111	813733	73029	938989	589507	462963	2000	290323	52		_	_		_	_	_				_	_			_			_	_	_	_	_	_	_			_		
810-1850	848-85	3337	68-4982	1282910	27.047	128937	102941	2772	982-120	-	_	_		_	_	_								_			_	_	_	_	_	_				_		
1730-1800 1	77.93	770.94						ĸ	22		22	-	_	_	-	_								_		_	_	_	_	_	_	_	_	_	_	_	-	
1 02100	921369	5477	_	1834561	_	_	=	=	_	0 29999r											В													B			0 1	
1630 1700 1700 1750	۲	-	=	6.8718 0.4	=	÷	=	=	19	1875 0.	0	8		0	8	0	-		-	0	0			0	-		0	0	0	0	0	0	0				0 1	
	۲	_	_	90,8338 0.6	_	_	=	_	_	_	0			0		0	=		=	0	0		-	0	=		0	0	0	0	0	0	0					0.0
1530 1600 1800 1650	3508 0.8	54091 0.7	_	_	-	_	-	293578 0.3	-	1006060		0	-		0		=	-	=			-			=	-										-	-	
1309-1350 155	35333 BB	72224 8.7	_	_	_	331166 05:	_	29-018 0.2	_	8 8	9.0	В		0	В	0	=		=	0	0		-	0	=		0	0	0	0	0	0	0				-	
1450-1500 1300	-	-	716946 0.72	_	÷	÷	384615 0.48	-	_	303 0.16	0.759231 0						=		=			-			=												-	
1409-1-60 1-430	9	_	72093 0.71	_		_	_	_	_	1369014 0 30303	Ï	7.0	в	0		0	=		=	0	0	=	-	0	=	в	0	0	0	0	0	0	0				-	
	-	_	_	_	_	_	=	_	÷	÷	÷	0	-	0	0	0	=	-	=	0	0	0	-	0	=	-	0	0	0	0	0	0	0				-	
1339-1400		36 0.774036	24 0.7216S3	_		_	=	0.307116	=	24 0.1-5039	_			0		0	=	-	=	0	0	=	-	0	=		0	0	0	0	0	0	0				-	
300 1300 1350	98 0.81321	S2 0.8011	_	_	Ĕ	Ť	Ĕ	Ĕ	_	18 0.395524	_	37 8.2	=				=		=		0	-	=		=												-	
250 1230-1300	F	÷	=	35 0.681319	=	-	=	=	=	=	27 B.113636	=	=	0		0	=		=	0	0	-	-	0	=		0	0	0	0	0	0	0				-	
1200-1250	F	÷	6 0.746803	_	_	9 0.55(862	_	_			1 0.182927	_	10				=		=			-	-		=												-	
50 1130-1200	-	=	0.713326	=	_	_	=	_	=	_	7 835534	=	0 18				_		_				-		=												-	
1030-1100 1100-1150	0.88785	_	0.746401	_	Ĕ	Ť	=	Ĕ	-	_	10.216867	-	_				=		_				-		=												-	
	10	_	_	0.713143	_	=	=	_	=	_	_	=	_				_		_			-	_		_													
1000-1050	0.83-632	0.803922	97050	B.78-62	0.65(11)68	0.57836	0.48873	0.62253	0.373851	_	0.144772	-	_	=	0.136364		_		_			-	-		_												-	
950-1000	0816649	0.79823	0.778828	0.722747	0.6566-48	0578595	0503736	0.459893	0362308	0.28-258	0.221135	0.12-514	0.0814815	0.0-405-405			_		_			-	. =		_													
056-006	987694	0.809981	0.763183	B.71337E	96736	12,002,00	0.524616	0.439182	0371585	0.293674	0238666	0.140187	_	_	0.02-6502		_		_			-	. =		_													
N50-900	1198621	0.823840	0.784038	0.71437	0.647185	0.58894	1532462	0.455387	0378892	0.299735	0.239492	0.177469	0.106557	0.0883721	92011900	200	_		_				-		_													
800-850	8/292/0	0.816514	10527.0	0.788637	961-199	0.601285	0.542526	0.465472	0.404237	0.312566	0.232889	0.207	708210	0.074184	0.0347222	0.0131579	_		_				. =		_													
250-810	288	177708.0	0.757746	0.713192	0.675-0	0.617109	0.5-6332	0.485242	8229070	0.324022	0.2383-57	0.153939	0.133188	0.0654045	0.04586	0.0121212	_		_				. =		_													
200 230	0.773438	0.828302	95522	1127121	0.67305	0.61255	0537204	1512734	0.42(8(8	033975	027558	9188606	0.135764	0.08-6.225	88952468	0.0234114	0.00546541																					
659 700	18-6-455	180404	1786013	175131	2011691	1638694	535136	12	3,410,602	13-28/2	0.269592	1214009	2281210	10883686	0.044898	0030135	_	_	_	_	_	-	-	_	_	_	_	_	_	_	_	_	_	_	_	_		
029-009	8.2975	841185	784336	-	681216	296259	57336	=	=	_	0.260561	_	_	_	_	_		_					_	_		_	_	_	_	_	_	_			_	_		
8	8829	2196	57402	8228	3245	86861	98	9882	38772	777	8649	310165	31-69	29918	83498	21873	-	-	_	0					_	_							0		_	_		0.0
0.550	816092 0	824047	79547	0.73062 0.7	0.00007	200221 0	66298	445 G3 B	35271	3787	220634 0	165696 0	109863	0.74586 0	1809081	0.41686 0				0	0						0	0	0	0	0	0	0			_		
450-300	0 0:091	0 20199	22122	8363	67.53	13-469	0.455	976	9229	9322		16142			018006 0	0 1228571				0	0			0			0	0	0	0	0	0	0			_		
Ė	2:0 9:8	878	9.0	8122 118	252 8.5	342.4 8.5	293	889 0.3	828 838	549 875	20 122	5702 0.3	_		0.00045756 0.0	8.8	=		=	В	В	-	-	В	=	-							0				-	
E1-00- Ot	822	150 152	3167 0.58	102 052	0.49	223 0.44	982 038	3465 032	B74 127	173194 822	75807 0.17	1102629 0.12	_		000		=		=	0	0				=												= 1	
220-400	578 528	11 0.490	529 0.475	937 0.460	223	783 0356	517 8328	223 0260	3	=	=	0.0555327 0.103		9016			=		=			-			=												-	
10 300-350	715.0	182 0.404	0.334	302 0389	374 8328	88 0284	350 0540	3GB 0.215	=	_	=		0.024				=	-	=	0	0	-	-		=												-	
250-310	9 0.260	1 0.218182	8 831	1 12.00	3 0.22037	4 0.2E	_	_	_	_	_	0.0115123		0		0	=	-	=	0	0	=	-	0	=		0	0	0	0	0	0	0				-	
200.230	0.25925	02468	4 0.15694	3 0.16-56	5 0.13-64	_	_	_	0.0376749	0.0185547	0.00241935		0			8	=		=	8	8	-	-	8	=	0								В	В	_	0 1	
130,200	0.09375	0.123457	0.0280374	0.0696203	0.0396825	0.0340615	0.0128535	0.00618357									_		_			-	-		_													
00 100 130		0							В	0			0			8	=		=	8	8	-	-	8	=	0								В	8	_	0 1	
001-02 2/Aar	0																=		=			-	-		=									0				
2 Valvage Red Rev	0				-		=		n			-			-		=		=					Dag B	Down II	0 200	D No.	D No.	D No.	D No.	Dag B	Dag B	Dag B	0 %00	0 200	9 mg	Deg B	0 th
New NACES	8-45 Deg	45-9 Deg	9-135 Drg	135-18 Deg	18-22.5 De	225.27 De	27-31 5 De	315.36 De.	36-40.5 De	485-45 Do	45-85 De	49554 De	5438 5 De	585.63 De	63.675 Do	675-72 De	72.75 Do	765.81 Dr.	81-85.5 Do	855.90 De	98-94-5 De	945.49 De	99-103.5 D	1055181	1 108-1125 L	112.5-117 Day	117-1215 Dag	12151361	126-1395 Deg	13054351	135 1395 1	135.144 Deg	1441485 Day	148 5-133 Day	153-1575 Day	1575-1621	162 1665 1	1865.12 Day
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_

Table 4: Table for 2D Histogram for New NM-Berger-Sehgal

_																																						
1950-2001	0.83333	0.29-521	0.083030	0.62711	D.55747	0.4545	0.306835	0.137931	0.181818	_																		_			_							
1900-1950	1111160	8061620		0.621212				0,258065																		_	_		_					_		_		
1850-1900	0.79.872	0.751905	0.00338		0.573342	0.406599		0.25																				_			_		_		_	_		
800-1850	1111797	62087		0.007573	0.525.0				0,333333		_	_	_	_	_	-	_	_	-	-	-	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
1730-1800	.810324	7,0003		52,029.1	_		_		0.214286		_	_															_	_			_				_	_		
1 120	0 25,35		0.700381 0		0.52568 0	_	_	_	_	_	_	_	0	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_		_	_		_				0.0
1620-1700 17	0.0	7,0657		1637168 0.0	_		_	_	_	12	_	_	_					0										_	-		_				_	_		0 0
1600-1620 16		17286-68 0.7	-	Ē	_	_	÷	=												0			0			0	0	_		0	_	0				_	0	
1000	20 202		Ī	-	0.087136 0.05	_	-		130 25 OF 1	26216 0.4	2323 0.2	0	-	0	0	0	0	-	0	0	0	0	0	-	-	0	0	_		0	_	0	_	0	_	_	0	
1550 1551	9230 910	651 0.74	7182-40 0.71	-	-	_	_	_	8130 878	002	4286 1(33232		-	-	-	-	0	-					-	-	-			-	-	0	-	0			-	_		0 0
1450-1500 1500-1550	280 268	320	65	578 0.668863	Ť	_	_	_			0.0714286								0	0	0		0			0	0		0			0		0			0	
420 1.50	0.823			51 0.64.678	Ť	Ī	_	_			64 0.25				0	0	0		0		0		0									0					0	0 0
00 1406-1450	4 0.85	4 0.75-206	0.75333	0.059251		_	_				7 0.136364							_						_	_			_			_		_		_	_		
1350-14	0.80873	0.73299	0.728296			÷	÷				T 0.166567		_	-	-	-		_					-	_	_			_			_		_		_	_		
1300-1320	0.783133	0.813655	0.708861	0.08321		0.529101	0.431408	_	_		0.083333		_									_		_	_			_	_		_							
1230-1300	0.78022	0.775033	0.737759	0.681775	0.579366	0.524138	0.426-556	0.405963	0.38785	0.189-474	0.0388235	0.0909091	0.111111									_						_	_		_				_			
1200-1250	2580	0.784204	0.731677 0.730769	720.67	0,60516	1525667	0.474736	0.405551	0300654	0.233918	0.16201.4	0.0759231														0												
120-1200	820281	0.814336	0.717308	0.683544	-	0.528678	0.44332	0397764	71120211	0.214286	0.138879	0.097361	0.0909091							0						_	_	_	_		_				_			
1100.1130	1813725	1804694	0.733735	1700101	16-0335	1557530	1475631	1437118												0	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
1020-1100	8	0.82-20-55		_	Ī	Ī	÷	-	-	_	_							_			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
10000-1020	982	0.235	TOTOLS II	703531 0	_	Ť	_	_						0 233333	714286 0	_	_	-	-		0	_	_	_	_	_	_	_	_		_	_	_	_	_	_		0.0
920-1000 10	31 21814	22657 0	70384 0.	683544 07	÷	÷	Ť	÷					108844 0.0	2	0.0-0.057	-	0	-					-	-	-	0	0	_	-	0	_	0				_	0	-
200-020	П	_		0.702804 0.6	_	_	=	_			0.215613 0.2			0 92/82	27273 0.0		0					0				0	0	_		0	_	0						
000	8333 (18)	5.448 0.8	_	Cu some	_	581746 0.5	Ξ	_						.103825 0.0	57.8 00	0.25	0			0					-	0	0	_			_	0		0	-	_	0	0 0
				G.0 780117.0		_	_	_	_	-	-	Ť	_	Ť.											-			_		0	_							
00 800 820		80211 0.80		722-403 0.71	÷	_	_	494202 0.47549	_	_	-	_	_	1850500 850080	31221010 SE1322-0.		0.0123457 0	-	0		0			-	-	-	-	-	-	0	-		-		-	_		
720.800	Г	0	0	0	_	_	_	_	<u></u>	0	_	_	82 0.118211	_	_	0 788	=							-	-			_			_				-			
200-220	_	0.814286	_	0.752791	Ť	_	_	0.479326	_	_	Ť	_	Ť	0.006020	0.0363636	7 0.0120482	=											_			_							
			-	_	0.677215	0.615-445	0.054788	0.402452	0.425845	-	-	-	0.135583	0.000000	0.039656	_	0.00280805	_						_	_			_			_		_		_	_		
000 000	0.880435	0.805305	0.8007204	0,730152	1687027	0.629688	0.554264	0.488325	0.419254	03-25679	0.261.486	0.203203	0.120520	00780856	0.0367232	0.0195072		_					-	_	_			_	-		_		_		_	_		
220.600	0.807002	0.817109	0.738354	0.736387	0.002351	0.637657	0.354316	0.472959	0.387414	0.319185	0.244851	0.187424	0.118058	0.0796216	0.0436238	0.0114213								_	_			_	_		_				_			
200-220	0.893522	0.811448	0.78089	0.22545	20292970	0.578365	0.406522	0.486444	0.356159	0.30273	0.24064	0.169353	0.115633	0.0534228	0.0364188	735037-00.0																						
	128827	_		63504		1521715	105.001	381784	27:00	272-452	(212341	9258-13	102659	10268421	10242028	10070046																						
T	2	581731	-	566116 0.6	7113 0.5	7548 0.5	5361 0.4	3753 R3	_	_	Ť	126234 0.1	Ť	0.0457205 0.0	00745573 0.0	070	0		0	0	0	0	0			0	0	_	-	0	_	0				-		
00 400 E0	-	_	-		15.0 BIG	265 0.44	105 0.35	494 0.33	<u></u>	ď	_	_	_	0.0128637 0.04	0.00													_			_							
	Ē	0.478261		0.448833	0.41500	0.38126	0312100	0.28349	4 023358	Ť	_	_	_				0		0	0	0														-	-	0	
300.350	0.4222	0,30600	0.35296	0,33023	0.31846	0.32187	0.25338	0.21827	0.17016	0.129657	0.0040285		_	0.00071-286	0	0	0		0	0	0		0			0	0	_		0	_	0						
220-310	1325	0.235849	0.223329	0200736	0.228145	0.197461	0.172859	0.1-2252				0.011.01.21	0.000-55620-4				0									0	0					0						
П	0.05/2013	12:21	912213	131222	10973451	10930736	0.0857143	71221800	0.032338	0.01391.47	291-122007	_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_				
0.200						0.0254025 0.	0.00485618 0.	n 7545700.	_	_	_	_	_					2						_	_	_	_	_	_	_	_	_	_	_	_	-	_	
00-150 15.	-	9.0	9.0	9.0	0.0	0.6	0.0	0.6	-	-	-	-					-									-	-	-	_	-	-	-	-	-	-	_	-	-
9.10	Ē	-	-	-		-	-	-	-	-	-	-	_	-	-	-	-						-	_	_	-	-	_	_	_	_	-	_	_	_	_		_
30 MeV/c																																						
MBS B	0 201	0 00	9.135 Deg 0 0 0.05075	1 Deg	O Maria	7 Dog 0	O Dog	1 Dog 0	O Dog	O Dog	O Dog	1 Dog 0	O Mari	1 Deg	Deg 0	Deg 0	Dog 0	Deg o	Dog	Deg 0	Dog	Dog 0	5 Day	0 2 Dog 0	2.5 Dag 0	0 200 Z	1.5 Dag 0	26 Dag 0	0.5 Dag 0	35 Day 0	9.5 Dag 0	44 Deg 0	8.5 Dag 0	63 Deg 0	0 No 92	62 Deg 0	6.5 Day 0	166.5171 Day
New N	0-45 D	450 D	94351	135.18	18-22.5	225.27	27.33.5	31.5.36	36.40.5	40.5-45	45.49.5	49.5.54	54.58.5	585.63	63.67.5	676-72	20.00	765.81	81.85.5	85.5.90	98.94.5	945.99	99-103,	103.5-1	108.112	112.5-1	117-121	121.5-1.	135.13	130.5.1	135-132	130.5.1	144-1-6	148.51	133.15,	157.5.1	163.18	1 1 1

Table 5: Table for 2D Histogram for Old NM-Rein-Sehgal

_	_																																							
1950-2000	_	183	0.708333	0.365217	0.722222	0.33333	17	12																																
1908-1950	$\vdash$	0.533333		12	_	7	-	789989																																
1859-1900 1	H	-	_	_	_	0.45	52	_	_	_			-	_	-		_													_	_	_			_	_	_			_
1800-1850 183	_	85743 88				0.015385 0.4		0 2999		-						0	=									0 0	0 0	= 0	= 0	-	_		0		0		-	0		
1730-1800 180	H	_	820187 0.95	_	_	_	_	_									=	-	-	-	-	-	-				= 0	= 0	= 0	=						-				
	F	H 82317	_	Ť	_	_	=	29 8.22222									=				-						= 0	= 0	= 0	=										
178-178	-	-	_	89 0	7 0.625	_	-	0.571429	-								-			-							= 0	= 0	= 0	=	=						_			
0 1630-1700	_	823529		0.63625	0.42307	=	0.454545	-	0.285714								-				-									=							_			
1638-1630	2999980	88	0.553333	0.612245	0.461538	0.44444	0.384615	0.35294	2	20							_	-	-		-	-					- 0	- 0	- 0	=			В					В		
1550-1600	12	182	0.804651	0.653846	971769	2	70	0.533333	0.33333								_				-									-										
1500-1550	0.909091	0.781905	0.688889	738685.0	0.368182	7,000,000	0.380552	0.3125	0.33333								_				-									_										
1459-1500	1833333	833333	3,755556	9962390	1,7118333	377778	1347826	2999921	2999911	_		_	_	_	_		_	_	_	_	_	_	_							_	_	_	_	_	_		_	_	_	
1400-1450		23	2289990	- 89	625	9268969	11111	- 70	1124	77	333333																										_			
1359 1400 1	52		_	_	_	_	_	23895 0	_	E		-	-	-	-		-									0 0					_	_		-		_	_			_
1300-1350 13	F	0.923077 0.8	_	_	_	1578947 0.0	_	0.54667 0.3	8471 0.5	20						0	=									0 0	0.0	0.0	0.0	=			Φ					Φ	0	
1230-1300 130	_	186111 0.95	÷	1,715882 0.67	-	152 0.57	-	_	_	125 83	_		0.33333				=				-		-				0 0	0 0	0 0	=				0			_		0	
	۲	Ē	÷	÷	-	-	÷	÷	=	357148 0.25	=	0	0.33	0	0	0	-	-	-	-	-	-	-					-		=		0		0	0	-	_		0	-
200 1200 1230	13	_	_	_	_	8 0.633465	=	_	=	_	0.083						=	=	-		-	-					= 0	= 0	= 0	=										
50 1150-1200	2	Ē	÷	Ť	Ĕ	Ť	=	0.376812	=	0.208333	10	0	0	0	0	0	-	-	-		-	-					0 0	0 0	0 0	=					0				0	
1100-1150	-	0.815789	_	Ť	=	0.303817	=	_	=	=	=	0.14285				0	-	-	-		-	-					= 0	= 0	= 0	-					0				0	
1959-1100	0.77778	0807892	0.851064	0.688889	0.641176	0.594937	123	0.465909	0397486	0.318182	0217391	0.1					_	-	-			-								=										
1009-1650	80	2999980	0.747253	0.70068	0.639535	0529801	_	=	0317308	0.146351	878	0.0714285					_	-	-		. =	-					- 0	- 0	- 0	=							_			
959-1000	0.789231	0.810811	_	0.735099	0.616505	0.654054	0.49359	799905-0	0.360825	0.25757B	0.285714	0.129032	0.357143	0.125			_				-									_						_	_			_
900-950	_	0.722453	0.746988	0.78115	0.651961	0.614213	0.513228	42	0.407407	0.29661	834222	0.236364					_													-						_				
850.90	0.77778	0.782609	0.863979	0.753425	1925291	0.593458	0.524887	0.474654	0.394737	0.28777	0.259259	0.122807	0.131579	799911667			_																В					В		
800830	1606060	0.65	0.744681	0.711656	912599	49190	0.592437	0.493878	0.409692	0.337423	0.273973	0.142857	176471	0.15625		1111111	_				-										_	_		_			_		_	
20.800	2280	187750	7062270	821917.0	7815690	0.659574	1216981	0.462838	0376518	0316742	0270115	0.15942	0.13233	71745700	0.0689655																									
92.00	97777	1,74359	1,732538	1.71229	3.705069	632939	1500781	1,456311	1,422222	136534	706721	1220238	28698F	1.0487805	_		_	_	_		_	_	_							_	_	_	_	_	_		_	_	_	
659.70	17	90	2002	92.82	98339	668224	244061	2101950	82458	340557	285136	220588	13878	0787402	0869365	022222																					_			
9 029 009	85743	_	_	_	_	1628291	_	1540881 0	_	291545 0	1263699 0	_	0.10291 0	_	_	0 689100	_														_	_	В			-	_	В		
530 600	8	-	_	=	=	818189	=	=	=	=	=	0.169381 0.2	_	=	=	-	=							0 0	0 0	0 0	= 0	= 0	= 0	D										
500 350 53	1 29990	14782 0.8	=	26925	=	÷	=	2222 0.7	=	=	2331 8.5	70588 0.0	-	=	_	9.0	-	-	-			-					0.0	0.0	0.0	-		0	0	0	0			0	0	
Г	9.0	989	67 187	562 0.61	9.0	969 0.61	188	70 LP	396 0.3	873 823	322 8.2	144 0.1	463 0.10	2556 0.0	1379 0.0	0.00338983 0	-	-	-		-	-	-				0.0	0.0	0.0	-						-	-			-
0 50 500	-	67 0.608696	0.642857	22 0.63095	15 E.S.	97 0.455	99-0	30 E324	0.36	28 0.271	62 0.159	19 0.133	513 0.107	354 0.653556			-	-	-	-	-	-	-			0 0	0.0	0.0	0.0	-					0					
400-50	-	1999970 8		1 0.621622	0.47427	4 0.5187	9 3008	13.65	7 0.2826	7 0.2562	76 0.163462	27 0.1608	95 0.4520513	32 0.0397354	0.0056	В	-				-							0.0		-			В					В		
350-400	F	0.470588	0.451613	0.5000	0.56513	0.3461	0.29625	0.29050	8.2355	0.16101	71 B1417	4 0.0816327	0.0238095	0.0129032			=				. =		. =				= 0	= 0	= 0	=					0					
330.350	23	0.285714	0.28571	0363854	0.242424	0.280488	9.16	0.24444	_	_	3 0.0843882	9 0.0636364	0.023522				=	-	-		-	-					= 0	= 0	= 0	=										
239 300	20	17	_	_	=	_	0.1373	287	0.133333	0.0869565	0.0434783	9698010 0					_	-	-		-	-								=						_	_			
200.250	0.33333			57	0.0689655		0.0615385	0.097561	10.0								_				-									_						_	_			
150.20				8271280	0.0571439	7999tb00	0.0217391	72591800													. =														0					
100-150																	_	-	-		-	-								-										
00 00 a/c				0				0									-	-	-		=	-	-				= 0	= 0	= 0	=	=					0	0			0
B-30 MeV/c		В															_	-	-			-											В				_	В		
MARS	3 Dag	45.9 Day	9-13.5 Deg	13.5-18 Deg	18-225 Deg	22.5-27 Dag	27-31.5 Lbg	31.5-35 Deg	40.5 Deg.	45 Deg	45-49.5 Day	St Day	85 Deg 1	G Deg	75 Day 1	72 Day 1	35 Dec 1	-81 Dec	55 Dec 1	85.5-91 Dev	42 De	on Dec	99-11G 5 Dov	108 5.408 Dec	IR.112 5 Dec	Total Control	The series of the series	Margarette.	121 3-120 Day	Mar care	130 5-135 Deg	.139.5 Deg.	5-144 Deg	144 148 5 Day	5-153 Deg 1	1575 Deg 1	157.5-162 Deg	166.5 Deg.	5-171 Deg 1	17-1755 Day
e	Ŧ.	9	9-13	13.5	185	22.5	25	12	36-4	9	4	69.5	245	28.0	63-6	67.5	25	15	8	17	9	100	Ē	100	1	1		ì		9	é	ġ	é	7	*	ġ	157	162	é	Ę

Table 6: Table for 2D Histogram for New ANM-Rein-Sehgal

_	_		_	_	_	_			_																									_	_	_	_	_	_
1930-2000	0.715882	8812125	0.688778	0.635714	218150	0321430	2	2 -									- 0		= 0		= 0	= 0														-		_	
1909-1950	0.884658	0.74335	0.719368	0.648275	0.635394	0.6.5759	210												= 0																			_	
1850-1900	0.783784	0.759991	0.698052	0.615385	0.525533	0.307244	8												= 0																				
1800-1830	0.808989	1734882	0.711409	8-86090	0.56923	0.49000	030360	90																														_	_
730-1800	835356	786323	578579	1648734	522	47222																																_	_
00-1750	1836538	9646	191461	81818	DB52	49,438.2	_												-																			_	_
20-170	_	_	_	0 28880	509091	1515385 0	-	2322			0.0	0.0	0 0	0.0	0.0	0.0			= 0	0 0			0 0	0 0														_	_
600-1650 16	_	_	=	677-81 0.6	Ξ	-	_	-	-	60	0.0	0.0		0.0	0.0	0.0	0.0		= 0	0.0	0.0	0.0	0.0	-	-									0.0				_	_
1910	_	_	_	200237 0.6	57381		-	-	0 0		0.0	0.0		0.0	0.0	0.0	0.0		= 0	0.0	0.0	0.0	0.0											0.0		-		_	_
1531	_	_	_	63144 0.39	59-65	-	-	-	-	-		0.0		0.0	0.0	0.0	0 0		= 0	0.0	0 0	0 0	0.0													-		-	-
1500 1500	Ξ	_	_	631818 0.63	598 059	488.872 0.48	-	-	-	000	2	0 0		0 0	0 0	0 0	= 0		= 0	0 0	= 0	= 0	0 0											0 0	-	=		-	-
-68 1430	-	_	_	_	53 0.597					1						0.0	0.0	-	= 0	0 0	0.0	0.0	0 0																
1400-1-6	_	_	_	72 0.65009-4	14 0.568153			-	-		1	0.0	- 0	0.0	0.0	0 0	= 0	= 1	= 0	0 0	= 0	= 0	0 0													-			
E 1358-1	_	6 0.790123	Ï	9 0.680972	4 058200	10232011	-	_	_	1000000		0 0		0 0	0 0	0 0	= 0	-	= 0	0 0	= 0	= 0	0 0												-	=		_	_
HH 1301-135	_	_	_	3 0.673329	=				3 0				- 0				0.0		= 0		0.0			0 0												-		В	В
1258-13	0.829352	0.809324	0.736299	0.482373	0.596.406		-	-	3 0	0 0	-		- 0	?		0 0			= 0	0 0			0 0																
1200-125	0.82(846	Φ	_	0.700468	2985190	05337	0.4008.38	0.37394	0.2002.10	_	-	1000	2 .						= 0																	-		_	_
1150-120	0.821212	0.806396	0.736423	0.693381	0.628975	0.555710	487885	0.374753	0.000	102010	20000	1000000							= 0																			_	_
1108-1158	8220280	0.79485	0.759636	0.68934	0.52265	0.564074	0.5383	933846	2000	0.01230		0.000000	The second					-	= 0																			_	_
1050-1100	0.835	0.780172	0.756151	0.69735	0.6-49693	0.5553	0.50302	0.420597	E 00 2 0 0	0.000	0.0000	0101010	0.01010						= 0																				
1000-1030	1811494	0.88895	0.764215	0.718415	0.646345	0.583082	0.408.878	0.425238	0.000	0.011000	200000	100000	270	200000				-																				В	В
929-1000	181-53	0.79961	0.777286	6705178	0.656827	0.57303	051302	0.439291	0.0000	0.00000	200000	0110170	01100						= 0																			_	_
000 930	0849776	0.816404	0.769928	0.699029	0.646262	0.588134	0.500007	1738421	0.07410.4	0.000000	0.01.00.0	0150007	7700000	0.022200	0105	7		-	= 0																				
K0.90	0.846791	0.803672	788-77-0	0.723165	0.653591	0.580724	0.525033	0.457878	1000	0.00000	2000000	0.126264	0.105369	0.0424240	0.000000																							_	_
E8-83	83-821	833534	6.778919	16224.0	1161291	785587	92.2.20	200	000000	0.000000	0.2220.4	170000	020270	0.0100000	195	7																						_	_
00800	780028.0	180314	1,772.47	1,736591	57,679,73	51000	250,057	45252	00100	2000000	2000	10000	190001		1000	7		_																		_	_	_	_
. B-220	846154	1328	280804	22938	565302	61161112	23826.4	82.00	0.000	1000	000000	102301	1001001	_	-	1	T GEORGE																					_	_
9299	1697	807565	7394 0	23165	685785	631820	200	2000			3000	2001	13000	0.000000	0.000000	2004	93.770		-																			_	_
630	_	812763 0.8	Ï	746173 0.7	20125	-	0 0	0 0			-	0.0		-	-		_		= 0		-	-		0 0										0 0				_	_
20.60	_	_	_	12411	70024 0.6	-	-		-	0 0		0 0	0.0	-		-	_	= 1	= 0	0 0	0 0	0 0	0 0				0	0	0 0	0 0	0 0	0	0 0			-		_	_
-	0.812332 0.80	_	_	=	DE9713 0.70	-	_	-	0.00	-	-	-	0.102000 0.10	-	_	-	_	= 1	= 0	0 0	- 0	- 0	0 0	- 0										0 0	-	-	-		
	_	_	_	_	578125 0.969	-		0 0	_	_		_	_	-	2.0	_	į.	-	= 0					0.0											-	-			
100-30	_	_	_	18 0.642908	19 0.578		_				0 0				-	-	= 0	= 1	= 0	0 0	= 0	= 0	0 0	-											-				
400-42	_	4 0.611916	Ť	0.529018	20020	98298 0			-	-			-	-	1	-			= 0	0 0			0 0													-			
330-400	_		_	8 0.527	0.2098	0.001005				0 0	0.0		10 0 00 000		9 0			= 1	= 0																				
300-320	0.398693	0.351406	0.359365	0.372408	0.32415	0.384053	0.23220	0.21823	10000	0.101000	0.000.0	0.000.00	1					-	= 0																				
230 300	0.225352	0.254881	0.272864	0.254642	0.240964	0.186729	0.157462	140781	0.107249	0.000000	0.0073000	0 01 4405	0.0011400						= 0																				
200-250	0.173913	179-87	0.175824	0.162912	0.135068	0.197303.9	and a			0.000000	10000000								_																			- 0	- 0
56-200	202200	0338922	92128	030269	10.67253	0.756881	78220	2505900																														_	_
100-150 15	0	0	0	0		-							20			2 0	20	2 1	20	2 0	20	20	2 0	20												-		8	8
c 56-100					_								- 0						= 0		-	-																	
0.50 MeV/c					_					_	_	_		_	_				=																				
ANMERS	0-45 Dag	ř.	9-135 Deg	135-18 Deg	8-22 5 Dec	295.97 Dec	The same	315,36 Dec		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1000000	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	100 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1000	100 CO CO CO CO	No. of the last	District Co.	2000	Mark Deg	1 TO 1 TO 1 TO 1	and a	20 24 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	100 5 100 Day	108 1175 Dec	112 5.117 Dec	17,1215 Day	27 5.1% Dec	136,130 5 Day	a STS Day	15,1395 Day	13 5144 Dec	44-148 5 Day	16 C 100 Day	133 1575 Day	157.5 162 Dev	162 1665 Drg	200 E15 80	17-17-5 Day
New.	27-	6.57	943	1354	18-22	2250	22.0	2	20.00	9	9 9	9	600	200	200	0 10	2 6	4	6 5	0.00	000	6 6	7 00	2000	3 2	110	12	5	18	F	1	B	1	9	125	197.5	162.1	1855	17.7

Table 7: Table for 2D Histogram for New ANM-Berger-Sehgal

Column   C	e		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
Column   C	1939-200	0.809524	182122	ucoson o concor	0.65139	-	-	-		В		_																						В		_	_		
Column   C	Н	0.823329	0.452378	0.00000	-	-	÷	=																															
Column   C	1830-1900	1602270	0.776824	200000	1/20/00	10000	0.33333	3333				_			В								_													_	_		
Column   C	1800-1850	6812180	0.08623	0.000000	0.535641	0.467073	136364	1							В																								
Column   C	1730-1800	0.797297	0.77323	0.027.000	1000000	0.4670	0.381070	1	0			_																									_		
Column   C	30-120	1/2/10/1	81,7829	200000	20100	442300	14	7,66667		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_	_	_	
Column   C	021-059	82238	1000	2000	200000		445800					_											_													_	_		
Column   C	300-1650	871795	6/8/9	00 400 4	100100	ACOUST D	W3880			- 17	_	_	_		Ε.								_	_	_	_	_	_	_				0		_	_	_		0.0
1	1 0991-02		-	-		_			_			_	_	-	8	-				-		_	_	_	_	_	_	_	_	_					-	_	_	-	0.0
Column   C	00-1550	823129 0	2000	- 0	0.0	3 0	( )		_			_			0	0	0	0	0		0	0	_								0	0	0			_	_		
Column   C	30-15m 1;	0	200	-	-	0.0	0 0	-	-	_	-	_	8	_	0						В	8	_	8	8	8	8	8	8	8						_	_	-	0.0
Colored   Colo	10-1-60	0	= :	<u>.</u>	0 0	0 0				85714		_			В						В		_										0			_	_		
Column   C	0.1400 1.4	0 0	= 1	-	_	0.0	-	_	_	÷	-	_			0								_											-		_	_	-	
1	0-1350 135	0 1	= :	0 0	0.0	0 0	2			_	17	0				0	0	0	0	0	В		0								0	0	0	В	0	0	0		
Column   C	130 130		_	_					_	Ť	3333 0	-	0							8	0	0	0	0	0	0	0	0	0	0			0	0	8	0	0	-	
Colored Colo	1250 1230	0	= 1				0 0		_	_	_					0	0	0	0		0	0									0	0	0	в					
Colonia   Colo	F	-	=	-	-	-		_	_	Ť	Ť	_		-																							_	-	
1   1   1   1   1   1   1   1   1   1	150 1130	-				3 0			-	_	_	22			В								_													_	_		
1	1901 001	-	7	-	-			250	_	_	_	÷	231																				0				_	-	
State   Stat	50 1030-1	23	= 1	0.0	0.0	0.0	0 0	8 0.3946		_	Ť	Ť	0.0769		В																			в					
1	100010	3 0	-	-	-			-		_	_	_	- -																								_		
State   Column   Co	0.026	-	=	_		_	9 6			-	0	_	=																										
State   Stat	9998	0	= :	- 0	-		0 0	0 0		d	d	_	<del>-</del>	889 0.125	19 8						В		_													_	_		
1	820-30	0	20	0 0	- 0		0 0			_	_	_	_	=	=	0	0	0	0												0	0	0				_	-	
1   10   10   10   10   10   10   10	800 E	B :	= 1	0.0	0.7314		0 0		-	74 0.3118	12 0.2540	_	_		-		133 B						_													_	_		
1   1   1   1   1   1   1   1   1   1	8-62	0 2	- 1		0.037	0 0	0 0			0.3207	0.2593	_	_	_		_	0.0833						_													_	_		
1   10   10   10   10   10   10   10	70.75	E2280	= 1	-	0.000		0 0			_	_	Ť	_	_	Ï	-	0 1-																						
1   10   10   10   10   10   10   10	659-70	J0611-8 0	SPER	0.0	0 0	0 0	0 0			0.348088	_	_	_	=		_	0.035696						_													_	_		
1	600-650	834184	1826	10000	0.00017	00000	0.561587	0.500339	7428140	0.337145	0.232955	0.189664	0.1-6609	_	_	_																				_	_	-	
1   10   10   10   10   10   10   10	920-600	DIC962:0	18122-6	0.750837	0.500.00	0.00000	052820	0.4768	0.400137	0.316995	0.236639	0.194822	0.128324	0.080-4312	0.0497512	0.0041-93							_													_	_		
1   1   1   1   1   1   1   1   1   1	20053	1980380	0.797824	0.707000	0.72030	0000000	0512134	0.431607	0353491	0300615	0.229012	0.1691.49	0.120419	0.0719951	0.0325901	0.00373399																							
1   1   1   1   1   1   1   1   1   1	GB 200	2400001	1,022694	9000	10000	2000	465%4	342.423	327259	252047	21244	1.148848	1.10-4636	1659-2262	1,0215545	101829KB	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		_	_	_	_	_
1   10   10   10   10   10   10   10	Ė								_	_	Ť	Ē	Ť	-						8	0	0							0	0					8			-	20
1   1   1   1   1   1   1   1   1   1	1.400		= 1							_	_	-	_	=	70					0												0	0		0			-	
1   1   1   1   1   1   1   1   1   1	336		= 1	0 0		- 1			-	_	_	_	=	8	0																							-	
Seguing 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	=		= 1	-	0.0	0 0	0.0	-				=	8.8		0																							-	
900 C	Ë	2	= 1	-	-	-	-			_	9.85	0.0																										-	
	Ī	-	=	_	_	-			100	0.42					0																							-	
	e e	0.093	910	0/00	1000	100	0.00																																
	н		= 1	-								_	_	-		_						_	_	_	_	_	_	_	_	_						_	_	= :	_
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	9	9		_ 0	_ 0	- 0					-	-	_	2	9	9	9	9	9	9	2	2	9	_	_	_	_	_	_	2	9	9	9	0	9	9	-	-2.1	
14.9 FO 14.9 F	MBS 05	0 2			200			- 0	100	200	0 200	0 %uc	0 %uc	0 25c	D No.	0 Mag	0 Mag	0 Mag	0 Mag	0 Mag	0 %uc	0 %uc	Deg D	8 Deg B	1 Deg 1	7 Deg B	1 Deg 1	i Deg II	1 Deg 1	1 Deg 1	Dog B	1 Deg B	Dog B	1 Dog 0	Deg 0	1 Deg 1	Deg 0	260	Des de la companya de
	NewAN	0.45 De	1 6 7 P	107.00	100000	000000	27.015.01	315.36 D	36-40.5 D	405-45 E	45-85 E	495-54 L	54851	585 63 L	63-67-5 L	675-72 L	72.H5.L	765-81 L	81.K5.5 L	855.90 L	98.9451	945.99 L	99-103.5	1635.1fb	108-1125	112.5-11,	117-1215	1215-126	126 138 5	135.135	135-1395	135.14	1441485	148 5 153	153-1575	1575-162	162-1665	185517	13.518

These are the corresponding tables to the 2D Efficiency plots for CC-Coherent Pion events.

Table 8: Table for 2D Histogram for New CC-Coh Pion NM-Rein-Sehgal

E 700	08/2	88888	2		_			_					_					_			_					_					_	_	_		
1 0251-0		_	0.000007	_				_	۷	٥	۷		٥		_		٥	٥	٥	٥	_ 0					_ 0			_	_	_	٥	_ 0		
1900 190	0.0			0		0 0	0	-	0	_	_	0	_	-	=	0	0	_	_	_	0 0	-	-		-	- 0		-	-	-	_		0 0	-	
1830 183	_	_	0.77	0	00	-	0	-	0	_	-	0	_	-	=	0	0	_	_	-	- 0		-		-	- 0		-	-	-	-	-	- 0	-	
180	_	98 088880	8	-	00	-		-	-	_	_	-	_	-	=		_	_	_	_	- 0	-	-		-	- 0		-		-		-	- 0	-	
1.004	=	7 0.714286	69	-	0				0					- 1	=						- 0	-										_	- 0	-	
14051	_	999999	2	0.666667	00			0							=		_				- 0											_	- 0		
16501.0	69	_	0.8%	2				_		_	_		_	-	=			_	_	_						- 0					_				
EXCHESS 1532-100   100-100   100-100   100-100   100-100   100-100   100-100   100-100   100-100   100-100	98889	0.805565	_	8				_		_	_		_	-	=			_	_	_					_						_				
15504000	88	0.82857	0.61111	_									0		-			0																	
5004550	0.816154	0.365277	0.539412	0.335353				_		_	_		_	-	_			_	_	_					_	_						_			
50450	90000	836/35	0785714	19																															
1 65140	2000000	-	-	7	20000	2 0		_	0	0	0	٥	_	-	_	٥	0	_	0	0		2 0			۰		2 0			0	0	_		2 =	0.0
Б	_	SEGRE 0.73	90 982391 100	3	8.		0		_	_	-	_	_	0 1	=	-	_	_	_	-	0 0	-	-			_		-	-	0	_		0 0	-	
130	_	_	_	90	8 0			_	-	-	-	-	_	-	=	-	-	_	-	-	-		-		_	- 0					_		-	-	
30 130	_	_	_	N 0.00005	c		0							0 1	=		_	_			0 0					- 0							0 0		
1221	0.913043		90		0.714286									0 1	=	0	_				0 0					- 0				0			0 0		0.0
SERGOD (SOUSSO) SERVICIO (1000-100) 1000-100 (100-130) 1130-130) 120-130 (120-130) 1300-130)			_	-	/999900/										=			_								- 0									
113012	_		92'0		70			_	_	-	-		_	-	_		_	_	-	-	- 0				_	- 0						_	- 0		
1100113	_	0.848485		_	0.77778					_			_	-	=		_	_		_						- 0									
10501100	08/0008	923550	0.813953	0.73556	0.77778								_					_							_	-									
0914001	S1B67.0	82Z80	760607	ogulos:	occaste	2 -	_	_	_	_	_	_	_	_	_	_	_	_	_	_			_	_	_				_	_	_	_			
000100	19	1813725		_	0633333	8.			-	-	-	_	_		-	_	-	_	-	-					_						-	_			
8606	œ.	0.838883	0.752888			0.53555	Ĺ	_		-			_		_		_	_							_				-	-	_	_			
8698	0.710536	0.858824	0.78162	0.770833	0.655172 0.73707 0.5000c 0.7170c		_	_	_	_	_		_		·		_	_	_	_				_	_						_	_			
	0.80662	80	0.755102	ß		778500		_	_			_	_	_	_	_	_	_			_	-			_	_	_		_			_	_		
(508II) SIDES	088880	83538	_	16,0030	ocodils 0.72				_	_	_	_	_	_	-	_	_	_	_	_		_			_	_						_			
005/00	074468	0789474	0752	100	OGDGD/S	Days.	2	8		_	_		_		_		_	_	_	_	-	-			_						_	_	-		
639-00	98998	987380	39080	90000	Stone	2000	12	68	_	-	-	_	_		-	_	_	_	-	-												_			
60.00		0.84375	0.790303 0.828125	0.736579	10,704		7	_					_	-	-		_	_			-	-			_	_	-					_	-		
23000	0.814815	0.717949	0.794303	0.748252	0.677905 0.704	2000	0.467407	0.142857	0.333333	_	_	_	_	-	_	_	_	_	_	_					_						_	_			
DESCRIPTION OF THE PERSON	8	1885714	0.792453	0.736	0.67222 0.	o descri	0.458333	0.368421	0.454545	0.10eee	_	_		_	-	_	_		_	_															
		0.854545	0.000000		0.63483	100000	270	0.22222	100000			_	_	_	_	_	_	_			_	-			_	_	_		_			_	_		
400450 450500	87189	1982390			0.481132	0.000	187	0.290K3	0.173913	0230489	128	_	_	_	_	_	_	_	_	_		_			_	_						_			
30.00	54	0.30000	0.6265	0.494845	0.4	10.000.01	14769 0.94118 0.32284 0.42 0.458333 0.407	2307547	32000	0.0434783	0.0855533	_	-	_	-		-	-	-	-					_		_				-				
311520	13137	_	_	_	341772	244 86	0.14769	0.0483871 0.190476 C	182	ucassle .	- September 1	9	_	_	_	_	_	_	-	_					_					_	_	_			
	pleaser 0.	0	230052 0.	_	224286 0.75673	Toward or	130435	HSSST 0	58825	desis 0	occesida occitisas	_	_	-	_	_	_	_	-	-	ے د	20		_	_		2 0		_		_		ے د	2 6	
2020 20	÷	0.235.94 0.1	_	÷	0.25882 02	- 10		8	8	Z.	z z	_		0 1	=	-	_		_	-	0 0	0 0				0 0	0 0				_	0	0 0		
		0.076823 0.2	_	00   10000000	21.5	3 8	œ		-	-	-	-		-	=	-	-		-	-	- 0	-	-			- 0				0	0	-	- 0	-	
130		000	_	900	00		0.0298	_		-	-		_	-	=		_	_	-	-	0 0		-			- 0					-		0 0		
9100 100	-		_	-	_	-		_	_	-	-	_	_	-	=	-	_	_	-	-	-		-		-	-	-	-	-	0	-	_	-	-	-
0.50 MeV/c 50:100 100:150 150:200	0	_	_	0				_	0	0	0	٥	_	-	-	٥	0	_	0	0		2 0					20			0	0	_		2 0	
RS 050	_	-	-	0 20	9 5	9 0	0 00		0 20			0 20		= 1 E°	=	0	0 20				e d	200	o d	Des	D Se	200	9 4	e e	Dec	Des	Deg	Deg	200	88	88
New NACKAS	045 Dg	450 Dg	9435 Deg	135-18 D.	18225Dg	And Spirit	315.35 De	36405 De	40.5-45 D.	45-49.5 D.	495.54 D.	545851A	38563D	636/5Lk	17.00	727651A	765.81 D.	81855D	85590 D.	90945 D.	10000	Institute Institute	Ins.1125.	1125117	1174215	1215126	lane lan	135 1305	1305141	14414851	1485133	1531575	15/5/6	1665171	171.175 Deg
_	_	_	_		_	_		_	_	_	_	_	_	=		_	_	_	_	_	_	_	_	_	_	_	_	_	_	Ť	_	_	_	_	

Table 9: Table for 2D Histogram for New CC-Coh Pion NM-Berger-Sehgal

Ī	900	)900																																				
	ã	ã	_	0	_	_	_	_		-	-	=	0 1	_	_	_	_	0	_	0	0			0		_	_	0	_	0	_	_	_	0	_	_	0 1	= 0
	0.7 1 0.833333 0.75 1 0.040066 0.041 0.5 1 1 0.040067	-	_		_	_	_	_		=	-	=	-	_	-	_	0	-	_	-	_	_	_	_	_	_	_		_		0	_	0	_	_		-	= 0
		_	_	_	_	_	_	_		_	-	-		_		_	0		_		_	_	_	_	_	_	_		_		0	_	0	_	_			= 0
	2.	_	_	_								_							_												0		0					= 0
		987	2		Ė	Ė	•									_	_	_		_	_	Ť	Ť	•	Ť	_	Ė		•		_	_	_	ė	•			
	20 /90	3	2		_	_	_	_		=	0 1	=	0 1	_	_	_	0		_		_						_		_		0		0	_	_	-	0 1	= 0
	88. 5	-		0	_	_	_	_		-	-	=	-	_		_	0		-		_	_	_	_	_	_	_	0	_	0		_		_	_	0	-	= 0
	9999	086	/900000	_						=		=			_				_		0																	= 0
		=	_	_		_	_	_		=	_	=	-	_	_		_		_		_							_	_	_					_		-	= 0
		7987	.,																																			
		ď	ď	_	_	_	_	_		-	-	-	-	_	0	_	0		-		_	-	-	_	-	_	_	0	_	0	0	_	0	_	_	0	-	= 0
	12	88	2	_					00	=	-	=	0 1		-		0		_		0	0	0		0						_		_			-	0 1	= 0
	0888	9	_	_					0.0	=		=			_				_																			= 0
	_	20	0.333333	_					_	=		=							_																			- 0
	-	9888	083888		_																						_											
	6		_	_	-	_	_	_	0.0	-	-	=	-	_	_	_	0	-	-	-	_	_	_	_	_	_	_	0	_	0	_	_	_	_	_	_	-	= 0
	88.		872		2			_	0.0	_	-	=	-	_			0		_								_							_			-	= 0
	2	0.81818	0.833333	910	_	_	_	_		_	-	=		_	_				_		_						_	_	_	_					_			= 0
Tables 1239131 1319131	0.00000	0.82105	0.333333	_								-							-																			
			J. S. C. C. C.		Ė	Ė			_						İ		_	_		_	_	_	_	•	_		Ė				_		_	Ė	•	İ		
TOPICI INTERIOR HONOR				20	_	_	_	_	0.0	=	-	=	-	_	0	_	0		-		_	_	_	_	_	_	_	0	_	0	0	_	0	_	_	0	-	= 0
	988	28	2.0	22 0.73	_				00	-	-	=					0		_		0										0		0					- 0
			280		2					=		=			_				_															_				= 0
SOME THEFT	12	0.73913	53	0.714286	_	_						-							-																			
	2000	77000	0.857143 0.084818 0.8 0	14286	•	_	_	_		_		_		_	_	_	_		_			_	_	_	_	_	<u>-</u>	_	_	_	_	_	_	<u>-</u>	_	_		
Bank	123	ď	34848 D.	14286				-	-	-	-	-	0	_	0		0		-		0			_			-	0		0	0		0	_		0		= 0
	8	2027 08	7143 0.6	8	4286 0.5		-	-	0.0	=	-	=	-	-	0		0		-		0			_			_	0	-	0	0		0	_	-	0	-	= 0
	'	3714286 0877	889	99			_	-	00	=	-	=	-	-	-	_	0		_	-	_	_	_		_	_	_	0	-	0	0	_	0		-		-	= 0
DESCRIPTION OF STREET	_	8714	15 15 15 15 15 15 15 15 15 15 15 15 15 1	95 058333	8	2	-	-	00	-	-	=	-	-	-		0		-								_		-					_	-	-	-	= 0
	2 0.7	8	7 0.7149 C	7 0.7368	20 02	_			0.0	=		=			_				_																			= 0
DAME IN COLUMN	0.83333	_	0.82am7				_	_		_	-	-		_	_	_			_		_	_	_	_	_	_	_		_			_		_	_			- 0
	_	0.000	_		0.62C3TS	0.77778			_										_																			
	2000	1882280	188	17897	29000	1999	244444	_		_	_	_	_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
	310007	1000	0.0000007 0	1000	18	19000		_		_		_	_ 1	_	_	۷	٥		-		_	٥	٥	٥	٥	۷	۷	_		_	٥	۷	٥			_	_ 1	_ 0
	0.855 1 0.94 0.057	55318 O.	ã	GE354 0.	39112 0.	9048	2014			-	-	=	-	_	0	_	0	-	-	-	_	_	_	_	_	_	_	0	_	0	0	_	0	_	_	_	-	= 0
	-	2318 D.G.	3867 0.8	90	3867 0.5	9867 0.65	점	3.0		=		=	0 1		_	0	0	0	_	0	0	0	0	0	0						0		0			0	0 1	= 0
	80.		0.600067				2	2.	- c	=	-	=	-	_	-	_	0	-	=	-	_	_	_		_	_	_		_		0	_	0	_	_		-	= 0
	0.77778	9.9	04375 0.5	33	g 0.355217	33 B.338462	3	B 0.355553				=		-	_		0		-								_		-					_	_			= 0
	8	3	04375	65789			1 Si	1000	3 0	8	9	=	-	-		_	0		_							_	_		_		0	_	0	_	_		-	= 0
	2	0.5555	23		0.193548			908	0.181818	9	-	_		_	_				_					_			_							_				= 0
		í		eg.	00768231				0												_																	
	-		13 -				ulsisis u		GUNESCE			_	_		_	۷	٥		-		_	٥	٥	٢	٥	۷	_					۷				_	_	_ 0
	-	-	널	_	_	3	3	á	86	1	-	=	-	-	-		0	0	-	0	_	-	-	_	-		_	0	-	0	0		0	_	-	0	-	= 0
	-		_	0	011111	_	-	-		=	-	=	-	-			0	-	-	-	_	-		_			_	0	-	0	0		0	_	-		-	= 0
	-	-		0		-	-	-		=	-	=	-	-			0		-					-				0	-	0	0			_	-		-	= 0
	_	-	_		_	_	_	_		=	-	=	-	_	-	_	0	-	=	-	_	_	_		_	_	_		_		0	_	0	_	_		-	= 0
DESCRIPTION OF THE PROPERTY OF	_		_	-	_	_	_	_		-	_	_	_	_	-	_		_	_	_	_	_	_	_	_	_	_	-	_	-		_		_	_	0	_	_
THE PART INC	ď.	287	5 Deg	SDeg	SDeg	2 Deg	S Deg	2479	S Deg	9370	2 FE	Sar re	S Deg	2012	5 Deg	2Deg	5 Deg	d Deg	5 Dec	ag De	5 Deg	9 Deg	35 Dg	JOS Deg	125 Dg	H7Dg	21.5 Dg	126 Dg	302 Dec	132 Dg	395 Deg	144 Deg	485 Deg	133 Dg	57.5 Deg	162.LVS	665 Deg	1
	2	9	9.135	1327	183	8	ri Ki	13 15	9 5	į.	9	8	89	8	89	67.57	72.76	765.8	8	55	909	95.59	30 IC	1685	1087	1125	1171	5	ď.	5	Ĭ	199	7	1485	ď.	12/0	Ž,	8 5
	ğ.	305	9-13.5 I	135-181	182251	125.02	27.03	8	20,403	9	9	1000	250	9	636753	67.572	727651	765.81	8,855	8550	909153	915.00	991035	1035-10	108.112	1125.11	117.121	121512	8	130513	135.130	130514	144.148	1485-15		in the	157516	157.5.162.Deg

Table 10: Table for 2D Histogram for Old CC-Coh Pion NM-Rein-Sehgal

~	_	_	_	_		_	_	_	_	_	_	_			_	_	_	_	_	_	_	_		_	_	_	_	_	_	_			_	_
		_	_	_			_	_	_	_	-							_	_	_	_			_	_	_	_	_	_	_	_			_
İ		_	_			_	_	_	_	Ī		_					_	_	_	_				_	_	_	_	_	_	_	_		_	Ī
=	-	_	_	- 0	-		_	_	_	_	-	-			-	-		_	_	-	-	= 0	0.0	-	_	_	_	_	_	_	_	00	-	_
				-	-				_		-	- 0				-					-	= 0									-	00		-
	0.833333	_		- 0								- 0									-	- 0					_		_		_			_
		_	_				_	_	_	_								_	_	_				_	_	_	_	_	_	_	_			_
ľ	7900000	_	_			_	_	_	_	_	_						_	_	_	_					_	_	_	_	_	_	_			_
-		0	_	-	-	0	0	0	_	_	-	- 0		0	-	-	0	0		_	-	= 0	-	0	0	_	_	_	_	_	_	00	-	_
	0.857148	_		-	-						-	- 0									-	= 0												
-	0.000007	_										- 0									-										_			
I				ıq.																														
	-	_	_			-	_	_	_	_	-	-					-	_	_	_	-	-	-	_	_	_	_	_	_	_	_			_
	13	-		-	-		0	0	_	_	-	- 0				-		0			-	= 0	-		0		_	_	_	_	-			_
		_	92	- 0																		- 0												_
	. 60	799990																																
İ	0.875	10																									_							
-		_	-		-				_	0	0 1	0 0			0 0	-					0 1	= 0	-		0		_	_	_	_	_			_
		870		- 0					_		-	- 0				-					-	= 0						_		_	_	0.0		
	0,600007	_		_					_	_												= 0					_			_	_			_
	80	520	is.																															
ľ		20	_			-	_	_	_	_	_	-	-				-	_	_	_		-	-	_	_	_	_	_	_	_	_	-	_	_
-		_	-		- =				_	0	-	- 0				-					-	= 0	-		0		0	_		_	-			_
=	080	89	_	0			0	0	_	_	-	- 0				-		0			-	= 0			0		_	_	_	_	-			-
	80	673	127	0																		= 0												
		799990	/9999	7900E/																														
ŀ	1 222	Æ	_	8,			_	_	_	_	-	-	-		0.0			_	_	_	0 1	= 0	-		_	_	_	_	_	_	_	0.0		_
-	15	_	5 0.75	0	-		0	0	_	_	-	- 0			-	-		0			-	= 0			0		_	_	_	_	_	0.0		-
	-		_	98			0	0	_	-	-	- 0				-		0			-	= 0	-	0	0		-	_	-	_	_		-	_
	60	700000 15	33 0.08V5	91	3 -						-	- 0				-					-	= 0								-			0	-
	0.7142	2000	0.7333	131	9 -				_	_	-	- 0									-	= 0					_	_	_	_	-			-
-				790000					_	_		- 0									-	= 0					_	_		_	-			_
	0,33333		1923290	0.7778	0.00																										_			
İ		8135	7383	0.5	2000		20																											
F	-	2999	124	0.684211		Ī	_	_	_	_	_	-	-				_	_	_	-	_	-	-	-	_	_	_	_	_	_	_	-	_	-
ŀ	-	2	23	38					_		0 1	- 0			-					_	-	= 0			0	0	_	_	_	_	_			_
-	-	0.8	714 0.81	0.065385 0.05	9 5	60	1 /990		_	0	-	-				-					-	= 0	-		0			_		_	-			_
=	0			2000	9 68 6	90 49	9970		_		-	- 0				-					-	= 0	-				_	_	_	_	-			_
		-	200	A00000			5 0.5		_			- 0				-					-	= 0						_		_	_	0.0		
		_	3,44444 0,727273 0,3625	0.533333 0.5625	3 =	0.214286 0.285714	0.460067 0.22222	625	20		-	_ 0						_		_		- 0		_			_	_		_				_
ŀ			0.44444	0,55555	0.25	0.214286	0.469957	7999900	27	_			= =			_ =				_		= 0	-				_		_				_ =	_
Į.	_	-		8	9 _	22	128X14	0.142857 0.125	928X14		_	_			_	_	_	-		_	_	_	_	_	-			_		_			_	_
f		_	_	8.	286714	Ī	_	142857	100007	_	_					_	_	_	_	_	_			_	_	_	_	_	_	_	_			Ť
ŀ		_		0 0	100	-	0	0	<u>.</u>	_	-	-						_		_	-	= 0		-	0		_		_	_	0	0.0		_
ŀ	-	0	_	-	- 2		0	0	_	0	0 1	0 0		0		-		0			-	= 0		0	0	_	0	_	0	_	0	0.0	0	-
ŀ	0 0	-		- 0	-		-	-	_	_	-	- 0		0		-		-		_	-	= 0		-	-		_	_	_	_	0	00	-	_
ŀ	0 0	-	_	-	-		-	-	_		-	- 0			-	-		-	_	_	-	= 0		-		_		_		_		00		_
ŀ	-	0		-	-		0	0	_		0 1	0 0						0			-	= 0		0	0			_		_	0			-
				- 0			_	-	_	_	-	- 0						_		_	- 1	= 0	0.0	-	-		_	_	_	_	0	00		_
ŀ	200	5 Deg	JS Deg	25 Deg	202	36 Dec	35 Deg	45 Deg	95 Deg	54 Deg	So De	2 0 0 1	20 DE	1	8 0 0 8	200	S D D	to Deg	30 Deg	Sp Deg	SIGN Deg	HZ5 Deg	117.121.5 Deg	3126 Deg	130.5 Deg	SISSDeg	139.5 Deg	S144Deg	148.5 Deg	SISSDeg	157.5 Deg	5162 Deg	MI De	175.5 Deg
E	1 13	5	13.5	20 6	3 2	15 155	8	9	7	9	3	888	3 6	8	9 19	2	12	ĝ	36	ã	ğ.	4	15	12	138	130	ij	ĕ	7	148,	á	15 8	18	Ē

Table 11: Table for 2D Histogram for New CC-Coh Pion ANM-Rein-Sehgal

66200	31882	12	571239																																
11980	9	2000	9	0 1	= 1		-	_	_	_	_		_	_	_	_		_	_	_	_	-					_	_	-	-	-			_	0.0
1981	1 120	90	88	-	= 1			_		_	_		-	_	-	_	-	_	_	-	_	-		-		-	_	_	- 0	-	- 0	-		_	0.0
1830	0.75534 0.80634 0.75 0.90537 0.94738 0.76906 0.76931 1 0.818.92	500		-	= 1						_		_	_		_			_	-	_	-						_	- 0	-	- 0				00
18001	0.76190	0.7083	0.6666		= 1					_								_	_			-					_	_	- 0	-	- 0			_	
1720180	SE2160	0.727273	88	-	-						_		_	_	_	_			_	_	_	-						_		-					
7001730	15550	82158	12	19.				_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			_	_	_	_						_	
8,198	150	100	1218						_	_	_	_	_	_	_	_	_	_	_	_	_					_	_	_						_	
1630	6624	988	189 00 1	-	= 1		0.00	_	_	_	0		_	0	_	0		_	0	-		-					_	0	- 0		- 0			_	0.0
ion Ton	301	318 0.72	150	-	= 1			_		_	_		_	0	_	_		_	_	-		-		-			_	_	- 0	-	- 0	-		_	0.0
931				-	= 1		0 0				_		_	_	_	_			_	-	-	- 1		-				_	0 0	- 1	- 0	-	0		00
1500159	۳	0848485	0735394	3	= 1						_		_		_	_			_	_		-						_		-					00
1430.130	0821429	0701731	0654545	g,	- 1						_		_		_	_		_	_	_		-						_	- 0	-	- 0				
14001450	0897435	182238	0818182	022220	-																														
001103	922	0838181	3,000,00	1681539 1									_		_			_																	
1300 1350 1350 1400 1400 1450 1450 1500	1939	ĭ	Ť	369048	-		-	_	_	_	_		_	_	_	_		_	_	_	_	-			-		_	_	-	-	-		-	_	0.0
130	918485 0.9	SECRE 0.2	_	=	3.	- 0	0		_		_		_	_	_	_		_	_	_	-	-	= =	-				_	0 0	- 1	- 0		0	0	
921	ã	_	٥	1 0.652/07	9 .		-	-		_				0			0	-	-			0 1		-			_	-	- 0	-	- 0			-	
1301	ä	Ĭ	0.652174	10808511				_		_			_					_		-		- 1					_		- 0	- 1	- 0			_	0.0
1130.12	0.774194	0.888383	0.725562	0.71864	E-Seption	2 -			_	_	_		_	_	_	_		_	_	_	-	-						_		- 1				_	
11001130	0.811463	0.83642	0.39575	0.727273	9 1	3 .																													
166110 H0H30 H30120 126120	885714	73380	768736	12	- Director 1																														
1 0301-0	10127 0	20178	Series	74906		100	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-	-		_	_	_	_	-		-		-	-	
Solon lonker	ĕ	ö			of the same	==		0	_		_		_	_	_	_	-		_	_	_	-	-			-	0	_	0 0	-	0 0		0	0	0.0
056 05606	ä	7998 08	9	_	-	0.57857	-	_		_	0	0	-	0		0	0	_	_		0	0 1		-		0	_	-	- 0	-	- 0	-		_	00
	3979 0.90	Ξ	_	_	_	290		-		-	0		_	0	-	0		_	0	-		-		-			-	0	- 0	= 1	- 0	-		_	00
88	287 0.86	257 0.83	629	350	100	2		-		0	_		_	_		_			_	-	_	-	0 0					_	0 0	-	0 0		0		0.0
SHAID SIDES SHOW	60 083	080	SGS 0.77	25 B	E 1	9 5	19	99		_	_			0		_		_	_	_		-		-			_	_	- 0	-	- 0	-		_	0.0
Н	111 087	283 086	304 8700	523 0732	100	9 (9 8 n	367 0.545455	999990		0	0		_		_	0				-		-		-					0 0	- 1	0 0		0		0.0
90.00	1980 20	14 0.833	78 0.779	56 0.78832	0.46	2000					_		_	_	_	_			_	-	-	-	0 0					_	0 0	- 1	0 0		0		00
002000	53 0.81812	21 0.83E84	3 0.777	0.745276	20 0	100000		16 0.44444		-	_					_		_	_	_	_	-		-			-	_	- 0	-	- 0	-		-	
029008	0.8210	23 0.837	37 0.7880	0.7488	2 5	200	11 0.354877	as ocassio		_			_		_			_		_	_	-		-			_		- 0	-	- 0	-		_	
20000	4 6738	07362	87948	2 0.7648	9 1	2000	6 04842	5 023258	0		_		_	_		_		_	_	-		- 1		-			_	_	- 0	- 1	- 0	-		_	0 0
9898	21	_	_	_	9 1	4 0 57773	-	-	0.391394	_	_		_	_	_	_		_	_	_	_	- 1		-			-	_	- 0	- 1	- 0			-	
18.8	1 0.71951	5 0.7008	0.513835 0.09774	0.513965 0.638243	0	S CONTRACTOR	0.4388	0.355833	910	컴	0							_	_			-					_	_	- 0	-	- 0			_	
951901	0.56923	0.61338	0.51263	_	9	CONTROL DATE OF THE PERSON OF	825	0.1987as 0.2987 0.3558	0.2387	8	0.14285		_	_		_		_	_	_	_	-					_	_	- 0	-	- 0			_	00
330400	970	0.54166	0.520025	0.437888	INCOME.	900	1300		0.18790	0.255	0.15384							_	_								_	_	- 0	-	- 0			_	0.0
30,30	70	032	7,26,50	0.3588.0		30050		0.18566	0.187097	0.07462B	0.058/002																		_	-	_				
220300	38	leggst	1314961	28574	107	oleans oleans	1620H	odastas	0004467	0043F83	0.052381																								
20020 2	P	_	_	100	_	O CANAGE OF THE PARTY OF THE PA			21212 0	_	_	_	_	_	_	_	_	_	_	_	_	-	-		-	_	_	_		-			_	_	
Н	92	slost 0.13	onesists of 52cc	OUGGERS 0-1				80	8	_	0			0		0		_	0		_	-	= =	-			_	0	- 0	-	- 0	-	-	_	00
100130 130300	-	0.08	000	000	3	4 5		_	_	_	0	_	_		_	0	_	_		_	_	-		-		_	_		- 0	- 1	- 0				0.0
100	-	_	0	-	= 1		0			0														-			-		0 0	-	0 0	-	-	_	0.0
MAY C 30 LOD	-	_	_	-	= 1	= =		_	_	_	0		_	0	_	0		_	0	_	_	-	= =	-			_	0	- 0		- 0	-		_	00
New ANA R S   0.30 MeV	-	_	_	-	= 1			_	_	_		_	_		_		_	_		_	_	<u> </u>	50 5	0 0		B		0 50	= 0	= 1	= 0		0 00		
TANK TO	045 Dg	No Dec	135 Deg	3548 Dg	842228	7.31 5 Dec	315.35 Dec	3640.5 Deg	10.5-45 Deg	5495 Dg	19554 Deg	54585 Dg	38.5463.Drg	63675 Deg	67.572 Deg	72,765 Deg	15.81 Deg	SI-SS 5 Drg	SATERS	90.945 Deg	S4166516	S010351Ag	R 175 P.	251F D	7125 D	1.513 D.	41305 D	119150	135135155	100	1441485 Deg	ALC: D	7516 D	2165 P	100517 Dg
É	Ē	7	7,	22.	9.8	418	78	8	9	4	9	7Š	88	8	8	63	92	76	18	8	ಹ	8.	2.5	É	Ξ	27	ם.	2.	2.5	2 :	7.3	1 5	12	9.	913

Table 12: Table for 2D Histogram for New CC-Coh Pion ANM-Berger-Sehgal

11128
_
_88000000000000000000000000000000000000
् ६ 
(1857 - 177
16.25.25.25.25.25.25.25.25.25.25.25.25.25.
8 8 43
图 <sub>2</sub> 26
18 18 19 19 19 19 19 19 19 19 19 19 19 19 19
10 10 10 10 10 10 10 10 10 10 10 10 10 1
200
258r
0.771.023 0.771.
285252 2857272 28574 285
10 10 10 10 10 10 10 10 10 10 10 10 10 1
112 123 123 123 123 123 123 123 123 123
Lan
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
10 10 10 10 10 10 10 10 10 10 10 10 10 1
0.000000000000000000000000000000000000
10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
7,588.64 89.118. 89.118. 89.11. 10. 10
2.53.8578 2.54.465 2.
200 See 1
28222883-0000000000000000000000000000000
1763 1 17
10.53823 0.089 0.089 0.085 0.0
00000000000000000000000000000000000000
10.000.000.000.000.000.000.000.000.000.
是据报程的 <b>自</b> 2012
3 0.8771 0.875781 0.8757
10.05.03.4
10 10 10 10 10 10 10 10 10 10 10 10 10 1
14.2.0 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 14.2.2 Dig. 15.2.2
医克洛克克氏征 经基本条件 经保险股份 医紫斑 黑黑 里面最后的 电电话电话 电电话 电电话