# SciBooNE Charged-Current Coherent Pion Production Acceptance Study Technical Note

Jonathan Asaadi<sup>1</sup> and Zachary Williams<sup>1</sup>

<sup>1</sup>Department of Physics, The University of Texas at Arlington

June 7, 2017

# 1 Introduction

This document is intended to serve as a reference for the acceptance study performed for the SciBooNE charged-current coherent pion production (CC-Coh  $\pi$ ) re-analysis, as well as provide documentation of the code used in this study (in the event anything needs to be revisited in the future). The code resides in the github repository labeled as and linked here: SciBooNE-MC. The corresponding ROOT files that were used in this acceptance study can be downloaded from here: SciBooNE-MC-ROOTFiles.

The paper is structured such that Section 2 outlines Monte Carlo (MC) samples used in this study, Section 3 describes the SciBooNE detector as it was simulated in this study, Section 4 describes the various event samples that were used to both validate and generate the acceptance studies for the CC-Coh  $\pi$  sample. Section 5 gives a high level summary of the results including the event-reduction tables as well as the CC-Coh  $\pi$  acceptance results and Section 6 gives the conclusion to our results of the acceptance study.

The appendix is left to explain the details of the macros. The appendix also details the important plots that each macro produces that play a role in making the plots shown in Section 5 (the Results section).

#### 1.1 Goal of the Re-Analysis

The goal of the re-analysis is to examine the acceptance modeling for the SciBooNE results in the presence of modern neutrino generators and updated models in order to hopefully shed light on why SciBooNE did not observe charged-current coherent pion production at low neutrino energy.

This study is intended to examine the effects of the acceptance modeling for a sample of CC-Coh  $\pi$  interactions inside the SciBooNE detector and compare what these would have been for various CC-Coh  $\pi$  production models. We utilize a simple, but robust, simulation of the SciBooNE detector and the NEUT neutrino generator to select and classify these neutrino events.

# 2 Samples

Five different samples were used in this study, three samples were generated in neutrino mode ( $\nu$ -mode) and two samples in antineutrino mode ( $\bar{\nu}$ -mode.)<sup>1</sup> Table 1 summarizes these samples. Details on these samples can be found in the Appendix.

<sup>&</sup>lt;sup>1</sup>All of these samples were generated by Callum Wilkinson (Thanks, Callum!)

#### **Summary of Samples**

Mode	NEUT version	Pion-Model	Number of simulated events
$\overline{\nu}$	5.3.6	Rein-Sehgal	1,000,000
$\overline{\nu}$	5.3.6	Berger-Sehgal	1,000,000
$\overline{\nu}$	5.0.1	Rein-Sehgal	100,000
$\bar{\bar{\nu}}$	5.3.6	Rein-Sehgal	1,000,000
$\bar{\bar{\nu}}$	5.3.6	Berger-Sehgal	1,000,000

Table 1: Summary of the samples used to build the acceptance model for this study.

#### 3 Detector Simulation

This section is intended to detail the detector simulation done in this acceptance model, and to describe the assumptions made in order to accomplish accurate classifications of simulated events as charged-current coherent pion production.

#### 3.1 The Detector

For the purposes of this acceptance study, the SciBooNE experiment is composed of two subdetectors. The first (and the more upstream) of the sub-detectors, is the Scintillator Bar Tracker (SciBar) which was originally conceived and constructed to function as the near detector for the K2K experiment. The second (and more downstream) of the sub-detectors, is the Muon Range Detector (MRD), which is the detector designed and constructed specifically for SciBooNE for measuring the momentum of muons produced from charged-current neutrino interactions up to  $1.2 \; GeV/c$  by using the observed range of the trajectory of the muon. The coordinante system used throughout this study, and illustrated in Figure 1, puts the origin in the lower corner of the SciBar detector, has z along the beam direction, y opposite to gravity, and x to beam left.




Figure 1: Representation of the SciBooNE detector and the coordinate frame used in this study

#### 3.1.1 The Scintillator Bar Tracker (SciBar)

The Scintillator Bar Tracker (SciBar) sub-detector is a scintillator detector which was used to identify neutrino interactions within SciBooNE. The dimensions of the SciBar detector used in this simulation are 0 < x < 3.0 m, 0 < y < 3.0 m, and 0 < z < 1.7 m. This simulation models the scintillator materials as having a constant energy deposition per unit length (dE/dx) for both

muons and pions of 2.04 MeV/cm based on previous SciBooNE analyses and on mean values for typical particle momentum listed in the particle data group (PDG).

### 3.1.2 The Muon Range Detector (MRD)

The Muon Range Detector (MRD), depicted in Figure 2, is located 0.55 m downstream of SciBar in the z-direction, and is a composition of two sets of thirteen alternating slabs of steel-scintillator layers, where the scintillator layers alternate between being horizontally oriented or vertically oriented, in the xy-plane. The steel layers have a z-direction thickness of 5.08 cm and the scintillator layers have a z-direction thickness of 0.6 cm. Combining all the layers of the different alternating materials results in 26 scintillator layers that "sandwich" twenty five steel layers inbetween and gives a total z-direction dimension of being 1.37m. The xy-plane is modeled as a square again (as was the case with SciBar, too) with dimensions in the x-direction and the y-direction of 2.6 m. The energy deposition per unit length (dE/dx) of a muon penetrating the scintillator layers is assumed to be a constant 2.04 MeV/cm while the energy deposition for the muon in the steel layers is assumed to be a greater value of 11.43 MeV/cm. Both values are typical for muons at the energy range produced in SciBooNE and taken from the PDG.




Figure 2: Depiction of the Muon Range Detector (MRD) which consists of alternating layers of horizontal scintillator (shown in pink) steel slabs (shown in grey) and vertical scintillator (shown in purple)

#### 4 Event Selection

Two main samples are used in this study to generate the acceptance tables. The first is a charged-current inclusive (CC-Inclusive) sample which requires a muon was created in the neutrino interaction and this muon intersects the MRD. This sample is described in Section 5.1 and is used to validate the building of the acceptance model by comparing it to previous SciBooNE analyses. The second sample is the charged-current coherent pion (CC-Coh  $\pi$ ) sample which requires a muon and charged pion are created in the neutrino interaction exclusively (e.g. no other final state particles in the event). This sample is described in Section 5.2.

Both of these samples are selected using NEUT MC-truth flags which ensure we are treating pure samples which are classified by the neutrino generator as belonging to the appropriate sample.

Whether or not the event identified by our selection makes it into the final sample used in the acceptance study depends on the behavior of the muon with respect to the MRD. A muon which enters the MRD from a neutrino interaction will either come to stop in the MRD, exit out the back of the MRD (assuming the muon's momentum is great enough), or exit out the side of the MRD. In the subsections below, these classifications are explained further.

# 4.1 Muon Stops within the MRD ("Stopped")

The requirement to classify a neutrino interaction as a "stopped" event requires the muon from the interaction to have reached the MRD, penetrated at least three layers of steel (giving activity in four layers of scintillator), and to then have deposited all of its remaining energy prior to reaching a boundary of the MRD. An illustration of a CC-Coh  $\pi$  event that would be classified as "stopped" is shown in Figure 3.




Figure 3: Depiction of an event that was classified as "stopped".

These events allow for complete reconstruction of the muon's momentum based on the number of layers which the muon penetrated and the muon's incident angle.

#### 4.2 Muon exits out the back of the MRD ("Out-the-back")

The classification of a neutrino interaction as "out-the-back" requires the muon from the interaction to have reached the MRD, and to have had sufficient energy such that the muon exited out the back face of the MRD without stopping. An illustration of such an event is given in Figure 4.




Figure 4: Depiction of an event that was classified as "out-the-back".

The exact momentum of muons which pass completely through the MRD could not be made in reconstruction, so these events were classified as having the minimum energy required to penetrate all the steel and scintillator layers of the MRD.

# 4.3 Muon exits out the side of the MRD ("Out-the-side")

The classification of a neutrino interaction as "out-the-side" requires that the muon from the interaction reached the MRD, penetrated at least three layers of steel, and then to have exited out the side of the active volume of the MRD (excluding the very back face) without stopping. Events which are classified as "out-the-side" are excluded from this study because no accurate reconstruction of the muon's momentum can be made when the muon exits out the side of the MRD. An illustration of such an excluded event which exits out the side of the MRD is given in Figure 5.




Figure 5: Depiction of an event that was classified as "out-the-side".

# 5 Results

The results of this acceptance study can be broken down into two different classification schemes of events. Those that met the conditions to qualify as CC-Inclusive events, and those that met the

conditions of classification as CC-Coh  $\pi$  events. The former is used to validate the acceptance modeling and detector simulation in order to reasonably reproduce previously published CC-Inclusive studies from SciBooNE, while the latter is used for the re-analysis.

#### 5.1 Charged-Current Inclusive Events

Here we define the CC-Inclusive sample which we use to validate our acceptance model against previously performed simulation studies from SciBooNE.

#### 5.1.1 $\nu$ -mode Charged-Current Inclusive Events

Table 2 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the  $\nu$ -mode MC.

Mode de Inclusive Event Reduction			
Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal
Total Sample	1,000,000	1,000,000	100,000
CC-Inclusive Interaction	725,730	727,278	69,363
$\mu$ + n-other particles in SciBar			
Muon enters the MRD	263,698	262,608	24,250
Muon enters the MRD and	231,089	230,054	21,001
penetrates $\geq 3$ layers of steel			
"Stopped"-Events	177,406	175,799	16,062
"Out-the-back"-Events	15,389	15,952	1,421
"Out-the-side"-Events	38,294	38,303	3,518
Good CC-Inclusive Events	192,795	191,751	17,483

 $\nu$ -mode CC-Inclusive Event Reduction

Table 2: Event reduction table for a sample of  $\nu$ -mode CC-Inclusive events simulated in the Sci-BooNE geometry.

The first quantity that is calculated for the different events is the momentum of both the muon and the pion, which are both found from the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{1}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{2}$$

where  $|\vec{p}_{\mu}|$  represents the magnitude of the momentum for the corresponding particle, and  $P_{\mu_a}$  (where a can be x, y, z, or E) represents the component of the four momentum for the corresponding particle. The momentum is reported in units of MeV/c.

The next quantity calculated is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle in spherical coordinates. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the momentum magnitudes of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1} \left( \frac{\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}}{P_{\mu_z}} \right) \tag{3}$$

$$\theta_{\pi} = tan^{-1} \left( \frac{\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}}{P_{\pi_z}} \right) \tag{4}$$

The angles are reported in units of °, and should run from 0° to 180°.




Figure 6: Muon Momentum (left) and Muon Angle (right) for  $\nu$ -mode CC-Inclusive interactions for all three models included in this study. These samples kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive.

Figure 6 shows the momentum and angular  $(\theta_{\mu})$  distribution for the sample of  $\nu$ -mode CC-Inclusive events passing all of our requirements for all three models considered in this study (NEUT v5.3.6 Rein-Sehgal, NEUT v5.0.1 Rein-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 7: One-dimensional efficiency plots for the  $\nu$ -mode CC-Inclusive sample.

Figure 7 represents the one-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events for

this study using all three different models compared to results derived from Hiraide's thesis <sup>2</sup> using the full SciBooNE Monte Carlo simulation. A few reference points are illustrated using dashed lines to guide the readers eye. A few percent difference is seen, but overall agreement between the two simulations hold.

Below are the two dimensional efficiency plots for the CC-Inclusive events in  $\nu$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section of the Appendix under the Charged-Current Inclusive Sample Efficiency Tables subsection.




Figure 8: Reference plot provided by Morgan (left), and the two-dimensional efficiency plot for CC-Inclusive events in  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal MC (right).

Figure 8 shows the two-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events in the NEUT v5.3.6 Rein-Sehgal MC. The left hand side is a reference plot provided by Morgan and the right hand side is for the NEUT v5.3.6 Rein-Sehgal MC used in this study. Table 6 is the efficiency table that corresponds to the two-dimensional efficiency plot on the right of Figure 8.




Figure 9: Reference plot provided by Morgan (left), and the two-dimensional efficiency plot for CC-Inclusive events in  $\nu$ -mode NEUT v5.3.6 Berger-Sehgal MC (right).

Figure 9 shows the two-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events in the NEUT v5.3.6 Berger-Sehgal MC. The left hand side is a reference plot provided by Morgan and the right hand side is for the NEUT v5.3.6 Berger-Sehgal MC used in this study. Table 7 is the efficiency table that corresponds to the two-dimensional efficiency plot on the right of Figure 9.

<sup>&</sup>lt;sup>2</sup>Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide\_dt.pdf




Figure 10: Reference plot provided by Morgan (left), and the two-dimensional efficiency plot for CC-Inclusive events in  $\nu$ -mode NEUT v5.0.1 Rein-Sehgal MC (right).

Figure 10 shows the two-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events in the NEUT v5.0.1 Rein-Sehgal MC. The left hand side is a reference plot provided by Morgan and the right hand side is for the NEUT v5.0.1 Rein-Sehgal MC used in this study. Table 8 is the efficiency table that corresponds to the two-dimensional efficiency plot on the right of Figure 10.

#### 5.1.2 $\bar{\nu}$ -mode Charged-Current Inclusive Events

Similar to before, Table 3 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the  $\bar{\nu}$ -mode MC.

#### $\bar{\nu}$ -mode CC-Inclusive Event Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal
Total Sample	1,000,000	1,000,000
CC-Inclusive Interaction	699,239	704,327
$\mu$ ( $\mu$ + n-other particles in SciBar)		
Muon enters the MRD	380,362	380,869
Muon enters the MRD and	336,373	337,979
penetrates $\geq 3$ layers of steel		
"Stopped"-Events	288,289	288,206
"Out-the-back"-Events	7,608	7,857
"Out-the-side"-Events	40,476	41,916
Good CC-Inclusive Events	295,897	296,063

Table 3: Event reduction table for a sample of  $\bar{\nu}$ -mode CC-Inclusive events simulated in the Sci-BooNE geometry.




Figure 11: Muon momentum (left) and muon angle (right) for  $\bar{\nu}$ -mode CC-Inclusive interactions for both models included in this study. These samples' kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive.

Figure 11 shows the momentum and angular distribution for the sample of  $\bar{\nu}$ -mode CC-Inclusive events passing all our requirements for both models considered in this study (NEUT v5.3.6 Rein-Sehgal, and NEUT v5.3.6 Berger-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 12: One-dimensional efficiency plots for the CC-Inclusive events in the  $\bar{\nu}$ -mode MCs. Muon momentum is on the right and the muon angle is on the left.

Figure 12 represents the one-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events for this study. No similar reference sample exists to be compared directly against, however we note that the shape and magnitude of the acceptance is nearly unchanged between  $\bar{\nu}$  and  $\nu$ -mode samples (as expected).

Below are the two dimensional efficiency plots for the CC-Inclusive events in  $\bar{\nu}$  mode. The tables that correspond to the plots can be found in the Efficiency Tables section of the Appendix under the Charged-Current Inclusive Sample Efficiency Tables subsection.




Figure 13: Two-dimensional efficiency plot for the CC-Inclusive events of the  $\bar{\nu}$ -mode NEUT v5.3.6 Rein-Sehgal MC.

Figure 13 shows the two-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events in NEUT v5.3.6 Rein-Sehgal MC. Table 9 is the efficiency table that corresponds to the two-dimensional efficiency plot in Figure 13.




Figure 14: Two-dimensional efficiency plot for the CC-Inclusive events of the  $\bar{\nu}$ -mode NEUT v5.3.6 Berger-Sehgal MC.

Figure 14 shows the two-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events in NEUT v5.3.6 Berger-Sehgal MC. Table 10 is the efficiency table that corresponds to the two-dimensional efficiency plot in Figure 14.

#### 5.2 Charged-Current Coherent Pion Production Events

Here we define the CC-Coh  $\pi$  sample which will be used in the re-analysis. The momentum and angle equations that were used above for the CC-Inclusive events were also used for the CC-Coh  $\pi$  events.

#### 5.2.1 $\nu$ -mode Charged-Current Coherent Pion Events

Table 4 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi$  events from the  $\nu$ -mode MC.

u-mode	CC-Coherent	Pion Event	Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal
Total Sample	1,000,000	1,000,000	100,000
CC-Coherent Pion Interaction	12,186	2,576	1,320
$(\mu + \pi + \varnothing \text{ in SciBar})$			
Both muon and pion are	8,535	1,845	884
forward going			
Muon enters the MRD and	7,407	1,592	767
penetrates $\geq 3$ layers of steel			
"Stopped"-Events	6,448	1,350	669
"Out-the-back"-Events	530	150	56
"Out-the-side"-Events	429	92	42
Good Coherent Pion Events	6,978	1,500	725

Table 4: Event reduction table for a sample of  $\nu$ -mode CC-Coh  $\pi$  events simulated in the SciBooNE geometry.





Figure 15: Muon momentum for all of the muons of the events that made it to the MRD (left) and muon angle for the muons of the events that made it to the MRD (right) for  $\nu$ -mode CC-Coh  $\pi$  interactions of all three models used in this study. The "Total" classification means that all three classifications for the CC-Coh  $\pi$  events ("stopped", "out-the-side", and "out-the-back") are included.






Figure 16: Muon momentum of both the "stopped" and the "out-the-back" samples (left) and muon angle of both the "stopped" and the "out-the-back" samples (right) for  $\nu$ -mode CC-Coh  $\pi$  interactions of all three models included in this study. The "Good" classification means that only the "stopped" and the "out-the-back" CC-Coh  $\pi$  events that the muon penetrated at least four layers of scintillator of the MRD are included for these histograms.

The last two quantities that are calculated are the two different types of four-momentum transfers specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{5}$$

This equation is in four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$

$$(6)$$

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{7}$$

This equation is in four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(8)

|t| is reported in units of  $(MeV/c)^2$ .

 $\nu$ -mode |t| and  $Q^2$  plots are below.




Figure 17: The |t| momentum transfer for the "stopped" and the "out-the-back" events (left) and  $Q^2$  momentum transfer for the "stopped" and the "out-the-back" events (right) for  $\nu$ -mode CC-Coh  $\pi$  interactions for the three models included in this study.




Figure 18: These two plots are the one-dimensional efficiency plots of the muon information. The left plot is the muon angle efficiency plot and the right is the muon momentum efficiency plot for  $\nu$ -mode.

Below are the two-dimensional efficiency plots for the CC-Coh  $\pi$  events in  $\nu$ -mode for all three models used in this acceptance study (NEUT v5.3.6 Rein-Sehgal, NEUT v5.3.6 Berger-Sehgal, and NEUT v5.0.1 Rein-Sehgal). The tables that correspond to the plots can be found in the Efficiency Tables section of the Appendix within the Charged-Current Coherent Pion Production Efficiency Tables subsection.




Figure 19: Two-dimensional efficiency plot for the CC-Coh  $\pi$  events of the  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal MC. Table 11 is the efficiency table for this two-dimensional efficiency histogram.




Figure 20: Two-dimensional efficiency plot for the CC-Coh  $\pi$  events of the  $\nu$ -mode NEUT v5.3.6 Berger-Sehgal MC. Table 12 is the efficiency table for this two-dimensional efficiency histogram.




Figure 21: Two-dimensional efficiency plot for the CC-Coh  $\pi$  events of the  $\nu$ -mode NEUT v5.0.1 Rein-Sehgal MC. Table 13 is the efficiency table for this two-dimensional efficiency histogram.

#### 5.2.2 $\bar{\nu}$ -mode Charged-Current Coherent Pion Events

Similar to before, Table 5 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi$  events from the  $\bar{\nu}$ -mode MC.

<i>ū</i> -mode	CC-Coherent	Pion	Event	Reduction
ν-mouc	CC-Concrenc	1 1011	L v CII t	ited action

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal
Total Sample	1,000,000	1,000,000
CC-Coherent Pion Interaction	36,669	7,790
$(\mu + \pi + \varnothing \text{ in SciBar})$		
Both muon and pion are	24,675	5,477
forward going		
Muon enters the MRD and	20,445	4,517
penetrates $\geq 3$ layers of steel		
"Stopped"-Events	18,935	4,203
"Out-the-back"-Events	372	82
"Out-the-side"-Events	1,138	232
Good Coherent Pion Events	19,307	4,285

Table 5: Event reduction table for a sample of  $\bar{\nu}$ -mode CC-Coh  $\pi$  events simulated in the SciBooNE geometry.

Below are the plots for CC-Coh  $\pi$  events for  $\bar{\nu}$ -mode. The layout of the rest will be very similar to  $\nu$ -mode, and the equations used previously for the momentums and angles are the same equations used for the variables of the histograms that follow.




Figure 22: Muon momentum (left) and muon angle (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi$  interactions for both the NEUT v5.3.6 Rein-Sehgal and the NEUT v5.3.6 Berger-Sehgal samples included in this study.

The structure of the plots in Figure 22 very closely resemble the same plots in  $\nu$ -mode above. The rest of the plots in this section have that same characteristic.




Figure 23: Muon momentum (left) and muon angle (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi$  interactions for both the "stopped" and "out-the-back" classification of events in both the NEUT v5.3.6 Rein-Sehgal and the NEUT v5.3.6 Berger-Sehgal samples.

 $\bar{\nu}$ -mode |t| and  $Q^2$  plots are below. They also have the same overall shape as the plots for  $\nu$ -mode above.




Figure 24: The |t| momentum transfer for the "stopped" and the "out-the-back" events (left) and  $Q^2$  momentum transfer for the "stopped" and the "out-the-back" events (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi$  interactions for both models included in this study.




Figure 25: These two plots are the one-dimensional efficiency plots of the muon information. The left plot is the muon angle efficiency plot and the right is the muon momentum efficiency plot for  $\bar{\nu}$ -mode.

Below are the two-dimensional efficiency plots for the CC-Coh  $\pi$  events in  $\bar{\nu}$ -mode. The tables that correspond to the plots can be found in the Efficiency Tables section of the Appendix in the Charged-Current Coherent Pion Production Efficiency Tables subsection.




Figure 26: Two-dimensional efficiency plot for the CC-Coh  $\pi$  events of the  $\bar{\nu}$ -mode NEUT v5.3.6 Rein-Sehgal MC. Table 14 is the efficiency table for this two-dimensional efficiency histogram.




Figure 27: Two-dimensional efficiency plot for the CC-Coh  $\pi$  events of the  $\bar{\nu}$ -mode NEUT v5.3.6 Berger-Sehgal MC. Table 15 is the efficiency table for this two-dimensional efficiency histogram.

# 6 Conclusion

The purpose of this acceptance study was to reproduce the acceptance model that was used in SciBooNE's original study, and to see the differences that resulted from multiple CC-Coh  $\pi$  models with a more modern NEUT generator (the old NEUT version originally used in the SciBooNE study is believed to be NEUT v5.0.1 and the modern version used in this study is NEUT v5.3.6). The two-dimensional efficiency histograms for CC-Inclusive events were made and are shown in Figures 8-10, Figure 13, and Figure 14 with the corresponding tables being Tables 6-10. The two-dimensional efficiency histograms for CC-Coh  $\pi$  were also made and can now be used in the re-analysis as they were intended with Figures 19-21, Figure 26, and Figure 27 being the associated CC-Coh  $\pi$  two-dimensional efficiency histograms. The tables that correspond to the CC-Coh  $\pi$  two-dimensional efficiency histograms are Tables 11-15.

While working with CC-Inclusive events we were able to reproduce the results of a previous study, which is evident in the close agreement between the one-dimensional efficiency histograms for the muon's momentum and angle with Hiraide's thesis in Figure 7. Though our model is simplistic, this agreement is a strong indication that our simplistic model is accurate and reasonable. In comparison with the two-dimensional efficiency histogram that was provided by Morgan for CC-Inclusive events, our two-dimensional efficiency histograms fail to corroborate what Morgan provided, as seen in Figures 8-10. This is a result that we do not understand, but we do realize that our acceptance study is being compared with two different studies, so we do not know for certain what this implies about our acceptance study.

The results for CC-Coh  $\pi$  events showed a strong difference between the Rein-Sehgal models and the Berger-Sehgal model in the four-momentum transfer variables: |t|, and  $Q^2$  (evident in Figure 17 for  $\nu$ -mode events and in Figure 24 for  $\bar{\nu}$ -mode events). This result has striking implications for the way that the data is sliced, but looking at the results of the two-dimensional efficiency histograms for CC-Coh  $\pi$  events (Figures 19-21 for  $\nu$ -mode and Figures 26-27 for  $\bar{\nu}$ -mode) there is no clearly distinguishable difference in the two-dimensional efficiency histograms between the different models. This implies that the detector is equally sensitive to both models, and that the issue of SciBooNE's non-observance of CC-Coh  $\pi$  is not a result of the detector not being sensitive to different models,

but is due to the slicing of the data.

The results above strongly indicate that our simplistic model is reasonable, and they also indicate that the SciBooNE detector is equally sensitive to the different models for CC-Coh  $\pi$  used in this acceptance study. This gives the clear implication that the cuts made for selecting CC-Coh  $\pi$  in data are model dependent, and the selection cuts would depend on which kinematic variables are used to distinguish a CC-Coh  $\pi$  event in the data. Future studies could be done on the differences in other kinematic variables between the different models, and there could also be a study that looked only at the events with a "stopped" muon and what their corresponding acceptance results would be.

# A Appendix: Sample Details

Appendix on samples

# A.1 $\nu$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$ -mode interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for charged-current coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

# A.2 $\nu$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$ -mode interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for charged-current coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack\_NEW.root

found at the following link (SciBooNE-MC-ROOTFiles).

### A.3 $\nu$ -Mode Rein-Sehgal NEUTv5.0.1

A sample of  $100,000 \nu$ -mode interactions were simulated using the NEUT generator (v5.0.1, believed to be the version used by the SciBooNE collaboration in the original publication) and the corresponding older Rein-Sehgal model for charged-current coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

#### A.4 $\bar{\nu}$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$ -mode interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for charged-current coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack.root

found at the following link (SciBooNE-MC-ROOTFiles).

#### A.5 $\bar{\nu}$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$ -mode interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for charged-current coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack\_NEW.root

found at the following link (SciBooNE-MC-ROOTFiles).

#### A.6 Vertex Distributions

The events were all given a random initial point that was generated with the goal that the vertex distributions of this simulation would closely match the vertex distributions that Hiraide <sup>3</sup> showed in his thesis. This was done using the code below and a set of histograms for one of the samples is shown below the code.

```
TRandom3 *randX = new TRandom3();
TRandom3 *randY = new TRandom3();
TRandom3 *flat = new TRandom3();
randX->SetSeed(jentry/2);
randY->SetSeed(jentry*jentry);
flat->SetSeed(jentry*jentry*jentry);
double Xpos = randX->Gaus(1.5,1.3);
while (Xpos<0 || Xpos>3.0) { Xpos = randX->Gaus(1.5,1.3); }
double Ypos = randY->Gaus(1.5,1.05);
while (Ypos<0 || Ypos>3.0) { Ypos = randY->Gaus(1.5,1.05); }
double Zpos = flat->Uniform(0,1.7);
```




Figure 28: Vertex distributions of the events in the  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal sample.

#### A.7 NewNMReinSehgal.C

This file is the macro that corresponds to the "NewNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms). When this macro is run (which can take a while), it also plots a few different histograms. The histograms that are plotted are the ones shown in the figures below with descriptions included with the corresponding figures. The order that the histograms appear in this paper is the same order they will be shown when this macro is run in root.

<sup>&</sup>lt;sup>3</sup>Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide dt.pdf




Figure 29: Vertex distributions of the events in the  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal sample.




Figure 30: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Inclusive events in the NEUT v5.3.6 Rein-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Inclusive.




Figure 31: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Coh  $\pi$  events in the NEUT v5.3.6 Rein-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Coh  $\pi$ .

The NewNMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{9}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{10}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle in spherical coordinates. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{11}$$

$$\theta_{\pi} = tan^{-1} \left( \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2} / P_{\pi_z} \right) \tag{12}$$

The angles are reported in units of  $^{\circ}$ , and should run from  $0^{\circ}$  to  $180^{\circ}$ .

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{13}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(14)

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}|$$
(15)

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(16)

|t| is reported in units of  $(MeV/c)^2$ .

# A.8 NewNMBergerSehgal.C

This file is the macro that corresponds to the "NewNMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack\_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 32: Vertex distributions of the events in the  $\nu$ -mode NEUT v5.3.6 Berger-Sehgal sample.




Figure 33: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Inclusive events in the NEUT v5.3.6 Berger-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Inclusive.




Figure 34: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Coh  $\pi$  events in the NEUT v5.3.6 Berger-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Coh  $\pi$ .

# A.9 OldNMReinSehgal.C

This file is the macro that corresponds to the "OldNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoOBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 35: Vertex distributions of the events in the  $\nu$ -mode NEUT v5.0.1 Rein-Sehgal sample.




Figure 36: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Inclusive events in the NEUT v5.0.1 Rein-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Inclusive.




Figure 37: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Coh  $\pi$  events in the NEUT v5.0.1 Rein-Sehgal sample in  $\nu$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Coh  $\pi$ .

#### A.10 NewANMReinSehgal.C

This file is the macro that corresponds to the "NewANMReinSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 38: Vertex distributions of the events in the  $\bar{\nu}$ -mode NEUT v5.3.6 Rein-Sehgal sample.




Figure 39: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Inclusive events in the NEUT v5.3.6 Rein-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Inclusive.




Figure 40: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Coh  $\pi$  events in the NEUT v5.3.6 Rein-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Coh  $\pi$ .

# A.11 NewANMBergerSehgal.C

This file is the macro that corresponds to the "NewANMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack\_NEW.root". This file performs the main

analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 41: Vertex distributions of the events in the  $\bar{\nu}$ -mode NEUT v5.3.6 Berger-Sehgal sample.




Figure 42: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Inclusive events in the NEUT v5.3.6 Berger-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Inclusive.




Figure 43: These are the two-dimensional histograms that when the left is divided by the right it returns the two-dimensional efficiency histogram for the CC-Coh  $\pi$  events in the NEUT v5.3.6 Berger-Sehgal sample in  $\bar{\nu}$  mode. The left only contains events that the muon "stopped" or went "out-the-back", penetrated the front face of the MRD, and passed through at least four layers of scinitillator before stopping. The right contains all of the events that were classified as CC-Coh  $\pi$ .

#### A.12 NMCombinedPlots.C

This is the file that performs the main plotting operations for the  $\nu$ -mode samples using the muon's information. All of the muon efficiency plots for  $\nu$ -mode are made with this file.




Figure 44: These are the  $\nu$ -mode one-dimensional efficiency plots of the muon information for CC-Inclusive events. The left is the angle efficiency plot and the right is the momentum efficiency plot for the muon in all three  $\nu$ -mode samples.




Figure 45: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal CC-Inclusive sample.




Figure 46: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.3.6 Berger-Sehgal CC-Inclusive sample.




Figure 47: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.0.1 Rein-Sehgal CC-Inclusive sample.




Figure 48: These are the two plots for muon angles in  $\nu$ -mode that are used to make the onedimensional efficiency plot for the angles in CC-Coh  $\pi$  events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either "stopped" or went "out-the-back" face, and penetrated at least four layers of scintillator of the MRD. The right plot is the muon angles for all of the events that were classified as CC-Coh  $\pi$ . The left divided by the right gives the efficiency plot for the muon angles in CC-Coh  $\pi$  events for all three  $\nu$ -mode samples.




Figure 49: This is the one-dimensional muon angle efficiency plot for the CC-Coh  $\pi$  events from the  $\nu$ -mode samples.




Figure 50: These are the two plots for muon momentums in  $\nu$ -mode that are used to make the one-dimensional efficiency plot for the momentums in CC-Coh  $\pi$  events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either "stopped" or went "out-the-back" face, and penetrated at least four layers of scintillator of the MRD. The right plot is the muon momentums for all of the events that were classified as CC-Coh  $\pi$ . The left divided by the right gives the efficiency plot for the muon momentums in CC-Coh  $\pi$  events for all three  $\nu$ -mode samples.




Figure 51: This is the one-dimensional muon momentum efficiency plot for the CC-Coh  $\pi$  events from the  $\nu$ -mode samples.




Figure 52: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.3.6 Rein-Sehgal CC-Coh  $\pi$  sample.




Figure 53: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.3.6 Berger-Sehgal CC-Coh  $\pi$  sample.




Figure 54: Two-dimensional efficiency plot for the  $\nu$ -mode NEUT v5.0.1 Rein-Sehgal CC-Coh  $\pi$  sample.

# A.13 NMPionPlotting.C

This is the file that performs the main plotting operations for the  $\nu$ -mode samples using the pion's information. All of the combined pion plots for  $\nu$ -mode are made with this file.




Figure 55: These are the angles of the pions in CC-Coh  $\pi$  events in the  $\nu$ -mode samples. The left plot is the angles for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for all three  $\nu$ -mode samples. The right plot is the pion angles for all of the events that were classified as CC-Coh  $\pi$  for the three  $\nu$ -mode samples.




Figure 56: These are the momentums of the pions in CC-Coh  $\pi$  events in the  $\nu$ -mode samples. The left plot is the momentums for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for all three  $\nu$ -mode samples. The right plot is the pion momentums for all of the events that were classified as CC-Coh  $\pi$  for the three  $\nu$ -mode samples.




Figure 57: These are the energies of the pions in CC-Coh  $\pi$  events in the  $\nu$ -mode samples. The left plot is the energies for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for all three  $\nu$ -mode samples. The right plot is the pion energies for all of the events that were classified as CC-Coh  $\pi$  for the three  $\nu$ -mode samples.

#### A.14 NMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for  $\nu$ -mode.




Figure 58: The |t| momentum transfer for the "stopped" and the "out-the-back" events (left) and  $Q^2$  momentum transfer for the "stopped" and the "out-the-back" events (right) for  $\nu$ -mode CC-Coh  $\pi$  interactions for the three models included in this study.

#### A.15 ANMCombinedPlots.C

This is the file that performs the main plotting operations for the  $\bar{\nu}$ -mode samples using the muon's information. All of the muon efficiency plots for  $\bar{\nu}$ -mode are made with this file.




Figure 59: These are the  $\bar{\nu}$ -mode one-dimensional efficiency plots of the muon information for CC-Inclusive events. The left is the angle efficiency plot and the right is the momentum efficiency plot for the muon in both  $\bar{\nu}$ -mode samples.




Figure 60: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode NEUT v5.3.6 Rein-Sehgal CC-Inclusive sample.




Figure 61: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode NEUT v5.3.6 Berger-Sehgal CC-Inclusive sample.




Figure 62: These are the two plots for muon angles in  $\bar{\nu}$ -mode that are used to make the one-dimensional efficiency plot for the angles in CC-Coh  $\pi$  events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either "stopped" or went "out-the-back" face, and penetrated at least four layers of scintillator of the MRD. The right plot is the muon angles for all of the events that were classified as CC-Coh  $\pi$ . The left divided by the right gives the efficiency plot for the muon angles in CC-Coh  $\pi$  events for both  $\bar{\nu}$ -mode samples.




Figure 63: This is the one-dimensional muon angle efficiency plot for the CC-Coh  $\pi$  events from the  $\bar{\nu}$ -mode samples.




Figure 64: These are the two plots for muon momentums in  $\bar{\nu}$ -mode that are used to make the one-dimensional efficiency plot for the momentums in CC-Coh  $\pi$  events. The left is the plot for the events that had a muon make it to the front face of the MRD and the muon either "stopped" or went "out-the-back" face, and penetrated at least four layers of scintillator of the MRD. The right plot is the muon momentums for all of the events that were classified as CC-Coh  $\pi$ . The left divided by the right gives the efficiency plot for the muon momentums in CC-Coh  $\pi$  events for both  $\bar{\nu}$ -mode samples.




Figure 65: This is the one-dimensional muon momentum efficiency plot for the CC-Coh  $\pi$  events from the  $\bar{\nu}$ -mode samples.




Figure 66: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode NEUT v5.3.6 Rein-Sehgal CC-Coh  $\pi$  sample.




Figure 67: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode NEUT v5.3.6 Berger-Sehgal CC-Coh  $\pi$  sample.

### A.16 ANMPionPlotting.C

This is the file that performs the main plotting operations for the  $\nu$ -mode samples using the pion's information. All of the combined pion plots for  $\nu$ -mode are made with this file.




Figure 68: These are the angles of the pions in CC-Coh  $\pi$  events in the  $\bar{\nu}$ -mode samples. The left plot is the angles for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for both  $\bar{\nu}$ -mode samples. The right plot is the pion angles for all of the events that were classified as CC-Coh  $\pi$  for both  $\bar{\nu}$ -mode samples.





Figure 69: These are the momentums of the pions in CC-Coh  $\pi$  events in the  $\bar{\nu}$ -mode samples. The left plot is the momentums for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for both  $\bar{\nu}$ -mode samples. The right plot is the pion momentums for all of the events that were classified as CC-Coh  $\pi$  for both  $\bar{\nu}$ -mode samples.







Figure 70: These are the energies of the pions in CC-Coh  $\pi$  events in the  $\bar{\nu}$ -mode samples. The left plot is the energies for the pions where the muon made it to the MRD and either "stopped" or went "out-the-back" face of the MRD, and penetrated four layers of scintillator of the MRD for both  $\bar{\nu}$ -mode samples. The right plot is the pion energies for all of the events that were classified as CC-Coh  $\pi$  for both  $\bar{\nu}$ -mode samples.

### A.17 ANMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for  $\bar{\nu}$ -mode.






Figure 71: The |t| momentum transfer for the "stopped" and the "out-the-back" events (left) and  $Q^2$  momentum transfer for the "stopped" and the "out-the-back" events (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi$  interactions for both models included in this study.

# B Efficiency Tables

This is the section of the appendix that contains the efficiency tables of the two-dimensional histograms that were seen throughout the technical note. This section is broken into two subsections: one for the CC-Inclusive two-dimensional efficiency histograms, and the other for the CC-Coh  $\pi$  two-dimensional efficiency histograms.

### B.1 Charged-Current Inclusive Sample Efficiency Tables

These are the corresponding tables to the 2D Efficiency histograms for CC-Inclusive events.

Table 6: Efficiency Table of 2D Histogram for  $\nu$ -mode CC-Inclusive NEUT v5.3.6 Rein-Sehgal

			_					_	_	_		_	_	_	_					_	_	_	_						_	_	
1930-2000	0.78574	0.713615	0.5625	0.453185	0.380932	133333								В			_		= 0						= 0						
1908-1920	78887.0	0.623653	0.530655	0.55932	12	0.1													= 0						= 0					0	
B81-881	813733	62620	509507	3833	299323	125			_		_	_	_	_	_	_	_	_				_	_	_				_	_	_	
800-1850	3835	684982	517647	100	272	921-58	71		_													_	_								
30.1800	198	2000	_	416667	12	22		2	_				_	_	_	_		_				_	_						_		
H	- 0	634561	-	28291		_	0 299991		_								0		= 0				_		= 0	-					0.00
H		53938 0.5 6.8718 0.6	_	24058		12	2810		_			-	-						= 0	0 0			_		= 0	0 0	0 0	0			0 00 1
900-1650 163	8 8	618935 0.7		1.63674 0.5		=	187		_					-					= 0				_		= 0	- 0					0 00 1
E	0.0	697572 0.7	_	301577 0.4	_	_	1936060	0.00	_			-	-				0		= 0		-		-	-	= 0	-				0.0	
	0.0	227468 0.69		1511166 051 1584875 035	-	929		0 0	_										= 0		-		_	-	= 0	= 0			В		
H		_	_		8	-	36803 0.16	2	_			-	-						= 0		-			-	= 0						
8	9		-			-		-	_										= 0		-		_	-	= 0						
H	66	2002	-	100000	_	_	20 0.16914 40 0.1747	_	_	-			-						= 0		-		_	-	= 0						
89	- 9	27 0 56146	-	1 0 2 6 8 8 8 8 7 1		_	14 014703												= 0	0 0					= 0	0 0					
E		0.735724	_	0.55685	_	_	0.39524												= 0						= 0	- 0					
0 123013	0.78982	0.730867	0.60652	0.458274		_	0.204918	-	0.33333			. =		В	В				= =						= 0				В		
H	0.3947	31880	-	9899	_	-	0.22(238	_	7														_		= 0						
1130-120	0.72272	0.713326	0.627978	0.553719		0.316279	0.191255	0.00	_										= 0				_		= 0				В		
1100 115	0.8385	0.746401	0.652881	0.56.628	0.62373	0.352837	0.27247	0.145161	0.432258										= 0						= 0						
1020-1100	0.828974	0.789633	0.632723	0.56538	0.403536	0.336748	0.2837	0.178295	7			. =							= 0				_		= 0						
1000 1020	0.83822	0.74026	890090	0.57886	0.62253	0.573851	26292	611010	0.457439	0.19096.4													_		= 0						
629-1000	0.79873	0.776828	0.656648	0578595	0.459893	0.362308	0.284258	0.12-614	0.0814815	0.0-465-465									= 0						= 0						
000 030	087664	0.75888	98796	0.524616	0.439182	0.371585	029367	7310110	0.122172	0.0840121	4000.400												_								
850-900	0.798611	12,540,80	06-71K5	05894	0.455387	1378892	129973	017748	0.106557	0.06010721	100			_				_					_		= 0						
800-850	816514	7867	961-199	542536	7465472	3,404237	312586	3267	12807	1974184	0212101			_	_	_	_	_				_	_	_					_	_	
28.830	877	713176	16745-0	54532	1.4862-22	3,416728	324022	115393	11318	0.0654045	01010101		_	_	_	_	_	_				_	_	_					_	_	
0.730	27.24.08 82.03.02	27.0396	067305	1537264	1512734	0.42(B(B	033975	18866	133764	188-625	10234114	0054641											_							_	
F		75513	-	536149		_	342862	_	=	0.083686	-	-			В				= 0	0 0			_		= 0	0 0		0 00	В	-	0.01
Ħ		1250	- 10	52336		=	331916	-	=	0751828 0.0	-	_					0		= 0				_		= 0	-				0	0 00 1
۳		787402 0.7		53869 0.5		_	3142	-	=	0.002000	0 0	-			0				= 0						= 0	-				0.0	0.001
12	0.816082 0.82 0.824047 0.73	0 0	-	280321 0.63	_	_		-	_	074586 0.08					0				= 0				0	-	= 0						0.001
H	<del>-</del> 2			B B B B B B B B B B B B B B B B B B B		_			_	0557781 0.07	Б	-			0				= 0				0		= 0						
7			-			~		0.00	_	-	0 0			В	В				= =	0 0			В		= 0	0 0					
H		67 0582	_	23 0.442424 82 0.381593	_	_	94 022549		_	922 0.0418139 0.000645726	0		0	0	8	0			= 0	0 0			0		= 0	0 0					
320-400		20 0479167	- 12	S 035622	-	_	8 0.17194		ᆖ	0.0165922									= 0	- 0				-	= 0	= 0					
3111-321		0.334426	-	0.28478		=	86 0129648		0.0248484					0	В				= 0	0.0			8		= 0						
250-310	0.218182	0.22900	0.23037	0.20088	_	_		0.015123											= 0						= 0				В		
200.230	0.25923	0.150948	_		_	0.0376749	0.018547	04004034											= =	- 0					= 0	- 0					
139 200	0.023457	0.0280374	0.0396825	0.034615	75281900.0			_ =											= -						=			_			
100-130						_													-			_							_		
V/c 59-100																			= 0	0.0					= 0	0.0					
WANTER BRIME															В					D 0			0 1	-	= 0				_		001
ew NATRS	0.45 Deg 459 Deg	435 Day	25 Deg	22.5.27 Day 27.31.5 Day	5.36 Deg	3-40.5 Drg	15-45 Day	15.54 Day	1.38.5 Deg	5-63 Day	7.72 Day	A S Dov	5.81 Deg	Sci Dog	15.90 Deg	90-94.5 Drg	45.99 Deg	99-103-5 Deg	10 3 10 Day	19 5 117 Day	17 1215 Dev	121 5 126 Deg	126-1305 Dry	130 5.135 Day	20 1395 Day	141 165 Day	148 5-4 50 Day	153-1575 Deg	1575-162 Day	162-1665 Day	71.175 Deg
	17	3 2	2 2	21 15	31	36	F 4	2 5	Š	25 5	3 12	13	ř	8	62	8	75	66	100	1 51	: =	12	12	g !	2 5	1 1		2	137	91 9	151

Table 7: Efficiency Table of 2D Histogram for  $\nu$ -mode CC-Inclusive NEUT v5.3.6 Berger-Sehgal

Column   C	
1,000   1,00	
1982   1981   1985   1982   1982   1982   1983	
77.20 (2012) (20	
1,200   1,20	
7783 (2012) (2012) (2017) (201	
1,242   1,24	
7720	
7790 (2011) (202	
7.720	
7.720	
1982 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
100 100 100 100 100 100 100 100 100 100	
77707 (1970)	
1.77 (2.2) (	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
100   100	
1.7.7.2.2.2.1.1.1.1.1.1.1.1.1.1.1.1.1.1.	
10 10 10 10 10 10 10 10 10 10 10 10 10 1	
20 10 20 20 20 20 20 20 20 20 20 20 20 20 20	
100 100 100 100 100 100 100 100 100 100	
20	
100 100 100 100 100 100 100 100 100 100	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
10.10 (10	
10 10 10 10 10 10 10 10 10 10 10 10 10 1	
202744222222222222222222222222222222222	
m b	
1	
10.000 10	
1401-500 10.201350 10.201350 10.201350 10.201350 10.2014	
1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1	
30.0021 10.00221 10.00221 10.00221 10.00221 10.00221 10.00221 10.00222 10.0022	
2015.00 201	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
12 12 12 12 12 12 12 12 12 12 12 12 12 1	
10.000 (10.000	
	17.17.5 Deg
(2.0 D ) (2.	762

Table 8: Efficiency Table of 2D Histogram for  $\nu$ -mode CC-Inclusive NEUT v5.0.1 Rein-Sehgal

-	
1 156 200	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1
1 0.533333	0.77.778 0.4 1111 0.2 10.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
1850-1900 0.8 0.882353	0.98420.5 0.70 0.70 0.825 0.82
1806-1850 0 0857143	0.055533 0.0555
750 1800 1 566667 1 923077	0.47 0.47 0.47 0.47 0.47 0.47 0.47 0.47
8 .	2.7.12.98 2.7.2.38 2.7.2.38 2.7.2.38 3.7.2.48
	6 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
1 65667 1 66667 1	7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7
0.16m	20 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1550 155 9001 0.77 1905 0.87	200
1450-1500 1500 1835533 0.50 1835533 0.70	15735556 1585 1585 1585 1585 1585 1585 1585
52	
1408 1408 1	further many in
858 1359.1 0 8333 177 0 825	18-18-18-18-18-18-18-18-18-18-18-18-18-1
300 1300-135 0.6 11 0.923077	25 75 75 75 75 75 75 75 75 75 75 75 75 75
1230-130	23 2 4 7 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
0.75 0.75 0.74859	10.771.05.0 771.05.0 771.05.0 771.05.0 771.05.0 771.05.0 771.05.0 771.0
0.75 0.75 0.8	0.000
1108-115	0.775374 0.755746 0.75576 0.75
0.777778 0.777778 0.807092	18851864 188588864 18941170 184717 1845181 185618 185618 18
98 98 98 98 98	0.7478.53 0.7208.53 0.7208.53 0.7308.53 0.74708.88 0.74708.88 0.74708.88 0.74708.88 0.74708.88 0.74708.88 0.74708.88 0.74708.88
950-1000 0.709231 0.810811	0-778187 0-6155165 0-6155165 0-6155165 0-6155165 0-615517
900-950 1 0.792-453	0.7351985 0.512085 0.513238 0.51328 0.5132
850 900 0.777778 0.782609	0.55257 0.575257 0.555555 0.555555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.55555 0.5555 0.55555 0.55555 0.555
0000	17.15881 64.4216 64.4216 64.62837 64.62
1-00	0.779179 0.659974 0.559874 0.51688 0.31672 0.3
0.77778 0.77778 0.74859	1,312,135 1,312,136 1,312,136 1,512,
650-70 8.7 8.8	10.1741.77 10.174
0.857143 0.857143	0.821853 0.821853 0.861823 0.861823 0.861823 0.861823 0.861823 0.86183
550-600 1 0.8	0.773.199 0.673.199 0.673.199 0.673.199 0.6771-149 0.67
500-550 0.505667 0.904762	1,500,577 1,500,577
50 500 0.5 0.608696	0.628.577 0.537 0.537 0.537 0.534 0.534 0.534 0.534 0.535
_	1921 152 1921 152 1921 152 1935 153 1935 153 193
700	A 451 61 13
000	12.85(25) 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
_	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
208-258 25 8-323323 8-2 8-323323 8-2	1,270,231 1,270,
28.28	14.7 (2.58)
108-158 158 8 8 8	
31 - 180 0 0 0	
B-SI MeV/c	
45.9	10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

Table 9: Efficiency Table of 2D Histogram for  $\bar{\nu}$ -mode CC-Inclusive NEUT v5.3.6 Rein-Sehgal

E		
1930-200	3/15/12/16/16/16/16/16/16/16/16/16/16/16/16/16/	
1900-1950	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
856-1900	100 100 100 100 100 100 100 100 100 100	
E81-90	2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2	
81 880 18	22.4.2.2	
21 821	20 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
E1 E21		
650 1650		
1600-1	7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7	
13281	2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
1200-1	2	
1420-13	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
1:488-1:5	1	
1359-140	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
13@-133	10 (1998) 10 (19	
1250-1310	2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
1200-1230	12.00 (19	
1130-12E		
1100-1150	6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	
020-1100	2	
0501-00	2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	
60-1000	120 120 120 120 120 120 120 120 120 120	
10.93	18.54.00 18.54.	
06.05	28-27-28-27-28-27-28-27-28-27-28-27-28-27-28-27-28-27-28-28-28-28-28-28-28-28-28-28-28-28-28-	
8 058 08	20 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
98 008 OE	25 25 25 25 25 25 25 25 25 25 25 25 25 2	
120	2 (2) (2) (2) (3) (3) (4) (4) (4) (4) (4) (4) (4) (4) (4) (4	
200		
E 650		
(B) (B) (F)		
19982	#	
500-550	2	
100 E	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
E1-80	10,000,000,000,000,000,000,000,000,000,	
330-400	13.10.24. 13.10. 13.10.24. 13.10. 13.10.24. 13.10.24. 13.10.24. 13.10.24. 13.10.24. 13.10.24. 13.10. 13.10. 13.10. 13.10. 13.10. 13.10. 13.10. 13.10	
300-220	8.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 14.0 15.0 15.0 15.0 15.0 15.0 15.0 15.0 15	
239.300	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Ī
20 250 25	17.73 (2.73	
150 200 20	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
100-150 150		
59 100 10		
0.50 MeV/c		
ew ANM R S   0	6.45 by 9.85 b	17 175 Day 0
New Al	1, 1, 2, 3, 19, 19, 19, 19, 19, 19, 19, 19, 19, 19	12.12

Table 10: Efficiency Table of 2D Histogram for  $\bar{\nu}$ -mode CC-Inclusive NEUT v5.3.6 Berger-Sehgal

Column   C	_																																							
Column   C	1930-2000	0.809524	0.68367	0.613924	67178140	0.482730	033333																								_		_							
The control of the	1900-1950	0.823329	0.716827	0.450888	0.552632	0.405405	0.555556	2																							_		_							_
	1830-1900	160220	120070	0.628571	0.527530	0.297297	0.33333	0.33333	_					-	-		-	-											-	-	_	-	_					_		_
	1800-1850	0.815789	0.008905	0.627737	0.525641	0.442623	0.136364	_																							_		_							
	1720-1800	7623620	277488	1682191	122331	0.46783	389932	_			_	_	_	_	_	-															_		_					_		
	921-98	1.824074	01 (BE	1452068	531915	0.443299	7.	299991	_	_	_	_	_	_	_	_	_	_	-			_				_	_	_	_	_	_	_	_		_			_	_	
	H21-0591	1825248	1502030	1654737	58333	1,477,273	0.446809	133294	- 2	_	_	_	_	_	_	_	_	_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
	091-00	871785	31236	18848	802208	216697	263889	33333	_		_	_	_	_	_	_	_	_				_				_	_	_	_	_	_	_			_			_	_	
Column   C	530-1600	825397	2002	1838201	788m8	760828	330189		799991	-	_	_	_	_	_	_	_	_											_	_		_						_		
	300 1350	823129	20046	88.0.98	381722	301502	=	77		_	_	_	_	_	_	_	_	_				_				_	_	_	_	_	_	_			_			_	_	
	450-150	299988	7,4634	997.739	5701-8	E988	407609	33333	157856	_	_	_	_	_	_	_																						_		-
	1408-1-50	1304	1982	883662	803365	318738	401487	32307	25224	285714	_	_	_	_	_	_																						_		-
The color of the	330.1400	-	-	Ē	÷	÷	485531	30303	_	Ī	17	_	_	_	_	-	-												-	-								_		-
	310-1350 1	819005 0	Z21674 II	Ĕ	_	12	-	_	_	_	0	0	0	0	0	_						-									_	0			-			_		
	330-13HI I.	0 0		_	_	_	Ť	_	Ť	_	33333	_	0	0	0	0	0	0											0	0		0								
	300-1250 1:	867133 0	72/66	=			Φ.		Φ.	_	285714 0	0	0	0	0	0																								
	30-1200	-	-	Ĕ	Ť	Ť	Ĕ	Ť	_	Ť	Ť	0	0	0	0	0	0	0											0	0		0						_		0
Column   C	00-H50 D	0.0	-	_	_	_	_	_	_	Ï	Ï	142857 0																												
	30-11-00	-		Ĕ	_	_	0	_	_	Ĕ	Ĕ	=	1759231																			0						_		-
Column   C	00-1020 10	8 0	0 0	_	_	_	301982 0.5			_	_	_	2			0															=							0		
	01 0001-0	0 0	-	_	_	_	_	_	Φ.	-	_	=	285714 0			0																0								-
	900 900	0	-	_	_	_	_	_	=	_	_	_	15	1125	-	-	-	-				_							-	-	_		_		-	-		_		- 10
	828-900	22092	795267	723615	98991	1.6(25.48	1525974	3,455385	332032	1253836	1228395	1202572	_	Ξ	9.67619	_	_	_	-	-	-					_	_	_	_	_	_	_	_	-		_		_	_	
1	820	8-8428	75586	231489	99229	613815	538056	762/097	3983	311813	254042	7108817	2998-11	=	10552381	_	_	_				_				_	_	_	_	_	_	_			_			_	_	
1,000,000,000,000,000,000,000,000,000,0	0.08-022	186324	1918	1,7337	1673-84	345687	1545725	1265	1229871	1320774	1259352	1210306	1149296	1.0851064	1.0933333	_	1.0833333	_	_	_	_	_			_	_	_	_	_	_	_	_		_	_		_	_	_	
1,000,000,000,000,000,000,000,000,000,0	052-00	7062281	1000	99-92	699989	80-4035	539545	-56883	201704	1297.021	982380	176355	12-426	18946372	187847	9869810	_	_											_	_								_	_	
1	99-300	200110	78362	7.8866	689239	629528	538263	298967	4229-6	3.8088	1912	203957	1.8563	2986180	0397554	01652486	786667																					_		
Company   Comp	9 629 6	834184 0	75014	74735	_	_	Φ.	500339	418747	337145 0		_	_	_	_	_																						0		
1,000,000,000,000,000,000,000,000,000,0	9	0.0	16837		_	_	_	_	_	16955	92936	_	_	_	_	=																								-
	230	_		-	-	_	÷	=	÷	÷	_	_	_	_	=	Ξ							-		-											-	-	0		
1	200	180		5	9.6	1980	2	70	2	. 8	87	13	3	8	8	8			-				-			-	-	-			=		=					0		-
		8.0				_	572 B.445	200 120		192	13 0.215	Ť	Ť	-	-	1000	0	0	-	-	-				-	-	-	-		0	=	0	=	-		-	-	0		
	9-901-00	1 20	- 24	9	721 0.4592	21 0.44%	25 B 3575	3335 B 3335	925 B.286b.	68 0.2352	22 B128	_	_		0.0075				-	-	-					-	-	-			=		-	-		-				
1802.01 (1800.00) 1802.01 (180	330-4		5 6	_	725 0.4237	67 0.3857	64 0.3226	61 82756	B 0.2345.	0	-	-	-	0.0156					-	-	-					-	-	-			=		-	-		-				
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	300 35	0.0	0 0	_	_	_	_	-		_			0.0216						-	-	-					-	-	-			=		-	-		-				
	T 230 36			_	_	_	_	_			0.0355	0.0115							-	-	-					-	-	-			=		-	-		-				0
	200.23	_	-	_	-	_	_	0.6525.	0.0441	0.0210							0	0	-	-	-					=	=	=	0	0	=	0	=	-		-		0		0
	50 139 200	0.09302	0.01000	0.04013.	0.02777	0.01964	0.00315.																								=		=			_				-
																			-	-	-					-	-	-			=		=	-		-				0
		0		0		В	0	В	0	0	0	0	0	0	0	0	0	0											0	0		0								
14.15   1.00   1	(BS 0.20	0			-	0 26		0 26	25							0 20	-	-						0	Doc 1	Dev	Dev	Dev	Deg	Deg	Dov	Deg	Dov	Down	Deg	Down	Doc 1	Deg 0	Dog	Day 10
	New ANA	8-45 Deg	4.25 Leg	135.18 De	18-22.5 Dt	225.27 Dt	27-31.5 Dt	315.36 Dt	36-40 5 D.	40.5-45 D.	45-85 Dt	495.54 Dt	54-28.5 D.	585.63 Dt	63-67-5 Di	675-72 D.	72.R5.Dt	765.81 De	81-85.5 De	855.90 De	98-94.5 De	945.99 De	00,103 5 L	105 5.108	168-1125	1125417	1171215	1215-126	126-1305	135.135	135-1395	1355144	1441485	1485453	133,1575	1575,189	162,1665	1665474	17.1755	13,5381

## B.2 Charged-Current Coherent Pion Production Efficiency Tables

These are the corresponding tables to the 2D Efficiency histograms for CC-Coh  $\pi$  events.

Table 11: Efficiency Table of 2D Histogram for  $\nu\text{-mode}$  CC-Coh  $\pi$  NEUT v5.3.6 Rein-Sehgal

199-200	0,00	0.00	9	2							_												_		_	_								_	_				_
	1 1/20/0			è	-			_	_		_		_		_		_					_	_		_	_						_		_					_
1831-1900 1900-1950	COSSESS OF		90			-		=	0	0	0	0	_	-	-	-	-	-	-	-	0	_	_		_	-	0 0		0 0		-	-	-	_	-	-	0.0	0	_
$\vdash$	100	-	-	3 -	-	0 0	= 1	=	0	0	0	0	_		-	-	-	-	-	-	-	0							-	-	-	-	-		-	5 0	-	-	_
170 lan 180 lan	9		-	8 -		-	-	=		0	0	0	_	-	_	-	-	-	-			_	_		_	-			-	-	-	-	-		- 0	-	-		_
1.501	orestro o	-		3_		- 0	-	=		0	0	0			_		_	-	-											-		_			-	- 0			
1,004730		0		Common			- 1	=		0					_		_	-	-											-		_			-				
1650,700	90	9 6	86	0 0	3 .		- 1	=					_		_		_		. =			_			_	_				. =		_			- 0				
16001650	000000	O COLUMN	neecon.	. 20	9		-	-							_		_															_							
0.001023	0.0	-		11110				_	-	_	_	_	_		_		_	-		-		_	_	_	_	_						_		_	_				_
022100	platet	20000		2000				=	_	_	_	_	_		_	_			_			_			_	_				_	_		_						_
1501301	200000		- 0	-		0 0		=	0	0	_	0	0		-		-	-		-	0.0	0	0	0	_	_	00		0 0			-		0	-	-	0 0	0 0	0
_	ť		-	-	1	8	-	=	0	0	0	0	_	-	-	-	-	-	-	-	0 0	_			_	-	0 0		0 0	-	-	-	-		-	-	0.0	0	_
E719091	2000000		,	_			- 1	=		0	0	0	_	-	-	-	-	=	-			_	_	_	_	_			-	-	-	-	-	_	-	- 0	-		_
97997 83	100000	-	-	-	8 1	0	-	=	0				0		_	0	-					0			0	0					0	-			- 0	- 0			. =
	0.7007	out the	0 0	-	-	- 0	- 1	=			_				_		_						_		_	_						_				- 0			
5	che loo	00000		0.714385		9	-	=			_		_		_		_		. =			_	_		_	_				. =		_			-				
170	0.70	O TOP	2000	o ereste	1			_		0	0	0			_		_															_							
8	2010	10000	9	00000			_	_	_	_	_	_	_	_	_	_	_	_	_	_		_	_	_	_	_				_	_	_	_	_	_			_	_
8	60000	- Stoles	2000	78 Sat	-	0		_	_	_	_	_	_	_	_	_	_		_			_	_		_	_				_	_	_	_	_					-
8198	000000	0 0000		-	_	9	- 1	=	_	_	_	_			_		_				-		_	_	_	_	0.0					_		_	-	-	-	-	_
1001400	00110	-	0		9 1	own of	4 1	=	0	0	0	0	_		-		-	-	-			0			_				-	-		-				0			_
=	000	400		off of the	_	200	2	=	0	0	0	0	_		-		-	-	-			_	_		_	-	0 0		-	-		-			-	- 0	-	0	_
	0.00	or o	9	9.0	2 1		9	1 535555	0	0	0	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-			-	-	-	-	-	-	-	= 0		-	-
88	000	0 0	- 0	j		0.00 mm	3	_	-	0	0	0	_		-	-	-	-	-			_	_	_	_	_	0 0		-	-	-	-	-	_	-	- 0		0	=
882	20 69500	3 0	96	i c	2		_	000 / MOSO	0	0	0	0	0		_		_	-	-		0	0					0 0			-		_			- 0	- 0	0.0	0	_
2	000000000000000000000000000000000000000	-	0 0	901		0 0	-	960	-	-	0	0	_		-		-	-	-		-	_	-	-	_	-			-	-		-		-	- 0	= 0		-	_
B (2)	1100		- 0	200		000	2 1	=	0	0	0	0			_		_	-	-		0				_		00			-		_			- 0	- 0	0.0	0	=
_	200			200		Stored over		920	2	8	0	0	_	-	-	-	-	-	-			_			_	_	0 0			-	-	-	-		-	- 0		0	=
3	90	. 6	0000	200			2	9	6.6	68	0	0	_	-	-	-	-	-	-	0		_	_		_	_	0 0			-	-	-	-		-	- 0		0	=
	Con plotter	Titolo godin			1	COUNTY DOWN		9	40/40/ 0.44	0.142857 0	0.333333	0	_	-	=	-	-	-	-	0		_	_		_	_	0 0			-	-	-	-		-	- 0		0	=
Heart Southfull	f	-	9 0				2	Ė	_	_	ŭ	0 29000	_	-	-	-	-	-	-	-	-	-	-	-	-	-			-	-	-	-	-	-	-	= 0			-
n	200	0 0	-	5 0		8 8	2	ì	â.	ä	â	ö	_	-	-	-	-	-	-	-	-	-	-	-	_	-			-	-	-	-	-	-	-	= 0		-	_
2	action of the	_	_		-	A female a conference		USSS00 USS041	032281 0.42	3RS 022	0.173913 0.352911	999		_	-		-	-	-		-	_	-	-	_	-			-	-		-		-	- 0	= 0		-	_
7	f	-	-	-	-	-	_	_			315 O.17	V83 023	533 0.12	-	_		_	-	-		0 0						00			-		_			- 0	- 0	-	0	_
3	2000		0.0000		-	24 034	_	_	_	.9		316 0.0434783	SS DUKKKER		-	0	-	-	-		-	0			-				-	-	0	-	0		- 0	- 0			_
3165	t	9 9	_	-	_	-	-	_	=	7 0.190476	22 0.1855	38 notestade	occide occides	89	_		_	-	-			0	_			0				-		_		_	- 0	- 0			-
200	2000	_	-	-	1 9	-		Ξ.	=	0048387	00588285	ordestas	Dressed		_		_	-					-		_							_			- 0	- 0			-
4	20.00		1000		_	1000		ngs/sp	0.115385				_		_		_					_	_	_	_	_						_		_	_				
9		Dog Conflored	1000000	longrapol o	-			-	0.029418																														
								=							_		_	-					-		_	_						_			_		-		_
	ļ		-	-	-	-	-	_	_		_		0		_		_		-		-	0	_	_	_	_	-		-	-		_		_		-	-	-	
BORDON C SMILE THEFT			- 0				-	-					0		_		_					0			0	0						_							
New YORKS	-	1		Passage New	9	200	9	200	201782	15 Deg	405-45 Deg	15 Deg	495.54 Deg	54585 Deg	SDE	63-67.5 Deg	72 Dec	72765 Dec	S Dec	STOR	85.500 Dec	(5 Deg	90 De	B5Dg	SIGN Deg	108-1125 Dvg	1 P.	19.0	1251315 Dec	135 Dec	205 Deg	144 Dec	485 Deg	1485133 Deg	1531575 Deg	801	less 17 De	755 Dec	MB Deg
New.	0.40	2 .	7 5	6 T	2 .	168	6 1	20.7	15 25	35.55	9	55.65	(5.0)	15	8	929	100	72.78	76.55	100	1 75	1606	35	30 FE	1035	168	177	12	185	130.5	Ţ.	0.00	144	1485	B	1 5	1 2	E	1755

Table 12: Efficiency Table of 2D Histogram for  $\nu$ -mode CC-Coh  $\pi$  NEUT v5.3.6 Berger-Sehgal

_	_						_	_														_													
90.300	7300000		_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_							_	_	_			
09670			_			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_					_	_	_	_				
1900			=		0		_		_	0		_	0	0	_	0			_	0	0			0 1	0 0		-	-		0	0	01	= 1	= 0	-
1880			-					_			_											_	_	-			-	-		0		0	= 1	= 0	-
1800-18	2-		-				_	_	_	_	_		_	_	_				_	_		_	_					. =					= 1	= 0	
Pariso	0.4	2	_																														-		
001780	299990																																		
71 00/14	200000	790000	=			-	_	_	_	_	_	_	_	_	_	0	_	-	_		0	_	_	0 1	0 0			_	-	_	0	0 1	= !	0.0	
620 1650	999	80.	-		0	_	_	0	_	_		_	_	_	_	0	_	_	_	0	0	0		0 1	0 0		-	-		_	0	0 1	= !	= 0	-
10001	٥-	-	=					_			_									_		_	_	- 1				-				-	= 1		
1550100	10,0007	2.	-					_			_											_	_					-					-	= 0	
300-1550	<i>1</i> 2 a	2 12				_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_							_	_	_			
005100	0.833338				_	_	_	Ē	_		Ī	_			_			_	_	Ī		Ē	Ī						_	_	Ī				
1450 114	80 8	100	-	0.0		_	_	_	_	_	_	_	_	_	_	0	_	_	_	_	0	_	_	-						_	_	-	= 1	= 0	_
140	- 8		-					_			_		-			0			0	_	0	_	_	-			-	-	0	0	_	-	= 1	= 0	-
1350.1	0.7	0.838									_												_	- 1	-			-					= 1	= 0	
130013	0.875	1872	_	2.			_	_	_	_	_		_	_	_					_		_	_					-					- 1	= 0	. =
2301300	050	088883	2			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_							_	_	_			
1987-1987   1987-1987   1787-28   1788-198   1388-198   1388-198   1488-198   1588-198	200000		_			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_					_		_	_				
120012	000000	_	_			_	_	_	_	0	_	_	_	0	_	0	_	_	_		0	_	_	0 1	0 0			-	-	-	0	01	= 1	= 0	
EU US	F .	6	3			0	0	_	0		_				0	0				_	0	_	_	- 1			-	_		0	_	-	= 1	= 0	-
11001	0.81882		g																					-	-			-					-		
	I ordente	G.	188888	2.			_	_	_		_	_	_		_					_		_	_	-				-					- 1	= =	. =
10004050	5.50 5.00 5.00 5.00 5.00 5.00 5.00 5.00	g.	11128		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_							_	_	_			
0001400	290000	8	9			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_				_	_	_	_	_				
6 06-06	0857143	8	9		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-		_		_	_	_	_	-	-	-	
80.900	0.77777		Ξ.	98777		_	_	_	_	_	_	_	_	_	_	0	_	_	_	_	0	_	_	-						_	_	-	= !	= 0	
80.80	1 020170		1000000	30	-	0	_	_	_	0	_	_	_	0	_	0	0	_	0		0	_	_	-			-	-		0	0	-	= 1	= 0	-
750-800 BIC	-10		200	88			_	_	_	0		_	_	0	_	0			_	0	0	_		0 1	0 0		-	-		0	0	01	= 1	= 0	-
$\vdash$	20 88	0	3	8	-	0	0	_	0	0	_			0	0	0		0		_	0	_	_	- 1			-	_		0	_	-	= 1	= 0	-
0.70070	0.833333	Sans		28 CO						_			_	_					_					-	-		-	-				-	= 1	- 0	
002020	0.000	8	=	S COLUMN	-	_	_	_	_	_	_	_	_	_	_				_			_	_	-			-	-		_		-	= 1	= 0	
0.090.00	0.92307		_	0.016067				_		_	_	_	_	_					_			_	_	-				-					= 1	= 0	
330600	0.916667		10000	2 0.55 0.628364	20	_		_			_					0				_	0	_	_	-	-			-			_		= 1	= 0	
200-230	1 0 or 0 ave	8		1000	0.2857	191	_	_	_		_	_			_		_	_				_	_	-			. =	. =		_			= 1	= 0	
98-8P	0.7cm/8 0.85	0.00000	3	200000			_	_	_	-	_	_	-	-	_		_	_		_		_	_	-				_			_	-	= 1	= -	_
400-620	0.75/7/08	2	9	0.788462		0.335353	0.142857	_	_	-	_	_	-	-	_		_	_	-			_	_	-	-		_	_	_	_	_		-	-	
30,400	¥3-	e)		9 50 50		935353	_	500000	200000		_	_			_		_		-	_		_	_	_				-	_	_		_			
300300	95			13038	77	_	_	oleess?   c	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_				_	_	_	_	_				_
2030 3			_	00000031	-	oliiii o	÷	olesse 0.		0	_	0	_	0					_	_		_	_	0 1	2 0						0	-	2 1	20	
$\vdash$	- 3		_		_	_	22	_	_	0	-	_	0	0	_	0	-	_	0	_	0	_	-	-		-	-	-	-	_	_	-	= 1	= 0	-
30.20		집.		0.133333	GISL	0.142	888	0111111	_		_	_	-		_	0	_	_	0	_	0	_	_	-			_	-		_		-	= 1	= 0	-
10-131 130-201		-	_			_	_	_	_		_	_	_		_		_	_	0	_		_	_	-			-	_		_		-	= 1	= 0	-
1001			=				_	_	_		_	_	-		_							_	_	-			-	-		0		-	= 1	= 0	-
0.50 MeV c 50 too		-	=			_	_	_	_	-	_	_	-	-	_	0	_	_	0		0	_	_	- 1	-	-	-	-	-	_	_	0 1	= 1	= 0	-
0-SOME			=					_					_							_		_											-		
New NM BS	2 Deg	9-13.5 Drg	Saran.	225.77 Deg	Z7315Deg	36 Deg	95 Deg	40.5-45 Deg	95 Deg	StDeg	54585 Deg	28.5463 Deg	75 Deg	72 Deg	72765 Deg	SI Deg	81-85.5 Deg	Salbe	90945 Deg	and one	RS5 Deg	SIGNOR	1125 Deg	112511/Deg	135 Dec	Parshe	JED De	Data Dec	SIMDE	1485 Deg	SISBDE	153-157.5 Deg	200	100000	175.5 Dec
New	14.5	8	2	2 1	27.3	の で	ž	9	ij	99	古	88	8	6	ĕ	765	8	12	88	3	ã	ğ	9	Zį į	15	2	Ē	1	18	144	148;	ń	5	1 8	Ē

Table 13: Efficiency Table of 2D Histogram for  $\nu\text{-mode}$  CC-Coh  $\pi$  NEUT v5.0.1 Rein-Sehgal

-	
7 α α	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
250000	
0.827.15	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
0.000	
3	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
1 (1777-177) 1 (17	1
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
8	-
1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
20 (20 (20 (20 (20 (20 (20 (20 (20 (20 (	
8 8 c	-
_ D=0	
at DU L	
C00000000000000000000000000000000000	
1 0 0 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
10000000000000000000000000000000000000	
25 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
100 100 100 100 100 100 100 100 100 100	
### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73 ### 10.727.73	
1 0.25 1 0.25	
224-310 10.25 10.10 10.1	
	58,
10.00 Mg 10.	171.175.5 Deg
> P + o 0 M M M 9 + + + 0 M M M D M M M M H H H H H H H H H H H H	

Table 14: Efficiency Table of 2D Histogram for  $\bar{\nu}\text{-mode}$  CC-Coh  $\pi$  NEUT v5.3.6 Rein-Sehgal

Column   C		
Column   C	950200	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Column   C	0510	
Column   C	61 0051	
Column   C	80 183	
Column   C	180	
Column   C	ᆮ	
Column   C	1700173	다 가 년 경기 : 1년 당기 : 1년
Column   C		88.21 88.21
Column   C		
Column   C	1001	
Column   C	F	
Column   C	$\vdash$	
Column   C		1886년 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
Column   C	1400-1450	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Column   C	201400	18181818181818181818181818181818181818
Column   C		
Column   C		
Column   C	1	
Columbia   Columbia		<u> </u>
Columbia   Columbia	1	19,888.89 (2,738
Columbia   Columbia	1100-1130	8. 14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15
Columbia   Columbia		23 28 28 21 28 28 28 28 28 28 28 28 28 28 28 28 28
Columbia   Columbia		b. 00.00 10
Columbia   Columbia	$\vdash$	87.87.48.4
Columbic of the party   Colu	$\vdash$	
Columbia   Columbia	-	
Columbia   Columbia	г	
Columbic of the part   Part   Columbic of the part   Columbic of t	$\vdash$	
Columbic Chief   1175   125	E	
Columbia   Columbia	F	
Columbia   Columbia	$\vdash$	
Columbic 2001   University   Columbic 2002	-	
1		22223 883 8
Columbic State   1975	80.68	
1	98.89	
1	0S)+00p	10,000 Sept. 10,00
1	330400	15 - 16 - 17 - 17 - 17 - 17 - 17 - 17 - 17
1	Г	
1	Г	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Н	10 - N N
	$\vdash$	[중하다 등학 주대형 프로젝 건강한 경험 전 경 경 경   '' 표 및 기계 등학 기
March   Marc	150200	100 100 100 100 100 100 100 100 100 100
Windows   Wind	0 10013	
	010S 0/	
March   Marc	D-SD MEV	
图 医皮肤 化硫酸 化二甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基	ANTES	10 P P P P P P P P P P P P P P P P P P P
	New	医安森氏征 化单位性 经未未存款 经现代的 医多种 电电子电话 电电话电话 医克里克氏

Table 15: Efficiency Table of 2D Histogram for  $\bar{\nu}$ -mode CC-Coh  $\pi$  NEUT v5.3.6 Berger-Sehgal

1981-2011	_	_	_		_															_				_				_			_							_							_			
1/201800 1800 1800 1800 1900 1900 1900 1900		0.714286								_		_							_																								_			_	_	
100	_	0.000007	_	-	_				_	_	_	-	-		-	_	_		_	_		=	_			-	_				=	-	-	-	0.0			=	0	_		-	_	-		_	10	
1001	-	90	_	0	_			0.0	-	0	0	0	-	0.0	-	0	-		0	=	0 0	=	0	_		-	_	_	-	-	=	0	0	-	0 0	0 0		=	0	-		-	0	0	_	_	-	0 0
100	-	0	_	_	_				=	_	0	0	0		=	0	_		_	_		=	0	_		=	_	_	-		=	0	0	-	0	0	= 1	=	0	_		-	-	0	_	_	10	-
	-	89		0	_				-	-	0	0	c		=		_		-	_	0 0	=		_	1 0	=	_	_	-		=		0	-	0 0	0 0	- 1	=	0	_		-	-		_	_		
THE PERSON	9.9	082		0	_				-		0	0			=		_	,		_		=		_		=		_			=		0	_			-	=	0	_		-			_			
	88/148	British			_				-	_					_		_	,	_	_		=		_		=		_			=			_			- 1	=		_		-	_		_			
DESCRIPTION.	_	100	100		_										-					_		-		_		-		_			=			-			-	=							_			
LANGE OF THE	_	1983830	033333	_	_					_	_								_	_		_		_				_			_		_				-	_					_		_			
DEFECT	0888888	98	6	_						_	_	_							_													_	_										_	_		_	_	
A LINE L	88		_	_	_		-		-	=	_	_	_		-	_	-	_	=	=		-	_	-		-	_	-			=	_	_	-	-			=	_	-		-	-	_	-	-		
DESCRIPTION OF STREET	_	0947BS 08	_	0	_			0.0	-	0	0	0	-	0.0	-	0	-		0	_	0 0	=	0	_		=	_	_			=	0	0	-	0.0	0 0		=	0	-		-	0	0	_	_	10	0 0
Terr Terr	à	_	- H	-	=			0.0	=	_	_	0	0	0.0	_	0	_		_	_		=	0	_		_	0	_		0 1	=	0	_	0	0		= 1	=	0	_		-	_	0	_	_		-
100	_	-	983636	99	_			0.0	-	_	0	0		0.0	-	0	_		_	_	0	=	0	_		-	0	_			=	0	0	-	0 0	0	-	=	0	_		-	_	0	_	_	10	
1205120 1235130 1305130 1335140	_	3 0.708333	-	_	_				-		0	0			-	0	_		-	_		=	0	_		-	0	_	-		=		0	-			-	=	0	_		_			_	-		
143041		0.795103	6.75	90	_				_	_		0			_		_		_	_		=		_		=	_	_			=			-			-	=		_		_	_		_			
	0.846.54	0.851852	69	89	Despera					_					_		_		_	_		_		_		_		_			=			-			-	=		_			_		_			
113141211	7		0.77.27		_												_			_		-		_				_			=			_			-	=		_					_			
	29766	0.882533	0.785714	0.727273	_					_	_					_			_	_			_	_			_	_			_		_				-	_	_				_		_	_	_	
	7,099,00	0.88565	-	14	Demon.																																									_		
TOTAL T	_		_	ogsts 0	n Dames	-	_		-	-	_	_	_		-	_	-	_	-	-		-	_	_		-	_	_			=	_	_	-				=	_	-		-	-	_	_	-	_	
1			200080	_	_			0.0	-	_	_	0	0	0.0	_	_	-		_	=		=	_	_		-	_	_		0 1	=	0	_	-			= 1	=	-	-		_	_	0	_	_		0 0
200	8	989	167857 B	8	0.00571 n				-	_	_	0	-		_	0			_	_		=	0		1 0	-	_			-	=		_	-	0 0		-	=	_			-	_			_	10	
No.			0.000017	0.038800	n.7dars n.				_	_	_	_	0		_	0	_		_	_		=	0		1 0	-	_			-	=		_	-	0 0			=	_	_		-	_			_	10	
OTTO DE				730167 0.	WESTS D.		2222		_	_	_	_	0		_	0	_		_	_		=	0		1 0	-	_			-	=		_	-				=	_	_		_	_			_	10	
1000	98376	85855	3601	75077 0	0 22227	_			_	_	_	_	-		-	_	_	_	_	_		=	_			-	_				=	_	_	-				=	_	_		-	_	_		_	-	
DOM:	_	_	80198	0.000	Some?	O STOR	8		_	_	_	_	0		_	0	_		_	_		-	0			-	_			-	=		_	-				=	_	_		_	_			_	-	
1000	_	_	0.781600 0.			-	-			_	_	_	0		_	0	_		_	_		=	0	_		-	_				=	-	_	_				=	_	_		_	_	-	_	_	-	
100	_			20000	Case o				,	_	_	_	-		-	_	_	_	_	_		=	_	_		-	_				=	_	_	-				_	0	_		-	_	_	_	_	10	
			0.78685			Very or			9	9	_	-	_		-	_	_	_	-	=		=	_	-		-	_				=	_	-	-	-			=	-	_		-	-	_	-	-	_	
000	0.88665									=	_	-	-		-	_	-	_	-	=		=	_	-		-	_				=	_	_	-	-			=	_	_		-	-	_	-	-	_	
no no	928888	0.788822	0.702703 0.79845	9,999	0.525	15				28310	_	_	_		-	_	_	_	=	=		=	_	_		-	_				=	_	_	-				=	_	_		-	_	_	_	-	_	
1		0.62675 0	0.51855	0.520trs 0	- 5	6	1			8181	2	99	-		-	_	_	_	_	_		=	_	_		-	_	_			=	_	_	_				=	_	_		_	_	_	_	_	10	
1	_	_	-	33338	047257 n	_		_	-	1988897	_	0.142857 0	-		-	_	_		_	_		_	_	-		_	٥	-		- 1	_	٥	_	-		. 0	21	_	-	_		_		٥	-	£	_	
STREET, ST	=	_	_	13528 02	131578n n.d		-	3 2		=			0	0.0	-	0	-		_	_		=	0	_		-	0	_		-	=		_	-	0 0		-	=	_	-		-	_		_	-		0.0
2000	-	_	_	33500	Dates.	-	-	-	_	m	olossa o	d	0	0.0	-	_	-		_	=		=	_	_		-	_		-	0 1	=	0	_	-	0		-	-	-	-		-	_	0	_	_	10	
7	_	olosas 02	_	1500l 03	-	0150802	_		_	192/9270	70	0	0	0.0	-	-	-		_	=		=	-	_		-	_	_	-		=	0	0	-	0	0		=	0	-		-	_	0	_	_	-	-
2000	1782	970	ors	_	OFF 0.14	210	1	100	á	8	_	0		0.0	_	_	_	-	_	_	0 0	=	_	_		=	_	_			=	0	0	-	0 0	0 0		=	0	_		-	0	0	_	_		-
ALL PARTY OF THE P		0	0	900	not	0.0			5	-	_		c		-		_	,	-	_		=		_		=	_	_			=	0	0	_			-	=	0	_		-	-		_	=	10	-
200	-			-	_				-	_	-	-	0		-	0	_		_	_		_	0	_		=	_	_	-		=		-	-			-	=	-	_		_	_		_	_		
1000		0	0	0	-				_	_	_	_	-		-	_	_		_	_	0 0	=	_	_		=	_				=	0	0		0.0	0 0		=	0	_		_	0	0	_	_	10	-
100	0			0	-			0.0	-	-	0	0	-	0.0	_	0	_		-	_		=	0	_		_	_	-	100	0	- 25	0 86	Dex 0	0 04	0		9	- 25	Dex 0	0 00	2		0 24	0 86	0 20	100	10	9.2
DELL'ANNE DELL'	045 Lbg	450 Dg	9435 Deg	135-18 Deg	18:22.5 Dec	20 5.77 Dec	27.91 C Par	of Care Day	diam'r.	36-40.5 Lbg	40.5-45 Deg	45-49.5 Deg	to c c t Day	A Se S Day	340000	SSS D1	92.67 5 Dec		6/5/21Ag	72765 Dec	Torsel D	977 1000	81-85.5 Deg	SSOnDe	Tools De	STORES.	212801kg	solins De	Instinct.		102112	1125.117 Deg	1174215 L	121 S.13t Dec	Tak Land D	d and a second	10101	13513501	1305141 L	144,1485 L	1485-178 Dec		153-15/5 Deg	157.51G2 L	Inclosed L	DESIG L	121 1766 D	1755181 Dec
ك	_		-11	_	_			. 0	_	- 2	-				_		_	_	_	-	-	÷	_	_			-1		_		-		_	-	_	_		•	_	=	_		-		_	=	Ť	_