# SciBooNE Charged-Current Coherent Pion Production Acceptance Study Technical Note

Jonathan Asaadi<sup>1</sup> and Zachary Williams<sup>1</sup>

<sup>1</sup>Department of Physics, The University of Texas at Arlington

June 2, 2017

# 1 Introduction

This document is intended to serve as a reference for the acceptance study performed for the SciBooNE charged current coherent pion production (CC-Coh  $\pi^{+/-}$ ) re-analysis, as well as provide documentation of the code used in this study (in the event anything needs to be revisited in the future). The code resides in the github repository labeled as and linked here: SciBooNE-MC. The corresponding ROOT files that were used in this acceptance study can be downloaded from here: SciBooNE-MC-ROOTFiles.

The paper is structured such that Section 2 outlines Monte Carlo samples used in this study, Section 3 describes the SciBooNE detector as it was simulated in this study, Section 4 describes the various event samples that were used to both validate and generate the acceptance studies for the CC-Coh  $\pi^{+/-}$  sample. Section 5 gives a high level summary of the results including the event-reduction tables as well as the CC-Coh  $\pi^{+/-}$  acceptance results.

The appendix is left to explain how the code is run and the details of the scripts within. The appendix also details the order in which the macros should be run in, and the important plots that each macro produces that play a role in making the plots shown in Section 5 (the Results section).

#### 1.1 Goal of the Re-Analysis

The goal of the re-analysis is to examine the acceptance modeling for the SciBooNE results in the presence of modern neutrino generators and updated models in order to hopefully shed light on why SciBooNE did not observe charged-current coherent pion production at low neutrino energy.

This study is intended to examine the effects of the acceptance modeling for a sample of coherent pion interactions inside the SciBooNE detector and compare what these would have been for various coherent pion production models. We utilize a simple, but robust, simulation of the SciBooNE detector and the NEUT neutrino generator to select and classify these neutrino events.

# 2 Samples

Five different samples were used in this study, three samples were generated in neutrino mode ( $\nu$ -mode) and two samples in antineutrino mode ( $\bar{\nu}$ -mode.)<sup>1</sup> Table 2 summarizes these samples. Details on these samples can be found in the Appendix.

<sup>&</sup>lt;sup>1</sup>All of these samples were generated by Callum Wilkinson (Thanks, Callum!)

#### **Summary of Samples**

Mode	NEUT version	Pion-Model	Number of simulated events
$\overline{\nu}$	5.3.6	Rein-Sehgal	1,000,000
$\overline{\nu}$	5.3.6	Berger-Sehgal	1,000,000
$\overline{\nu}$	5.0.1	Rein-Sehgal	100,000
$\bar{\bar{\nu}}$	5.3.6	Rein-Sehgal	1,000,000
$\bar{\bar{\nu}}$	5.3.6	Berger-Sehgal	1,000,000

Table 2: Summary of the samples used to build the acceptance model for this study.

#### 3 Detector Simulation

This section is intended to detail the detector simulation done in this acceptance model, and to describe the assumptions made in order to accomplish accurate classifications of simulated events as charged-current coherent pion production.

#### 3.1 The Detector

For the purposes of this acceptance study, the SciBooNE experiment is composed of two sub-detectors. The first (and the more upstream) of the sub-detectors, is the Scintillator Bar Tracker (SciBar) which was originally conceived and constructed to function as the near detector for the K2K experiment. The second (and more downstream) of the sub-detectors, is the Muon Range Detector (MRD), which is the detector designed and constructed specifically for SciBooNE for measuring the momentum of muons produced from charged-current neutrino interactions up to  $1.2 \ GeV/c$  by using the observed range of the trajectory of the muon. The coordinante system used throughout this study, and illustrated in Figure 3.1, puts the origin in the lower corner of the SciBar detector, has z along the beam direction, y opposite to gravity, and x to beam left.




Figure 3.1: Representation of the SciBooNE detector and the coordinate frame we use in this study

#### 3.1.1 The Scintillator Bar Tracker (SciBar)

The Scintillator Bar Tracker (SciBar) sub-detector is a scintillator detector which was used to identify neutrino interactions within SciBooNE. The dimensions of the SciBar detector used in this simulation are 0 < x < 3.0 m, 0 < y < 3.0 m, and 0 < z < 1.7 m. This simulation models the scintillator materials as having a constant energy deposition per unit length (dE/dx) for both

muons and pions of 2.04 MeV/cm based on previous SciBooNE analyses and on mean values for typical particle momentum listed in the particle data group (PDG).

#### 3.1.2 The Muon Range Detector (MRD)

The Muon Range Detector (MRD), depicted in Figure 3.1.2, is located 0.55 m downstream of SciBar in the z-direction, and is a composition of two sets of thirteen alternating slabs of steel-scintillator layers, where the scintillator layers alternate between being horizontally oriented or vertically oriented, in the xy-plane. The steel layers have a z-direction thickness of 5.08 cm and the scintillator layers have a z-direction thickness of 0.6 cm. Combining all the layers of the different alternating materials results in 26 scintillator layers that "sandwich" twenty five steel layers inbetween and gives a total z-direction dimension of being 1.37m. The xy-plane is modeled as a square again (as was the case with SciBar, too) with dimensions in the x-direction and the y-direction of 2.6 m. The energy deposition per unit length (dE/dx) of a muon penetrating the scintillator layers is assumed to be a constant 2.04 MeV/cm while the energy deposition for the muon in the steel layers is assumed to be a greater value of 11.43 MeV/cm. Both values are typical for muons at the energy range produced in SciBooNE and taken from the PDG.




Figure 3.1.2: Depiction of the Muon Range Detector (MRD) which consists of alternating layers of horizontal scintillator (shown in pink) steel slabs (shown in grey) and vertical scintillator (shown in purple)

#### 4 Event Selection

Two main samples are used in this study to generate the acceptance tables. The first is a charged current inclusive (CC-Inclusive) sample which requires a muon was created in the neutrino interaction and this muon intersects the MRD. This sample is described in Section 5.1 and is used to validate the building of the acceptance model by comparing it to previous SciBooNE analyses.

The second sample is the charged current coherent pion (CC-Coh  $\pi^{+/-}$ ) sample which requires a muon and charged pion are created in the neutrino interaction exclusively (e.g. no other final state particles in the event). This sample is described in Section 5.2.

Both of these samples are selected using NEUT MC-truth flags which ensure we are treating pure samples which are classified by the neutrino generator as belonging to the appropriate sample.

Whether or not the event identified by our selection makes it into the final sample used in the acceptance study depends on the behavior of the muon with respect to the MRD. A muon which enters the MRD from a neutrino interaction will either come to stop in the MRD, exit out the back of the MRD (assuming it's momentum is great enough), or exit out the side of the MRD. In the next sections we explain this classification further.

# 4.1 Muon Stops within the MRD ("Stopped")

The requirement to classify a neutrino interaction as a "stopped" event requires the muon from the interaction to have reached the MRD, penetrated at least three layers of steel (giving activity in three layers of scintillator), and to then deposit all of its remaining energy prior to reaching a boundary of the MRD. An illustration of a CC-Coh  $\pi^{+/-}$  event which would be classified as "stopped" is shown in Figure 4.1.




Figure 4.1: Depiction of an event that was classified as "Stopped."

These events allow for complete reconstruction of the muon's momentum based on the number of layers which the muon penetrated and the muons incident angle.

#### 4.2 Muon exits out the back of the MRD ("Out-the-back")

The classification of a neutrino interaction as "out-the-back" requires that the muon from the interaction to have reached the MRD and to have had sufficient energy to have exited out the back face of the MRD without stopping. An illustration of such an event is given in Figure 4.2.




Figure 4.2: Depiction of an event that was classified as "out-the-back".

The exact momentum of muons which pass completely through the MRD could not be made in reconstruction, so these events were classified as having the minimum energy required to penetrate all the steel and scintillator layers of the MRD.

### 4.3 Muon exits out the side of the MRD ("Out-the-side")

The classification of a neutrino interaction as "out-the-side" requires that the muon from the interaction reached the MRD, penetrated at least three layers of steel, and then to have exited out the side of the active volume of the MRD (excluding the very back face). Events which are classified as "out-the-side" are excluded from this study because no accurate reconstruction of the muons momentum can be made when the muon exits out the side of the MRD. An illustration of such an excluded event which exits out the side of the MRD is given in Figure 4.3.




Figure 4.3: Depiction of an event that was classified as "Out-Side."

# 5 Results

The results of this acceptance study can be broken down into two different classification schemes of events. Those that met the conditions to qualify as CC-Inclusive events, and those that met the

conditions of classification as Charged-Current Coherent Pion events. The former is used to validate the acceptance modeling and detector simulation to reasonably reproduce previously published CC-Inclusive studies from SciBooNE, while the latter is used for the reanalysis.

#### 5.1 Charged-Current Inclusive Events

Here we define the charged current inclusive sample (CC-Inclusive) which we use to validate our acceptance model against previously performed simulation studies.

#### 5.1.1 $\nu$ -mode Charged-Current Inclusive Events

Table 5.1.1 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the neutrino mode ( $\nu$ -mode) Monte Carlo.

u-mode CC-Inclusive Event Reduction				
Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal	
Total Sample	1,000,000	1,000,000	100,000	
CC-Inclusive Interaction	725,730	727,278	69,363	
$\mu$ ( $\mu$ + n-other particles in SciBar)				
Muon enters the MRD	263,698	262,608	$24,\!250$	
Muon enters the MRD and	231,089	230,054	21,001	
penetrates $\geq 3$ layers of steel				
"Stopped"-Events	177,406	175,799	16,062	
"Out-the-back"-Events	15,389	15,952	1,421	
"Out-the-side"-Events	38,294	38,303	3,518	
Good CC-Inclusive Events	192,795	$191,\!751$	17,483	

ν-mode CC-Inclusive Event Reduction

Table 5.1.1: Event reduction table for a sample of  $\nu$ -mode CC-Inclusive events simulated in the SciBooNE geometry.

Figure 5.1.1 shows the momentum and angular ( $\theta$ ) distribution for the sample of  $\nu$ -mode CC-Inclusive events passing all our requirements for all three models considered in this study (NEUT v5.3.6 Rein-Sehgal, NEUT v5.3.6 Berger-Sehgal, NEUT v5.0.1 Rein-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 5.1.1: Muon Momentum (left) and Muon Angle (right) for  $\nu$ -mode CC-Inclusive interactions for all three models included in this study. These samples kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive

Figure 5.1.1 represents the one-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events for this study using all three different models compared to results derived from Hiraide's thesis <sup>2</sup> using the full SciBooNE Monte Carlo simulation. A few reference points are illustrated using dashed lines to guide the readers eye. A few perecent difference is seen, but overall agreement between the two simulations hold.




Figure 5.1.1: One-dimension efficiency plots for the  $\nu$ -mode CC-Inclusive sample.

Figure 5.1.2 shows the two-dimensional efficiency for selecting  $\nu$ -mode CC-Inclusive events. The left hand side is a reference plot provided by Morgan and the right hand side is for the Rein-Sehgal MC used in this study.




Figure 5.1.2: Two-dimensional efficiency plots for the  $\nu$ -mode Rein-Sehgal CC-Inclusive sample.

<sup>&</sup>lt;sup>2</sup>Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide dt.pdf




Figure 5.1.2: Two-dimensional efficiency plots for the  $\nu$ -mode Berger-Sehgal CC-Inclusive sample.




Figure 5.1.1: Two-dimensional efficiency plots for the  $\nu$ -mode Old Rein-Sehgal CC-Inclusive sample.

#### 5.1.2 $\bar{\nu}$ -mode Charged-Current Inclusive Events

Similar to before, Table 5.1.2 goes through the event selection criteria for selecting a sample of CC-Inclusive events from the antineutrino mode ( $\bar{\nu}$ -mode) Monte Carlo.

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal			
Total Sample	1,000,000	1,000,000			
CC-Inclusive Interaction	699,239	704,327			
$\mu$ ( $\mu$ + n-other particles in SciBar)					
Muon enters the MRD	380,362	380,869			
Muon enters the MRD and	336,373	337,979			
penetrates $\geq 3$ layers of steel					
"Stopped"-Events	288,289	288,206			
"Out-the-back"-Events	7,608	7,857			
"Out-the-side"-Events	40,476	41,916			
Good CC-Inclusive Events	295,897	296,063			

 $\bar{\nu}$ -mode CC-Inclusive Event Reduction

Table 5.1.2: Event reduction table for a sample of  $\bar{\nu}$ -mode CC-Inclusive evnets simulated in the SciBooNE geometry.

Figure 5.1.1 shows the momentum and angular distribution for the sample of  $\bar{\nu}$ -mode CC-Inclusive events passing all our requirements for both models considered in this study (NEUT v5.3.6 Rein-Sehgal, and NEUT v5.3.6 Berger-Sehgal). The distributions have been normalized to the same area and show no strong differences between them.




Figure 5.1.2: Muon Momentum (left) and Muon Angle (right) for  $\bar{\nu}$ -mode CC-Inclusive interactions for all three models included in this study. These samples kinematics are, unsurprisingly, very similar for the sample of CC-Inclusive

Figure 5.1.2 represents the one-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events for this study. No similar reference sample exists to be compared directly against, however we note that the shape and magnitude of the acceptance is nearly unchanged between  $\bar{\nu}$  and  $\nu$ -mode samples (as expected).




Figure 5.1.2: One-dimension efficiency plots for the  $\bar{\nu}$ -mode CC-Inclusive sample. Muon's Momentums is on the right and the Muon's Angles is on the left.

Figure 5.1.2 shows the two-dimensional efficiency for selecting  $\bar{\nu}$ -mode CC-Inclusive events for this study compared to results derived from Morgan's reference sample (need more words here about this....see email)




Figure 5.1.2: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode Rein-Sehgal CC-Inclusive sample.




Figure 5.1.2: Two-dimensional efficiency plot for the  $\bar{\nu}$ -mode Berger-Sehgal CC-Inclusive sample.

#### 5.2 Charged-Current Coherent Pion Production Events

Here we define the Charged-Current Coherent Pion Production sample (CC-Coh  $\pi^{+/-}$ ) which we use to validate our acceptance model against previous simulation studies which were done.

#### 5.2.1 $\nu$ -mode Charged-Current Coherent Pion Events

Table 5.2.1 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi^{+/-}$  events from the neutrino mode ( $\nu$ -mode) Monte Carlo.

The first quantity that is calculated for the different events is the momentum of both the muon and the pion, which are both found from the equations:

 $\nu$ -mode CC-Coherent Pion Event Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal	NEUT v5.0.1 Rein-Sehgal
Total Sample	1,000,000	1,000,000	100,000
CC-Coherent Pion Interaction	12,186	2,576	1,320
$(\mu + \pi + \varnothing \text{ in SciBar})$			
Both muon and pion are	8,535	1,845	884
forward going			
Muon enters the MRD and	7,407	1,592	767
penetrates $\geq 3$ layers of steel			
"Stopped"-Events	6,448	1,350	669
"Out-the-back"-Events	530	150	56
"Out-the-side"-Events	429	92	42
Good Coherent Pion Events	6,978	1,500	725

Table 1: Event reduction table for a sample of  $\nu$ -mode Charged Current Coherent Pion events simulated in the SciBooNE geometry.

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{1}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{2}$$

where  $|\vec{p}_{\mu}|$  represents the magnitude of the momentum for the corresponding particle, and  $P_{\mu_x}$  represents the component of the four momentum for the corresponding particle. The momentum is reported in units of MeV/c.

The next quantity calculated is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1} \left( \frac{\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}}{P_{\mu_z}} \right) \tag{3}$$

$$\theta_{\pi} = tan^{-1} \left( \frac{\sqrt{P_{\pi_x}^2 + P_{\pi_y}^2}}{P_{\pi_z}} \right) \tag{4}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of charged-current coherent pion production, the angle should never be larger than 90°.




Figure 1: Muon Momentum for all of the muons of the events that made it to the MRD and penetrated at least three layers (left) and Muon Angle for the muons of the events that made it to the MRD and penetrated at least three layers (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study. The "Total" classification means that all CC-Coh  $\pi^{+/-}$  events are included in these histograms.

Here will be the plots for CC-Coh Pion with the good momentum efficiencies and the angle efficiencies!




Figure 2: Muon Momentum of both the "stopped" and "not-stopped" samples (left) and Muon Angle of both the "stopped" and "not-stopped" samples (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study. The "Good" classification means that only the stopped and not-stopped CC-Coh  $\pi^{+/-}$  events are included for these histograms.

The last two quantities that are calculated are the two different types of four-momentum transfers specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{5}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$

$$(6)$$

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{7}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
(8)

|t| is reported in units of  $(MeV/c)^2$ .

 $\nu$ -Mode |t| and  $Q^2$  plots are below:




Figure 3: The |t| Momentum Transfer for the "stopped" and "not-stopped" events (left) and  $Q^2$  Momentum Transfer for the "stopped" and "not-stopped" events (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for the three models included in this study.




Figure 5.2.1: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.




Figure 5.2.1: Two-dimensional efficiency plot for the new NEUT  $\nu$ -mode Berger-Sehgal CC-Coherent sample.




Figure 5.2.1: Two-dimensional efficiency plot for the old NEUT  $\nu$ -mode Rein-Sehgal CC-Coherent sample.

# 5.2.2 $\bar{\nu}$ -mode Charged-Current Coherent Pion Events

Similar to before, Table 5.2.2 goes through the event selection criteria for selecting a sample of CC-Coh  $\pi^{+/-}$  events from the anti-neutrino mode ( $\bar{\nu}$ -mode) Monte Carlo.

Below are the plots for CC-Coh  $\pi^{+/-}$  Events for  $\bar{\nu}$ -mode. The layout of the rest will be very similar to  $\nu$ -mode, and the equations used previously are the same equations used for the plots below.

#### $\bar{\nu}$ -mode CC-Coherent Pion Event Reduction

Events Selection	NEUT v5.3.6 Rein-Sehgal	NEUT v5.3.6 Berger-Sehgal
Total Sample	1,000,000	1,000,000
CC-Coherent Pion Interaction	36,669	7,790
$(\mu + \pi + \varnothing \text{ in SciBar})$		
Both muon and pion are	24,675	5,477
forward going		
Muon enters the MRD and	20,445	4,517
penetrates $\geq 3$ layers of steel		
"Stopped"-Events	18,935	4,203
"Out-the-back"-Events	372	82
"Out-the-side"-Events	1,138	232
Good Coherent Pion Events	19,307	4,285

Table 2: Event reduction table for a sample of  $\bar{\nu}$ -mode Charged Current Coherent Pion events simulated in the SciBooNE geometry.




Figure 4: Muon Momentum (left) and Muon Angle (right) for  $\nu$ -mode CC-Coh  $\pi^{+/-}$  interactions for all three models included in this study.

The structure of the plots in Figure: 5.2.2 very closely resembles the plots for the  $\nu$ -mode above, and the rest of the plots in this section have that same characteristic.




Figure 5: Muon Momentum (left) and Muon Angle (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi^{+/-}$  interactions for both the "stopped" and "not-stopped" samples of events.

 $\bar{\nu}$ -mode |t| and  $Q^2$  plots are below. They also have the same overall shape as the plots for  $\nu$ -mode above.




Figure 6: The |t| Momentum Transfer (left) and  $Q^2$  Momentum Transfer (right) for  $\bar{\nu}$ -mode CC-Coh  $\pi^{+/-}$  interactions for both of the models included in this study which are the "stopped" and "not-stopped" events.




Figure 5.2.2: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Rein-Sehgal CC-Coherent sample.




Figure 5.2.2: Two-dimensional efficiency plot for the new NEUT  $\bar{\nu}$ -mode Berger-Sehgal CC-Coherent sample.

# A Appendix: Sample Details

Appendix on samples

### A.1 $\nu$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$  interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack.root

found at the following link (put link to sample here).

#### A.2 $\nu$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\nu$  interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample correspond to the file labeled

SciBooNE\_numu\_coh\_RooTrack\_NEW.root

found at the following link (put link to sample here).

#### A.3 $\nu$ -Mode Rein-Sehgal NEUTv5.0.1

A sample of 100,000  $\nu$  interactions were simulated using the NEUT generator (v5.0.1, believed to be the version used by the SciBooNE collaboration in the original publication) and the corresponding older Rein-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root

found at the following link (put link to sample here).

# A.4 $\bar{\nu}$ -Mode Rein-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$  interactions were simulated using the NEUT generator (v5.3.6) and the Rein-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack.root

found at the following link (put link to sample here).

# A.5 $\bar{\nu}$ -Mode Berger-Sehgal NEUTv5.3.6

A sample of 1,000,000  $\bar{\nu}$  interactions were simulated using the NEUT generator (v5.3.6) and the Berger-Sehgal model for coherent pion production. This sample corresponds to the file labeled

SciBooNE\_numubar\_coh\_RooTrack\_NEW.root

found at the following link (put link to sample here).

#### A.6 Vertex Distributions

The events were all given a random initial point that was generated with the goal that the vertex distributions of this simulation would closely match the vertex distributions that Hiraide <sup>3</sup> showed in his thesis. This was done by... etc.

```
TRandom3 *randX = new TRandom3();
TRandom3 *randY = new TRandom3();
TRandom3 *flat = new TRandom3();
randX->SetSeed(jentry/2);
randY->SetSeed(jentry*jentry);
flat->SetSeed(jentry*jentry*jentry);
double Xpos = randX->Gaus(1.5,1.3);
while (Xpos<0 || Xpos>3.0) { Xpos = randX->Gaus(1.5,1.3);}
double Ypos = randY->Gaus(1.5,1.05);
while (Ypos<0 || Ypos>3.0) { Ypos = randY->Gaus(1.5,1.05);}
double Zpos = flat->Uniform(0,1.7);
```




Figure 7: Vertex distributions of the events in the new Rein-Sehgal sample in  $\nu$ -mode.

#### A.7 NewNMReinSehgal.C

This file is the macro that corresponds to the "NewNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms). When this macro is run (which can take a while), it also plots a few different histograms. The histograms that are plotted are the ones shown in the figures below with descriptions included with the corresponding figures. The order that the histograms appear in this paper is the same order they will be shown when this macro is run in root.

<sup>&</sup>lt;sup>3</sup>Hiraide's thesis can be found here: http://www-he.scphys.kyoto-u.ac.jp/theses/doctor/hiraide dt.pdf




Figure 8: New  $\nu$ -Mode Rein-Sehgal X-Y vertex distributions for muons that made it to the MRD and penetrated at least to the third layer of steel.




Figure 9: New  $\nu$ -Mode Rein-Sehgal Z vertex distributions for the interactions.




Figure 10: New  $\nu$ -Mode Rein-Sehgal Y vertex distributions for the interactions.




Figure 11: New  $\nu$ -Mode Rein-Sehgal X vertex distributions for the interactions.




Figure 12: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 13: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".




Figure 14: This histogram shows the amount of muons that embedded (or "Stopped") in a corresponding layer of steel in our simulation.




Figure 15: This is a 2D histogram for the momentum and angle of the pion in the total CC Coh Pion events.

#### Total CC Coh Events Muon Information Muon Momentum (MeV) Muon Angle (Degrees)

Figure 16: This is a 2D histogram for the momentum and angle of the muon in the total CC Coh Pion events.

The NewNMReinSehgal.C macro also calculates many different quantities for the generated simulation of the events and saves the information in histograms that are later called upon through the plotting macros (which are after all of the analysis macros). The first quantity that is calculated for the different vertexes is the momentum of both the muon and the pion, which are both calculated using the equations:

$$|\vec{p}_{\mu}| = \sqrt{P_{\mu_x}^2 + P_{\mu_y}^2 + P_{\mu_z}^2} \tag{9}$$

$$|\vec{p}_{\pi}| = \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2 + P_{\pi_z}^2} \tag{10}$$

The momentum is reported in units of MeV/c.

The next quantity that is calculated in the macro is the angle from the beam-direction for both the muon and the pion, which are labeled as either  $\theta_{\mu}$ , or  $\theta_{\pi}$ , respectively. The angle from the beam-direction is the same as the angle from the z-direction, and this angle is known as the azimuthal angle. The calculation of the azimuthal angle is slightly more involved than the simple calculation used for finding the magnitude of the momentum of the two particles, and is calculated using the equations:

$$\theta_{\mu} = tan^{-1}(\sqrt{P_{\mu_x}^2 + P_{\mu_y}^2}/P_{\mu_z}) \tag{11}$$

$$\theta_{\pi} = tan^{-1} \left( \sqrt{P_{\pi_x}^2 + P_{\pi_y}^2} / P_{\pi_z} \right) \tag{12}$$

The angles are reported in units of °, and should run from 0° to 180°. In the case of Charged-Current Coherent Pion Production, the angle should never be larger than 90°.

The last two quantities that this analysis macro calculates are the two different types of four-momentum transfers specific to this interaction, which are  $Q^2$  and |t|. The  $Q^2$  corresponds to the four-momentum transfer from the neutrino and muon to the nucleus and pion, and is calculated using the equation:

$$Q^2 = |(P_{\nu_{\mu}} - P_{\mu})^2| \tag{13}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute  $Q^2$ :

$$Q^{2} = |(P_{\nu_{\mu,x}} - P_{\mu_{x}})^{2} + (P_{\nu_{\mu,y}} - P_{\mu_{y}})^{2} + (P_{\nu_{\mu,z}} - P_{\mu_{z}})^{2} + (P_{\nu_{\mu,E}} - P_{\mu_{E}})^{2}|$$
(14)

 $Q^2$  is reported in units of  $(MeV/c)^2$ .

The |t| corresponds to the four-momentum transfer from the neutrino, muon, and pion to the nucleus, and is calculated using the equation:

$$|t| = |(Q - P_{\pi})^{2}| = |(P_{\nu_{\mu}} - P_{\mu} - P_{\pi})^{2}| \tag{15}$$

This equation is the four-momentum notational form. The code follows the equation below in order to compute |t|:

$$|t| = |(P_{\nu_{\mu,x}} - P_{\mu_x} - P_{\pi_x})^2 + (P_{\nu_{\mu,y}} - P_{\mu_y} - P_{\pi_y})^2 + (P_{\nu_{\mu,z}} - P_{\mu_z} - P_{\pi_z})^2 + (P_{\nu_{\mu,E}} - P_{\mu_E} - P_{\pi_E})^2|$$
 (16)

|t| is reported in units of  $(MeV/c)^2$ .

# A.8 NewNMBergerSehgal.C

This file is the macro that corresponds to the "NewNMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_RooTrack\_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 17: New  $\nu$ -Mode Berger-Sehgal X-Y vertex distributions for muons that made it to the MRD and penetrated at least to the third layer of steel.




Figure 18: New  $\nu$ -Mode Berger-Sehgal Z vertex distributions for the interactions.




Figure 19: New  $\nu$ -Mode Berger-Sehgal Y vertex distributions for the interactions.




Figure 20: New  $\nu$ -Mode Berger-Sehgal X vertex distributions for the interactions.




Figure 21: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 22: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".!




Figure 23: This histogram shows the amount of muons that embedded (or "Stopped") in a corresponding layer of steel in our simulation.




Figure 24: This is a 2D histogram for the momentum and angle of the pion in the total CC Coh Pion events.




Figure 25: This is a 2D histogram for the momentum and angle of the muon in the total CC Coh Pion events.

# A.9 OldNMReinSehgal.C

This file is the macro that corresponds to the "OldNMReinSehgal.h" file, which connects with this file: "SciBooNE\_numu\_coh\_OLDNEUT\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoOBS.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 26: Old  $\nu$ -Mode Rein-Sehgal X-Y vertex distributions for muons that made it to the MRD and penetrated at least to the third layer of steel.




Figure 27: Old  $\nu$ -Mode Rein-Sehgal Z vertex distributions for the interactions.




Figure 28: Old  $\nu$ -Mode Rein-Sehgal Y vertex distributions for the interactions.




Figure 29: Old  $\nu$ -Mode Rein-Sehgal X vertex distributions for the interactions.




Figure 30: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 31: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".

# Steel Layer the Muon Stops Inside

Figure 32: This histogram shows the amount of muons that embedded (or "Stopped") in a corresponding layer of steel in our simulation.




Figure 33: This is a 2D histogram for the momentum and angle of the pion in the total CC Coh Pion events.




Figure 34: This is a 2D histogram for the momentum and angle of the muon in the total CC Coh Pion events.

# A.10 NewANMReinSehgal.C

This file is the macro that corresponds to the "NewANMReinSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoRSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 35: New  $\bar{\nu}$ -Mode Rein-Sehgal X-Y vertex distributions for muons that made it to the MRD and penetrated at least to the third layer of steel.




Figure 36: New  $\bar{\nu}$ -Mode Rein-Sehgal Z vertex distributions for the interactions.




Figure 37: New  $\bar{\nu}$ -Mode Rein-Sehgal Y vertex distributions for the interactions.




Figure 38: New  $\bar{\nu}$ -Mode Rein-Sehgal X vertex distributions for the interactions.




Figure 39: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 40: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".




Figure 41: This histogram shows the amount of muons that embedded (or "Stopped") in a corresponding layer of steel in our simulation.




Figure 42: This is a 2D histogram for the momentum and angle of the pion in the total CC Coh Pion events.




Figure 43: This is a 2D histogram for the momentum and angle of the muon in the total CC Coh Pion events.

### A.11 NewANMBergerSehgal.C

This file is the macro that corresponds to the "NewANMBergerSehgal.h" file, which connects with this file: "SciBooNE\_numubar\_coh\_RooTrack\_NEW.root". This file performs the main analysis for this generated sample, and then organizes the information into many different histograms. The histograms are then written to a file titled "totalmuoninfoBSBar.root" inside the "ROOTFILES" directory. The "ROOTFILES" directory is included in the SciBooNE-MC repository (it is absolutely pertinent that this directory be located where the macro files are located due to how the calls of the combined data macros reference the now saved histograms).




Figure 44: New  $\bar{\nu}$ -Mode Berger-Sehgal X-Y vertex distributions for muons that made it to the MRD and penetrated at least to the third layer of steel.




Figure 45: New  $\bar{\nu}$ -Mode Berger-Sehgal Z vertex distributions for the interactions.




Figure 46: New  $\bar{\nu}$ -Mode Berger-Sehgal Y vertex distributions for the interactions.




Figure 47: New  $\bar{\nu}$ -Mode Berger-Sehgal X vertex distributions for the interactions.




Figure 48: This is a 2D histogram for the momentum and angle of the pion in the CC Coh Pion events that met the qualification of being "good".




Figure 49: This is a 2D histogram for the momentum and angle of the muon in the CC Coh Pion events that met the qualification of being "good".




Figure 50: This histogram shows the amount of muons that embedded (or "Stopped") in a corresponding layer of steel in our simulation.




Figure 51: This is a 2D histogram for the momentum and angle of the pion in the total CC Coh Pion events.




Figure 52: This is a 2D histogram for the momentum and angle of the muon in the total CC Coh Pion events.

### A.12 NMCombinedPlots.C

This is the file that performs the main plotting operations for the neutrino mode samples using the muon's information. All of the muon efficiency plots for neutrino mode are made with this file.




Figure 53




Figure 54




Figure 55




Figure 56




Figure 57




Figure 58




Figure 59




Figure 60




Figure 61




Figure 62




Figure 63




Figure 64




Figure 65




Figure 66




Figure 67




Figure 68




Figure 69




Figure 70




Figure 71




Figure 72




Figure 73




Figure 74




Figure 75




Figure 76




Figure 77

# ${\bf A.13} \quad {\bf NMPionPlotting.C}$

This is the file that performs the main plotting operations for the neutrino mode samples using the pion's information. All of the pion efficiency plots for neutrino mode are made with this file.




Figure 78




Figure 79




Figure 80




Figure 81




Figure 82




Figure 83




Figure 84




Figure 85




Figure 86




Figure 87




Figure 88




Figure 89

## A.14 NMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for neutrino mode.




Figure 90




Figure 91




Figure 92




Figure 93




Figure 94




Figure 95

### A.15 ANMCombinedPlots.C

This is the file that performs the main plotting operations for the anti-neutrino mode samples using the muon's information. All of the muon efficiency plots for anti-neutrino mode are made with this file.




Figure 96




Figure 97




Figure 98




Figure 99




Figure 100




Figure 101




Figure 102




Figure 103




Figure 104




Figure 105




Figure 106




Figure 107




Figure 108




Figure 109




Figure 110




Figure 111




Figure 112




Figure 113

### A.16 ANMPionPlotting.C

This is the file that performs the main plotting operations for the anti-neutrino mode samples using the pion's information. All of the pion efficiency plots for anti-neutrino mode are made with this file.




Figure 114




Figure 115




Figure 116




Figure 117




Figure 118




Figure 119




Figure 120




Figure 121




Figure 122




Figure 123




Figure 124




Figure 125

#### A.17 ANMFourSquaredPlotting.C

All of the four-momentum transfer (both |t| and  $Q^2$ ) combined plots are made with this file for anti-neutrino mode.




Figure 126




Figure 127




Figure 128




Figure 129




Figure 130




Figure 131

### B Steps for Running the Code

The instructions on how to run the code and the order the files need to run in so that there are no resulting error messages, or other issues while running the code, are detailed in this section.

- Step 1: This is the first step. (Run the NewNM macros and the NewANM macros and the OldNM macro.)
- Step 2: This is the second step. (Run the combined plotting macros.)
- Step 3: This is the third step. (Run the Pion Plotting macros.)
- Step 4: Etc. (Run the FourSquaredMomentum macros.)

# C Efficiency Tables

These are the corresponding tables to the 2D Efficiency plots for CC-Inclusive events.

Table 3: Table for 2D Histogram for New NM-Rein-Sehgal

al	p .		- L	_		10	1 55	_	F1	-	_			_	_	-	_	_	-	_	_	-		_	_	_	_	_	-	-	_	-	_	-	-	-	-	-	-	_	_	-	-	-	-	_
H	7 082352	-	_	_	_	-	-	-	28892	0.33333		=		=			=	=		-		- 1	=			=		0 0	- 1	=	=				=	-		0 0	=				=	-		
Н	7899280	0.0	-	-	-	-	0.25074	-	-	_		2		_			=	=		-		-	=			_		0 0	-	=	=				_	-		0 0	=				_	=		
	1111980	- 0	000000	0.000000	Medic II	0.462963	0.38388	3	=	-	2 1	=		_			=	_		_		= 1	=			_			= 1	=	=				_	_			=				_			
1810-182	0.848485	0.000	10000	1,455	197,047	12000	0.405941	0.000	2272	0.571-286		3		_			=			-			=			_		0 0		=	=				_	-			=				_	-		
1730-1800	198	0.777	0.00000	1539531	10.550.00	H-42857	733311.0	70000000	2,63	0.25		=	0.25	_			=	=		_		-	=			_			-	=	=				_	_			=				_			
1701120	0.521359	10.04(1)	1000000	150-501	1 332(b)	1,447,41	0.355001	0.000	0.272727	12		1.455th		_						_			=			_				=	=				_	_			-				_			
1631-170	0.921569	0.490.400 0.700.00	0.00000	07/9/07	A-09-0-11	530065	0.34058	070.70	0.32857	0.238095		0.4873	В	_			=	Ξ.	В	_			=	В	8	_				=	=		В	В	_	_						В	_			
1600 1650	80	0.774004	0.000000	0.010300	15011633	0.63674	0.0.463.4	0.000.000	9239	0.127273		0.4873		_				=					=	В		_				=	=		В		_								_	-		
1530-1600	0.9355B	150,000	0.000000	0.004030	11 212 11 22	0.513616	0.3605.77	o companie	0.293578	0.191489		1936360	9.0	_									=			_				=	_				_								_			
13061350	0.783333	0.745424	0.00000	0.00000	1.0000	0.511166	0.48,675	0.00.00	8.29.4118	0.8656		9.16		_					-			-	=			_			-	=	=			-	_								_			
1450-150	200	0.744.00	2000000	and and a	192025	0.4978.35	2199810	2000	19325191	125		136613	0.0759231	-					_											_	_			_	_								_			_
1:00-1:50	9.53386	0.0000	o contrary	1570000	926979	n Shinni	0.48953	200000	0.081702	0.351194		1.163014	0.11.75.47	_									_							_	_				_								_			
330.140	82546	2007	20000	991-100	160000	58828	383868	magar	9307116	1182-422		97,179,79	9.1917	_			_	_	_			_	_	_	_	_			_	_	_	_	_	_	_				_	_	_	_	_	_		-
13001350	0.81321	DOMESTIC OF	100000	0.694-702.0	0.269138	589955	0.455487		=	0.282509	_	1.20(02.0	1919-980	12					-				_							_	_	_		-							_	-			-	_
230-1388	362198	- Cooper	100000	600000	100000	1,498274	1 474016	0.00.00	3711783	13333	-	-	9292111	11.2857	000000		_	_	_					_	_		_			_	_	_	_	_	_		_	_		_	_	_	_	_		-
200 1250	0.897592 TOTAL	1000000	20000		884459	1.55 FM 62	58259	- PERSON	879848	0.293333	-	122(238	0.182927	0.08	-			_	_	_			_	_	_	_			-	_	_	_	_	_	_	_				-	_	_	_	-		_
130-120	25272	2424	200000	10000	927379	533719	445302	4000	-	-	-	191739	=	07070	-			_	_					_						_		_	_	_						_	_	_				
100-1150	28.85	Suite Ca	3 6	=	188208	354828 B	481938	00000	2222	352837	2 1	=	216867	145161	1021661			Ξ.					_	В		_				=	=	_	В		_		0.0						_		_	-
	822174		_	-	Ē	56538		-	10336	335748	-	=	168675	1782%	-				-				=							=	=	_		-	_						_	-	_		-	-
9	884522	= 0	0.00	=	=	57036	~	0	=	_		=	14572	0 61101.0	9		=	136364			0.0		=					0.0		=	=	В					0		=	0	В					-
Н	3815649		-	-	thomps a	Ξ	-	Ì	=	362308 0.	-	=	221135 0.	12-514 n.	-	3 6	=						=	0	0			0.0		=	=	8	0				0.0	0.0	-	0	В				-	-
1881	87564 04			-	_	570674	_	_	_	371585 02	_	=	23866	140187 0.		-	-	0.548902			0 0		=	8	0			0.0		=	=	8	8				0.0	0.0	-	0	В				-	-
8 8 8	798611			=	2	18894	20,460	-	455387 D.	378892	_	=	239462	177400 0.	-	-	Ξ.	9201	0.02		0 0	-	=	8	8			0.0	-	=	=	В	8		=		0 0	0 0	-	0	В		=		-	
0.850 850	2007.48			=	=	601285 n.		ė	46.472	404237	1	=	232889 02	202	13007	-	=	=	0131579 04		0 0	= 1	=			=		0.0	= 1	=	=	В			=		0 0	0.0	-	0	В		=			-
0.810	100	2000	0 0	20101	=	6171m		-	_	416728 0.	2 1	=	2383-5	163939			2	_	0121212 0.		0 0		=	0						=	=		0						-	0						-
12	950	9 5	-	_	=	161255	_		_		1	=	2755	1188606			=		0234114 0 1	00540541			=	В	В	_		0.0		=	=		В	8	_		0.0	0.0	-			В	_			
98	8645		- 0	=	-	628604 0.61	00130	200	2	2090	-	=	269592 0.27	214009 0.18			=	_	1020135 0.00	0	5 0		=	0	0	=		0 0		=	=		0	0	=	=	0 0	0 0	=			0	=	-	-	_
650	842975	5 0	- 0	=	481216 U.D.	3967 1953	52336 055	220	2	0.40		=	200561 026	138172 021		3 0	_	=	9173-0 995	-	0 0		=	0	0	=		0.0		=	=		0	0	=	-	0 0	0 0	=	0		0	=	-	-	
Ø	828829 0.84	<u> </u>	0 0	=	=	53.0 888918		ė	18882 B20	399772 0.4	2 1	=	248649 0.26	÷	10160 010	3 0	=	=	0121873 0.01	-	0 0		=			=		0 0		=	=				=	-		0 0	=				=	=		_
27	816092 0.82	0 0	0 0	oute alle	=	200221 0.61		-	=	5527 0.39	-	=	220634 0.24	-	-	3 6	Ξ.	-	m41686 0.01	-	0.0		=			=		0 0		=	=				=	-		0 0	=				=	=		_
17	7160-49 0.81	0 0	0 0	=	=	513469 0.50		-		335716 0.35	2 1	=	200127 0.22	35.14.22	-		-	ó	00228571 0.00	-	0.0		=			=		0 0		=	=				=	-		0 0	=				=	-		_
4		= 0	0.0	=	=			0	_	=		=		_		2 6	=	0.00045756 0.018	0.002	=	0 0	= 1	=	в		=		0.0	= 1	=	=		в		=	-	0 0	0.0	=				=	-		_
	9223840		10000	10224	U-498232	223 0.44242424	187 0381503	100	465 0.322889	174 0.273659		=	202 B1222	=	-	_	Ξ.	9000		-	0 0	= 1	=			=		0 0	= 1	=	=	0			=	-		0 0	=		0		=	=	0	
320-40	27		1000	127	3	NS 0.3562.	17 03280	Table In	.53 02604	75 02289		=	1688 0.12990	5227 0.102639	-	-	U-DATE.	=		_	0 0	= 1	=			=			= 1	=	=				=	_			=				=	-		
300.00	1 1 1 1 1 1	5 0	10000	10000	3	128478	12457	0	68 02155	14 0.1678.25	,	_	908 0.0851638	7255327			=			-		-	=	В		=				=	=		В		=	-		0 0	=				=	-		
-	0.20087	-	1000	=	0.2203	4 n.2ms.	0.146300	=	=	20 0.1043-0	2 1	=	=	0.0115123		0 0	=	=		_	0 0	-	=	В		=			-	=	=		В		=	_		0 0	=				=	-		
200 230	025939	-	-	_	=	5 0.120514	_	-		0.0335749		140,000,000	0.00241935	=			=			_		= 1	=			=			= 1	=	=				=	_			=				=	-		
F	0.09373	100000	1000000	0.4050203	0.46596825	0.78-0.615	0.00555	0.00+0.00	0.400618357	_		=		_			=			-			=			_				=	=				_	-							_			
E1-001	-			=	=	_		-		=		=		=			=			_		= 1	=			=			= 1	=	=				=	-			=				=	-		
TeV/c 50-100	_	0.0	0 0	=	=	_				=		=		=			=			-			=			=		0 0		=	=				=	-		0 0	=				=	-		-
WANTER BRIMEVO	-	= 0		=	=	_			=	=		=		=			=	=	-	-		= 1 	=	-	-	=		1	100	n 2500	neg I	0 org. 0	9 and	Dog B	Down II	U AND	-		0 %00	D No.	0 pg 0	Dog B	Down II	0 400	0 200	
انتم	0-45 Deg	San San		100348.00	18-22 a Day	27 Dr.	22.0 5 Dec	1	36 Da	36-41.5 Dev		Allo-to Deg	45-85 Drg	495.54 Dec	4 50 C Day		103 205	63-67.5 Dry	375-72 Day	TARREDO.	10000	101	N-No Deg	90 De	98-945 Day	445.49 Dov	00 102 C Day	See and a see	1 917	18-1125 Deg	12 3-417 D/g	17-1215 Deg	121 5-126 Day	126-1305 Day	1315-135 Dev	55,1395 Day	130 C 141 Day		1 1001	148 5-133 Deg	153-1575 Drg	157.5 162 Dry	162-1665 Dev	165.47 Dev	7 1755 Day	10000

Table 4: Table for 2D Histogram for New NM-Berger-Sehgal

_	_																											_			_							
1550-300	0.83333	0.794521	0.083030	0.622711	D.55747	0.5555	0.306835	0.137931	0.181818	_																			_			_	_	_	_		_	
e						_		0.258065		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		_	
1850-1900	27.872	761905	0.00338	299997	0.573342	66290		0.25																				_			_							
1800-1850 1		_	0.081967	Ī	÷	÷			0,333333 0	- 2	_	_	_	_	_	_	-	0	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	0 1	_	0.0
1730 1800 18	F	770053 0.7	0.0	_	_		÷			0.165657 0.5			_	_	_	_	_	_		_			_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	0.0
170.170 173	P	_	_	_	Ť	Ť	Ť	÷	÷	_	-		-	-	-	-												_	-	-	_	-	-			-		0 0
	F	_	23 0.70381	~	-	127 0.435681	_	÷	÷	22	-	0	0	0	0	0	0	0	0	0	-	0	0	0		0	0	_		0	_					-		0 0
53 1630 1700	F	Ť	0.747-23	Ť	_	_	÷	_	_	_	_	_	-	_	_	_	_											_	-		_	-	-	-		-	-	
n 16m-1630	Н	_	_	_	_	_	_	_	181818	_	_		-	-	-	-					-							_	-		_					-		
0 12201600	F	Ť	0.717833	0.66548	0.587336	22	_	_	7:08ID	0.0526316	_																	_			_					-		
1500-1530	0.873016	0.793631	0,7182-0	0.668863	0.590654	0.484043	0.433735	0.277372	0.2321-63	0.2	0.0714280																	_			_							
1.50.150	765528.0	0.790036	0.75089	0.64678	0.3835	0.335335	0.403145	0.39333	0.304819	0.157855	0.25																	_			_					-		
14001420	98.0	90,25.20	0.753835	0.659251	611602.0	0.49993	162131	0.381395	0.348624	0.22222	0.136364		0.0															_			_						_	
1350 1400	г	_	0.728296	12180.	612863	209404	0.455169	0.33354	_	_	299																											
1300 1350 1	F	813055	-		Ť	÷	÷	Ť	_	_	0.0832323	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		_	
1230-1300 13	Р.	-	(730769 0.	0	_	_	_	_	0.38785 0.	_		0 150505	11111				0	0					0	0	0	0	0	_	_	0	_					-		0 0
1200-1250 125	П	1784704 0.7	731677 0.7		60516 0.5	535057 0.5	4736 0.4	(406331 0.4	_	_		0.0752231 0.0	0.1		0	0	0	0		0	-		0	0	0	0	0	_	0	0	_	0	0	0	0	-	0	
Ē	F	÷	_	-	_	=	1332 0.47	Ĭ	Ť	÷	_	-	0 16061	0	0	0				0			0					_			_					-	0	
	Н	_	_	_	÷	Ī	÷	÷	_	_	_	_	=							0								_			_					-		
0211:0011:00	Н	10311031	8 R733735	_	Ť	÷	÷	÷	_	-	_	_	E4 0.0344828	_	- 2	-	-	-	0	-		0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	=	
20 1020-1100	H	-	8 0770286	_	_	_	_	_	_	_	_	1 0.133838	34 0.085105		36 0.166667													_			_					-		
2	Н	0.80-255	0.766738	0.703534	0.653195	0.571275	÷	Ť	0.342320	÷			0.0617284	0.03333														_			_					-		
920 1000	0.814815	0.72267	0.720384	0.683544	0.634515	0.50278	0.518353	0.451835	0.368859	_	_	0.151292	0.108844	_	0.0-0.0567	_					_							_	_		_	_	_	_	_			
020000	7800280	0.82270	0756598	0,702804	0.65554	0.571899	0.517520	0.461108	0.3699.42	0.312284	0.219613	0.142512	5757130	0.0428716	0.0227273													_	_		_	_						
820-200	0.833333	0.735.448	0.773.00	0.704508	0.661634	0.581746	0.521099	0.42864	0.400812	0.305906	0.254659	0.170744	0.118863	0.103825	0.0357148	0.03125												_			_							
830.830	11858333	1120.080	0.762235	750117.0	0.651835	0.60-238	0.5-48275	0.47549	0396230	0.323815	1,235071	0.161433	0.105568	0.0780781	0.0162162													_			_							
0.08.052	0.833333	0.800211	780287	0.732403	7491997	0.622306	0.55257	0.494205	0.392764	0.331052	0.255.423	0.197558	0.118211	0.0845538	0.0-26136		735210.0			_			_	_	_	_	_	_	_	_	_	_	_	_	_		_	
052 002	843373	.81.4286	782736	752731	689404	637363	1221201	479326	403834	330302	82882	2004002	.125182	(2000)	.0363636	0.0120482	0.06-2031																					
Г	F	_	0.4269	0 68-652	77210	0 2452 0	0.054788 0	.402452	0 250527	345987	_	Ť	135533	0 1606000	0 999691	_	0.280899 0			_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		-	
030,630	F	0.806305 0.8	Ť	_	Ť	_	Ť	Ť	_	_	÷	-	1120520 0.1	0.0780856 0.0	0.0367232 0.0	_	0.0			0	-							_			_					-	0	0.0
г	Н	181 601218	738354 1181		÷	÷	Ť	Ť	÷	÷	÷	÷	3.118058 R.15	0.0796216 0.07	Ξ	0.01.2213 0.01	0	0		0			0	0				_			_	0			0	-	0	
002022	F	Ť	_	_	_	_	_	_	_	_	÷	Ť	÷	_	_	1	0	0		0	-		0	0	0	0	0	_		0	_					-		0.0
2002	0.8933	0.8114	0.7808	0.727545	0.6567	0.5783	0.4065	0.436444	0.356159	0.3027	0.2406	0.1693	0.1196	1 0.0534228	_	456 0.00-E035												_			_					-		0 0
50,300	120827.0	0.773273	0.70245	0.63459.4	11582547	0.521715	0.465.01	0.381784	0.327/46	0.232-452	0.212341	0.1-18536	0.102659	0.035842	0.0242928	0.000700466												_			_					-		
400,50	0.615385	0.581731	0.5725	0.566116	0.517113	0.447548	0.355.351	0.333753	0.275212	0.22483	781.81.0	0.125334	0.0810065	0.0457305	0.00745573		0	0					0	0	0	0	0			0								
320-400	0.4915165	0.478261	0.4633-43	0,448833	0.415013	0.381265	0312100	0.283494	0.233583	8022610	0.141328	0.0958814	0.0552683	0.0128637													_	_			_							
300.350	2222	200004	62963	80233	18452	62812	55358	12281	70164	2962	282011-00	0.0531873	0.0199623	382-12000																								
Г	0.0	0	529 0.3	736 0.3	145 0.3	461 0.2	859 0.3	202 0.3	7963 0.1	_	_	0.0110121 0.0	0.000-256204 0.0	0.0	0	0	0	0	0	0		0	0	0	0	0	0	_	0	0	_	0	0	0	0	-	0	0.0
220.310	5 (1325)	1 0.235849	6 0.22372	2 (120)	51 0.228	36 0.197	43 0.172	17 0.142202	8 0.0817362	_	_	0.011	0000	-	-	-	-	-	0		-	0						-	-		-	-	-	-	-	-		
310.250	0.1578	175710	0.135716	-	0.0973451			_	0.032338	0.01391.47	701-152000										-							_			_					-		
150,200	0	0.0625	0.09875	0.0382166	0.030404	0.0254025	0.00485618	0.00754717			_																	_			_							
0 100 120	0		0								_									0								_			_					-		
New NATES   0.20 MeV/c   20.100   100.150   150.200	-			-							-		-	-	-	-					-							_	-		_	-				-		
0.20 Me)	0	0	0								_					0				0								-			-						0	
WANTES	C Deg	4.50 Deg	35 Deg	135-18 Dag	22.5 Deg	5-27 Day	31.5 Deg	5-36 Deg	40.5 Day	5-45 Day	40.5 Day	5-54 Day	28.5 Day	5-63 Day	67.5 Day	5-72 Day	76.5 Day	5-81 Day	85.5 Day	5-90 Day	94.5 Day	5-09 Day	1035 Dag	3.5-108 Deg	4412.5 Dag	35-117 Dag	7421.5 Day	.5 126 Day	1430.5 Dag	1.5-135 Deg	1430.5 Day	7.5-144 Drg	4-1-8-5 Dag	3.5-153 Deg	3-157.5 Dag	7.5-162 Day	2-100.5 Day	16.5 171 Day
É	Ē	4	5	Ĕ	ž	220	14	31.	36	ŧű.	4	49.0	í	585	63.1	225	ř	ř	81	855.	8	940	99	163	108	112	117	Ē	138	100	H	139	7	2	9	6	162	9 5

Table 5: Table for 2D Histogram for Old NM-Rein-Sehgal

1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
18-19-19-10-18-19-18-19-18-19-18-19-18-19-18-19-18-19-18-18-18-18-18-18-18-18-18-18-18-18-18-
No. 1 (1982)
200 - 200 -
1711-1710-1710-1710-1710-1710-1710-1710
18.00.00 (19.00.00 (19.00.00 (19.00.00 (19.0
18 18 18 18 18 18 18 18 18 18 18 18 18 1
10 10 10 10 10 10 10 10 10 10 10 10 10 1
10 10 10 10 10 10 10 10 10 10 10 10 10 1
20 (1997)
14-10-14-15-15-16-15-16-16-16-16-16-16-16-16-16-16-16-16-16-
25 25 25 25 25 25 25 25 25 25 25 25 25 2
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
100 100 100 100 100 100 100 100 100 100
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0
11 11 11 11 11 11 11 11 11 11 11 11 11
2
18   19   19   19   19   19   19   19
1.757-211 115 12 12 12 12 12 12 12 12 12 12 12 12 12
10 - 10 - 10 - 10 - 10 - 10 - 10 - 10 -
7. 17. 17. 17. 17. 17. 17. 17. 17. 17. 1
10 10 10 10 10 10 10 10 10 10 10 10 10 1
18.70.00 to 18.70.
2.00 (1997) 1.00 (
6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
88.07.67 8.05.77 8.05.
231 den 1 1 3 1 3 1 3 1 3 1 3 1 3 1 3 1 3 1 3 1
1000 000 000 000 000 000 000 000 000 00
10 10 10 10 10 10 10 10 10 10 10 10 10 1
440-441-441-441-441-441-441-441-441-441-
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
13 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
12   12   12   12   12   12   12   12
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
N. A.
12.2 Mar. 12.2 M
医库尔克氏氏试验检检验测验检验检验测验验 医克里曼 医克里曼 医格里氏病 医皮肤炎 化

Table 6: Table for 2D Histogram for New ANM-Rein-Sehgal

ANALIES HOUNES			_	-				_			=				_					_		=		_		00 100 E Day		The street	18-112-5 Deg   0	125-117 Deg   0	117-121.5 Dov. 0	121.5-126 Dev   0	36,1305 Dov. 0	131 S.1 S. Dov. 10	2 1305 Doy	135 5144 Dec	41 40 5 Day		THE CALLS LINE II	133 1575 Deg   0	0	162 1665 Deg 0	166.5-171 Deg B	17 1755 Day 8
V/C SEPTING			_	_				_			_	_			_					_		=		_					=		_	_							=	_			_	_
100-120	_	_	_					_	-		_	_		_	_					_		_		_					_	_	_	_	-	-					=	_		_		
Ï	0.037037 0	0.0538922 0	0.081226	0.00000		Total Maria	1825681	10135747	-			_	_	_	_				_	_		_	_	_			_	_	_	_	_								_	_	_	_	_	_
٦	Ť	0 179-87 0	0.175824 0	0 16291.0		_	=	n service in	-		_	0.0164031 0	9	_				_	9	_	-	_	9	_					_	_								-			9		-	-
III SHE	225352	264831	272864	26,46,47	0.00000	.000	186722	157462	140781	0.10701.00	70000	185222	047888	011465	710554017			_										_											_	_				
SHIP-SOIL	0.398693	0.351406	0.369365	8000000		7	-284672	1.237722	0.21821	100000	) diagona	9731676	0.099-6-4	0.0658882	0.0170428	o mimic	-		_			_	_						_	-									_	_	_	-		
330-400	0.510204	0.322654	72620	5030	0.40000	200	28182	33.8172	0.302653	000000	7 (57777)	0.200825	0.13113	0.103003	0.0537772	0.00000	0.00100107	Townson o																						_			- 0	- 0
Alle-Par	0.583034	961190	0.563628	8100020	O'LOUGH O	D-100000	0.46546	0.3833-0	0343622	10000	100-62-0	02267B	117744	0.127891	1258800	0.0400544	000000000000000000000000000000000000000	Decise Don																										
"		0.722011 0	0.680187	0 642008 0	-	-	_	B-65377 B	-		_	_	_	0.149639 0	0.116889	0		-	9	_		_	9	_					_	9	_	_								-	9	9	9	9
7	1.812332	1,822749	1,778963	13.000	0 1000	2000	. 28783	1,52,7443	000000	1000	1001001	1828	1,222,933	1372282	0.122696	1000000	00200	-	1.00515-464	_			_	_	_					_	_	_	_	_		-	_		_	-	_	_	_	_
SOU BILL	_	0.806912 0.	0.784609 0.	0.55.511		-	0.641457 0.	0.566528 0.	0.48047	-	=	0.320273	Ξ.	0.1785.6	0.12-613 0.			_	0.0163693 0.	_		_	9	_				-		-									2	9	9	-	-	-
#10-63fl	=	1812763 04	175534	0.748173	-	-	Φ	1573181	-	-	_	=	_	338367 0.	11-6336 0.	-	-	Ξ.	10144928 04	_	 2 1	_	-	-			2 0	2 1	_	-	-	-						2 1	_	_	-	-	-	-
0.00	_	807565 0.	77094 n	73165		-	630832	590777			-	_	_	390678 B.	1390-68		-	-	0.0071-285 0.			=	8						=	8							0.0		=		8	8		
ì	Ť	8777 08	280804 0.7	20028 0.7	-	-	ó	538564 0.5	-	0 0	=	_	_	185291 0.1	133106 0.1	-		_	0.89835	-		=		-			0 0		=		=	=	-	-			0.0		=					
_	<u>-</u>	80314 0.80	772247	-		_	610015 0.58	550457 0.53		0 0	_	_	25,756 0.23	18963 0.17	13/564 0.12	7				=		=		=	-		0 0		=		=	=	-	-					=				0	0
٦	_	803654 0.805	773010 0.77	7201 1000	-	_	\$5767 B.38.	535736 0.525		0.00	-	_	_	175556 0.136	121076 0.103	_	-	-		=		=		=	-		0 0		=		=	=							=	-				
1	_	803672 0.816	774887 0.769	22/2165		_	Ĕ	525933 0509	-	_	-	=	=	136364 0.15E	105263 0.054	-		Top or		=		=		=			0 0	= 1	=		=	=	-	-					=					
1	9	816404 0.79961	255928 0.772	1211211 0210031		-	Ï	50907 0513	-	0 0	-	Ĕ	-	1156327 0.116279	82-11-08					=		=		=	-		0 0		=		=	=	-	-					=	_			0	
1000	_	961 0.808856	7285 0.754215	THE REST		2	303 0.5KS0.	51202 0.49878	-	2 0	=	_		_	285 0.157895		0 0			=		=		=			3 0	= 1	=		=	=							=					
1996.1	_	22.032.032	115 0,756151	15 060785		-	100000	178 n.5n.2n.2	_	3 0	-	-	_	0.181818	- 9			-		_		=		=					=		=	=	-	-					=					
110011	_	72 0.73495	-	5 080%		Ė	_	2 0.5382	-	-	-	=	_	18 0.0833333	_		0.0			=		=		=				-	=		_	_							=					
38 1136-126	-	0.805395	0.756423	-		-	_	12 0.487885		9 0	=	_	0.307692	E	=				В	_		=	В	_				-	=		=	=							=		В			
12001-123	_	0.799636	3 0.745802	2000000		-	Ĕ	5 0.49 R.38	-	-	-	Ĕ	2 0233294	120	_					_		=		_					=		_	_							=					
1		0.809324	0.236299	0.082003		-	_	0.554704	-	3 0	=	0.22222			2					_		=		_				-	=		_	_							=					
1388-1336	0.86385	0.739766	0.725705	0.623330	72000	-	=	0.467192	0.320121	000	-	0.111111	7.0	_	_			-	В	_		=	В	_				-	=		_	_							=	=	В			
1338-1400	0.785124	0.790123	0.729168	0.680072	_	n-disedura	0.5332H	0.42857	0.342857	-	_	0.272727			_					_		_		_					_		_	_							=				_	_
1-99-1-99	905+88-0	0.793201	0.731919	0.05000.4	0 200120	2000	6901050	0.405405	2000	0.000	ULBS200	7			_					_		=		_					=		_	_							_					
1450-1300	0.820896	0.770154	0.712513	818189	0000000	-	=	0.464286	-	-	-			_	_							-							_	_	_	_							=			_		
Sulli-toon 1.	_	0.7907.8	10000000	063144 03		-	=	0.381679 0.0	-		-	0.53533	-	-	_				-	-		=	-	-	-			-	_	-	_	_	-	-					-	-	-	-	-	-
SINTERIO 16	_	9008	20017 0.3	20037 0.6		-	160605	Jan 23162 10.5			700007						0.0			=		=					0.0		=										=		0			
6HII-165H	÷	702-84 87	724876 0.7	677.481	-	-	=	346154 0.3	-	-	3		В	8				-	В	=		=	В	=	-		0 0		=	8							0.0		=		В	8		
17.1	-	5.733146 0.73	T21537 1.3	89.0 038.691		-	=	1348877 11.35	-		=				=		0 0			=		=		=	-		0 0		=		=	=	-	-			0.0		=				0	0
-	_	50 SEAS	13461 0.71	681818	_			258774 0.55	-		=	=	В	0	=		0.0	-	В	=		=	В	=			0 0		=	0	=	=							=	=	В	0		
3	Ŧ	786323 0.73	718579 0.71	÷	E E	-	_		_	2	=				=		0 0	-		=		=		=	-		0 0	= 1	=		=	=	-	-					=					
H	_	123.082 0.75091	7114m 0.098052	16098-48 0.615385		-	-	0307F02 0.25	-		=		0	0	=		0 0		0	=		=	0	=			0 0		=	0	=	=							=		0	0	0	0
	84 0.884GE	91 0.7703	52 0.719368	85 0 6.8275	-	-	44 0.647159	0.125		0 0	=				=		0 0			=		=		=			0 0	= 1	=		=	=							=					
	200		8	К	1	5	60				Ī																																	

Table 7: Table for 2D Histogram for New ANM-Berger-Sehgal

2	
2	
2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
1	
2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	
2 4 4 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
2	
88867 17472 17472 17472 17472 17472 17472 17472 17472 17472 17785	
8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
2. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	
2 (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)	
2 (2) (2) (2) (3) (3) (3) (3) (3) (3) (3) (3) (3) (3	
21 2 3 2 7 3	
25 25 25 25 25 25 25 25 25 25 25 25 25 2	
8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
2	
2 (2) (2) (2) (3) (3) (3) (3) (3) (3) (3) (3) (3) (3	
2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2	
2 (1987)	
2 (1972) 1 (	
200 (200 (200 (200 (200 (200 (200 (200	
15 of 18 15	
(8) (8) (8) (8) (8) (8) (8) (8) (8) (8)	
(1) 173 173 4 (1) 173 173 173 173 173 173 173 173 173 173	
1, 10, 10, 10, 10, 10, 10, 10, 10, 10, 1	
0.000 1 0.000	
1,59,600 P. 1,59,6	
2. 21. 33. 4 2. 21. 33. 4 2. 21. 21. 21. 21. 21. 21. 21. 21. 21. 2	
25 15 16 16 16 16 16 16 16 16 16 16 16 16 16	
1147.77 1147.77 1161.93 1161.9	
20 00 00 00 00 00 00 00 00 00 00 00 00 0	
14-15 Dec. 15-15 Dec.	185.17 Deg
	975

These are the corresponding tables to the 2D Efficiency plots for CC-Coherent Pion events.

Table 8: Table for 2D Histogram for New CC-Coh Pion NM-Rein-Sehgal

	_		_				_						_		_									_				_		_			
9	98/2	088888	35	_	_	_			_	_		_		-	-	-		_	_		_		_	-		_	_		-		_	_	
894	F	-	799990	_		_	_		_	_			_	_	_	_			_	_	_		_						_	_		_	
81	0.833333 0.	_	_								-		-	_	_	_	-		-	_	_	0.0	-	0	= 0	-	-	_	_	-	-	0.0	0.0
1820	_	_	_		-		_	_	_	_	_		-	0	0	0	- 0			-	0	0.0		-	= 0		-	0	_	-		0.0	
181	0.6523E	286 0.88	88	-	-	-			-	-	- 0						- 0		-		-	0 0	-	-	= 0	-	-		_	-		0.0	00
STATES ST	1990	10	69	-	0	-		-	-	_	- 0	-			_		- 0	-		_	-	00		-	= 0			0	-	-			
8	_	0.680	20	0,660	- 0			-	_	_	- 0				_	-	- 0		-	-	-		-	-	= 0	==	-			-	- 0		
9	Н	_	0.85	2							- 0											0 0			= 0							0.0	0.0
10016	Н	0.885565	_	8	_															_				-	<b>=</b> 0					-			
999	1	18883	0.61111	_	- 0			_									- 0			_				-	- 0					-			
	19191810	0.365217	0.539412	0.335353						_	- 0				_	_									- 0								
981984	200000	0836735	0.785714	673																													
14001430 14504300	200000	-	_	_	200000		_	_	-	-			_	_	_	-				_	-							-	_	-	_		
977	Н		9												_	_				_	_					. 4			_				
	0	_	÷	3000000																												0.0	0.0
130	0.913043   0.78	_	_	-	92																									-		0.0	0.0
9			0.7005 0.6	846 D.71												_	-		0.0		-		0.0	-	= 0					-			
120			_													0	- 0			_		0.0		-	= 0				_	-		0.0	0.0
8	0	5 0.885565		97 000,300,0																								0				0.0	0.0
1050-100   1100-150   1150-150   120-120   120-130   130-130	29/30/20 8	_	_	0.7858	9.7777	- =		_	-	_	- 0						- 0			_		0.0			= 0					-			0.0
1000116	8200280	_	_	0.75	0.77778	- =				_	- 0														- 0								0.0
30-100   100-100	811167.0	082285	700000		0.653846	3 _			-	_															= 0								
_	0.75	0.813725	_		900000	8											-			_				-		-			_		-		
888	0.89(802	0.838333	0	0.700067	0.78707	0.777	_			_				-	_						_												
89	0.71626	0.858824		0.70833	0.65372	9 5									_						_			_									
8	0.806422	-	0.755102	_	600ths 0.72	0.02377		_	-		- 0						- 0								- 0								
0.000	0.888889	0.83638	<u> </u>	_			_		-		- 0						- 0								- 0					-			
90	0	0.789474	_	_	06960/8	-	_	2	-	-	- 0													-	- 0								0.0
8	200080	0857380	0.830645	00000	0.69015	-	15	솀		-	- 0														- 0								0.0
8	989	0.8855	1 08/8125	0.736576	0.704		10.44		_	_	- 0						- 0							-	= 0				_				
8	0.81481	1 0.717946	3 0.79(333	174825	2 067906	1000		1 0.12857	5 033333		- 0				0	0	- 0			_					= 0			0	_	-			0.0
_	880 /	5 0.88574	6	0.736	3 067222	0.4887	0.45833	2 0.368[2]	0.45454	9999	- 0				0	0	_			_					= 0				_	-			0.0
999	9 064285	1 0854545	552KS 0.00000	6 0.711Hz	2 0.6348	0.5553	9 0.294118 0.322381 0.42 0.	3 0.2222	3 0.35.294	0 6	- 0				0	0	_			_					= 0				_	_			0.0
	0.571429	0.653.64	: 0.552R	0.5055	0.48113	1000	0.3228	0.200R	01730	8 02306	27				_		- 0				-			-	= 0								
30050 300400	0.3625	0.3000	0.6265	0.494845	0.4	2 20	0.291118	0.307547	0.3803.5	0.0034/8	OURSES.				_					_	-			-	= 0				_	-			
38830	03125	870	0.350877	0310345	0.34772	0.24486	0.147659	0.00483871 0.190076 0	0.1855	0.0545316	DESIGN OF STREET													_									
98	200010	ods	020023	램	82128	denot.		Doless71	00588285	0.0463138	0000000																			-	-		
	220	1235301	0.166667	6192700	0.25882	UNKAR	0.115385		-					-	-	-								-	_								
98	F	0.0768231		Joseph			0.029018		_	_			_		<u>-</u>						_	_			_				_		_		
57	9 6	-	-	0	0.0	-	10	-	-		0.0			2	0	0	0.0			-		0.0		-	-			10	0	-		-	
8	0	-	_	_	_			_	-	_				-	_	_				_	_			-	- 0			-	_	-			
New YALKS   0-30 MeV/c   30-100   100-150   150-200   200-200				_	_	_		_	_	_	_						_		_	_	_	_	_	_	_	_				_	_	_	_
MKS 0	9 Sac	25	Deg	SDg 0	2 de 1	200	i Dec	5Deg 0	o Spe	o sare	200	2	De C	3 Deg 0	5 Deg 0	Deg 0	# d	2	i d	Selesting 0	0 Ser 193	105 L25 L95	15 Deg 0	121 5 125 Deg 0	200	2 D 2 D 2 D 2 D 2 D 2 D 2 D 2 D 2 D 2 D	4 Deg	S Deg 0	3 Deg	75 Dec	200	166517 Dg	171.1755 Deg 0
New N	045 Dg	4501	9435	135-ls	18.225 Deg	1 12	315.35 Drg	35-40.5 De	40.5-45 Deg	\$100 PT	1000	200	63675 Deg	67.5.72 Day	72765	76.58l Dep	SI SES DE	100	945.00 Deg	99 IES	1529	122	11712	12.51	200	120	1887	144.14	14851	191	10.01	100	1221

Table 9: Table for 2D Histogram for New CC-Coh Pion NM-Berger-Sehgal

Ī	900	)900																																				
	ã	ã	_	0	_	_	_	_		-	-	=	0 1	_	_	_	_	0	_	0	0			0		_	_	0	_	0	_	_	_	0	_	_	0 1	= 0
	0.7 1 0.833333 0.75 1 0.000006 0.000006 0.4 0.5 1 1 0.000006	-	_		_	_	_	_		=	-	=	-	_	-	_	0	-	_	-	_	_	_	_	_	_	_		_		0	_	0	_	_		-	= 0
		_	_	_	_	_	_	_		=	-	-		_		_	0		_		_	_	_	_	_	_	_		_		0	_	0	_	_			= 0
	2.	_	_	_								_							_												0		0					= 0
		987	2		Ė	Ė	•		_							_	_	_		_	_	Ť	Ť	•	Ť	_	Ė		•		_	_	_	ė	•			
	20 /90	3	2		_	_	_	_		=	0 1	=	0 1	_	_	_	0		_		_						_		_		0		0	_	_	-	0 1	= 0
	88. 5	-		0	_	_	_	_		-	-	=	-	_		_	0	-	-	-	_	_	_	_	_	_	_	0	_	0		_		_	_	0	-	= 0
	9999	086	/900000	_						=		=			_				_		0																	= 0
		=	_	_		_	_	_		=	_	=	-	_	_		_		_		_							_	_	_					_		-	= 0
		7987	.,																																			
		ď	ď	_	_	_	_	_		-	-	-	-	_	0	_	0		-		_	-	-	_	-	_	_	0	_	0	0	_	0	_	_	0	-	= 0
	12	88	2	_					00	=	-	=	0 1		_		0		_		0	0	0		0						_		_			-	0 1	= 0
	0888	9	_	_					0.0	=		=			_				_																			= 0
	_	20	0.333333	_					_	=		=							_																			- 0
	-	9888	083888		_																						_											
	6		_	_	-	_	_	-	0.0	-	-	=	-	-	_	_	0	-	-	-	_	_	_	_	_	_	_	0	_	0	_	_	_	_	_	_	-	= 0
	88.		872		2			_	0.0	_	-	=	-	_			0		_								_							_			-	= 0
	2	0.81818	0.833333	910	_	_	_	_		_	-	=		_	_				_		_						_	_	_	_					_			= 0
Tables 1239131 1319131	0.00000	0.82105	0.333333	_								-							-																			
			J. S. C. C. C.		Ė	Ė			_						İ		_	_		_	_	_	_	•	_		Ė				_		_	Ė	•	İ		
TOPICI INTERIOR HONOR				20	_	_	_	_	0.0	=	-	=	-	_	0	_	0		-		_	_	_	_	_	_	_	0	_	0	0	_	0	_	_	0	-	= 0
	988	28	2.0	22 0.73	_				00	-	-	=					0		_												0		0					- 0
			280		2					=		=			_				_								_							_				= 0
SOME THEFT	12	0.73913	53	0.714286	_	_						-							-																			
	2000	77000	0.857143 0.684818 0.8 0	14286	•	_	_	_		_		_		_	_	_	_		_			_	_	_	_	_	<u>-</u>	_	_	_	_	_	_	<u>-</u>	_	_		
Bank	125	ď	34848 D.	14286				-	-	-	-	-	0	_	0		0		-		0		0	_	0		-	0		0	0		0	_		0		= 0
	8	2027 08	7143 0.6	8	4286 0.5		-	-	0.0	=	-	=	-	-	0		0		-					_			_	0	-	0	0		0	_	-	0	-	= 0
	'	3714286 0877	889	99			_	-	00	=	-	=	-	-	-	_	0	-	_	-	_	_				_	_	0	-	0	0	_	0		-		-	= 0
DESCRIPTION OF STREET	_	8714	15 15 15 15 15 15 15 15 15 15 15 15 15 1	95 058333	8	2		-	00	-	-	=	-	-	-		0		_								_		-					_	-	-	-	= 0
	2 0.7	8	7 0.7149	7 0.7368	20 02	_			0.0	=		=			_				_																			= 0
DAME IN COLUMN	0.83333	_	0.82am7				_	_		_	-	_		_	_	_			_		_	_	_	_	_	_	_		_			_		_	_			- 0
	_	0.000	_		0.62C3TS	0.77778			_										_																			
	2000	1882280	188	17897	29000	1999	244444	_		_		_	_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
	310007	1000	0.0000007 0	1000	18	19000		_		_		_	_ 1	_	_	۷	٥		-		_	٥	٥	٥	٥	۷	۷	_		_	٥	۷	٥			_	_ 1	_ 0
	0.855 1 0.94 0.057	55318 O.	ã	GE354 0.	39112 0.	9048	2014			-	-	=	-	_	0	_	0	-	-	-	_	_	_	_	_	_	_	0	_	0	0	_	0	_	_	_	-	= 0
	-	2318 D.G.	3867 0.8	90	3867 0.5	9867 0.65	점	3.0		=		=	0 1		_		0		_		0	0	0	0	0						0		0			0	0 1	= 0
	80.		0.600067				2	2.	- c	=	-	=	-	_	-	_	0	-	=	-	_	_	_		_	_	_		_		0	_	0	_	_		-	= 0
	077778	9.9	04375 0.5	33	g 0.355217	33 B.338462	3	B 0.355553				=		-	-		0		-								_		_					_	_			= 0
	8	3	04375	65789			1 Si	1000	3 0	8		=	-	-		_	0		_							_	_		_		0	_	0	_	_		-	= 0
	2	0.5555	23		0.193548			908	0.181818	9	-	_		_	_				_					_			_							_		_		= 0
		í		eg.	00768231				0												_																	
	-		13 -				ulsisis u		GUNESCE			_	_		_	۷	٥		-		_	٥	٥	٢	٥	۷	_					۷				_	_	_ 0
	-	-	널	_	_	3	3	á	86	1	-	=	-	-	-		0	0	-	0	_		-	_	-		_	0	-	0	0		0	_	-	0	-	= 0
	-		_	0	011111	_	-	-		=	-	=	-	-			0	-	-	-	_	-		_			_	0	-	0	0		0	_	-	0	-	= 0
	-	-		0		-	-	-		=	-	=	-	-			0		-					-				0	-	0				_	-		-	= 0
	_	-	_		_	_	_	_		=	-	=	-	_	-	_	0	-	=	-	_	_	_		_	_	_		_		0	_	0	_	_		-	= 0
DESCRIPTION OF THE PROPERTY OF	_		_	-	_	_	_	_		-	_	_	_	_	-	_		_	-	_	_	_	_	_	_	_	_	-	_	-		_		_	_	0	_	_
THE PART INC	ď.	287	5 Deg	SDeg	SDeg	2 Deg	S Deg	2479	S Deg	9370	2 FE	Sar re	S Deg	2012	5 Deg	2Deg	5 Deg	d Deg	5 Dec	ag De	5 Deg	9 Deg	35 Dg	JOS Deg	125 Dg	H7Dg	21.5 Dg	126 Dg	302 Dec	132 Dg	395 Deg	144 Deg	485 Deg	133 Dg	57.5 Deg	162.14g	665 Deg	1
	2	9	9.135	1332	183	8	ri Ki	13 15	9 5	į.	9	8	89	8	89	67.57	72.76	765.8	8	55	909	95.59	30 IC	1685	1087	1125	1171	5	ď.	5	Ĭ	199	7	1485	ď.	12/0	Ž,	9 5
	ğ.	305	9-13.5 I	135-181	182251	125.02	20.00	8	20,403	9	9	1000	250	9	636753	67572	727651	765.81	8,855	8550	909153	915.00	991035	1035-10	108.112	1125.11	117.121	121512	8	130513	135.130	130514	144.148	1485-15		ja ja	10,000	157.5.162.Deg

Table 10: Table for 2D Histogram for Old CC-Coh Pion NM-Rein-Sehgal

Table 11: Table for 2D Histogram for New CC-Coh Pion ANM-Rein-Sehgal

966200	0.818182	571429		_			_	_				_		_	_	_		_	_	_			_	_	_				_			_	
9819	-	1	_	_			_						_	_	_				_					_	_							_	
198	27		-	_	0.0		_		0 1		-		0	_	_					-		-		_	_	= 0	0	= 1	= 0	0.0	-	_	
1850	0.76905 0.76	9	_	_		0	_		-	0 0	-		0	_	_	-					- 0	0.0		_	_	= 0	- 0	= 1	= 0		0.0	0	
188			_	_			_		-		-		0	_	_	- 0	-			-		-	_	_	_	= 0	- 0	= 1	= 0		-	_	
(A)	897760		_	_	0.0		_		-					_	_	- 0			_					_	_	= 0	- 0	= 1	= 0			-	
170017	0.9087	§ 2	12				_		- 1																	= 0	- 0	= 1	= 0			_	
1650170	92	0.68421	_				_		-						_	_			_					_	_	= 0		- 1	= 0			_	
Society 1500-100 (society 1600-170 (700-170 (170-180 (180-180 (190-190 (1900-190 (190-190-190 (190-190 (190-190 (190-190 (190-190 (190-190)(190-190 (190-190)(190-190 (190-190)(190-190)(190-190 (190-190)(190-190)(190-190)(190-190)(190-190)(190-190)(190-190-190)(190-190-190)(190-190-190-190-190-190-190-190-190-190-	1253180																									= 0		-					
001000	163371	80.08		_			_	_	_			_	_	_	_			_	_	_			_	_	_							_	
83.00	57143	2000	_	_			_	_				_	_	_	Ī			_	Ī				Ī	_	_							_	
88	877		-	_	0.0	0.00	_	_	-	-	-	-	0	_	_	-		-	_	-	= 0	-	_	_	_	= 0		2	= 0		-	-	
57	-		93	-		-	_	Ξ.	0 1				0	_	_	_			_	-	= 0	-		_		= 0	-	= 1	= 0	- 0	-	_	
1400			99 055550	-					0 1	- 0						- 0	-		0	-	- 0	0 0				= 0	- 0	= 1	= 0		0 0		
8		S CARGO	4s 0.6315/9	_			_		-		-			_	_		-			-			_	_	_	= 0	- 0	= 1	= 0			-	
1908	0.0000	3 2	7 0.6500	98					0 1				0			- 0			0							= 0	- 0	= 1	= 0	-			
12043	-	224.38	0.6830	910			_		-	- 0			0	_	_	- 0				-				_	-	= 0	- 0	= 1	= 0				
99-100 100-100 100-110 110-120 130-120 120-130 130-130 130-130 130-130 130-140 140-140 140-140	082580	0.66234	OSISSII	0.0857			_		0 1				0	_	_	- 0			0					_	-	= 0	- 0	- 1	- 0				
1150-120	0.774194	29.262	0.71864	0.5555	2.											_										= 0		-	= 0				
81100	1811463	2000	3,727,273	90	7.		_		_	_			_	_	_	_			_		_			_			-		_			_	
818	1128	900	-	0.676471			_	_	_			_	_	_	_			_	_				_	_	_							_	
0900	10127	9 2	7,4996 0.		10000	-	_	-	-	-		-	_	_	_	-		-	_	-	- 0	-	_	_	_	= 0	-	= 1	= 0		-	_	
8	80	3 2 2	50 07		0.6619 0.6		_		-	00	-		0	_	_	-	-				= 0	0.00		_	_	= 0	-	= 1	= 0	-	0.0	-	
8	19814 0.7	92	15054 0.0		0.58883 0.4	-		0	0 1		0 0	0	0	0	0	- 0		0		0 1	= 0	-	0	0		= 0	0 0	= 1	= 0	0 0	-	-	
70 STO STORED STORED 900 STORED	6.00		7.12189 0.7		7 055		_		0 1				0	0	0	- 0	-			0	= 0	-	0	0		= 0	-	= 1	= 0	0 0	-	-	
22	3667 08	100	35192 0.7	108704 0.66	553816 0.6 76		-		0 1				0	0	0	- 0			0	0	= 0	-	0	0		= 0	-	= 1	= 0	0 0	-	-	
8	80 08		32730	4709 0.7	0000	1515155	000000	-	-	00			0	_	_	00	0 0		_	-	0 0	0.0	-	_	_	= 0	-	= 1	= 0	- 0	0.0	0	
	61111 08		58523 07	10001	20 12/130	_	_	0	-	-			0			- 0				0	= 0	-	0			= 0	-	= 1	= 0	-	-	-	
8	0.81802 0.8	100	48276 0.7	799677 0.5	1559		0.44444 0.2		-	00	-		0	_	_	- 0				0	= 0	-	0	_		= 0	-	= 1	= 0	-	-	_	
600 650 700 700 700 700 700 700 700 700 700 7	S20053 02	2017 SWG 0.7	748800 0.7	300187 0.7	57.082 11at 02	33.0877 0.5	33836 0.X		-	-			0	_	_	- 0			0	-	= 0		_	_	_	= 0	-	= 1	= 0			_	0 0
2000	180		764858 0.7	ã	GE3852 05		123258 0.5	_	-	0.0			0	_	_	- 0	-		0	-	= 0		_	_	_	= 0		-	= 0			_	0 0
19	0.789474 0.7	385	0.07	_	od otes 157773		<u> </u>	1901301	-	0 0			-	0		-			-	0 1	- 0	0.0		0		= 0	-	= 1	= 0	-	0.0	-	
	0.719512 0.	·	_	_		8	13	_	21	00	-		0	_	_	00		-	_	-	- 0	0.00	_	_	_	= 0	-	= 1	= 0	-	0.00	0	0.0
20-fe   40-fe   40-fe	0.569231 0.4	15.25 OPT	J. 137888 0.513006 0.638243	154555 062864	0.38465 0.457.43 0.5565 0.20ms 0.40778 0.44444	35281	obese obsess over over	20378 0.	9	)9			0	_	_	00			_	-	- 0	0.00		_	_	= 0	-	= 1	= 0		0.00		0.0
9	365		\$7888 D.	458lol 03	Solds 0.	150	198538	187919 0.	2549 0.	0 0			0	_	_	0.0			_	-	- 0	0.00	-	_	_	= 0		= 1	= 0		0.00		0.0
800	6	3000	328820		2005	200	200	S/00/2	1,000	0 0 0 0			-	0	0				0	0 1				0		= 0	0.0	= 1	= 0	0.0		_	00
	10		8574 03	_	0.0	12	odesies ode	10 Jan	100	3 c	-	-	0	_	_	-	-	-	0	-		0.0	-	_	-	= 0	-	= 1	= 0	-	0.0	-	00
20030	31	3 2		25 023enl	tes others	72 0165	150 El 32	222	98	ij.	-		0	-	_	- 0			-	-		-	-	_	-	= 0	- 0	= 1	= 0	-	-	-	
New ANM R.S. 0-20 MeV c   30-100   100-120   150-20   200-20	0.2302.0	100	3 oleni.	9 0.1372	oofstas olasta o	0.08837.3	0.0016422	0.012	-				-	_	_	- 0								_	_	= 0	- 0	= 1	= 0	- 0			
150.20		nnesisie.	DICTOR	0.04764	0.0233		_		0.1				0			_			0					_	_	= 0	- 0	= 1	= 0			_	
10015			0	_			_		0 1					_	_	- 0			0	-			_	_		= 0	- 0	- 1	= 0	-		-	
W. C. S.			0	_			_		0.1	0 0				_	_	- 0			0	-	- 0	0.0		_	_	= 0	- 0	= 1	= 0	0.0	0.00		00
W E-B V			0				_		0.1				0								= 0		_	_	_	= 0	- 0	= 1	= 0			_	00
wANG RE	5 Deg	S Dec	NB Deg	225 Deg	225.27 Deg	San	36-40.5 Deg	545 Deg	5-49.5 Lbg	49554 Deg	Section 1	63675 Drg	572 Deg	765 Deg	765-81 Deg	825Deg	mod 5 Dec	200 Deg	loss Deg	SAIR DE	51125 Lig	11712 5 Deg	.5135 Deg	Hans Deg	SIZE De	100	Series Inc.	200	10.00	20 CO 10	HES De	Sel Pies	17.17.55 Dg 17.5.5180 Dg
New	i.	\$ <del>2</del>	135	185	ลัธ	176	Ä.	9	J.	97	5 8	8	63	25	765	8	á	95	8	<u> </u>	4 5	H	2	123	<u> </u>	9 8	2 :	3	2 5	37	19	8	EE

Table 12: Table for 2D Histogram for New CC-Coh Pion ANM-Berger-Sehgal

	30-300																																			
1   1   1   1   1   1   1   1   1   1	$\vdash$	_	98				-	_	-	_	_		_	0.0		-			-	_	_	-	-		-	_	_	_	_	-	_	_	_	-	-	-
1   1   1   1   1   1   1   1   1   1	-1900 1900	-	_			-		-	-	_	-	-	=					-		_	-	-		-			0			-	-	0	_	- 0	= =	
1		_	#B			-				-	0	-	=			= 0		-	0	0	-	-		-		0					_		_	- 0		
1   1   1   1   1   1   1   1   1   1		_				-				_	_		-			= 0		-			-						_		_		_		_	- 0		
Column   C	Enlar	Ļ	89										_																							
Column   C	001730	15	6																																	
Column   C	1 00/10	27.48	8		0 0			_	_	_	_		-				0 0		-	_	_	-				_	_	_	_	_	_	_	_	-	==	-
Column   C	E	88	129	-		-		_	0	_	0	00	=	0.0		= 0		-		_	-	-			0	0		0	_	0	_		_	- 0		
Column   C	-		_			-				_	0	- 0	=			= 0		-		_	-	-									_		-	- 0		
1   1   1   1   1   1   1   1   1   1	F	_	063636	9								-	-			= 0				_		-											_	- 0		
Column   C	Smilan	083333	9	Ŷ.									_			= 0					_															
Column   C	15015m	877777	90																																	
1   1   1   1   1   1   1   1   1   1	1 61430	F	8					_	_	_	_		_			= 0				_	_	-	-		0.00	_	_	_	_	_	_	_	_	- 0		
Column   C	14m 14c	716		i -		-	-	_	-	_	0	-	=	0.0		= 0		-		_	0	-			-	-	_	0	_	-	0		_	- 0		000
1		Ť	÷	9 2	3 -	_		_	_	_			=			= 0		_		_		-				_	_	0	_	_	0	_	_	- 0		
Column   C	3 13mb1	F	-	g						_		- 0	=							_		-				_		0	_		0		_	- 0		
1	1250130	/900000	0.795103	9 9	3 -						_		-								_										_					
1	20120	184654	1851852	2 2	Description 1	ĺ	_	_	_	_		_	_	_		_	_	_	_	_		_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
Column   C	1120	Ē	Ξ	_	,			_	_	_	_		_							_	_						_	_	_	_	_	_	_			
	HS0 H	-	_	-	-	. =	-			_	_	0.0	=	0.0		= 0				_	_	-	= 0	0.0	0			_			_		_	= 0	0.0	-
Column   C	100	P	_	-			0			_	_	- 0	=			= 0		-		0	_	0 1	- 0	0 0	0			_			_			0 0		
1	ㄷ	99990	÷		-	-					_	- 0	=			= 0				_	_	-	= 0					_			_		_	- 0		
1	1000-100	1970	OSES	200	0.0000					_			-			= 0				_		-											_			
1	9804000	0.78923	0.8103022	2		3 =					_		_							0	_															
1	000-000	0.77778	88	) S (S)	1.00cm								_								_															
1	000	90	763830	1000	769.6	75000		_	_	_	_	_	_						_	_	_	_	-	-		_	_	_	_	_	_	_	_	_		
Column   C	Н	1.794872	813333	200	O'Dare	9	333333	_	_	_	_	_	_		_			_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_		
	Ξ	9623-16	85865	100	CCCCC.			_	_	_	_		_						_	_	_	_				_	_	_	_	_	_	_	_			
1	2 05/0	8384 0	75257	200	2 James	2CTR		-		_	_	0.0	_	0.0		= 0				_	_	-	= 0		-	_	_	_	_		_	-	_	-	-	
1	90	9	ď	200		-	-	_	-	_	0	-	_		0 1	= 0				_	_	-			0.00	_	0	0	0		0	0	_	- 0		
1	_	-	-	3 6	0 0	1	_	88		_	0	0.0	=	0.0	0 1	= 0		-		_	0	0	= 0		0 0	_	_	0	_		0		_	0 0	0.0	
	Ē	ᆮ	8		-	_		2	1 200	_	0		=			= 0		_		_	0	-			-	-	_	0	_	-	0	-	_	- 0	==	00
1	Н		_		, ,	-			99			-	=	0.0	- 1	= 0		-				-	= 0	-	0 0			-		0	_			= 0		-
1	8		18 18 18 18 18 18 18 18 18 18 18 18 18 1				32	38 135		_		-	=			= 0		-		_		-				_	_	0	_	_	0	_	_	- 0		00
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	H		22 O.78TB	8 000	100	_					0	-	-			= 0		-				-		-				0			0			- 0		
	Н	Н	_	_	-	_	_	_	_	_	-	0-	-			= 0		. =		_							_	0	_		0		_	- 0		
1	330400	Ĕ	_									_	=		_			_	_	_	_	_				_	_	_	_	_	_	_	_	_		
	30830	8	0361702	082380 0845280	03550	Sec.	2	910	012905				-										-													
	2030	12	28574	1000	Select.	2018	d48148	deget	0.0555533	Hess	-	_	_						-	_	_	_				_	_	-	_	_	-	_	_	_		_
	П	-	_		-	-	_		_	٥	۷	ە ت	_			_ 0		_		_		ا ت	_ 0			_	٥	٥	٥		٥	٥	٥	_ 0		
	П	20	3					98	98	_	0	0.0	=	0.0		= 0		-		_	0	0	= 0	-		_	_	0	_		0		_	0 0		
		0		- 6	100	2		0	0			0 0	=			= 0						0		-	0 0					0	-			- 0		
		0		= =								0.0	=	0.0		= 0						0	- 0	-	0 0									0 0		
10   10   10   10   10   10   10   10	0	0	_	0.0		_		_		_	0	00	=	0.0	0 1	= 0		_		_	0	01	- 0	-		_	_	0	_				_	0 0	0.0	
100 M	S 0-30 Mi		_			. =		_		_			-			= 0		. =		_		-					_	0	_				_	- 0		
2 日本 日本 日本 日本 日本 日本 日本 日本 日本 日本 日本 日本 日本	AMIBS	5 Deg	9 de	Ja De	25 Dec	Z Dec	15 Deg	36 Deg	05 Deg	45 Deg	95 Dg	St De	Series.	75.00	2	2 P. C.	S De	25 Dec	Seg Des	45 Deg	2010	085 Deg	126 14	2117 Dec	1215 Dec	5125 Deg	1305 Deg	SIE De	1305 Deg	S14 Dg	1485 Deg	SIE De	1575 Deg	5162 Deg	20 PM	1755 Deg
	New	Ĭ	ij.	9 1	2	E)	27.53	100	78	9	Ą	97	8	98	3 (	36	182	200	120	8	250	8.	9 8	100	H	121	8	é	ń	Ŕ	144	148.	19	3 6	3 2	E

## ${\bf D} \quad {\bf Acknowledgements}$

Thank everyone who helped, and thank everyone who gave their inputs into your acceptance study. YOU NEED TO GIVE A HUGE AND SPECIAL THANKS TO DR. ASAADI RIGHT HERE! (He has been suuuuuuuper patient...)