

Problem 1 In class we proved the Euler-Maclaurin summation formula

$$\int_0^1 f(x)dx = \frac{1}{2} (f(0) + f(1)) + \sum_{m=1}^{\infty} b_m \left(f^{(2m-1)}(1) - f^{(2m-1)}(0) \right)$$

for some unknown constants b_m independent of f .

(a) Find a recursive formula for b_m by evaluating both sides for $f(x) = e^{\lambda x}$ where λ is a parameter.

(b) Compute $b_1, b_2, b_3, \dots, b_{10}$.

(c) Compound the formula to show

$$\int_0^n f(x)dx = \frac{1}{2}f(0) + f(1) + f(2) + \dots + f(n-1) + \frac{1}{2}f(n) + \sum_{m=1}^{\infty} b_m \left(f^{(2m-1)}(n) - f^{(2m-1)}(0) \right).$$

(d) Use the Euler-Maclaurin formula to show that

$$\sum_{j=1}^n j^k = P_{k+1}(n)$$

is a degree- $(k+1)$ polynomial in n . Example:

$$\sum_{j=1}^n j = \frac{n(n+1)}{2}.$$

(e) Show that the error in the trapezoidal rule satisfies

$$\begin{aligned} \int_0^1 f(x)dx - h \left(\frac{1}{2}f(0) + f(h) + f(2h) + \dots + f((n-1)h) + \frac{1}{2}f(nh) \right) \\ = \sum_{m=1}^{\infty} b_m h^{2m} \left(f^{(2m-1)}(1) - f^{(2m-1)}(0) \right). \end{aligned}$$

Problem 2 Write a matlab program `ectr.m` of the form

```
function w = ectr(n, k)
% n : quadrature points are 0 ... n
% k: degree of precision
```

which produces the weight vector (w_0, \dots, w_n) containing endpoint-corrected trapezoidal weights of even order $k = 2, 4, \dots, 10$: for given $n \geq 2k$. Your code should combine previous codes for the differentiation matrix, the Bernoulli numbers b_1 through b_{10} , and the Euler-Maclaurin summation formula. Use it to complete the following table of endpoint-corrected trapezoidal weights of even order $k = 2, 4, \dots, 10$:

k	w_0	w_1	w_2	w_3	w_4	w_5	w_6	w_7	\dots
2	$1/2$	1	1	1	1	1	1	1	\dots
4	$9/24$	$28/24$	$23/24$	1	1	1	1	1	\dots
6						1	1	1	\dots
\vdots									

Check the order of accuracy of the weights on

$$\int_0^1 e^x dx$$

for $h = 1/32, \dots, 1/1024$ and even $k = 2, \dots, 10$.

Problem 3 (a) Find an exact formula for the quintic polynomial $P_5(x) = x^5 + \dots$ such that

$$\int_{-1}^1 P_5(x)q(x)dx = 0$$

for any quartic polynomial q .

(b) Find exact formulas for the five roots x_1, x_2, x_3, x_4, x_5 of the equation $P_5(x) = 0$.

(c) Find exact formulas for the integration weights w_1, w_2, w_3, w_4, w_5 such that

$$\int_{-1}^1 q(x)dx = \sum_{j=1}^5 w_j q(x_j)$$

exactly whenever q is a polynomial of degree 5.

(d) Given any real numbers $a < b$, find exact formulas for points $y_j \in [a, b]$ and weights $u_j > 0$ such that

$$\int_a^b q(x)dx = \sum_{j=1}^5 u_j q(y_j)$$

whenever q is a polynomial of degree 5.

(e) Explain why each of the three factors in the error estimate

$$\int_a^b f(x)dx - \sum_{j=1}^5 u_j f(y_j) = C_{10} f^{(10)}(\xi) \int_a^b (y-y_1)^2 (y-y_2)^2 (y-y_3)^2 (y-y_4)^2 (y-y_5)^2 dy$$

is inevitable and determine the exact value of the constant C_{10} .

Problem 4 Write, test and debug an adaptive 5-point Gaussian integration code `gadap.m` of the form

```
function [int, abt] = gadap(a, b, f, r, tol)
% a,b: interval endpoints with a < b
% f: function handle f(x, r) to integrate
% r: parameters for f
% tol: User-provided tolerance for integral accuracy
% int: Approximation to the integral
% abt: Endpoints and approximations
```

Build a list $\mathbf{abt} = \{[a_1, b_1, t_1], \dots, [a_n, b_n, t_n]\}$ of n intervals $[a_j, b_j]$ and approximate integrals $t_j \approx \int_{a_j}^{b_j} f(x, r) dx$, computed with 5-point Gaussian integration. Initialize with $n = 1$ and $[a_1, b_1] = [a, b]$. At each step $j = 1, 2, \dots$, subdivide interval j into left and right half-intervals l and r , and approximate the integrals t_l and t_r over each half-interval by 5-point Gaussian quadrature. If

$$|t_j - (t_l + t_r)| > \mathbf{tol} \max(|t_j|, |t_l| + |t_r|)$$

add the half-intervals l and r and approximations t_l and t_r to the list. Otherwise, increment \mathbf{int} by t_j . Guard against infinite loops and floating-point issues as you see fit and briefly justify your design decisions in comments.

Problem 5 (a) Show that

$$\int_0^1 x^{-x} dx = \sum_{n=1}^{\infty} n^{-n}$$

- (b) Use the sum in (a) to evaluate the integral in (a) to 12-digit accuracy.
- (c) Evaluate the integral in (a) by `ectr.m` to 1, 2, and 3-digit accuracy. Estimate how many function evaluations will be required to achieve p -digit accuracy for $1 \leq p \leq 12$. Explain the agreement or disagreement of your results with theory.
- (d) Approximate the integral $\int_0^1 x^{-x} dx$ using your code `gadap.m`. Tabulate the total number of function evaluations required to obtain p -digit accuracy for $1 \leq p \leq 10$. Compare your results with the results and estimates for endpoint-corrected trapezoidal integration obtained in (c).

Problem 6 Implement, debug and test a MATLAB function `pleg.m` of the form

```
function p = pleg(t, n)
% t: evaluation point
% n: degree of polynomial
```

This function evaluates a single value $P_n(t)$ of the monic Legendre polynomial P_n of degree n , at evaluation point t with $|t| \leq 1$. Here $P_0 = 1$, $P_1(t) = t$ and P_n is determined by the recurrence

$$P_n(t) = tP_{n-1}(t) - c_n P_{n-2}(t)$$

for $n \geq 2$, where $c_n = (n-1)^2/(4(n-1)^2 - 1)$. Be sure to iterate forward from $n = 0$ rather than recurse backward from n , and do not generate any new function handles. Test that your function gives the right values for small n where you know P_n .

Problem 7 Implement a MATLAB function `gaussint.m` of the form

```
function [w, t] = gaussint( n )
% n: Number of Gauss weights and points
```

which computes weights w and points t for the n -point Gaussian integration rule

$$\int_{-1}^1 f(t)dt \approx \sum_{j=1}^n w_j f(t_j).$$

(a) Find the points t_j to as high precision as possible, by applying your code `bisection.m` to `p1eg.m`. Bracket each t_j initially by the observation that the zeroes of P_{n-1} separate the zeroes of P_n for every n . Thus the single zero of $P_1 = t$ separates the interval $[-1, 1]$ into two intervals, each containing exactly one zero of P_2 . The two zeroes of P_2 separate the interval $[-1, 1]$ into three intervals, and so forth. Thus you will find all the zeroes of P_1, P_2, \dots, P_{n-1} in the process of finding all the zeroes of P_n .

(b) Find the weights w_j to as high precision as possible by applying your code `gadap.m` to

$$w_j = \int_{-1}^1 L_j(t)^2 dt$$

where L_j is the j th Lagrange basis polynomial for interpolating at t_1, t_2, \dots, t_n .

(c) For $1 \leq n \leq 20$, test that your weights and points integrate monomials $f(t) = t^j$ exactly for $0 \leq j \leq 2n - 1$.