

# Applied Complex Analysis

Willie Rush Lim

2020



## Introduction

Denote the set of complex numbers by

$$\mathbb{C} = \{x + iy \mid x, y \in \mathbb{R}\}$$

where  $i = \sqrt{-1}$  is defined such that  $i^2 = -1$ .

Complex analysis is the study of functions of a complex variable. In the first few chapters, we shall explore some introductory concepts, such as basic properties of complex numbers and continuity of complex-valued functions. The main emphasis is the concept of *holomorphic* functions, i.e. complex-valued functions which are differentiable in a complex sense, and the many applications of their somewhat magical properties. I used the word 'magical' because holomorphicity is such a rigid condition that many of the results you will see are somewhat unintuitive yet true.

We will start with some motivation. Basic algebra tells us that the number of roots of a polynomial with real coefficients is at most its degree. For example,  $x^2 + c$  has two real roots if  $c < 0$ , one root if  $c = 0$ , and no roots if  $c > 0$ . Introducing the imaginary number  $i$  provides us with a more elegant way of formulating this idea.

**Theorem** (Fundamental Theorem of Algebra). *The field  $\mathbb{C}$  is algebraically closed, that is, any polynomial with coefficients in  $\mathbb{C}$  of degree  $d > 1$  has exactly  $d$  roots in  $\mathbb{C}$ , counting multiplicity.*

Many initial attempts of proving the theorem by prominent mathematicians D'Alembert, Euler, Gauss, Lagrange, and Laplace in 1700s were incomplete. In 1806, a Swiss accountant, Parisian bookstore manager and 'amateur' mathematician Jean-Robert Argand completed D'Alembert's ideas and hence became the first person to rigorously prove the fundamental theorem of algebra. We will in fact use properties of holomorphic functions to give 3 different proofs of the theorem, including D'Alembert and Argand's approach.

It is difficult to list the many applications of the fundamental theorem of algebra. The main idea is that the field of complex numbers is the perfect setting to solve equations!

A direct consequence in linear algebra is that every square matrix with entries in  $\mathbb{C}$  admits an eigenvalue. When a  $2 \times 2$  matrix has imaginary eigenvalues, it acts as a rotation of the plane rather than expansion or contraction in certain directions. In the study of continuous dynamics arising from mechanical systems, it is common to use complex numbers in order to capture oscillations in the system.

One of the direct applications of the study of holomorphic functions is contour integration. The integral of a complex function along a closed path is not dependent on the path itself but rather on certain values called *residues* of the function's singularities. This means that it is often easier to integrate a real function of a real variable by converting it into a problem involving a contour integral in the complex plane.

Fourier series and Fourier transforms are useful in decomposing functions into its frequency components. (Think of decomposing nice functions as a sum or an integral of different sine and cosine waves.) Fourier analysis can be easily formulated via complex analysis, and it comes up everywhere: in differential equations, probability, quantum mechanics, signal processing, etc.

Mechanical and electrical engineers as well as computer musicians also encounter complex variables in electrical circuits with alternating current. Digital filters are designed by looking at the locations of *zeros* and *poles* of rational functions called *transfer functions*, which essentially model a device's inputs and outputs.

Iterations of holomorphic functions have long been known to have many applications. Complex polynomials, for example, can be used to model the population of rabbits over time. Powerful basic results in complex analysis, many of which do not apply to generic real differentiable functions, make up one of the many reasons why the study of iterations of holomorphic functions (holomorphic dynamics) is very well developed compared to the other branches of the field of dynamical systems.

Conformal functions are holomorphic functions with strictly non-zero derivative. Such functions have an amazing geometric property of angle preservation at every point and are useful in transforming regions with complicated boundary to those of a much nicer shape (square, disk, etc). You may, for example, want to transform a mechanical problem on a complicated domain into an equivalent problem on a circular disk. In cartography, conformal maps are useful in creating a world map as well as local nautical charts using Mercator and stereographic projections. More recently, conformal functions are applied to the surface of the human brain for brain development study and diagnosis of Alzheimer's disease and schizophrenia.

# Chapter 1

## Complex Numbers

In this chapter, we will go through the basic algebraic and geometric properties of complex numbers.

### 1.1 The Algebra of $\mathbb{C}$

The set  $\mathbb{C}$  is equipped with the usual arithmetic operators, namely:

- addition  $+$ :  $(x + iy) + (a + ib) = (x + a) + i(y + b)$ ,
- multiplication  $\times$ :  $(x + iy) \times (a + ib) = (xa - yb) + i(xb + ya)$ .

Let's denote by  $\mathbb{C}^*$  the set of non-zero complex numbers  $\mathbb{C} \setminus \{0\}$ . This set is equipped with an additional operator:

- inversion of a non-zero number:  $(x + iy)^{-1} = \frac{x - iy}{x^2 + y^2}$ .

Similar to  $\mathbb{R}$ , the set of complex numbers  $\mathbb{C}$  is a *field*; it satisfies the following axioms:

1.  $(\mathbb{C}, +)$  is an abelian group:
  - $+$  is associative and commutative,
  - $0$  is the identity element of  $+$ , i.e.  $z + 0 = z$  for all  $z \in \mathbb{C}$ ,
  - Additive inverses exist, i.e.  $z + (-z) = 0$  for all  $z \in \mathbb{C}$ ;
2.  $(\mathbb{C}^*, \times)$  is an abelian group:
  - $\times$  is associative and commutative,
  - $1$  is the identity element of  $\times$ , i.e.  $z \times 1 = z$  for all  $z \in \mathbb{C}^*$ ,

- Multiplicative inverses exist, i.e.  $z \times z^{-1} = 1$  for all  $z \in \mathbb{C}^*$ ;
- 3.  $\times$  is distributive over  $+$ .

The set  $\mathbb{C}$  of complex numbers can be identified with the real vector space  $\mathbb{R}^2$  by the vector space isomorphism:

$$\begin{aligned}\mathbb{C} &\rightarrow \mathbb{R}^2, \\ z &\mapsto (\operatorname{Re} z, \operatorname{Im} z), \\ x + iy &\mapsto (x, y).\end{aligned}$$

Unlike  $\mathbb{C}$ , the real plane  $\mathbb{R}^2$  is only equipped with addition operator  $+$  but not a natural multiplication operator  $\times$ . Nonetheless, the mapping above allows us to geometrically represent complex numbers as points on the plane. This is typically known as *Argand diagram*.

## 1.2 The Geometry of $\mathbb{C}$

Every complex number  $z = x + iy$  comes with a unique *real part*  $x$  and *imaginary part*  $y$ . We shall denote them as follows:

$$\operatorname{Re} z = x, \quad \operatorname{Im} z = y.$$

Geometrically,  $\operatorname{Re}$  and  $\operatorname{Im}$  can be thought as functions  $\mathbb{C} \rightarrow \mathbb{R}$  acting as projections onto the real and imaginary axes respectively.

The *complex conjugate*  $\bar{z}$  of a complex number  $z = x + iy$  is  $\bar{z} = x - iy$ . Geometrically, the operation  $z \mapsto \bar{z}$  is a reflection over the real axis. The following identity can be thought of as a change of basis from  $(x, y)$  to  $(z, \bar{z})$ .

**Proposition 1.1.** *For any  $z \in \mathbb{C}$ ,  $\operatorname{Re} z = \frac{z+\bar{z}}{2}$  and  $\operatorname{Im} z = \frac{z-\bar{z}}{2i}$ .*

Another straightforward algebraic exercise also gives us the following basic properties.

**Proposition 1.2.** *For any  $z, w \in \mathbb{C}$ ,  $\overline{z+w} = \bar{z} + \bar{w}$  and  $\overline{zw} = \bar{z}\bar{w}$ . If  $z \neq 0$ ,  $\overline{z^{-1}} = \bar{z}^{-1}$ .*

The *absolute value / modulus* of a complex number  $z = x + iy$  is

$$|z| = \sqrt{x^2 + y^2}.$$

Pythagoras' theorem indicates that geometrically the modulus  $|z|$  of  $z$  is equal to the distance between 0 and  $z$ .

**Proposition 1.3.** *For any  $z, w \in \mathbb{C}$ ,*

- $|zw| = |z||w|$ ,
- $z\bar{z} = |z|^2$ ,
- $|z + w| \leq |z| + |w|$  (*Triangle inequality*).

The *argument* of  $z$ ,  $\arg(z)$ , is defined to be the counterclockwise angle (measured in radians) subtended by the positive real axis  $\mathbb{R}^+$  and the line segment joining 0 and  $z$ . See figure 1.1.

Notice that  $\arg$  is a multivalued function. For example, both  $\pi$  and  $3\pi$  are arguments of  $i$ . We can refine this by defining the *principal argument* of  $z$ ,  $\text{Arg}(z)$ , to be the unique argument of  $z$  lying in  $(-\pi, \pi]$ .

*Remark.* The interval  $[0, 2\pi)$  is also often chosen to be the codomain of the principal argument.

**Proposition 1.4.** *For any  $z, w \in \mathbb{C}^*$ ,*

- $\arg(zw) = \arg(z) + \arg(w)$ ,
- $\text{Arg}(zw) = \text{Arg}(z) + \text{Arg}(w) \bmod 2\pi$ .

**Example 1.** Let  $z = 1 + i$  and  $w = -1 + \sqrt{3}i$ . The modulus and arguments of  $z$  and  $w$  are:

$$|z| = \sqrt{2}, \quad |w| = 2, \quad \arg(z) = \frac{\pi}{4}, \quad \arg w = \frac{2\pi}{3}.$$

Then, the modulus and argument of  $(1 + i)(-1 + \sqrt{3}i)$  are  $2\sqrt{2}$  and  $\frac{11\pi}{12}$  respectively.

For any non-zero complex number  $z = x + iy$ , if  $r = |z|$  and  $\theta = \text{Arg}(z)$ , then basic trigonometry gives us the following change of variables:

$$x = r \cos \theta, \quad y = r \sin \theta.$$

The expression  $z = r(\cos \theta + i \sin \theta)$  from above is the *polar form* of  $z$ .

**Theorem 1.5** (Euler's formula). *For any  $\theta$ ,  $e^{i\theta} = \cos \theta + i \sin \theta$ .*

*Proof.* We will give two different proofs of the result - one with differential equations, and another with Maclaurin series. The expression  $e^{i\theta}$  is a non-zero complex number, so there is a unique  $r > 0$  and  $\hat{\theta}$  such that

$$e^{i\theta} = r(\cos \hat{\theta} + i \sin \hat{\theta}). \tag{1.1}$$




Figure 1.1: A point  $z = x + iy = re^{i\theta}$  on the Argand diagram

Here,  $r$  and  $\hat{\theta}$  are functions of  $\theta$ . When  $\theta = 0$ ,  $r(0) = 1$  and  $\hat{\theta}(0) = 0$ . Differentiating (1.1) with respect to  $\theta$ , we obtain

$$\begin{aligned} ie^{i\theta} &= \frac{dr}{d\theta}(\cos \hat{\theta} + i \sin \hat{\theta}) + r \frac{d\hat{\theta}}{d\theta}(-\sin \hat{\theta} + i \cos \hat{\theta}) \\ &= \frac{dr}{d\theta} \frac{e^{i\theta}}{r} + i \frac{d\hat{\theta}}{d\theta} e^{i\theta} \\ &= \left( \frac{dr}{d\theta} + i \frac{d\hat{\theta}}{d\theta} \right) e^{i\theta}, \end{aligned}$$

where the second equality above is obtained from (1.1). From above, we see that  $\frac{dr}{d\theta} = 0$  and  $\frac{d\hat{\theta}}{d\theta} = 1$ . By our initial conditions, we obtain  $r(\theta) \equiv 1$  and  $\hat{\theta} \equiv \theta$ .

Alternatively, recall the following Maclaurin series:  $e^{i\theta} = \sum_{n=0}^{\infty} \frac{(i\theta)^n}{n!}$ . Using the fact that  $i^n = (-1)^{n/2}$  if  $n$  is even, and  $i^n = (-1)^{n-1/2}i$  if  $n$  is odd,

$$\begin{aligned} e^{i\theta} &= \sum_{\text{even } n} \frac{(-1)^n \theta^n}{n!} + \sum_{\text{odd } n} \frac{(-1)^{n-1/2} \theta^n}{n!} \\ &= \left( 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots \right) + i \left( \theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \dots \right) = \cos \theta + i \sin \theta. \end{aligned}$$

□

**Example 2.** When  $\theta = \pi$ , we have Euler's identity:  $e^{i\pi} = -1$ .

The polar form of a complex number  $z$  can alternatively be written in the form of  $z = re^{i\theta}$ . This expression is particularly useful when performing multiplication of complex numbers as we can use the laws of exponent. One particular instance is the following.

**Theorem 1.6** (De Moivre's Theorem). *For any  $\theta$  and integer  $n \in \mathbb{Z}$ ,*

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta).$$

**Example 3.** To compute and simplify  $\left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)^{10}$ , we can use De Moivre's theorem. The term inside the bracket is essentially  $\cos \theta + i \sin \theta$  where  $\theta = \frac{2\pi}{3}$ . Then,

$$\left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)^{10} = \cos\left(\frac{20\pi}{3}\right) + i \sin\left(\frac{20\pi}{3}\right) = -\frac{1}{2} + \frac{\sqrt{3}i}{2}.$$

## 1.3 Complex Roots

Consider a complex number  $z_0$  and a positive integer  $n$ . A complex number  $w$  satisfying  $w^n = z_0$  is called an  $n^{\text{th}}$  root of  $z_0$ .

Suppose  $z_0 = 0$ . Regardless of  $n$ , there is only one root of 0, which is 0 itself. This is due to the fact that  $\mathbb{C}$  is an integral domain, i.e. for any two complex numbers  $z_1$  and  $z_2$ , if  $z_1 z_2 = 0$  then either  $z_1 = 0$  or  $z_2 = 0$ .

Suppose  $z \neq 0$  now, then surely any root  $w$  is also non-zero. Using their polar forms  $z = re^{i\theta}$  and  $w = se^{it}$ , then the equation becomes:

$$s^n e^{int} = r e^{i\theta}$$


Considering the modulus and the argument independently, we obtain two real equations  $s^n = r$  and  $nt = \theta \bmod 2\pi$ . There are therefore  $n$  different solutions to  $w$ :

$$w_k = r^{1/n} e^{i(\theta+2\pi k)/n}, \quad k \in \{0, 1, \dots, n-1\}.$$

In the expression above,  $w_0$  is called the *principal root* of  $z_0$ . On the complex plane, these roots are evenly spaced on the circle  $\{z \in \mathbb{C} \mid |z| = r^{1/n}\}$  of radius  $r^{1/n}$  centered at the origin.

When  $z_0 = 1$ , the  $n^{\text{th}}$  roots of 1 are called the  $n^{\text{th}}$  roots of unity. They all lie on the unit circle and are of the form  $e^{2\pi ik/n}$ , where  $k \in \{0, 1, \dots, n-1\}$ .

**Example 4.** The 3rd roots of unity are  $1$ ,  $e^{2\pi i/3}$  and  $e^{4\pi i/3}$ . The Cartesian forms of these roots are  $1$ ,  $\frac{-1+i\sqrt{3}}{2}$ , and  $\frac{-1-i\sqrt{3}}{2}$ .

Figure 1.2: 3<sup>rd</sup> roots of unity

## 1.4 The Topology of $\mathbb{C}$

An *open disk* of radius  $r > 0$  centred at a complex number  $a \in \mathbb{C}$  is a subset of  $\mathbb{C}$  of the form:

$$\mathbb{D}(z, r) = \{z \in \mathbb{C} \mid |z - a| < r\}.$$

The boundary of this disk is a circle of radius  $r > 0$  centred at  $a$ , denoted with a partial sign in front:

$$C(z, r) = \partial\mathbb{D}(z, r) = \{z \in \mathbb{C} \mid |z - a| = r\}.$$

If we include the boundary, we obtain a *closed disk* typically denoted with an overline:

$$\overline{\mathbb{D}(z, r)} = \{z \in \mathbb{C} \mid |z - a| \leq r\}.$$

**Example 5.** Let's consider the sets

$$A = \{re^{i\theta} \mid r = \sin \theta, \theta \in \mathbb{R}\}, \quad B = \{re^{i\theta} \mid 0 < r < \sin \theta, \theta \in \mathbb{R}\}.$$

If  $z = x + iy$  lies in  $A$ , then  $x = \sin \theta \cos \theta$  and  $y = \sin^2 \theta$  for some  $\theta$ . By double angle formulas,

$$\sin^2(2\theta) + \cos^2(2\theta) = (2x)^2 + (1 - 2y)^2 = 1.$$

This equation represents a circle of radius  $\frac{1}{2}$  centered at  $\frac{i}{2}$ . Therefore,  $A$  is the circle  $C(\frac{i}{2}, \frac{1}{2})$ . For points  $z$  on the set  $B$ , we only need to consider the case when  $\sin \theta > 0$ , or principally when  $0 < \theta < \pi$ . The set  $B$  is the open disk  $\mathbb{D}(\frac{i}{2}, \frac{1}{2})$ .

The geometric and topological properties of the complex plane  $\mathbb{C}$  are essentially the same as those of the real plane  $\mathbb{R}^2$  since we have the obvious identification  $x + iy \mapsto (x, y)$ . We will give a brief introduction of necessary topological terminology that we will use in the next few chapters.

**Definition 1.** A subset  $S \subset \mathbb{C}$  is:

- *open* if for every  $s \in S$ , there is some  $r > 0$  such that  $\mathbb{D}(s, r) \subset S$ ,
- *closed* if its complement  $\mathbb{C} \setminus S$  is open,
- *bounded* if there is some  $r > 0$  where  $S \subset \mathbb{D}(0, r)$ ,
- *compact* if  $S$  is closed and bounded.

**Example 6.** Below are some subsets of  $\mathbb{C}$  which we will commonly encounter.

1. The empty set  $\emptyset$  is trivially open and compact.
2. The complex plane  $\mathbb{C}$  is both open and closed, but not bounded.
3. The punctured plane  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$  is open, but not closed nor bounded.
4. The unit disk  $\mathbb{D} := \mathbb{D}(0, 1)$  is open and bounded, but not closed.
5. The closed unit disk  $\overline{\mathbb{D}}$  and its boundary  $\partial\mathbb{D}$  are compact.
6. The real axis  $\mathbb{R}$  is closed and unbounded.

**Definition 2.** An open/closed set  $S \subset \mathbb{C}$  is:

- *connected* if  $S$  cannot be expressed as a disjoint union of two open/closed non-empty subsets of  $\mathbb{C}$ ,
- *simply connected* if it is connected and it has no "holes", i.e. the complement  $\mathbb{C} \setminus S$  has no bounded connected component,
- *multiply connected* if it is connected but not simply connected.

We say that  $S$  is a *domain* if it is a non-empty open connected subset of  $\mathbb{C}$ .




Figure 1.3: Four connected subsets of  $\mathbb{C}$ . Solid boundary lines are included in the colored set, whereas dashed boundary lines are not included. The first (from the left) is a simply connected domain. The second is closed and simply connected. The third is a punctured disk, which is a multiply connected domain. The last is closed and multiply connected.

**Example 7.**

1.  $\emptyset, \mathbb{C}, \overline{\mathbb{D}}$  and  $\mathbb{R}$  are simply connected.
2. The punctured plane  $\mathbb{C}^*$ , the punctured unit disk  $\mathbb{D}^* := \mathbb{D} \setminus \{0\}$ , and the unit circle  $\partial\mathbb{D}$  are multiply connected.
3. The annulus  $\{z \in \mathbb{C} \mid r < |z| < R\}$  of inner radius  $r$  and outer radius  $R$  is multiply connected.

**Example 8.** Consider the set  $S = \{z \in \mathbb{C} \mid |\operatorname{Im}(\frac{1}{z})| < 1\}$ . In polar form  $z = re^{i\theta}$ , the inequality becomes

$$|\operatorname{Im}(r^{-1}e^{-i\theta})| = |r^{-1}\sin(-\theta)| = r^{-1}|\sin \theta| < 1$$

Therefore,  $|\sin \theta| < r$ . Similar to Example 5, this represents all the complex numbers lying outside two closed disks  $\overline{\mathbb{D}(\pm \frac{i}{2}, \frac{1}{2})}$ . The set  $S$  is illustrated in Figure 1.4; it is open, unbounded, and multiply connected.




Figure 1.4: The set  $S$ .

**Short Quiz 1**

1. Simplify  $\frac{1+i}{i-1}$ .
2. Find the modulus of  $(3+4i)(-4+3i)$ .
3. Find the argument(s) of  $\arg(-1+i)$ .
4. Express  $2e^{-2\pi i/3}$  in the form of  $x+iy$ .
5. What are the 3<sup>th</sup> roots of  $8i$ ?
6. Find the value of  $(1+i)^6$ .
7. If  $z \neq 0$ , express  $\text{Im}(\frac{z}{z+\bar{z}})$  in terms of  $\theta = \text{Arg}(z)$ .

Consider the following subsets:

$$\begin{array}{ll} A = \mathbb{D}(2, 2) \cup \mathbb{D}(-2, 2), & B = \overline{\mathbb{D}(i, 1)} \cup \overline{\mathbb{D}(-i, 1)}, \\ C = \mathbb{D}(2, 1) \cup \mathbb{D}(-2, 1), & D = C(i, 1) \cup \overline{\mathbb{D}(-i, 1)}. \end{array}$$

8. Identify subsets which are open.
9. Identify subsets which are connected.
10. Identify subsets which are simply connected.

Answers: 1.  $-i$ , 2. 25, 3.  $\frac{3\pi}{4} + 2\pi k$ , 4.  $-1 - i\sqrt{3}$ , 5.  $\pm\sqrt{3} + i$  &  $-2i$ , 6.  $-8i$ , 7.  $\frac{1}{2}\tan\theta$ , 8.  $A$  and  $C$ , 9.  $B$  and  $D$ , 10.  $B$ .



# Chapter 2

## Complex Functions

### 2.1 Convergence and Continuity

**Definition 3.** A sequence of complex numbers  $\{z_n\}_{n \in \mathbb{N}}$  converges to a limit  $z$  if and only if:

for all  $\epsilon > 0$ , there exists  $N > 0$  such that for all  $n \geq N$ ,  $|z_n - z| < \epsilon$ .

Convergence of a sequence  $z_n$  to  $z$  can be denoted by  $z_n \rightarrow z$ ,  $|z_n - z| \rightarrow 0$ , or  $\lim_{n \rightarrow \infty} z_n = z$ .

**Proposition 2.1.** *The limit of a convergent sequence is unique.*

*Proof.* Suppose for a contradiction that there are distinct limits  $z \neq w$  of a sequence  $z_n$ . Let  $\epsilon = \frac{1}{2}|z - w| > 0$ , then for all sufficiently high  $n$ ,  $|z_n - z| < \epsilon$  and  $|z_n - w| < \epsilon$ . However, by triangle inequality,

$$|z - w| \leq |z - z_n| + |z_n - w| < 2\epsilon = |z - w|.$$

We then have a contradiction. □

**Theorem 2.2.** *If  $z_n \rightarrow z$  and  $w_n \rightarrow w$ , then*

- $z_n + w_n \rightarrow z + w$ ,
- $z_n w_n \rightarrow zw$ .

*Proof.* Let's pick  $\epsilon > 0$ . There are some high  $N_1, N_2 \in \mathbb{N}$  such that  $|z_n - z| < \epsilon/2$  if  $n \geq N_1$ , and  $|w_n - w| < \epsilon/2$  if  $n \geq N_2$ . By triangle inequality, when  $n \geq \max\{N_1, N_2\}$ ,

$$|z_n + w_n - z - w| \leq |z_n - z| + |w_n - w| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

This shows that  $z_n + w_n \rightarrow z + w$ .

Set  $M = \max\{|w_1|, \dots, |w_{N_2}|, |w| + \epsilon\}$ . The sequence  $\{w_n\}_{n \in \mathbb{N}}$  is bounded because we have the following inclusions:

$$\{w_n\}_{n \in \mathbb{N}} \subset \{w_1, \dots, w_{N_2}\} \cup \mathbb{D}(w, \epsilon/2) \subset \mathbb{D}(0, M).$$

There are some  $N_3, N_4 \in \mathbb{N}$  such that  $|z_n - z| < \epsilon/2M$  if  $n \geq N_3$ , and  $|w_n - w| < \epsilon/2 \max\{1, |z|\}$  if  $n \geq N_4$ . Then, when  $n \geq \max\{N_3, N_4\}$ ,

$$\begin{aligned} |z_n w_n - zw| &\leq |z_n w_n - z w_n| + |z w_n - zw| = |w_n| |z_n - z| + |z| |w_n - w| \\ &< M \cdot \frac{\epsilon}{2M} + |z| \cdot \frac{\epsilon}{2 \max\{1, |z|\}} \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

This shows that  $z_n w_n \rightarrow zw$ . □

In particular, a sequence of complex numbers converges exactly when the real parts and the imaginary parts converge respectively.

**Corollary 2.3.**  $x_n + iy_n \rightarrow x + iy$  if and only if  $x_n \rightarrow x$  and  $y_n \rightarrow y$ .

*Proof.* The  $\Leftarrow$  direction is immediate from the previous proposition. The  $\Rightarrow$  direction comes from the following inequality:

$$\max\{|x_n - x|, |y_n - y|\} \leq \sqrt{|x_n - x|^2 + |y_n - y|^2} = |x_n + iy_n - x - iy|.$$

As  $|x_n + iy_n - x - iy| \rightarrow 0$ , sandwich rule forces both  $|x_n - x|$  and  $|y_n - y|$  to converge to 0 too. □

**Definition 4.** Let  $U$  and  $V$  be non-empty subsets of  $\mathbb{C}$ . A function  $f : U \rightarrow V$  is *continuous at  $a \in U$*  if and only if:

$$\begin{aligned} &\text{for all } \epsilon > 0, \text{ there exists } \delta > 0 \text{ such that} \\ &\text{if } z \in U \cap \mathbb{D}(a, \delta), \text{ then } f(z) \in \mathbb{D}(f(a), \epsilon). \end{aligned}$$

We say that  $f$  is *continuous* if it is continuous at every point in  $U$ .

**Proposition 2.4.** A function  $f : U \rightarrow V$  is continuous at  $a \in U$  if and only if for any sequence  $z_n$  in  $U$ , if  $z_n \rightarrow a$ , then  $f(z_n) \rightarrow f(a)$ .

*Proof.* Let  $f$  be continuous at  $a$  and pick the pair  $(\epsilon, \delta)$  in the definition of continuity. Suppose  $z_n \rightarrow a$ , then there is some high  $N \in \mathbb{N}$  such that if  $n \geq N$ ,  $|z_n - a| < \delta$ . By continuity, if  $n \geq N$ ,  $|f(z_n) - f(a)| < \epsilon$ . Therefore,  $f(z_n) \rightarrow f(a)$ .

Suppose for any sequence  $z_n$  converging to  $a$ ,  $f(z_n) \rightarrow f(a)$ . Suppose for a contradiction that  $f$  is not continuous at  $a$ , then there is some  $\epsilon > 0$  and sequence of points  $z_n \in U \cap \mathbb{D}(a, \frac{1}{n})$  for  $n \in \mathbb{N}$  such that  $|f(z_n) - f(a)| \geq \epsilon$ . Since  $|z_n - a| < \frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , then  $z_n \rightarrow a$ . The assumption implies that  $f(z_n) \rightarrow f(a)$ , but this cannot happen because  $f(z_n)$  is always at least  $\epsilon$  away from  $f(a)$ . This gives the contradiction. □

The statement above gives an equivalent way of defining continuity at a point. A shorter way of saying this is

$$\lim_{z \rightarrow a} f(z) = f(a).$$

The following is a direct consequence of 2.2.

**Proposition 2.5.** *Let  $f, g : U \rightarrow V$  and  $h : V \rightarrow W$  be continuous functions, then the sum  $f + g$ , the product  $f \cdot g$  and the composition  $h \circ f$  are continuous.*

**Example 9.** Constant functions  $f(z) = a$  are trivially continuous. The identity function  $\text{Id}(z) = z$  is also continuous. By taking products and sums, we can inductively obtain that every complex polynomial  $a_d z^d + a_{d-1} z^{d-1} + \dots + a_1 z + a_0$  is continuous.

**Example 10.** The modulus function  $m(z) = |z|$  is a continuous function on  $\mathbb{C}$ . Indeed, for any  $a \in \mathbb{C}$ , if  $z \rightarrow a$ , then by triangle inequality,

$$|m(z) - m(a)| \leq ||z| - |a|| \leq |z - a| \rightarrow 0.$$

By sandwich rule,  $m(z) \rightarrow m(a)$  too.

By the previous proposition, if  $f(z)$  is a continuous function on a subset of  $\mathbb{C}$ , then so is the composition  $|f(z)|$ .

**Example 11.** The functions  $\text{Re}$  and  $\text{Im}$  are continuous. (Refer to Corollary 2.3.) Since  $\bar{z} = \text{Re}(z) - i \text{Im}(z)$ , complex conjugation is also continuous.

Continuous functions behave nicely on compact subsets of  $\mathbb{C}$ .

**Theorem 2.6.** *Let  $f : K \rightarrow V$  be a continuous function and  $K$  be a compact subset of  $\mathbb{C}$ . Then,  $f$  attains a maximum and a minimum on  $K$ , i.e. there are points  $a, b \in K$  where  $|f(a)| \leq |f(z)| \leq |f(b)|$  for all  $z \in K$ .*

The theorem above is a consequence of a result from topology. In particular, the image  $|f(K)|$  of a compact set  $K$  under a continuous function  $|f|$  is compact, and any compact subset of  $\mathbb{R}$  contains maximum and minimum points.

## 2.2 Holomorphic Functions

We can define differentiability of complex-valued functions the same way as we define that of functions of one real variable. However, we will emphasise in the next few sections that complex differentiability is actually a much more rigid notion than the usual multivariable real differentiability.

**Definition 5.** Let  $U, V \subset \mathbb{C}$  be open and non-empty. A complex function  $f$  is (*complex*) *differentiable* at a point  $a$  if and only if the following limit exists:

$$f'(a) = \lim_{z \rightarrow a} \frac{f(z) - f(a)}{z - a}$$

If so, then  $f'(a)$  is called the (*complex*) *derivative* of  $f$  at  $a$ . The function  $f$  is said to be *holomorphic* on  $U$  if it is holomorphic at every point in  $U$ , and *entire* if additionally  $U = \mathbb{C}$ .

*Remark.* The term "analytic" and "complex differentiable" are often used interchangeably with "holomorphic".

**Example 12.**

1. Constant functions  $f(z) = a$  are entire with derivative 0 everywhere.
2. The identity function  $\text{Id}(z) = z$  is an entire function and its derivative is 1 everywhere.
3. The inversion function  $\tau(z) = 1/z$  is holomorphic on  $\mathbb{C}^*$  and its derivative is  $-z^{-2}$ . Indeed, if we choose any angle  $\theta$ , then if  $z = re^{i\theta} + a$ ,

$$\begin{aligned} \lim_{z \rightarrow a} \frac{\tau(z) - \tau(a)}{z - a} &= \lim_{r \rightarrow 0} \frac{\frac{1}{re^{i\theta} + a} - \frac{1}{a}}{(re^{i\theta} + a) - a} = \lim_{r \rightarrow 0} \frac{\frac{-re^{i\theta}}{a(re^{i\theta} + a)}}{re^{i\theta}} \\ &= \lim_{r \rightarrow 0} -\frac{1}{a(re^{i\theta} + a)} = -\frac{1}{a^2}. \end{aligned}$$

4. Complex conjugation  $f(z) = \bar{z}$  has no derivative at any point. If we choose any angle  $\theta$ , then using  $z = re^{i\theta} + a$ ,

$$\lim_{z \rightarrow a} \frac{f(z) - f(a)}{z - a} = \lim_{r \rightarrow 0} \frac{(re^{-i\theta} + \bar{a}) - \bar{a}}{(re^{i\theta} + a) - a} = e^{-2i\theta},$$

but the value of this limit is not the same if we choose different values of  $\theta$ . For example, the limit is 1 when  $\theta = 0$ , but it is  $-1$  if  $\theta = \frac{\pi}{2}$ .

**Proposition 2.7.** *Every holomorphic function is continuous.*

*Proof.* Let  $f : U \rightarrow V$  be holomorphic. If  $a \in U$ ,

$$\begin{aligned} \lim_{z \rightarrow a} f(z) - f(a) &= \lim_{z \rightarrow a} (z - a) \frac{f(z) - f(a)}{z - a} \\ &= \lim_{z \rightarrow a} (z - a) \cdot \lim_{z \rightarrow a} \frac{f(z) - f(a)}{z - a} \\ &= 0 \cdot f'(a) = 0, \end{aligned}$$

where the second equality follows from Theorem 2.2. Therefore,  $f$  is continuous at  $a$ . As  $a$  is arbitrary,  $f$  is continuous on  $U$ .  $\square$

The rules for differentiation of complex-valued functions is more or less the same as those of functions of one real variable.

**Proposition 2.8.** *Let  $f, g : U \rightarrow V$  and  $h : V \rightarrow W$  be holomorphic. Then,*

- (a) *the sum  $f + g$  is holomorphic and  $(f + g)'(z) = f'(z) + g'(z)$ ,*
- 1. *the product  $f \cdot g$  is holomorphic and  $(f \cdot g)'(z) = f'(z)g(z) + f(z)g'(z)$ ,*
- 2. *the composition  $h \circ f$  is holomorphic and  $(h \circ f)'(z) = h'(f(z))f'(z)$ .*

*Proof.* (a) follows immediately from Proposition 2.5. For (b),

$$\begin{aligned} (f \cdot g)'(a) &= \lim_{z \rightarrow a} \frac{f(z)g(z) - f(a)g(a)}{z - a} \\ &= \lim_{z \rightarrow a} \frac{f(z)(g(z) - g(a))}{z - a} + \frac{g(a)(f(z) - f(a))}{z - a} \\ &= f(a)g'(a) + g(a)f'(a). \end{aligned}$$

For (c), we use the fact that  $f$  is continuous:

$$\begin{aligned} (h \circ f)'(a) &= \lim_{z \rightarrow a} \frac{h(f(z)) - h(f(a))}{z - a} \\ &= \lim_{z \rightarrow a} \frac{h(f(z)) - h(f(a))}{f(z) - f(a)} \cdot \frac{f(z) - f(a)}{z - a} \\ &= \lim_{w \rightarrow f(a)} \frac{h(w) - h(f(a))}{w - f(a)} \cdot \lim_{z \rightarrow a} \frac{f(z) - f(a)}{z - a} \\ &= h'(f(a))f'(a). \end{aligned} \quad \square$$

### Example 13.

1. Every polynomial  $p(z) = \sum_{n=0}^d a_n z^n$  with complex coefficients  $a_n \in \mathbb{C}$  is an entire function. We can show this inductively by product rule above that every monomial  $a_n z^n$  is holomorphic with derivative  $a_n n z^{n-1}$  and by the addition rule,  $p$  is holomorphic.
2. Every rational function, i.e. a function of the form  $f(z) = p(z)/q(z)$  where  $p$  and  $q$  are polynomials, is holomorphic on  $\mathbb{C} \setminus \{z \in \mathbb{C} | q(z) = 0\}$ .

Every complex function  $f(z)$  admits a unique real-imaginary splitting  $f(x+iy) = u(x, y) + iv(x, y)$ , where  $u$  and  $v$  are real-valued functions defined on an open subset of  $\mathbb{R}^2$  given by:

$$u(x, y) = \operatorname{Re} f(x + iy), \quad v(x, y) = \operatorname{Im} f(x + iy).$$

We say that  $f$  is (*real*) *differentiable* if the partial derivatives of  $u$  and  $v$  with respect to  $x$  and  $y$  exist. When we are given  $u$  and  $v$ , we will see that holomorphic functions are precisely solutions of a system of partial differential equations.

**Theorem 2.9** (Cauchy-Riemann Equations). *Let  $f = u + iv$  be a complex function on an open set  $U \subset \mathbb{C}$ . Then,  $f$  is holomorphic if and only if  $u$  and  $v$  are continuously differentiable and  $u_x = v_y$  and  $v_x = -u_y$ .*

*Proof.* Let  $f$  be holomorphic at a point  $a = a_1 + ia_2 \in U$ , then

$$f'(a) = \lim_{h \rightarrow 0} \frac{u(a_1 + h_1, a_2 + h_2) - u(a_1, a_2)}{h} + i \frac{v(a_1 + h_1, a_2 + h_2) - v(a_1, a_2)}{h}.$$

The dummy variable  $h = h_1 + ih_2$  can converge to zero in various directions, but we will only consider two cases. Suppose  $h_2 = 0$ , then

$$\begin{aligned} f'(a) &= \lim_{h_1 \rightarrow 0} \frac{u(a_1 + h_1, a_2) - u(a_1, a_2)}{h_1} + i \frac{v(a_1 + h_1, a_2) - v(a_1, a_2)}{h_1} \\ &= u_x + iv_x. \end{aligned} \tag{2.1}$$

Similarly, if we consider the limit in the direction satisfying  $h_1 = 0$ ,

$$\begin{aligned} f'(a) &= \lim_{ih_2 \rightarrow 0} \frac{u(a_1, a_2 + ih_2) - u(a_1, a_2)}{ih_2} + i \frac{v(a_1, a_2 + ih_2) - v(a_1, a_2)}{ih_2} \\ &= -iu_y + v_y. \end{aligned} \tag{2.2}$$

Comparing (2.1) and (2.2), it is clear by taking the real and imaginary parts of  $f'(a)$  that the partial derivatives of  $u$  and  $v$  at  $a$  exist and satisfy  $u_x = v_y$  and  $v_x = -u_y$ . To show that these partial derivatives are continuous, we need continuity of  $f'$ . We will obtain this for granted in Corollary 4.4.

Conversely, suppose  $u$  and  $v$  are continuously differentiable satisfying  $u_x = v_y$  and  $v_x = -u_y$ . The Taylor series of  $u$  and  $v$  at  $a$  can be expressed as:

$$\begin{aligned} u(a_1 + h_1, a_2 + h_2) &= u(a_1, a_2) + u_x h_1 + u_y h_2 + |h|\psi(h), \\ v(a_1 + h_1, a_2 + h_2) &= v(a_1, a_2) + v_x h_1 + v_y h_2 + |h|\phi(h), \end{aligned}$$

for some functions  $\psi$  and  $\phi$  where  $\psi(h), \phi(h) \rightarrow 0$  as  $h \rightarrow 0$ . All partial derivatives of  $u$  and  $v$  are evaluated at  $(a_1, a_2)$ . Then,

$$\begin{aligned}
& \lim_{h \rightarrow 0} \frac{u(a_1 + h_1, a_2 + h_2) - u(a_1, a_2)}{h} + i \frac{v(a_1 + h_1, a_2 + h_2) - v(a_1, a_2)}{h} \\
&= \lim_{h \rightarrow 0} \frac{[u_x h_1 + u_y h_2 + |h|\psi(h)] + i[v_x h_1 + v_y h_2 + |h|\phi(h)]}{h} \\
&= \lim_{h \rightarrow 0} \frac{(u_x + iv_x)(h_1 + ih_2) + |h|(\psi(h) + i\phi(h))}{h} \\
&= u_x + iv_x.
\end{aligned}$$

As the limit converges to  $f' = u_x + iv_x$ ,  $f$  is indeed holomorphic.  $\square$

It is natural to consider a change of variables from  $(x, y)$  to  $(z, \bar{z}) = (x+iy, x-iy)$ . By using multivariable chain rule, we can obtain an expression for partial derivatives  $\frac{\partial}{\partial z}$  and  $\frac{\partial}{\partial \bar{z}}$  in terms of  $\frac{\partial}{\partial x}$  and  $\frac{\partial}{\partial y}$ .

**Definition 6.** The *Wirtinger derivatives* are defined as follows:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$$

It is straightforward to check that the following identities hold:

$$\frac{\partial z}{\partial z} = \frac{\partial \bar{z}}{\partial \bar{z}} = 1, \quad \frac{\partial \bar{z}}{\partial z} = \frac{\partial z}{\partial \bar{z}} = 0.$$

Given a complex function  $f = u + iv$  with differentiable real and imaginary parts  $u$  and  $v$ ,

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2}(f_x + if_y) = \frac{1}{2}((u_x - v_y) + i(v_x + u_y))$$

Clearly, the Cauchy-Riemann equations hold if and only if the expression above is 0. The equation

$$\frac{\partial f}{\partial z} = 0$$

is the complex form of the Cauchy-Riemann equations, as it allows us to deduce holomorphicity without having to know the real and imaginary parts  $u$  and  $v$ , but rather by the absence of the variable  $\bar{z}$ . Roughly speaking, if  $f$  is not a function of  $\bar{z}$ , then it is holomorphic!

If  $f$  is holomorphic, the notation for the complex derivative of  $f$  is also consistent since

$$\frac{\partial f}{\partial z} = \frac{1}{2}(f_x - if_y) = \frac{1}{2}((u_x + v_y) + i(v_x - u_y)) = u_x + iv_x = f',$$

where the last inequality follows from (2.1).

**Example 14.** Re and Im are differentiable but not holomorphic. (Refer to Proposition 1.1.)

## 2.3 Exponential and Logarithmic Functions

One of the many elementary functions we will commonly encounter is the exponential function

$$\exp(z) := e^z = e^x e^{iy}.$$

The real and imaginary parts are  $u = e^x \cos(y)$  and  $v = e^x \sin(y)$ , and we can easily check that the Cauchy-Riemann equations are satisfied. Thus, this is an entire function and its derivative is itself. Below are some of its properties:

1.  $|\exp(z)| = e^x$ ,
2.  $\arg(\exp(z)) = y + 2\pi ik$  where  $k \in \mathbb{Z}$ ,
3.  $\exp(z + 2\pi ik) = \exp(z)$  for any  $k \in \mathbb{Z}$ ,
4. The image of  $\exp$  on  $\mathbb{C}$  is  $\mathbb{C}^*$ .

We would now like to find the inverse of  $\exp$ . If we denote the inverse by  $\log$ , then

$$\log(z) := \ln|z| + i\arg(z)$$

for any  $z \neq 0$ . However, as  $\arg$  is multivalued,  $\log$  is multi-valued and therefore it is not a well-defined function. This is consistent with the fact that  $\exp$  is not injective due to property 3.

This problem can be fixed by using the principle argument  $\text{Arg}$ . Doing so will introduce a ray of discontinuity  $(-\infty, 0]$  corresponding to the points in  $\mathbb{C}$  with argument  $\pi$ . We can replace  $\arg$  with  $\text{Arg}$  and define the *principal value* of  $\log(z)$ , denoted by  $\text{pv } \log(z)$ . If we choose the codomain to be  $\text{Arg}(z) \in (-\pi, \pi]$ , we then have the *principal branch* of the logarithmic function

$$\text{Log} : \mathbb{C} \setminus (-\infty, 0] \rightarrow \mathbb{C}, \quad \text{Log}(z) := \ln|z| + i\text{Arg}(z).$$

The ray  $(-\infty, 0]$  is often called a *branch cut*. Using this choice of branch cut,  $\text{Log}$  is holomorphic on  $\mathbb{C} \setminus (-\infty, 0]$ , with image  $\{x + iy \mid x \in \mathbb{R}, -\pi < y < \pi\}$  and derivative  $\frac{1}{z}$ . (To verify this, compute the Wirtinger derivatives in polar coordinates.)

**Example 15.**  $\log(i)$  is purely imaginary and multivalued since

$$\log(i) = \ln(1) + i\arg(i) = \frac{\pi i}{2} + 2\pi ik, \text{ where } k \in \mathbb{Z}.$$

Its principal value is  $\text{Log}(i) = \frac{\pi i}{2}$ .




Figure 2.1: Various branch cuts of the logarithm

*Remark.* In general, the branch cut can be taken to be any unbounded curve from 0 which does not intersect itself. Straight rays of different angles are often used if necessary.

For a non-integer  $c \in \mathbb{C}$ , we define the *power function*  $z^c$  to be the multivalued function on  $\mathbb{C}^*$  given by

$$z^c := \exp(c \log(z)).$$

Similar to the logarithm, we can take the principal value of  $z^c$  to be

$$\text{pv } z^c := \exp(c \text{Log}(z)).$$

Again, this becomes a holomorphic function outside a chosen branch cut  $(-\infty, 0]$ .

**Example 16.**  $i^i$  is (perhaps surprisingly) real and multivalued since

$$i^i = e^{-\pi/2 - 2\pi k}, \text{ where } k \in \mathbb{Z}.$$

Its principal value is  $\text{pv } i^i = e^{-\pi/2}$ .

## 2.4 Trigonometric Functions

Euler's formula allows us to express  $\sin \theta$  and  $\cos \theta$  in terms of  $e^{\pm i\theta}$  as follows:

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}.$$

It is then natural to define trigonometric functions of a complex variable  $z$  using the exponential function:

$$\cos(z) := \frac{e^{iz} + e^{-iz}}{2}, \quad \sin(z) := \frac{e^{iz} - e^{-iz}}{2i}, \quad \tan(z) = \frac{\sin(z)}{\cos(z)}.$$

As  $\exp$  is entire, so are  $\cos$  and  $\sin$ . However, the function  $\tan$  is holomorphic everywhere except at points  $z$  such that  $\cos(z) = 0$ . You may check that the usual trigonometric identities still hold.

The generalisation of hyperbolic functions are also clear:

$$\cosh(z) := \frac{e^z + e^{-z}}{2}, \quad \sinh(z) := \frac{e^z - e^{-z}}{2}, \quad \tanh(z) = \frac{\sinh(z)}{\cosh(z)}.$$

The functions  $\cosh$  and  $\sinh$  can be viewed as the even part and the odd part of the exponential function respectively. Both are entire functions, but  $\tanh$  is only holomorphic everywhere except at points  $z$  such that  $\cosh(z) = 0$ . The following property can be easily deduced by definition.

**Proposition 2.10.** *For any  $z \in \mathbb{C}$ ,  $\cos(iz) = \cosh(z)$  and  $\sin(iz) = i \sinh(z)$ .*

A point  $w$  is a *zero* of a function  $f$  if  $f(w) = 0$ . It turns out that the zeros of trigonometric functions in  $\mathbb{C}$  are the same as the zeros of trigonometric functions in  $\mathbb{R}$ .

**Proposition 2.11.** *The zeros of trigonometric and hyperbolic functions are as follows:*

$$\begin{aligned} \{z \in \mathbb{C} \mid \sin(z) = 0\} &= \{n\pi\}_{n \in \mathbb{Z}}, \\ \{z \in \mathbb{C} \mid \cos(z) = 0\} &= \left\{ \frac{\pi}{2} + n\pi \right\}_{n \in \mathbb{Z}}, \\ \{z \in \mathbb{C} \mid \sinh(z) = 0\} &= \{in\pi\}_{n \in \mathbb{Z}}, \\ \{z \in \mathbb{C} \mid \cosh(z) = 0\} &= \left\{ i \left( \frac{\pi}{2} + n\pi \right) \right\}_{n \in \mathbb{Z}}. \end{aligned}$$

*Proof.* By addition rule and Proposition 2.10,  $\sin(x + iy) = \sin x \cosh y + i \sinh y \cos x$ . Thus, if  $z = x + iy$ ,

$$\begin{aligned} |\sin(z)|^2 &= \sin^2 x \cosh^2 y + \sinh^2 y \cos^2 x \\ &= \sin^2 x (1 + \sinh^2 y) + \sinh^2 y (1 - \sin^2 x) \\ &= \sin^2 x + \sinh^2 y. \end{aligned}$$

As such,  $\sin(z) = 0$  if and only if  $\sin x = 0$  and  $\sinh y = 0$ , and the latter occurs exactly when  $x \in \{n\pi\}_{n \in \mathbb{Z}}$ , and  $y = 0$ .

The zeros of  $\cos$  can be obtained from those of  $\sin$  using the identity  $\cos(z) = \sin(\pi/2 - z)$ , and the zeros of hyperbolic functions can be obtained from those of trigonometric functions by Proposition 2.10.  $\square$

Tgonometric and hyperbolic functions are definitely not surjective, but we are still able to find their local inverses on some restricted domains. Since they are made up of the exponential function, their inverses can be described in terms of logarithms.

**Proposition 2.12.** *The inverses of trigonometric and hyperbolic functions are multivalued functions:*

$$\begin{aligned}\sin^{-1}(z) &= -i \log(iz + [1 - z^2]^{1/2}), \\ \cos^{-1}(z) &= -i \log(z + i[1 - z^2]^{1/2}), \\ \tan^{-1}(z) &= \frac{i}{2} \log \frac{i+z}{i-z}, \\ \sinh^{-1}(z) &= \log(z + [z^2 + 1]^{1/2}), \\ \cosh^{-1}(z) &= \log(z + [z^2 - 1]^{1/2}), \\ \tanh^{-1}(z) &= \frac{1}{2} \log \frac{1+z}{1-z}.\end{aligned}$$

*Proof.* Let  $\sin^{-1}(z) = w$ , then  $z = \frac{e^{iw} - e^{-iw}}{2i}$ . This can be written in quadratic form:

$$(e^{iw})^2 - 2ize^{iw} - 1 = 0.$$

The quadratic formula gives us  $e^{iw} = iz + [1 - z^2]^{1/2}$ , and using logarithm,

$$w = -i \log(iz + [1 - z^2]^{1/2}).$$

The resulting function is multivalued since the square root and the logarithm are multivalued. Similar algebraic methods can be applied to obtain the inverses of other functions and will be left to the reader as an exercise.  $\square$

For each of the functions above, we can pick a branch cut of the square root and the logarithm in order to obtain a holomorphic branch of the function. However, finding a nice branch cut for the inverse of the function requires a more involved argument and we shall not attempt to find it.

**Example 17.** Let's find the solution for the equation  $\sin(2\pi z) = 2$ . By the proposition above,

$$\begin{aligned}z &= -\frac{i}{2\pi} \log(2i + (-3)^{1/2}) = -\frac{i}{2\pi} \log(i(2 \pm \sqrt{3})) \\ &= -\frac{i}{2\pi} \left[ \ln(2 \pm \sqrt{3}) + \frac{\pi i}{2} + 2\pi ik \right], \\ &= -\frac{i \ln(2 \pm \sqrt{3})}{2\pi} + \frac{1}{4} + k, \quad k \in \mathbb{Z},\end{aligned}$$

### Short Quiz 2

1. Does the sequence  $i^n$  converge as  $n \rightarrow \infty$ ? If so, what is the limit?
2. Does the sequence  $\left(\frac{i}{n}\right)^n$  converge as  $n \rightarrow \infty$ ? If so, what is the limit?
3. Find the limit of the sequence  $\frac{1}{n} + \left(1 + \frac{i}{n}\right)^n$ .
4. Which of the following functions are continuous on  $\mathbb{C}$ ?

$$z^3, \quad 1/z, \quad |z - 2| + |z + 2|, \quad \arg(z)$$

5. At which of the following points is  $\operatorname{Arg}(z)$  continuous?

$$1, \quad i, \quad -1, \quad -i, \quad 0$$

6. What are the solutions of the equation  $e^z = -\pi$ ?
7. What is the value of  $(i^i)^i$ ?
8. What are the solutions of the equation  $\tanh(iz) = 0$ ?
9. Which of these functions are holomorphic on a domain containing  $2\pi i$ ?

$$\sin z, \quad \sinh z, \quad \csc z, \quad \operatorname{csch} z, \quad \cot z, \quad \coth z$$

10. Which of these complex numbers correspond to  $\log(1)$ ?

$$0, \quad 1, \quad 2\pi i, \quad 1 + 2\pi i, \quad -2\pi i, \quad 1 - 2\pi i.$$

Answers: 1. No, 2. Yes, to 0, 3.  $e^i$ , 4.  $z^3$  and  $|z - 2| + |z + 2|$ , 5. 1,  $i$ ,  $-i$ .  
 6.  $\ln \pi + i(\pi + 2\pi ik)$ . 7.  $-i$ . 8.  $k\pi$ . 9.  $\sin z, \sinh z, \csc z, \cot z$ . 10. 0 and  $\pm 2\pi i$ .

# Chapter 3

## Contour Integration

### 3.1 Curves in $\mathbb{C}$

**Definition 7.** An *arc / curve / path* in  $\mathbb{C}$  is subset of  $\mathbb{C}$  parametrized by a continuous function  $\gamma : [a, b] \rightarrow \mathbb{C}$  defined on a closed interval  $[a, b] \subset \mathbb{R}$ .

A curve  $\gamma : [a, b] \rightarrow \mathbb{C}$ , can be expressed as  $\gamma(t) = u(t) + iv(t)$  where  $u$  and  $v$  are real-valued functions. We say that  $\gamma$  is *differentiable* when both  $u$  and  $v$  are differentiable on  $[a, b]$  as real functions. The derivative of  $\gamma$  at  $t$  is

$$\gamma'(t) = u'(t) + iv'(t).$$

For each  $t$ , when  $\gamma'(t) \neq 0$ ,  $\gamma'(t)$  represents a tangent vector of the curve at the point  $\gamma(t)$  of magnitude  $|\gamma'(t)| = \sqrt{u'(t)^2 + v'(t)^2}$ .

**Definition 8.** A curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  is:

- *closed* if  $\gamma(a) = \gamma(b)$ ,
- *simple* if  $\gamma$  is injective on the open interval  $(a, b)$ ,
- *smooth* if  $\gamma$  is differentiable and  $\gamma'(t) \neq 0$  for all  $t \in (a, b)$ ,
- a *contour* if  $\gamma$  is piecewise smooth, i.e.  $\gamma$  can be partitioned into finitely many smooth curves.

**Example 18.** A circle  $C(z, r)$  of radius  $r > 0$  centered at  $w \in \mathbb{C}$  is a simple closed smooth curve. It can be parametrised by  $\gamma(t) = z_0 + re^{it}$ ,  $0 \leq t \leq 2\pi$ .

**Example 19.** The function  $\sigma(t) = e^{it} \sin(2t)$ ,  $0 \leq t \leq 2\pi$ , parametrises the locus of the equation  $r = \sin(2\theta)$ . This curve is a closed and smooth but clearly not simple. See the leftmost curve in Figure 3.1.




Figure 3.1: A closed non-simple smooth curve, a non-closed non-piecewise-smooth curve (Devil's Staircase), and a closed piecewise smooth curve.

We will state without proof two results from topology.

**Proposition 3.1.** *Any domain  $U \subset \mathbb{C}$  is path-connected. That is, for any two points  $z$  and  $w$  in a domain  $U \subset \mathbb{C}$ , there is a smooth curve  $\gamma : [0, 1] \rightarrow U$  such that  $\gamma(0) = z$  and  $\gamma(1) = w$ .*

**Theorem 3.2** (Jordan Curve Theorem). *The complement of any simple closed curve in  $\mathbb{C}$  has exactly two connected components, exactly one of which is bounded.*




Figure 3.2: A simple closed curve splits its complement into a bounded domain  $U$  and an unbounded domain  $V$ .

*Remark.* Due to this theorem, simple closed curves are often called *Jordan curves*. You may think that the theorem seems very intuitive, but the proof is rather involved. The very first proof was out in 1910s (not by Camille Jordan), relying on heavy machineries such as the theory of homology groups

in algebraic topology. The shortest proof I know is by Maehara (1984), relying only on basic knowledge of topology, including the Brouwer fixed point theorem.

**Definition 9.** A point  $w$  or a subset  $S$  of  $\mathbb{C}$  is said to be *enclosed* by a simple closed curve  $\gamma$  if  $\{w\}$  or  $S$  is contained in the bounded connected component of the complement of  $\gamma$  in  $\mathbb{C}$ .

Every curve  $\gamma$  has two possible orientations. Given a curve  $\gamma : [a, b] \rightarrow \mathbb{C}$ , we can reverse its orientation to obtain  $\gamma^- : [a, b] \rightarrow \mathbb{C}$  defined by  $\gamma^-(t) = \gamma(a + b - t)$ .




Figure 3.3: Reversing the orientation of  $\gamma$ .

When  $\gamma$  is a simple closed curve, the orientation of  $\gamma$  is *positive* if for any point  $w$  enclosed by  $\gamma$ ,  $\gamma(t)$  runs anticlockwise with respect to the basepoint  $w$  as  $t$  increases. The orientation is *negative* if  $\gamma(t)$  runs clockwise. Unless stated otherwise, we always assume that every simple closed curve  $\gamma$  is positively oriented, and  $\gamma^-$  is negatively oriented.

## 3.2 Integration Along a Contour

Any closed interval  $[a, b]$  can be parametrized by a trivial curve  $\gamma(t) = t$ ,  $t \in [a, b]$ . The integral of a continuous function  $f$  along  $\gamma$  is taken to be  $\int_a^b f(z)dz$ . We shall introduce a way to generalise line integrals along an arbitrary contour.

**Definition 10.** Let  $f$  be a complex-valued continuous function defined on a smooth curve parametrized by  $\gamma : [a, b] \rightarrow \mathbb{C}$ . The *integral* of  $f$  along  $\gamma$  is

$$\int_{\gamma} f(z)dz := \int_a^b f(\gamma(t))\gamma'(t)dt.$$

If  $\gamma$  is a contour which is smooth on  $[a_{j-1}, a_j]$  for  $j = 1, \dots, N$  where  $a_0 = a$  and  $a_N = b$ , then the *integral* of  $f$  along  $\gamma$  is a finite sum of the integral over the smooth parts:

$$\int_{\gamma} f(z) dz := \sum_{j=1}^N \int_{a_{j-1}}^{a_j} f(\gamma(t)) \gamma'(t) dt.$$

*Remark.* When  $\gamma$  is a simple closed curve, it is common to denote the integral of  $f$  along  $\gamma$  by the notation

$$\oint_{\gamma} f(z) dz.$$

Two parametrizations  $\gamma : [a, b] \rightarrow \mathbb{C}$  and  $\sigma : [c, d] \rightarrow \mathbb{C}$  of a smooth curve are *equivalent* if there is a continuously differentiable bijection  $h : [a, b] \rightarrow [c, d]$  such that  $h(a) = c$ ,  $h(b) = d$ , and  $\gamma = \sigma \circ h$ .

Our definition of contour integral is robust because it is independent of our choice of parametrization of the curve. If two curves  $\gamma$  and  $\sigma$  are equivalent, convince yourself by change of variables  $t \mapsto h(t)$  defined above that the integral of  $f$  along  $\gamma$  and  $\sigma$  must be equal:

$$\int_{\gamma} f(z) dz = \int_{\sigma} f(z) dz.$$

If we reverse the orientation of  $\gamma$ , we can again apply change of variables  $t \mapsto a + b - t$  and check that the integral of a continuous function  $f$  on  $\gamma^-$  is

$$\int_{\gamma^-} f(z) dz = - \int_{\gamma} f(z) dz. \quad (3.1)$$

**Example 20.** Let's compute the integral  $I = \oint_{\gamma} f(z) dz$  of the function  $f(z) = 1/(z - z_0)$  along the curve  $\gamma(t) = z_0 + re^{it}$  where  $0 \leq t \leq 2\pi$ .

$$I = \int_0^{2\pi} \frac{\gamma'(t)}{\gamma(z) - z_0} dt = \int_0^{2\pi} \frac{ire^{it}}{(z_0 + re^{it}) - z_0} dt = \int_0^{2\pi} idt = 2\pi i.$$

Notice that the value  $I$  is independent of the radius  $r$  and the center  $z_0$ .

The *length*  $L(\gamma)$  of a smooth curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  can be computed by the following integral:

$$L(\gamma) = \int_a^b |\gamma'(t)| dt.$$

We can think of  $L(\gamma)$  as the integral of the continuous function  $f$  along  $\gamma$  defined by  $f(z) = |\gamma'(t)|/\gamma'(t)$  where  $z = \gamma(t)$ . As such, the length is invariant under parametrization and change of orientation.

**Example 21.** The length of any circle of radius  $r$  is  $2\pi r$ . Indeed, using the parametrisation  $\gamma(t) = z_0 + re^{it}$  where  $0 \leq t \leq 2\pi$ ,

$$L(\gamma) = \int_0^{2\pi} |ire^{it}| dt = \int_0^{2\pi} r dt = 2\pi r.$$

An equivalent parametrization that is commonly used is  $\sigma(t) = z_0 + re^{2\pi it}$ ,  $0 \leq t \leq 1$ . Can you show that the length computed using this parametrization is the same?

**Proposition 3.3.** *Let  $f$  and  $g$  be continuous functions on a contour  $\gamma$  and let  $\alpha, \beta \in \mathbb{C}$ . Then,*

- $\int_{\gamma} \alpha f(z) + \beta g(z) dz = \alpha \int_{\gamma} f(z) dz + \beta \int_{\gamma} g(z) dz$ , (linearity)
- $|\int_{\gamma} f(z) dz| \leq L(\gamma) \cdot \max_{z \in \gamma} |f(z)|$ . (ML inequality)

*Proof.* Linearity is trivial. The curve parametrized by  $\gamma$  is a compact subset of  $\mathbb{C}$  and therefore  $|f|$  always attains its maximum along  $\gamma$ . Let  $M := \max_{z \in \gamma} |f(z)|$  and pick a parametrization  $\gamma : [a, b] \rightarrow \mathbb{C}$ . Then,

$$\begin{aligned} \left| \int_{\gamma} f(z) dz \right| &\leq \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \leq \int_a^b |f(\gamma(t))| \cdot |\gamma'(t)| dt \\ &\leq \int_a^b M |\gamma'(t)| dt = M \int_a^b |\gamma'(t)| dt = ML(\gamma). \end{aligned}$$

The second inequality follows from the fact that for any continuous function  $h : [a, b] \rightarrow \mathbb{C}$ , we always have the inequality  $|\int_a^b h(t) dt| \leq \int_a^b |h(t)| dt$ . We shall prove this below. Let  $re^{i\theta}$  be the polar form of  $\int_a^b h(t) dt$ . Then,

$$\begin{aligned} \left| \int_a^b h(t) dt \right| &= e^{-i\theta} \int_a^b h(t) dt = \operatorname{Re} \int_a^b e^{-i\theta} h(t) dt \\ &= \int_a^b \operatorname{Re}[e^{-i\theta} h(t)] dt \leq \int_a^b |e^{-i\theta} h(t)| dt = \int_a^b |h(t)| dt, \end{aligned}$$

and we are done. □

### 3.3 Primitives

**Definition 11.** An *antiderivative / primitive* of a continuous function  $f$  on a domain  $U \subset \mathbb{C}$  is a holomorphic function  $F : U \rightarrow \mathbb{C}$  such that  $F' = f$ .

The existence of primitives makes computation of contour integrals much easier. Regardless of the shape of the contour, it turns out that the integral only depends on the endpoints of the contour.

**Lemma 3.4.** Suppose  $f$  is continuous on a domain  $U$  and has a primitive  $F$ . Then, for any contour  $\gamma : [a, b] \rightarrow U$ ,

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)),$$

*Proof.* Let  $\gamma : [a, b] \rightarrow U$  be a smooth curve, then

$$\int_{\gamma} f(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} F(\gamma(t)) dt = F(\gamma(b)) - F(\gamma(a)),$$

If  $\gamma$  is piecewise smooth, we can sum up the integral over all smooth parts to obtain a similar result.  $\square$

**Corollary 3.5.** If  $F$  is holomorphic on a domain  $U$  and  $F' \equiv 0$ , then  $F$  is a constant function.

*Proof.* For any two points  $u$  and  $v$  in  $U$ , we can apply the lemma above by setting  $f = 0$  to obtain  $f(u) = f(v)$  because the integral over  $f$  over any curve is 0.  $\square$

**Theorem 3.6.** Suppose  $f : U \rightarrow \mathbb{C}$  is continuous on a domain  $U$ . The following are equivalent:

- (a)  $f$  has a primitive  $F : U \rightarrow \mathbb{C}$ ,
- (b) For any closed contour  $\gamma$  in  $U$ ,  $\oint_{\gamma} f(z) dz = 0$ .

*Proof.* Suppose (a) is true. Let  $\gamma : [a, c] \rightarrow U$  be a closed contour, then  $F(\gamma(b)) = F(\gamma(b))$  because  $\gamma(b) = \gamma(a)$ . Then, Lemma 3.4 immediately gives us (a)  $\Rightarrow$  (b).

Suppose (b) is true. Pick a basepoint  $z_0 \in U$  and for each  $z \in U$ , let  $\gamma_z : [0, 1] \rightarrow U$  be a smooth curve from  $\gamma_z(0) = z_0$  to  $\gamma_z(1) = z$ . Let's define  $F$  by

$$F(z) = \int_{\gamma_z} f(w) dw.$$

To prove that  $F$  is a well-defined function, we must show that the value  $F(z)$  is independent of the choice of the smooth curve  $\gamma_z$ . Let  $\gamma_z$  and  $\sigma_z$  be two such curves, then reverse the orientation  $\sigma_z$  to obtain  $\sigma_z^-$ , a smooth curve from  $z$  to  $z_0$ . The two curves  $\gamma_z$  and  $\sigma_z^-$  glue together to form a closed contour  $\Gamma_z$ . By (3.1) and (b),

$$\int_{\gamma_z} f(w)dw - \int_{\sigma_z} f(w)dw = \int_{\gamma_z} f(w)dw + \int_{\sigma_z^-} f(w)dw = \oint_{\Gamma_z} f(z)dz = 0.$$

Therefore, the integrals of  $f$  along  $\gamma_z$  and  $\sigma_z$  coincide, proving that  $F$  is a function. The proof is complete once we show that  $F$  is holomorphic and  $F' = f$ . Pick any point  $z \in U$ , then  $U$  contains a small ball  $\mathbb{D}(z, \epsilon)$ . Pick  $h \in \mathbb{C}$  such that  $|h| < \epsilon$ , and define a line segment  $\alpha_h(t) = z + th$ ,  $0 \leq t \leq 1$ . By a similar argument as above, (b) implies

$$F(z + h) - F(z) = \int_{\alpha_h} f(w)dw.$$

Then, using the fact that  $\int_{\alpha_h} dz = h$  and  $L(\alpha_h) = |h|$ , by ML inequality,

$$\begin{aligned} & \left| f(z) - \frac{F(z + h) - F(z)}{h} \right| \\ &= \left| f(z) - \frac{1}{h} \int_{\alpha_h} f(w)dw \right| = \left| \frac{1}{h} \int_{\alpha_h} (f(z) - f(w))dw \right| \\ &\leq \frac{1}{|h|} L(\alpha_h) \max_{w \in \alpha_h} |f(z) - f(w)| = \max_{w \in \alpha_h} |f(z) - f(w)|. \end{aligned}$$

By continuity of  $f$ , as  $h \rightarrow 0$ ,  $\max_{w \in \alpha_h} |f(z) - f(w)| \rightarrow 0$ . The limit of the left hand side of the inequality above is also 0, thus proving holomorphicity of  $F$  near  $z$ .  $\square$

**Example 22.** The function  $f(z) = (z - z_0)^{-n}$  on  $\mathbb{C} \setminus \{z_0\}$  admits a primitive  $\frac{(z - z_0)^{-n+1}}{-n+1}$  when  $n \neq 1$ . As such, if  $n \neq 1$  and  $\gamma$  is any closed contour not passing through  $z_0$ ,

$$\oint_{\gamma} \frac{dz}{(z - z_0)^n} = 0.$$

**Example 23.** The derivative of  $\log : \mathbb{C} \setminus (-\infty, 0] \rightarrow \mathbb{C}$  is the function  $\tau(z) = 1/z$ , so then  $\tau$  has primitive  $\log$  in the domain  $\mathbb{C} \setminus (-\infty, 0]$ . This domain cannot be extended any further since  $\log$  becomes discontinuous on the branch cut. Alternatively, we can say that  $\tau$  has no primitive in  $\mathbb{C}^*$  because, from Example 20, the integral of  $\tau$ , along any circle centered at 0 is non-zero.

### 3.4 Cauchy-Goursat Theorem

Theorem 3.6 does not give a nice criterion for the existence of primitives because it can be rather troublesome to compute integrals over all possible closed contours.

**Theorem 3.7** (Cauchy-Goursat). *Let  $f$  be holomorphic on a simply connected domain  $U \subset \mathbb{C}$  and let  $\gamma$  be a closed contour in  $U$ , then*

$$\oint_{\gamma} f(z) dz = 0.$$

Cauchy proved the theorem using Green's theorem from multivariable calculus, whereas Goursat's proof, albeit rather lengthly, only uses continuity of partial derivatives. We will only present the proof using Green's theorem.

*Proof.* Let's assume first that  $\gamma$  is a simple closed contour and let  $V$  be the bounded domain enclosed by  $\gamma$ . Let  $f(x + iy) = u(x, y) + iv(x, y)$ . Assume that  $\gamma$  is positively oriented. By the usual change of variables  $z = x + iy$  and by Green's theorem,

$$\begin{aligned} \oint_{\gamma} f(z) dz &= \oint_{\gamma} f(x + iy) dx + i f(x + iy) dy \\ &= \iint_V i f_x - f_y dx dy = \iint_V (iu_x - v_x) - (u_y + iv_y) dx dy = 0, \end{aligned}$$

where the last equality follows from Cauchy-Riemann equations.

When  $\gamma$  is negatively oriented (clockwise), we can replace it by  $\gamma^-$  and the minus sign will not change the value zero. When  $\gamma$  is not a simple curve, there is a way to partition  $\gamma$  into a finite number of components consisting of simple closed curves and a degenerate closed curves, i.e. those which enclose a region of zero area. Taking the sum of the integrals over each of these components, we will still obtain 0.  $\square$

*Remark.* Simply connectedness is an essential criterion. If the domain  $U$  is multiply connected, there is always a contour  $\gamma$  such that the domain  $V$  enclosed by  $U$  is not contained in  $U$  and thus the integral of  $f$  over the region  $V \setminus U$  will not make sense. Example 20 shows that this theorem fails when the domain is multiply connected.

Combining this theorem with Theorem 3.6, we obtain the following.

**Corollary 3.8.** *Any holomorphic function on a simply connected domain has a primitive.*

**Theorem 3.9** (Deformation Theorem). *Let  $f$  be holomorphic on a domain  $U \subset \mathbb{C}$ , and let  $\gamma$  and  $\sigma$  be two simple closed contours such that  $\sigma$  lies in the domain  $V$  enclosed by  $\gamma$  and that both contours have the same orientation. If the annulus  $A$  enclosed by  $\gamma$  and  $\sigma$  is contained in  $U$ , then*

$$\oint_{\gamma} f(z) dz = \oint_{\sigma} f(z) dz.$$

*Proof.* Pick a simple contour  $\alpha$  in  $A$  joining a point on the outer boundary to another on the inner boundary. (See Figure 3.4.) Removing  $\alpha$  from  $A$  gives us a region  $A'$  and its boundary can be parametrized by the closed contour  $\Gamma$  obtained by gluing in order the curves  $\gamma$ ,  $\alpha$ ,  $\sigma^-$ , and  $\alpha^-$ . Then, by Cauchy-Goursat,

$$\begin{aligned} \oint_{\gamma} f(z) dz - \oint_{\sigma} f(z) dz &= \oint_{\gamma} f(z) dz + \oint_{\alpha} f(z) dz + \oint_{\sigma^-} f(z) dz + \oint_{\alpha^-} f(z) dz \\ &= \oint_{\Gamma} f(z) dz = 0. \end{aligned} \quad \square$$




Figure 3.4: Curve  $\alpha$  joining  $\gamma$  and  $\sigma$ .




Figure 3.5: The square  $\gamma$  can be replaced by the smaller circle.

**Example 24.** Let's compute the integral  $I$  of  $(z^2 - 4)^{-1}$  along  $\gamma$ , a simple closed contour parametrising the square of side length 4 centered at  $1 + i$ . The partial fraction decomposition of  $I$  is

$$I = \oint_{\gamma} \frac{1}{z^2 - 4} dz = \frac{1}{4} \oint_{\gamma} \frac{1}{z - 2} dz - \frac{1}{4} \oint_{\gamma} \frac{1}{z + 2} dz$$

Since the function  $1/(z + 2)$  is holomorphic on and inside  $\gamma$ , the second integral is zero by Cauchy-Goursat. The function  $\frac{1}{z-2}$  is holomorphic everywhere

except at 2, so then by deformation theorem, we can replace  $\gamma$  on the first integral with any circle centered at 2, e.g.  $C(2, 0.5)$ . (See Figure 3.5.) Example 20 immediately tells us that the first integral is equal to  $\frac{1}{4} \cdot 2\pi i$ . Therefore,  $I = \frac{1}{2}\pi i$ .

In fact, the deformation theorem can be applied to any pair of arbitrary closed contours in  $U$  which are *homotopic*. This means that one curve can be continuously deformed in  $U$  to the other curve. If the two contours are not closed, then they must be *homotopic relative to their endpoints*, i.e. we need the additional assumption that both have the same endpoints. (Lemma 3.4 hints at why fixing endpoints are important.) Below is a simple pictorial explanation of homotopic curves.




Figure 3.6: On the gray annular domain above, curves 1, 2 and 3 have the same endpoints but only 1 and 2 are homotopic relative to their endpoints. Among closed curves 4 - 7, the only pair of homotopic curves is 5 and 6.

### Short Quiz 3

1. What is the integral of  $1/z$  along the circle  $C(0, 2)$ ?
2. Compute the length of the curve  $\gamma(t) = \cos(t)e^{it}$  where  $0 \leq t \leq \pi$ .
3. Compute the contour integral of the function  $8z^3$  along an L-shaped contour which starts from 1 to  $2i$  and passes through 0.

Answers: 1.  $2\pi i$ , 2.  $\pi$ , 3. 30.

# Chapter 4

## Integration Formulas

### 4.1 Cauchy's Formulas

**Theorem 4.1** (Cauchy's Integral Formula). *Let  $f$  be holomorphic on a simply connected domain  $U$ ,  $\gamma$  be a simple closed contour in  $U$ , and  $z_0$  be a point lying in the domain enclosed by  $\gamma$ . Then,*

$$f(z_0) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z - z_0} dz.$$

*Proof.* We can assume that  $z_0 = 0$  without loss of generality because when  $z_0 \neq 0$ , we can replace  $f$  with the function  $f(z + z_0)$  on the domain  $\{z \in \mathbb{C} : z + z_0 \in U\}$  and the contour  $\gamma(t)$  with  $\gamma(t) - z_0$ .

By the deformation theorem, we can replace  $\gamma$  with  $\gamma_r$ , a contour parametrizing the circle  $C(0, r)$  for arbitrarily small radius  $r > 0$ . Recall that

$$\frac{1}{2\pi i} \oint_{\gamma_r} \frac{1}{z} dz = 1.$$

Then, by taking the limit as  $r \rightarrow 0$ ,

$$\begin{aligned} \left| \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z} dz - f(0) \right| &= \left| \frac{1}{2\pi i} \oint_{\gamma_r} \frac{f(z)}{z} dz - f(0) \right| \\ &= \left| \frac{1}{2\pi i} \oint_{\gamma_r} \frac{f(z) - f(0)}{z} dz \right| \\ &\leq \frac{1}{2\pi} \cdot L(\gamma_r) \cdot \max_{z \in \gamma_r} \left| \frac{f(z) - f(0)}{z} \right| \\ &= r \cdot \max_{z \in \gamma_r} \left| \frac{f(z) - f(0)}{z} \right| \longrightarrow 0 \cdot f'(0) = 0. \end{aligned}$$

As the term of the left hand side is independent of  $r$ , it is identically 0.  $\square$