

Chapter 2 Convex sets

Last update on 2025-02-24 14:15:08+08:00

Table of contents

Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

three groups of concepts

affine combination	convex combination	conic combination
affine set	convex set	convex cone
affine hull	convex hull	conic hull

Convex combination

convex combination of $x_1, \dots, x_k \in \mathbb{R}^n$: points of the form

$$\theta_1 x_1 + \dots + \theta_k x_k, \quad \text{where } \theta_1, \dots, \theta_k \geq 0 \quad \text{and} \quad \theta_1 + \dots + \theta_k = 1$$



line segment between x_1 and x_2 : the set of all convex combinations of x_1 and x_2

$$\{x = \theta x_1 + (1 - \theta)x_2 \mid 0 \leq \theta \leq 1\}$$

Convex set

convex set: $C \subseteq \mathbb{R}^n$ is convex if contains line segment between any pair of points in C



examples (one convex, two nonconvex)



Convex hull

convex hull of $C \subseteq \mathbb{R}^n$: the set of all convex combinations of points in C

$$\mathbf{conv} C = \{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_1, \dots, x_k \in C; \theta_1, \dots, \theta_k \geq 0; \theta_1 + \cdots + \theta_k = 1\}$$



facts

- ▶ the convex hull of C is the smallest convex set containing C
- ▶ if C is a convex set, then $\mathbf{conv} C = C$


Affine combination

affine combination of $x_1, \dots, x_k \in \mathbb{R}^n$: points of the form

$$\theta_1 x_1 + \dots + \theta_k x_k, \quad \text{where} \quad \theta_1 + \dots + \theta_k = 1$$

line through x_1 and x_2 : the set of all affine combinations of x_1 and x_2

$$\{x = \theta x_1 + (1 - \theta)x_2 \mid \theta \in \mathbb{R}\}$$



affine set: $C \subseteq \mathbb{R}^n$ is affine if it contains the line through any pair of points in C

example

- ▶ the solution set of linear equations $\{x \mid Ax = b\}$ is an affine set
- ▶ conversely, every affine set can be expressed as the solution set of a system of linear equations

affine hull of $C \subseteq \mathbb{R}^n$: the set of all affine combinations of points in C

$$\mathbf{aff} C = \{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_1, \dots, x_k \in C, \theta_1 + \cdots + \theta_k = 1\}$$

facts


- ▶ the affine hull of C is the smallest affine set containing C
- ▶ if C is an affine set, then $\mathbf{aff} C = C$

Conic combination

cone: $C \subseteq \mathbb{R}^n$ is a cone if $\theta x \in C$ for every $x \in C$ and $\theta \geq 0$.

conic combination of $x_1, \dots, x_k \in \mathbb{R}^n$: points of the form

$$\theta_1 x_1 + \dots + \theta_k x_k, \quad \text{where } \theta_1, \dots, \theta_k \geq 0$$



convex cone: $C \subseteq \mathbb{R}^n$ is a convex cone if it is convex and a cone


fact

C is a convex cone $\iff C$ contains all conic combinations of points in itself

Conic hull

conic hull of $C \subseteq \mathbb{R}^n$: the set of all conic combinations of points in C

$$\{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_1, \dots, x_k \in C; \theta_1, \dots, \theta_k \geq 0\}$$



facts

- ▶ the conic hull of C is the smallest convex cone containing C
- ▶ if C is a convex cone, then its conic hull is itself

Exercises

- ▶ Study the following concepts from text:

affine dimension, **relative interior.**

- ▶ Suppose that $C \subseteq \mathbb{R}^n$ is convex, then $\mathbf{int} C$ and $\mathbf{cl} C$ are also convex.
- ▶ Suppose that $C \subseteq \mathbb{R}^n$ is convex, then

$$\mathbf{int} C = \emptyset \quad \Longleftrightarrow \quad C \text{ is contained in a hyperplane.}$$

- ▶ Suppose that $C \subseteq \mathbb{R}^n$ is convex, and $\mathbf{int} C \neq \emptyset$, then

$$\mathbf{cl}(\mathbf{int} C) = \mathbf{cl} C.$$

) Look at $d + 1$, the largest number of affinely independent points from C . Let x_0, \dots, x_d one such affinely independent subset of largest size. Note that every other point is an affine combination of the points x_k , so lies in the affine subspace generated by them, which is of dimension d .

) If $d < n$ then this subspace is contained in an affine hyperplane.

If $d = n$, then C contains $d + 1$ affinely independent points. Since C is convex, it will also contain the convex hull of those $n + 1$ points. Now, in an n -dimensional space the convex hull of $n + 1$ affinely independent points has non-empty interior. So the interior of C is non-empty.

Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities


Dual cones and generalized inequalities

four types of elementary examples

- ▶ LP type: hyperplanes, halfspaces, polyhedra
- ▶ ball type: Euclidean balls, ellipsoids, norm balls
- ▶ cone type: second-order cone, norm cones
- ▶ matrix type: positive semidefinite cone

Hyperplanes


hyperplane: set of the form $\{x \mid a^T x = b\}$ ($a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R}$)



fact: hyperplanes are affine and convex

Halfspaces

halfspace: set of the form $\{x \mid a^T x \leq b\}$ ($a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R}$)



fact: halfspaces are convex


Polyhedra

polyhedron: solution set of finitely many linear inequalities and equalities

$$Ax \preceq b, \quad Cx = d$$

where $A \in \mathbb{R}^{m \times n}$, $C \in \mathbb{R}^{p \times n}$, \preceq is componentwise inequality

i.e. polyhedra are intersections of finite number of halfspaces and hyperplanes;



fact: polyhedra are convex (page 31)

Euclidean ball with center x_c and radius r : two equivalent representations

- ▶ set of the form

$$B(x_c, r) = \{x \mid \|x - x_c\|_2 \leq r\}$$

- ▶ set of the form

$$B(x_c, r) = \{x_c + ru \mid \|u\|_2 \leq 1\}$$

fact: Euclidean balls are convex

Ellipsoids


ellipsoid: two equivalent representations

- ▶ set of the form

$$\{x \mid (x - x_c)^T P^{-1} (x - x_c) \leq 1\} \quad \text{with} \quad P \in \mathbb{S}_{++}^n$$

- ▶ set of the form

$$\{x_c + Au \mid \|u\|_2 \leq 1\} \quad \text{with } A \text{ square and nonsingular}$$



fact: ellipsoids are convex

Norm balls

norm ball with center x_c and radius r : set of the form


$$\{x \mid \|x - x_c\| \leq r\}$$

fact: norm balls are convex

Norm cones

second-order cone:

$$\{(x, t) \in \mathbb{R}^n \times \mathbb{R} \mid \|x\|_2 \leq t\}$$



norm cone: for any norm $\|\cdot\|$ on \mathbb{R}^n

$$\{(x, t) \in \mathbb{R}^n \times \mathbb{R} \mid \|x\| \leq t\}$$


fact: norm cones are convex cones

Positive semidefinite cone

positive semidefinite cone

$$\mathbb{S}_+^n = \{X \in \mathbb{S}^n \mid X \succeq 0\}$$

fact: positive semidefinite cone \mathbb{S}_+^n is a convex cone



Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

Establishing convexity

Practical methods for establishing convexity of a set C

1. apply definition

$$x_1, x_2 \in C, 0 \leq \theta \leq 1 \quad \implies \quad \theta x_1 + (1 - \theta)x_2 \in C$$

2. reconstruct C from known convex sets by operations preserving convexity:

- ▶ intersection
- ▶ affine functions
- ▶ perspective function
- ▶ linear-fractional functions

Intersection

an arbitrary intersection of convex sets is convex

example

the positive semidefinite cone \mathbb{S}_+^n is convex

affine function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form

$$f(x) = Ax + b \quad \text{with } A \in \mathbb{R}^{m \times n} \text{ and } b \in \mathbb{R}^m$$

- ▶ the image of a convex set under f is convex

$$S \subseteq \mathbb{R}^n \text{ convex} \quad \implies \quad f(S) = \{f(x) \mid x \in S\} \text{ convex}$$

- ▶ the inverse image $f^{-1}(C)$ of a convex set under f is convex

$$C \subseteq \mathbb{R}^m \text{ convex} \quad \implies \quad f^{-1}(C) = \{x \in \mathbb{R}^n \mid f(x) \in C\} \text{ convex}$$

examples

- ▶ scaling and translation: if $S \subseteq \mathbb{R}^n$ is convex, $\alpha \in \mathbb{R}$ and $a \in \mathbb{R}^n$, then

$$\alpha S = \{\alpha x \mid x \in S\} \quad \text{and} \quad S + a = \{x + a \mid x \in S\}$$

are convex

- ▶ projection: if $S \subseteq \mathbb{R}^m \times \mathbb{R}^n$ is convex, then

$$T = \{x_1 \in \mathbb{R}^m \mid (x_1, x_2) \in S \text{ for some } x_2 \in \mathbb{R}^n\}$$

is convex

- ▶ sum: if $S_1, S_2 \subseteq \mathbb{R}^n$ are both convex, then

$$S_1 + S_2 = \{x + y \mid x \in S_1, y \in S_2\}$$

is convex

- ▶ solution set of linear matrix inequality

$$\{x \in \mathbb{R}^n \mid x_1 A_1 + \cdots + x_n A_n \preceq B\}$$

where $A_i, B \in \mathbb{S}^m$, is convex

proof

inverse image of the positive semidefinite cone under the affine function

$$f: \mathbb{R}^n \rightarrow \mathbb{S}^m, \quad f(x) = B - (x_1 A_1 + \cdots + x_n A_n)$$

► hyperbolic cone

$$\left\{x \in \mathbb{R}^n \mid x^T P x \leq (c^T x)^2, c^T x \geq 0\right\}$$

where $P \in \mathbb{S}_+^n$ and $c \in \mathbb{R}^n$, is convex

proof

inverse image of the second-order cone

$$\{(z, t) \mid z^T z \leq t^2, t \geq 0\}$$

under the affine function $f: \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$ given by $f(x) = (P^{1/2}x, c^T x)$

Perspective function

perspective function $P: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ given by

$$P(x, t) = x/t, \quad \mathbf{dom} P = \mathbb{R}^n \times \mathbb{R}_{++} = \{(x, t) \mid t > 0\}$$

- ▶ images of convex sets under perspective function are convex
- ▶ inverse images of convex sets under perspective function are convex
- ▶ prove it

Linear-fractional functions

linear-fractional function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by

$$f(x) = \frac{Ax + b}{c^T x + d}, \quad \text{dom } f = \{x \mid c^T x + d > 0\}$$



it is the composition of an affine function g and the perspective function P , where

$$g(x) = \begin{bmatrix} A \\ c^T \end{bmatrix} x + \begin{bmatrix} b \\ d \end{bmatrix}$$

- ▶ images of convex sets under linear-fractional functions are convex
- ▶ inverse images of convex sets under linear-fractional functions are convex

example

$$f(x) = \frac{x}{x_1 + x_2 + 1}, \quad \text{dom } f = \{(x_1, x_2) \mid x_1 + x_2 + 1 > 0\}$$



Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

two fundamental properties of convex sets


- ▶ separating hyperplane theorem
- ▶ supporting hyperplane theorem

Separating hyperplane theorem

Theorem

If C and D are nonempty disjoint convex sets, then there exist $a \neq 0$ and b such that

$$a^T x \leq b \text{ for all } x \in C, \quad a^T x \geq b \text{ for all } x \in D.$$



Remark $\{x \mid a^T x = b\}$ is called a **separating hyperplane**

Sketch of proof

Step 1. Theorem holds if there exists $c \in C$ and $d \in D$ such that

$$\|c - d\|_2 \leq \|u - v\|_2$$

for all $u \in C$ and $v \in D$.

Idea: take the perpendicular bisector of the line segment connecting c and d .

Step 2. If $C = \{0\}$, then there exists $a \neq 0$ such that $a^T x \geq 0$ for all $x \in D$.

Idea: if $0 \notin \text{cl } D$, apply Step 1 to $\{0\}$ and $\text{cl } D$; if $\text{int } D = \emptyset$, D is contained in a hyperplane; if $0 \in \text{cl } D$ and $\text{int } D \neq \emptyset$, shrink D by ε and let $\varepsilon \rightarrow 0$.

Step 3. Prove the general case.

Idea: apply Step 2 to $\{0\}$ and $S = \{y - x \mid x \in C, y \in D\}$.

Step 1. Define

$$f(x) = (d - c)^T \left(x - \frac{d + c}{2} \right) = (d - c)^T x - \frac{1}{2} \|d - c\|_2^2.$$

For any $v \in D$, we have $tv + (1 - t)d \in D$ for $t \in [0, 1]$, hence the function

$$g(t) = \|tv + (1 - t)d - c\|_2^2$$

satisfies $g(t) \geq g(0)$ for $t \in [0, 1]$, which implies $g'(0) \geq 0$. Since

$$g'(t) = 2(tv + (1 - t)d - c)^T (v - d)$$

we obtain $g'(0) = 2(d - c)^T (v - d) \geq 0$, therefore

$$\begin{aligned} f(v) &= (d - c)^T \left(v - \frac{d + c}{2} \right) = (d - c)^T \left(v - d + \frac{d - c}{2} \right) \\ &= (d - c)^T (v - d) + \frac{1}{2} \|d - c\|_2^2 \geq 0. \end{aligned}$$

Similarly we can show $f(u) \leq 0$ for any $u \in C$.

Step 2. We first check two simple cases, then prove the remaining cases.

Case 1. Assume $0 \notin \text{cl } D$.

Note that $\text{cl } D$ is convex, and there exists some $d \in \text{cl } D$ such that $\|d - 0\|_2 \leq \|y - 0\|_2$ for all $y \in \text{cl } D$ (why?). Applying Step 1 to $\{0\}$ and $\text{cl } D$, there exists $a \neq 0$ and b such that $a^T y \geq b \geq a^T 0 = 0$ for all $y \in \text{cl } D$.

Case 2. Assume $\text{int } D = \emptyset$.

In such a case D is contained in a hyperplane $\{z \mid a^T z = b\}$ for some $a \neq 0$ and b . Assume wlog $b \geq 0$, then $a^T y = b \geq 0$ for all $y \in D$.

Case 3. Consider the remaining cases.

For each sufficiently small $\varepsilon > 0$, the set $D_{-\varepsilon} = \{z \mid B(z, \varepsilon) \subseteq D\}$ is nonempty and convex, and $0 \notin \text{cl } D_{-\varepsilon}$ (why?). By Case 1 there exists $a_\varepsilon \neq 0$ such that $a_\varepsilon^T z \geq 0$ for all $z \in D_{-\varepsilon}$. Assume wlog $\|a_\varepsilon\|_2 = 1$. Choose any positive sequence $\{\varepsilon_i\}$ converging to 0, a subsequence of $\{a_{\varepsilon_i}\}$ converges to some \bar{a} with $\|\bar{a}\|_2 = 1$. Then $\bar{a}^T z \geq 0$ for all $z \in \text{int } D$ (why?) hence for all $z \in D$ (why?).

Step 3. The nonempty set

$$S = \{y - x \mid x \in C, y \in D\}$$

is convex (why?) and disjoint from $\{0\}$. By Step 2, there exists $a \neq 0$ such that

$$a^T(y - x) \geq 0 \quad \Longleftrightarrow \quad a^T x \leq a^T y$$

for all $x \in C$ and $y \in D$. It follows that

$$\sup\{a^T x \mid x \in C\} \leq \inf\{a^T y \mid y \in D\}$$

whose both sides are finite. Then any b in between satisfies

$$a^T x \leq b \leq a^T y$$

for all $x \in C$ and $y \in D$.

Remarks


- ▶ strict separation requires additional assumptions (e.g. point and closed convex set)
- ▶ converse separating theorem requires additional assumptions (e.g. one set is open)

Supporting hyperplane theorem

A **supporting hyperplane** to a set C at a boundary point x_0 is a hyperplane

$$\{x \mid a^T x = a^T x_0\}$$

where $a \neq 0$, such that $a^T x \leq a^T x_0$ for all $x \in C$



Theorem

If C is convex, then supporting hyperplane exists at every boundary point of C .

Sketch of proof

Step 1. Assume $\text{int } C \neq \emptyset$. (Idea: use separating hyperplane for $\{x_0\}$ and $\text{int } C$.)

Since $\text{int } C$ is also convex (why?), we apply separating hyperplane theorem to $\text{int } C$ and $\{x_0\}$ to conclude the existence of $a \neq 0$ and b such that $a^T x \leq b \leq a^T x_0$ for all $x \in \text{int } C$. Since $\text{cl } C = \text{cl}(\text{int } C)$ (why?) it follows that $a^T x \leq b$ for all $x \in \text{cl } C$. In particular $a^T x \leq b = a^T x_0$ for all $x \in C$ and the given $x_0 \in \text{bd } C$.

Step 2. Assume $\text{int } C = \emptyset$. (Idea: C is contained in a hyperplane.)

Then C is contained in a hyperplane $\{x \mid a^T x = b\}$ for some $a \neq 0$ and b (why?). It follows $a^T x = b = a^T x_0$ for all $x \in C$.

Tasks Study the following extensions of both theorems from the text

- ▶ strict separation theorem
- ▶ converse of separating hyperplane theorem
- ▶ converse of supporting hyperplane theorem

Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

Proper cones

proper cone: a cone $K \subseteq \mathbb{R}^n$ satisfying

- ▶ K is convex
- ▶ K is closed (contains its boundary)
- ▶ K is solid (has nonempty interior)
- ▶ K is pointed (contains no line, or equivalently, $\pm x \in K \implies x = 0$)

examples

- ▶ nonnegative orthant $K = \mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x_i \geq 0, i = 1, \dots, n\}$
- ▶ positive semidefinite cone $K = \mathbb{S}_+^n$

Generalized inequalities

generalized inequality on \mathbb{R}^n defined by a proper cone $K \subseteq \mathbb{R}^n$

$$x \preceq_K y \iff y - x \in K$$

$$x \prec_K y \iff y - x \in \text{int } K$$

examples (similar for \prec , \succeq , \succ)

- ▶ nonnegative orthant and componentwise inequality ($K = \mathbb{R}_+^n$)

$$x \preceq_{\mathbb{R}_+^n} y \iff x_i \leq y_i, \quad i = 1, \dots, n$$

- ▶ positive semidefinite cone and symmetric matrix inequality ($K = \mathbb{S}_+^n$)

$$X \preceq_{\mathbb{S}_+^n} Y \iff Y - X \text{ positive semidefinite}$$

these two types are so common that we drop the subscript in \preceq_K

properties (same for \prec , \succeq , \succ)

- ▶ many properties of \preceq_K are similar to \leq on \mathbb{R} ; for example:

$$x \preceq_K y, \quad u \preceq_K v \quad \implies \quad x + u \preceq_K y + v$$

- ▶ partial ordering (not always a total/linear ordering)

it could happen that $x \not\preceq_K y$ and $y \not\preceq_K x$

- ▶ pay attention to other subtleties

for example: $x \preceq_K y$ does not imply $x \prec_K y$ or $x = y$

- ▶ if $x \prec_K y$, then for u and v small enough, $x + u \prec_K y + v$

Minimum and minimal elements

$x \in S$ is the **minimum element** of S with respect to \preceq_K if for all

$$y \in S \quad \implies \quad x \preceq_K y$$

$x \in S$ is a **minimal element** of S with respect to \preceq_K if

$$y \in S, \quad y \preceq_K x \quad \implies \quad y = x$$

Remark Minimum element, if exists, must be unique.


property A point $x \in S$ is the minimum element of S if and only if

$$S \subseteq x + K.$$


property A point $x \in S$ is the minimal element of S if and only if

$$(x - K) \cap S = \{x\}.$$

example for $K = \mathbb{R}_+^2$



x_1 is the minimum element of S_1



x_2 is a minimal element of S_2

coming up

criteria for determining minimum/minimal elements

Affine and convex sets

Elementary examples

Operations preserving convexity

Separating and supporting hyperplanes

Generalized inequalities

Dual cones and generalized inequalities

dual cone of a cone $K \subseteq \mathbb{R}^n$

$$K^* = \{y \in \mathbb{R}^n \mid y^T x \geq 0 \text{ for all } x \in K\}$$

examples

$$K = \mathbb{R}_+^n \implies K^* = \mathbb{R}_+^n$$

$$K = \mathbb{S}_+^n \implies K^* = \mathbb{S}_+^n \text{ (example 2.24)}$$

$$K = \{(x, t) \mid \|x\|_2 \leq t\} \implies K^* = \{(x, t) \mid \|x\|_2 \leq t\}$$

$$K = \{(x, t) \mid \|x\|_1 \leq t\} \implies K^* = \{(x, t) \mid \|x\|_\infty \leq t\}$$

first three examples are **self-dual** cones

Properties of dual cones

- ▶ K^* is a cone, and is always convex, even when the original cone K is not convex
- ▶ K^* is closed and convex
- ▶ $K_1 \subseteq K_2$ implies $K_2^* \subseteq K_1^*$
- ▶ If K has nonempty interior, then K^* is pointed
- ▶ If the closure of K is pointed then K^* has nonempty interior
- ▶ K^{**} is the closure of the convex hull of K . (Hence if K is convex and closed, $K^{**} = K$)
- ▶ If K is a proper cone, then so is its dual K^* , and moreover, that $K^{**} = K$

Dual generalized inequalities

- ▶ assume K is a proper cone, then K^* is also a proper cone, and $K^{**} = K$
- ▶ in such a case K^* also defines generalized inequalities
- ▶ generalized inequalities with respect to K^* can usually be interpreted via K

examples

$$y \succeq_{K^*} 0 \quad \Longleftrightarrow \quad y \in K^* \quad \Longleftrightarrow \quad y^T x \geq 0 \text{ for all } x \succeq_K 0$$

$$y \succ_{K^*} 0 \quad \Longleftrightarrow \quad y \in \mathbf{int} K^* \quad \Longleftrightarrow \quad y^T x > 0 \text{ for all } x \succeq_K 0 \text{ and } x \neq 0$$

Dual characterization of the minimum element


x is the minimum element of S
with respect to \preceq_K \iff x is the unique minimizer of $\lambda^T z$
over $z \in S$ for each $\lambda \succ_{K^*} 0$

Geometrically, this means that for any $\lambda \succ_{K^*} 0$, the hyperplane

$$\{z \mid \lambda^T (z - x) = 0\}$$

is a strict supporting hyperplane to S at x .

Note that convexity of S is **not required**



Sketch of proof

(\implies) for each $z \in S$ and $z \neq x$

- ▶ x is minimum $\implies z - x \succeq_K 0$ and $z - x \neq 0$
- ▶ $\lambda^T(z - x) > 0$ for each $\lambda \succ_{K^*} 0 \implies x$ is the unique minimizer of $\lambda^T z$

(\impliedby) for each $z \in S$ and $z \neq x$

- ▶ x is the unique minimizer of $\lambda^T z \implies \lambda^T(z - x) > 0$ for each $\lambda \succ_{K^*} 0$
- ▶ continuity $\implies \lambda^T(z - x) \geq 0$ for each $\lambda \succeq_{K^*} 0 \implies z - x \succeq_K 0$


Dual characterization of a minimal element

There is a gap between the necessary and sufficient conditions. Compared to the minimum, there is no uniqueness in the condition.

x is a minimal element of S
with respect to \preceq_K



x minimizes $\lambda^T z$ over $z \in S$
for some $\lambda \succ_{K^*} 0$



x is a minimal element of S
with respect to \preceq_K
and S is **convex**



x minimizes $\lambda^T z$ over $z \in S$
for some nonzero $\lambda \succeq_{K^*} 0$

Sketch of proof

$$\begin{array}{ccc} x \text{ is a minimal element of } S & \Longleftarrow & x \text{ minimizes } \lambda^T z \text{ over } z \in S \\ \text{with respect to } \preceq_K & & \text{for some } \lambda \succ_{K^*} 0 \end{array}$$

(\Leftarrow)

- ▶ suppose that x is **not** minimal, i.e., there exists a $z \in S, z \neq x$, and $z \prec_K x$ (by def. 2.1 on page 45)
- ▶ Then $\lambda^T(x - z) > 0$ which is obtained from the second property relating a generalized inequality and its dual on page 53
- ▶ However, this contradicts our condition that x is the minimizer of $\lambda^T z$ over S

A point x can be minimal in S , but not a minimizer of $\lambda^T z$ over $z \in S$, for any λ (see Fig. 2.25 on page 56). This suggests that convexity plays an important role in the converse, which is correct.

Sketch of proof

$$\begin{array}{ccc} x \text{ is a minimal element of } S & & x \text{ minimizes } \lambda^T z \text{ over } z \in S \\ \text{with respect to } \preceq_K & \implies & \text{for some nonzero } \lambda \succeq_{K^*} 0 \\ \text{and } S \text{ is convex} & & \end{array}$$

(\implies)

- ▶ x is minimal $\implies ((x - K) \setminus \{x\}) \cap S = \emptyset$
- ▶ Applying the separating hyperplane theorem to the disjoint convex sets $((x - K) \setminus \{x\})$ and S , we conclude that there is a $\lambda \neq 0$ and μ such that $\lambda^T(x - y) \leq \mu$ for all $y \in K$, and $\lambda^T z \geq \mu$ for all $z \in S$
- ▶ From the first inequality, we have $\lambda^T y \geq \lambda^T x - \mu \geq 0$ since we take $z = x$ in the second inequality $\lambda^T z \geq \mu$ for all $z \in S$. We conclude $\lambda \succeq_{K^*} 0$ based on page 53.
- ▶ Since $x \in S$ and $x \in x - K$, we have $\lambda^T x = \mu$ on the separating hyperplane, so the second inequality implies that μ is the minimum value of $\lambda^T z$ over S .
- ▶ x is a minimizer of $\lambda^T z$ over S , where $\lambda \neq 0$, $\lambda \succeq_{K^*} 0$