



# Direct Estimation of Carbon Emissions from High Latitude Fires: The Adapted FREM Approach

Dr. Will Maslanka, Prof. Martin Wooster



Abstract



At high latitudes, global climate change is exacerbating conditions that impact and alter wildfires regimes

- Larger and more frequent fires, lengthening of fire season
- Warmer temperatures, regionally decreased precipitation<sup>[1]</sup>
- July 2023 “2<sup>nd</sup> warmest on record” – increase in wildfire news stories

Warming is accelerated in higher latitude regions, and increases the risk of high-latitude feedback mechanisms

- Release of carbon sequestered in high-latitude peatlands / permafrost
- Black carbon deposits on ice increasing rate of melting

Effective monitoring of high-latitude fires and emissions imperative for climate monitoring and reporting.

[1] IPCC 2019: Land-Climate Interactions in Special Report: Climate Change and Land



## Nova Scotia battles its largest wildfire on record

🕒 1 June

## Canada wildfire smoke leaves millions under air quality advisory

🕒 18 July

## Canada wildfires: At least 30,000 households in British Columbia told to evacuate

🕒 3 days ago

## Canada wildfire season is now the worst on record

🕒 29 June 2023

## Canada wildfires: Residents scramble to flee fires in Kelowna and Yellowknife

🕒 18 August 2023

# The FREM Approach: Current and Adaptation

One approach to estimate emissions from observations of Fires is through the top-down Fire Radiative Energy Emission (FREM) approach.

- Directly linking observations of FRP/FRE to TPM and (more recently) directly to CO to estimate emissions.
- Grouping these observations via biomes, can get a biome specific emission coefficient for species  $X$ ,  $EC_X^{biome}$
- Previous work has focused on Africa<sub>[2, 3]</sub> and SE Asia<sub>[4]</sub>, taking advantage of high temporal sampling with Geostationary Platforms.

This approach not valid for the high-latitudes, due to poor spatial resolution (if visible).

- For high-latitude fire prone areas, geostationary pixel area  $\approx 20\text{km}^2$
- High-latitude adaptation (aFREM): substituting Polar Orbiting platforms for Geostationary.

[2] Mota and Wooster 2018: 10.1016/j.rse.2017.12.016

[3] Nguyen et al 2023: 10.5194/acp-23-2089-2023

[4] Fisher et al., 2020: 10.3390/s20247075

# Data and High Latitude Region(s) of Interest

4

## VIIRS onboard Suomi-National Polar-orbiting Partnership (S-NPP)

- 375m / 750m spatial resolution
- Multispectral data, including derived fire radiative power (FRP) product
- Seen as the replacement for MODIS


## TROPOMI onboard Sentinel-5P

- 7.0 x 5.5km spatial resolution, trace gas products including CO
- Near-simultaneous observations with S-NPP

## Hourly Gridded FRP from locally ran “development” GFAS

- Using S-NPP FRP as input, rather than MODIS
- S-NPP capable of detecting far lower FRP fires than MODIS can

## ESA CCI Land Cover Map (2018, 300m resolution)





# aFREM Method – Excess “in plume” Mass of CO Estimation

5


Locating “well observed” (cloud-free) fires and smoke plumes

- Finding Plumes in RGB imagery (A) and AOD<sub>550</sub> (B)
- Digitize Plumes using RGB imagery (A) and AOD<sub>550</sub> (B)
- Extracting Excess CO from Sentinel-5P (C)
  - $\text{Excess CO} = \sum [\text{CO}_{\text{Obs}} - \text{CO}_{\text{Bgd}}]$

Locating these fire-plume “matchups” only needs to be done in the “setup” stage.

- Identified matchups provide the emission coefficients linking the total FRE emitted to total excess CO release.
- These coefficients can then be applied to all FRP data; not just those of the match up fires.

Potential for AI automation of plume mapping



# aFREM Method – Excess “in plume” Mass of CO Estimation

6


Locating “well observed” (cloud-free) fires and smoke plumes

- Finding Plumes in RGB imagery (A) and AOD<sub>550</sub> (B)
- Digitize Plumes using RGB imagery (A) and AOD<sub>550</sub> (B)
- Extracting Excess CO from Sentinel-5P (C)
  - $\text{Excess CO} = \sum [\text{CO}_{\text{Obs}} - \text{CO}_{\text{Bgd}}]$

Locating these fire-plume “matchups” only needs to be done in the “setup” stage.

- Identified matchups provide the emission coefficients linking the total FRE emitted to total excess CO release.
- These coefficients can then be applied to all FRP data; not just those of the match up fires.

Potential for AI automation of plume mapping



# aFREM Method – Fire Radiative Energy (FRE) Estimation


7

Locating “well observed” (cloud-free) fires and smoke plumes

- Finding Plumes in RGB imagery (A) and AOD<sub>550</sub> (B)
- Digitize Plumes using RGB imagery (A) and AOD<sub>550</sub> (B)
- Extracting Excess CO from Sentinel-5P (C)
  - $\text{Excess CO} = \sum [\text{CO}_{\text{Obs}} - \text{CO}_{\text{Bgd}}]$

Similar Process for FRE and Biome

- Using False Colour Composite (D) to highlight fire activity
- Digitize Fire Area using False Colour Composite (D) and Active Fire
- Extract Hourly FRP from GFAS
  - 06:00 → Sentinel-5P overpass time
- Finding all Active Fire pixels within Fire Polygon
- $\geq 50\%$  same biome → Dominant Biome






# aFREM Emission Coefficient ( $EC_{CO}^{biome}$ ) Calculation

8

$EC_{CO}^{biome}$  relates the rate of release of CO (g.s<sup>-1</sup>) to the rate of release of FRE (i.e. FRP, MW)

Grouping plumes by biome, can generate Biome Specific Emission Coefficients for CO,  $EC_{CO}^{biome}$







# CO Emission Estimation – Deciduous Needleleaf

9

aFREM produces similar CO Emissions for “Deciduous Needleleaf” forest fires to existing inventories

- ~86% w.r.t GFAS, ~88% w.r.t FEER





# CO Emission Estimation – Evergreen Needleleaf

10

aFREM produces similar CO Emissions for “Evergreen Needleleaf” forest fires to existing inventories

- ~74% w.r.t GFAS, ~60% w.r.t FEER





# CO Emission Estimation – Grassland (inc. high latitude Tundra)

11

aFREM generates CO Emissions lower for “High Latitude Grassland Fires” than existing inventories

- ~26% w.r.t GFAS, ~39% w.r.t FEER



# Summary and Next Steps

Introduced the aFREM approach, directly linking CO Emission rate to FRP for polar orbiters

- Digitizing Plumes and Fire Activity using VIIRS
- Calculating  $EC_{CO}^{biome}$  for Evergreen and Deciduous Needleleaf Forest, and Grasslands (Tundra)
- Estimates of Daily CO emissions compared those of FEER and GFAS

## Next Steps

- Completing Analysis of COI23 (193 additional plume observations)
- Full Timeseries Analysis (2019 – 2023)
- Impact of peat vs non-peat on  $EC_{CO}^{biome}$
- Full Carbon Emissions estimation from the CO values
- Monthly Comparison with GFED



Abstract



KCL Research  
Group

# Thanks for listening!



**National Centre for  
Earth Observation**  
NATURAL ENVIRONMENT RESEARCH COUNCIL

Will Maslanka | William.maslanka@kcl.ac.uk | NCEO King's College London

**KING'S**  
College  
LONDON