

Safe Envelopes for High-Dimension Reachability by Linear Hopf Solutions with Antagonistic Error

Will Sharpless

University of California, San Diego

November 6, 2024



Contents

- 1 Preliminaries - Hamilton-Jacobi Reachability
- 2 Preliminaries - Hopf Formula
- 3 Safe Envelopes by Linear Hopf Solutions
- 4 Limitations
- 5 Methods for Tighter Envelopes

Preliminaries - Hamilton-Jacobi Reachability

System

Consider a system,

$$\dot{x} = f_x(x, \tau) + h_1(x, \tau)u + h_2(x, \tau)d \triangleq f(x, u, d, \tau), \quad (1)$$

where $\tau \in [t, T]$, u and d are drawn from convex sets $\mathcal{U} \subset \mathbb{R}^{n_u}$, $\mathcal{D} \subset \mathbb{R}^{n_d}$, and (23) is Lipschitz continuous in (x, u, d) and continuous in t .

Differential Game Problem Statement

Suppose a game is defined by,

$$P(x, u(\cdot), d(\cdot), t) = J(x(T)) + \int_t^T L(u(\tau), d(\tau))d\tau. \quad (2)$$

where Player I (actions u) and Player II (actions d) compete by optimizing (2), in a zero-sum fashion.

Differential Game Problem Statement

Suppose a game is defined by,

$$P(x, u(\cdot), d(\cdot), t) = J(x(T)) + \int_t^T L(u(\tau), d(\tau))d\tau. \quad (2)$$

where Player I (actions u) and Player II (actions d) compete by optimizing (2), in a zero-sum fashion.

Reach game \triangleq (Player I, Player II) have objectives $(\min_u P, \max_d P)$

Avoid game \triangleq (Player I, Player II) have objectives $(\max_u P, \min_d P)$

(Note, we use \cdot, \cdot^- to define analogous *Reach & Avoid* game objects resp.)

Differential Game Problem Statement

Given a target set $\mathcal{T} \in \mathbb{K}\mathbb{R}^{n_x}$ to reach or avoid, let the terminal cost $J : \mathbb{R}^{n_x} \mapsto \mathbb{R}$ be defined by

$$\begin{cases} J(x) < 0 & \text{for } x \in \mathcal{T} \setminus \partial\mathcal{T} \\ J(x) = 0 & \text{for } x \in \partial\mathcal{T} \\ J(x) > 0 & \text{for } x \notin \mathcal{T} \end{cases}$$

such that it is convex and lower semicontinuous.

Differential Game Problem Statement

Given a target set $\mathcal{T} \in \mathbb{KR}^{n_x}$ to reach or avoid, let the terminal cost $J : \mathbb{R}^{n_x} \mapsto \mathbb{R}$ be defined by

$$\begin{cases} J(x) < 0 & \text{for } x \in \mathcal{T} \setminus \partial\mathcal{T} \\ J(x) = 0 & \text{for } x \in \partial\mathcal{T} \\ J(x) > 0 & \text{for } x \notin \mathcal{T} \end{cases}$$

such that it is convex and lower semicontinuous.

Running costs $L, L^- : \mathbb{R}^{n_u} \times \mathbb{R}^{n_d} \mapsto \mathbb{R}$ solely constrain the inputs,

$$\begin{aligned} L(u, d) &= \mathcal{I}_{\mathcal{U}}(u) - \mathcal{I}_{\mathcal{D}}(d), \\ L^-(u, d) &= \mathcal{I}_{\mathcal{D}}(d) - \mathcal{I}_{\mathcal{U}}(u), \end{aligned} \tag{3}$$

where $\mathcal{I}_{\mathcal{C}}(c) = \{0 \text{ if } c \in \mathcal{C}, +\infty \text{ else}\}$.

Game Cost and Value Properties

For trajectory $x(\cdot)$ satisfying the dynamics, if $u(\cdot) \in \mathbb{U} : [t, T] \mapsto \mathcal{U}$,
 $d(\cdot) \in \mathbb{D} : [t, T] \mapsto \mathcal{D}$,

$$P(x, u(\cdot), d(\cdot), t) \leq 0 \iff x(T) \in \mathcal{T}. \quad (4)$$

Game Cost and Value Properties

For trajectory $x(\cdot)$ satisfying the dynamics, if $u(\cdot) \in \mathbb{U} : [t, T] \mapsto \mathcal{U}$,
 $d(\cdot) \in \mathbb{D} : [t, T] \mapsto \mathcal{D}$,

$$P(x, u(\cdot), d(\cdot), t) \leq 0 \iff x(T) \in \mathcal{T}. \quad (4)$$

Consider the optimal values of the games $V, V^- : \mathbb{R}^{n_x} \times (-\infty, T] \mapsto \mathbb{R}$ defined

$$\begin{aligned} V(x, t) &= \sup_{d(\cdot) \in \Gamma(t)} \inf_{u(\cdot) \in \mathbb{U}} P(x, u(\cdot), d(\cdot), t), \\ V^-(x, t) &= \inf_{d(\cdot) \in \Gamma(t)} \sup_{u(\cdot) \in \mathbb{U}} P(x, u(\cdot), d(\cdot), t). \end{aligned} \quad (5)$$

Game Cost and Value Properties

For trajectory $x(\cdot)$ satisfying the dynamics, if $u(\cdot) \in \mathbb{U} : [t, T] \mapsto \mathcal{U}$,
 $d(\cdot) \in \mathbb{D} : [t, T] \mapsto \mathcal{D}$,

$$P(x, u(\cdot), d(\cdot), t) \leq 0 \iff x(T) \in \mathcal{T}. \quad (4)$$

Consider the optimal values of the games $V, V^- : \mathbb{R}^{n_x} \times (-\infty, T] \mapsto \mathbb{R}$ defined

$$\begin{aligned} V(x, t) &= \sup_{d(\cdot) \in \Gamma(t)} \inf_{u(\cdot) \in \mathbb{U}} P(x, u(\cdot), d(\cdot), t), \\ V^-(x, t) &= \inf_{d(\cdot) \in \Gamma(t)} \sup_{u(\cdot) \in \mathbb{U}} P(x, u(\cdot), d(\cdot), t). \end{aligned} \quad (5)$$

These analogously satisfy,

$$\begin{aligned} V(x, t) \leq 0 &\iff x \in \mathcal{R}(\mathcal{T}, t), \\ V^-(x, t) \leq 0 &\iff x \in \mathcal{R}^-(\mathcal{T}, t). \end{aligned} \quad (6)$$

Reach and Avoid Sets

where $\mathcal{R}(\mathcal{T}, t), \mathcal{R}^-(\mathcal{T}, t) \in \mathbb{KR}^{n_x}$ are the *Reach* and *Avoid* sets resp., formally given by

$$\begin{aligned}\mathcal{R}(\mathcal{T}, t) &\triangleq \{x \mid \exists u(\cdot) \in \mathbb{U} \ \forall d(\cdot) \in \mathbb{D} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}, \\ \mathcal{R}^-(\mathcal{T}, t) &\triangleq \{x \mid \exists d(\cdot) \in \mathbb{D} \ \forall u(\cdot) \in \mathbb{U} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}. \end{aligned} \tag{7}$$

Reachability versus Feasibility

where $\mathcal{R}(\mathcal{T}, t), \mathcal{R}^-(\mathcal{T}, t) \in \mathbb{KR}^{n_x}$ are the *Reach* and *Avoid* sets resp., formally given by

$$\begin{aligned}\mathcal{R}(\mathcal{T}, t) &\triangleq \{x \mid \exists u(\cdot) \in \mathbb{U} \ \forall d(\cdot) \in \mathbb{D} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}, \\ \mathcal{R}^-(\mathcal{T}, t) &\triangleq \{x \mid \exists d(\cdot) \in \mathbb{D} \ \forall u(\cdot) \in \mathbb{U} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}.\end{aligned}\tag{8}$$

In contrast, the feasible set $\mathcal{S} \in \mathbb{KR}^{n_x}$ is given by,

$$\mathcal{S}(\mathcal{T}, t) \triangleq \{x \mid \exists u(\cdot) \in \mathbb{U} \ \exists d(\cdot) \in \mathbb{D} \text{ s.t.} \\ x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\},\tag{9}$$

and captures all trajectories which might arrive at the target, irrespective of player objectives.

Reachability versus Feasibility

where $\mathcal{R}(\mathcal{T}, t), \mathcal{R}^-(\mathcal{T}, t) \in \mathbb{KR}^{n_x}$ are the *Reach* and *Avoid* sets resp., formally given by

$$\begin{aligned}\mathcal{R}(\mathcal{T}, t) &\triangleq \{x \mid \exists u(\cdot) \in \mathbb{U} \ \forall d(\cdot) \in \mathbb{D} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}, \\ \mathcal{R}^-(\mathcal{T}, t) &\triangleq \{x \mid \exists d(\cdot) \in \mathbb{D} \ \forall u(\cdot) \in \mathbb{U} \text{ s.t.} \\ &\quad x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\}.\end{aligned}\tag{8}$$

In contrast, the feasible set $\mathcal{S} \in \mathbb{KR}^{n_x}$ is given by,

$$\mathcal{S}(\mathcal{T}, t) \triangleq \{x \mid \exists u(\cdot) \in \mathbb{U} \ \exists d(\cdot) \in \mathbb{D} \text{ s.t.} \\ x(\cdot) \text{ satisfies (23)} \wedge x(T) \in \mathcal{T}\},\tag{9}$$

and captures all trajectories which might arrive at the target, irrespective of player objectives. This set is useful in this work because $\mathcal{S} \supset \mathcal{R}, \mathcal{R}^-$.

Hamilton-Jacobi-Bellman PDE

Theorem (Evans 84, (1))

The value function V defined in (5) is the viscosity solution to,

$$\begin{aligned} \frac{\partial V}{\partial t} + H(x, \nabla_x V, t) &= 0 && \text{on } \mathbb{R}^{n_x} \times [t, T], \\ V(x, T) &= J(x(T)) && \text{on } \mathbb{R}^{n_x} \end{aligned} \tag{10}$$

where the Hamiltonian $H : \mathbb{R}^{n_x} \times \mathbb{R}^{n_x} \times [t, T] \mapsto \mathbb{R}$ is

$$H(x, p, t) = \min_{u \in \mathcal{U}} \max_{d \in \mathcal{D}} p \cdot f(x, u, d, t). \tag{11}$$

Hamilton-Jacobi-Bellman PDE

Theorem (Evans 84, (1))

The value function V defined in (5) is the viscosity solution to,

$$\begin{aligned} \frac{\partial V}{\partial t} + H(x, \nabla_x V, t) &= 0 && \text{on } \mathbb{R}^{n_x} \times [t, T], \\ V(x, T) &= J(x(T)) && \text{on } \mathbb{R}^{n_x} \end{aligned} \tag{10}$$

where the Hamiltonian $H : \mathbb{R}^{n_x} \times \mathbb{R}^{n_x} \times [t, T] \mapsto \mathbb{R}$ is

$$H(x, p, t) = \min_{u \in \mathcal{U}} \max_{d \in \mathcal{D}} p \cdot f(x, u, d, t). \tag{11}$$

Additionally, the optimal control is given by,

$$u^*(t) = \arg \min_{u \in \mathcal{U}} \nabla_x V(x, t) \cdot h_1(t)u. \tag{12}$$

Hamilton-Jacobi-Bellman PDE

Theorem (Evans 84, (1))

The value function V defined in (5) is the viscosity solution to,

$$\begin{aligned} \frac{\partial V}{\partial t} + H(x, \nabla_x V, t) &= 0 && \text{on } \mathbb{R}^{n_x} \times [t, T], \\ V(x, T) &= J(x(T)) && \text{on } \mathbb{R}^{n_x} \end{aligned} \tag{10}$$

where the Hamiltonian $H : \mathbb{R}^{n_x} \times \mathbb{R}^{n_x} \times [t, T] \mapsto \mathbb{R}$ is

$$H(x, p, t) = \min_{u \in \mathcal{U}} \max_{d \in \mathcal{D}} p \cdot f(x, u, d, t). \tag{11}$$

Additionally, the optimal control is given by,

$$u^*(t) = \arg \min_{u \in \mathcal{U}} \nabla_x V(x, t) \cdot h_1(t) u. \tag{12}$$

This identically applies to V^- with $H^-(x, p, t) = \max_u \min_d p \cdot f(x, u, d, t)$.

Preliminaries - Hopf Formula

Forwards to Backwards change

For simplicity, we change the previous *final* value problem into an *initial* value problem by $\phi(x, t) \triangleq V(x, T - t)$,

Forwards to Backwards change

For simplicity, we change the previous *final* value problem into an *initial* value problem by $\phi(x, t) \triangleq V(x, T - t)$, s.t.

$$\begin{aligned} -\frac{\partial \phi}{\partial t} + H(x, \nabla_x \phi, t) &= 0 && \text{on } \mathbb{R}^{n_x} \times [0, t], \\ \phi(x, 0) &= J(x) && \text{on } \mathbb{R}^{n_x} \end{aligned} \tag{13}$$

with Hamiltonian

$$H(x, p, t) = \max_{u \in \mathcal{U}} \min_{d \in \mathcal{D}} -p \cdot f(x, u, d, T - t). \tag{14}$$

Forwards to Backwards change

For simplicity, we change the previous *final* value problem into an *initial* value problem by $\phi(x, t) \triangleq V(x, T - t)$, s.t.

$$\begin{aligned} -\frac{\partial \phi}{\partial t} + H(x, \nabla_x \phi, t) &= 0 && \text{on } \mathbb{R}^{n_x} \times [0, t], \\ \phi(x, 0) &= J(x) && \text{on } \mathbb{R}^{n_x} \end{aligned} \tag{13}$$

with Hamiltonian

$$H(x, p, t) = \max_{u \in \mathcal{U}} \min_{d \in \mathcal{D}} -p \cdot f(x, u, d, T - t). \tag{14}$$

For systems lacking state dependence $f(x, u, d, t) \equiv f(u, d, t)$, H also lacks state dependence $H(x, p, t) \equiv H(p, t)$. In this setting, there is a solution to the PDE ...

The Hopf Formula

Theorem (Rublev 00, (2))

Assume that $J(x)$ is convex and that $H(t, p)$ is pseudoconvex in p and satisfies (B.i-B.iii) in [(2)], then the minimax-viscosity solution of (13) is given by

$$\phi(x, t) = - \min_{p \in \mathbb{R}^{n_x}} \left\{ J^*(p) - x \cdot p + \int_0^t H(p, \tau) d\tau \right\} \quad (15)$$

where $J^*(p) : \mathbb{R}^{n_x} \rightarrow \mathbb{R} \cup \{+\infty\}$ is the Fenchel-Legendre transform (i.e. convex-conjugate) of a convex, proper, lower semicontinuous function $J : \mathbb{R}^{n_x} \rightarrow \mathbb{R}$ defined by

$$J^*(p) = \sup_{x \in \mathbb{R}^{n_x}} \{p \cdot x - J(x)\}. \quad (16)$$

The Hopf Formula

Theorem (Rublev 00, (2))

Assume that $J(x)$ is convex and that $H(t, p)$ is pseudoconvex in p and satisfies (B.i-B.iii) in [(2)], then the minimax-viscosity solution of (13) is given by

$$\phi(x, t) = - \min_{p \in \mathbb{R}^{n_x}} \left\{ J^*(p) - x \cdot p + \int_0^t H(p, \tau) d\tau \right\} \quad (15)$$

where $J^*(p) : \mathbb{R}^{n_x} \rightarrow \mathbb{R} \cup \{+\infty\}$ is the Fenchel-Legendre transform (i.e. convex-conjugate) of a convex, proper, lower semicontinuous function $J : \mathbb{R}^{n_x} \rightarrow \mathbb{R}$ defined by

$$J^*(p) = \sup_{x \in \mathbb{R}^{n_x}} \{p \cdot x - J(x)\}. \quad (16)$$

Remark. The minimax-viscosity and viscosity solutions coincide when $H(p, \tau)$ is convex in p for $\tau \in [0, t]$.

In an *Avoid* Game, minimax viscosity is a safe estimate

Remark. The minimax-viscosity solution given by the Hopf formula is always less than or equal to the viscosity solution [(3)].

Hence, in an avoid problem, ϕ^- (the Avoid Hopf value) guarantees a conservative lower bound of V^- for any H^- satisfying Thm. 2.¹

¹This yields a potentially non-convex objective, thus one should employ ADMM, known to solve similar non-convex problems [(4)].

Hopf with Linear Systems

Requiring “state-independence” is restrictive. It happens that, however, any linear system,

$$\dot{x} = A(\tau)x + B_1(\tau)u + B_2(\tau)d \quad (17)$$

may be transformed with the mapping $z(\tau) \triangleq \Phi(\tau)x(\tau)$ where $\Phi(\tau)$ defined by the *fundamental matrix* $\dot{\Phi} = -\Phi(\tau)A(\tau)$, $\Phi(0) = I$. If $A(\tau) \equiv A$ then $\Phi(\tau) = \exp(-\tau A)$.

Hopf with Linear Systems

Requiring “state-independence” is restrictive. It happens that, however, any linear system,

$$\dot{x} = A(\tau)x + B_1(\tau)u + B_2(\tau)d \quad (17)$$

may be transformed with the mapping $z(\tau) \triangleq \Phi(\tau)x(\tau)$ where $\Phi(\tau)$ defined by the *fundamental matrix* $\dot{\Phi} = -\Phi(\tau)A(\tau)$, $\Phi(0) = I$. If $A(\tau) \equiv A$ then $\Phi(\tau) = \exp(-\tau A)$.

In the \mathcal{Z} -space,

$$\begin{aligned} z(\tau) = \Phi(\tau)x(\tau) &\implies \dot{z} = \dot{\Phi}(\tau)x + \Phi(\tau)\dot{x} \\ &= \Phi(\tau)(-A(\tau)x) + \Phi(\tau)\dot{x} \\ &= \Phi(\tau)(B_1(\tau)u + B_2(\tau)d) \end{aligned} \quad (18)$$

and thus may be solved by the Hopf formula.

Hopf, Linear Systems, \mathcal{Z} -space

After the change(s) of variables, the *Reach* \mathcal{Z} -Hamiltonian takes the form,

$$\begin{aligned}
 H_{\mathcal{Z}}(p, t) &= \max_{u \in \mathcal{U}} \min_{d \in \mathcal{D}} -p \cdot \Phi(T-t) (B_1(T-t)u + B_2(T-t)d), \\
 &= \max_{u \in \mathcal{U}} \{p \cdot R_1(\tau)u\} - \max_{d \in \mathcal{D}} \{p \cdot R_2(\tau)d\} \\
 &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(\tau)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(\tau)p)
 \end{aligned} \tag{19}$$

where $R_i(\tau) \triangleq -\Phi(T-t)B_i(T-t)$.

Hopf, Linear Systems, \mathcal{Z} -space

After the change(s) of variables, the *Reach* \mathcal{Z} -Hamiltonian takes the form,

$$\begin{aligned}
 H_{\mathcal{Z}}(p, t) &= \max_{u \in \mathcal{U}} \min_{d \in \mathcal{D}} -p \cdot \Phi(T-t)(B_1(T-t)u + B_2(T-t)d), \\
 &= \max_{u \in \mathcal{U}} \{p \cdot R_1(\tau)u\} - \max_{d \in \mathcal{D}} \{p \cdot R_2(\tau)d\} \\
 &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(\tau)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(\tau)p)
 \end{aligned} \tag{19}$$

where $R_i(\tau) \triangleq -\Phi(T-t)B_i(T-t)$.

If \mathcal{U} and \mathcal{D} are defined by norms, then $\mathcal{I}_{\mathcal{U}}^*$ and $\mathcal{I}_{\mathcal{D}}^*$ are the corresponding dual norms. This is useful for fast computation and proving the convexity of H (*but may be ignored, if confusing*).

Safe Envelopes by Linear Hopf Solutions

Running Example (Van der Pol)

To illustrate theoretical results, we consider the system,

$$\dot{x} = \begin{bmatrix} x_1 \\ \mu(1 - x_1^2)x_2 - x_1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix}(u + d) \quad (20)$$

with $\mu = 1$, and inputs

$$u \in \mathcal{U} \triangleq \{|u| \leq 1\}, \quad d \in \mathcal{D} \triangleq \{|d| \leq 1/2\}, \quad (21)$$

and a target set defined by,

$$\mathcal{T} \triangleq \{||(x - c_{\mathcal{T}})||_{Q_{\mathcal{T}}} \leq 1\}, \quad (22)$$

for parameters $c_{\mathcal{T}} \in \mathbb{R}^{n_x}$, $Q_{\mathcal{T}} \in \mathbb{S}_{++}^{n_x}$. Note,

$$||(x - c_{\mathcal{T}})||_{Q_{\mathcal{T}}} = ||(x - c_{\mathcal{T}})^{\top} Q_{\mathcal{T}}^{-1} (x - c_{\mathcal{T}})||_2.$$

Comparing Nonlinear and Linear Systems

Recall, in general, the system takes the form,

$$\dot{x} = f_x(x, \tau) + h_1(x, \tau)u + h_2(x, \tau)d \triangleq f(x, u, d, \tau). \quad (23)$$

Comparing Nonlinear and Linear Systems

Recall, in general, the system takes the form,

$$\dot{x} = f_x(x, \tau) + h_1(x, \tau)u + h_2(x, \tau)d \triangleq f(x, u, d, \tau). \quad (23)$$

Consider a continuous, linear function

$$l(x, u, d, \tau) \triangleq A(\tau)x + B_1(\tau)u + B_2(\tau)d, \quad (24)$$

which will serve as an approximate model of (23) which we may solved with Hopf.

Comparing Nonlinear and Linear Systems

Recall, in general, the system takes the form,

$$\dot{x} = f_x(x, \tau) + h_1(x, \tau)u + h_2(x, \tau)d \triangleq f(x, u, d, \tau). \quad (23)$$

Consider a continuous, linear function

$$l(x, u, d, \tau) \triangleq A(\tau)x + B_1(\tau)u + B_2(\tau)d, \quad (24)$$

which will serve as an approximate model of (23) which we may solve with Hopf.

Across \mathbb{R}^{n_x} , their difference may be unbounded. But, in the feasible set \mathcal{S} , since $(\mathcal{T} \times \mathcal{U} \times \mathcal{D} \times [t, T])$ is compact, by Filippov's Lemma (5),

$$\delta^* \triangleq \max_{x \in \mathcal{S}} \|(f - l)(x, u, d, \tau)\| \quad (u, d) \in \mathcal{U} \times \mathcal{D} \quad (25)$$

is finite.

Comparing Nonlinear and Linear Systems

Hence, for $x \in \mathcal{S} \subset \mathbb{R}^{n_x}$,

$$\dot{x} = f(x, u, d) = l(x, u, d) + \varepsilon \quad ||\varepsilon|| \leq \delta^*, \quad (26)$$

and

$$H(x, p, t) = H_l(x, p, t) + p \cdot \varepsilon, \quad (27)$$

where $H_l(x, p, t) \triangleq \min_{u \in \mathcal{U}} \max_{d \in \mathcal{D}} \{p \cdot l(x, u, d)\}$.

Comparing Nonlinear and Linear Systems

Hence, for $x \in \mathcal{S} \subset \mathbb{R}^{n_x}$,

$$\dot{x} = f(x, u, d) = l(x, u, d) + \varepsilon \quad ||\varepsilon|| \leq \delta^*, \quad (26)$$

and

$$H(x, p, t) = H_l(x, p, t) + p \cdot \varepsilon, \quad (27)$$

where $H_l(x, p, t) \triangleq \min_{u \in \mathcal{U}} \max_{d \in \mathcal{D}} \{p \cdot l(x, u, d)\}$.

Although ε is nonlinear and state-dependent, we upper-bound this relationship to use of the Hopf formula, assuming the worst for safety.

Game with Antagonistic Error

Consider another antagonistic player with actions

$$\varepsilon \in \mathcal{E} \triangleq \{\varepsilon \in \mathcal{X} \mid \|\varepsilon\| \leq \delta^*\}. \quad (28)$$

Game with Antagonistic Error

Consider another antagonistic player with actions

$$\varepsilon \in \mathcal{E} \triangleq \{\varepsilon \in \mathcal{X} \mid \|\varepsilon\| \leq \delta^*\}. \quad (28)$$

The corresponding Hamiltonians for the *Reach* and *Avoid* games resp. are,

$$H_{\delta^*}(x, p, t) \triangleq H_l(x, p, t) + \max_{\varepsilon \in \mathcal{E}} p \cdot \varepsilon, \quad (29)$$

$$H_{\delta^*}^-(x, p, t) \triangleq H_l(x, p, t) + \min_{\varepsilon \in \mathcal{E}} p \cdot \varepsilon. \quad (30)$$

Remark. $H_{\delta^*} = \max_{\varepsilon \in \mathcal{E}} H$ and hence H_{δ^*} is the *envelope* of the Hamiltonian H in the parlance of optimization and PDEs (6).

Safe Upper & Lower Values for *Reach & Avoid* games

Lemma

For $V(x, t)$ defined in (5), arising from target \mathcal{T} and dynamics f with players confined to \mathcal{U}, \mathcal{D} , then

$$V(x, \tau) \leq V_{\delta^*}(x, \tau) \quad x \in \mathcal{S}, \tau \in [t, T] \quad (31)$$

where $V_{\delta^*}(x, t)$ is defined as the value of the Reach game with dynamics l and an additional antagonistic player, confined to \mathcal{E} , s.t. $V_{\delta^*}(x, t)$ is the solution to the PDE in (10) with Hamiltonian H_{δ^*} .

Safe Upper & Lower Values for *Reach & Avoid* games

Lemma

For $V(x, t)$ defined in (5), arising from target \mathcal{T} and dynamics f with players confined to \mathcal{U}, \mathcal{D} , then

$$V(x, \tau) \leq V_{\delta^*}(x, \tau) \quad x \in \mathcal{S}, \tau \in [t, T] \quad (31)$$

where $V_{\delta^*}(x, t)$ is defined as the value of the Reach game with dynamics l and an additional antagonistic player, confined to \mathcal{E} , s.t. $V_{\delta^*}(x, t)$ is the solution to the PDE in (10) with Hamiltonian H_{δ^*} .

In the Avoid game, analogously

$$V(x, \tau) \geq V_{\delta^*}^-(x, \tau) \quad x \in \mathcal{S}, \tau \in [t, T] \quad (32)$$

s.t. $V_{\delta^*}^-(x, t)$ is the solution to the PDE in (10) with Hamiltonian $H_{\delta^*}^-$.

Safe Upper & Lower Values for *Reach & Avoid* games

Proof.

From [(1)], the viscosity solution corresponds to

$$V(x, T) = \inf_x \left\{ J(x(T)) + \int_t^T \mathcal{L}(x(\tau), f(x(\tau))) d\tau \right\}$$

$$V_{\delta^*}(x, T) = \inf_x \left\{ J(x(T)) + \int_t^T \mathcal{L}_{\delta^*}(x(\tau), f(x(\tau))) d\tau \right\}$$

where $\mathcal{L}, \mathcal{L}_{\delta^*}$ are the Lagrangians of the system,

$$\mathcal{L} = \inf_x \{x \cdot p + H(x, p)\}, \quad \mathcal{L}_{\delta^*} = \inf_x \{x \cdot p + H_{\delta^*}(x, p)\}.$$

By definition $H \leq H_{\delta^*}$, so $\forall p, \mathcal{L} \leq x \cdot p + H(x, p) \leq x \cdot p + H_{\delta^*}(x, p)$, and thus $\mathcal{L} \leq \mathcal{L}_{\delta^*}$.

Safe Upper & Lower Values for *Reach & Avoid* games

Proof.

From [(1)], the viscosity solution corresponds to

$$V(x, T) = \inf_x \left\{ J(x(T)) + \int_t^T \mathcal{L}(x(\tau), f(x(\tau))) d\tau \right\}$$

$$V_{\delta^*}(x, T) = \inf_x \left\{ J(x(T)) + \int_t^T \mathcal{L}_{\delta^*}(x(\tau), f(x(\tau))) d\tau \right\}$$

and thus $\mathcal{L} \leq \mathcal{L}_{\delta^*}$. Then,

$$\begin{aligned} \forall x, V(x, t) &\leq J(x(T)) + \int_t^T \mathcal{L}(x(\tau), f(x(\tau))) \\ &\leq J(x(T)) + \int_t^T \mathcal{L}_{\delta^*}(x(\tau), f(x(\tau))). \end{aligned}$$

Hence, $V(x, t) \leq V_{\delta^*}(x, t)$.



Subsets & Supersets for *Reach & Avoid* games

Corollary

In the Reach game, at any time $\tau \in [t, T]$,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \subset \mathcal{R}(\mathcal{T}, \tau) \quad (33)$$

and for any $x \in \mathcal{R}_{\delta^}(\mathcal{T}, \tau)$, $u_{\delta^*}^*(\cdot)$ derived from (12) with V_{δ^*} will yield $x(T) \in \mathcal{T}$ under the true, nonlinear dynamics (23) despite any disturbances $d(\cdot) \in \mathbb{D}$.*

Subsets & Supersets for *Reach & Avoid* games

Corollary

In the Reach game, at any time $\tau \in [t, T]$,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \subset \mathcal{R}(\mathcal{T}, \tau) \quad (33)$$

and for any $x \in \mathcal{R}_{\delta^}(\mathcal{T}, \tau)$, $u_{\delta^*}^*(\cdot)$ derived from (12) with V_{δ^*} will yield $x(T) \in \mathcal{T}$ under the true, nonlinear dynamics (23) despite any disturbances $d(\cdot) \in \mathbb{D}$.*

Analogously, for the Avoid game,

$$\mathcal{R}_{\delta^*}^-(\mathcal{T}, \tau) \supset \mathcal{R}^-(\mathcal{T}, \tau) \quad (34)$$

and $\forall x \notin \mathcal{R}_{\delta^}^-(\mathcal{T}, \tau)$, the trajectory $x(\cdot)$ arising from $u_{\delta^*}^*(\cdot)$ solved from (12) with $V_{\delta^*}^-$ will yield $x(T) \notin \mathcal{T}$ under the true, nonlinear dynamics despite any $d(\cdot) \in \mathbb{D}$.*

Subsets & Supersets for *Reach & Avoid* games

Proof.

The crucial property of the game value (6) guarantees,

$$x \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \iff V_{\delta^*}(x, \tau) \leq 0 \implies V(x, \tau) \leq 0 \iff x \in \mathcal{R}(\mathcal{T}, \tau).$$

Subsets & Supersets for *Reach & Avoid* games

Proof.

The crucial property of the game value (6) guarantees,

$$x \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \iff V_{\delta^*}(x, \tau) \leq 0 \implies V(x, \tau) \leq 0 \iff x \in \mathcal{R}(\mathcal{T}, \tau).$$

By definition $x \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau)$ implies

$$\begin{aligned} &\exists u_{\delta^*}^*(\cdot) \in \mathbb{U}, \forall \varepsilon(\cdot) \in \mathcal{E}, \forall d(\cdot) \in \mathbb{D}, \text{ s.t.} \\ &x(T) \in \mathcal{T} \text{ and } \dot{x} = l(x, u_{\delta^*}^*, d) + \varepsilon. \end{aligned}$$

Subsets & Supersets for *Reach & Avoid* games

Proof.

The crucial property of the game value (6) guarantees,

$$x \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \iff V_{\delta^*}(x, \tau) \leq 0 \implies V(x, \tau) \leq 0 \iff x \in \mathcal{R}(\mathcal{T}, \tau).$$

By definition $x \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau)$ implies

$$\begin{aligned} \exists u_{\delta^*}^*(\cdot) \in \mathbb{U}, \forall \varepsilon(\cdot) \in \mathcal{E}, \forall d(\cdot) \in \mathbb{D}, \text{ s.t.} \\ x(T) \in \mathcal{T} \text{ and } \dot{x} = l(x, u_{\delta^*}^*, d) + \varepsilon. \end{aligned}$$

Assuming that \mathcal{S} is sufficiently large, for the trajectory $x(\cdot)$ arising from $u_{\delta^*}^*(\cdot)$,

$$\forall \tau \in [t, T], x(\tau) \in \mathcal{R}_{\delta^*}(\mathcal{T}, \tau) \subset \mathcal{R}(\mathcal{T}, \tau) \subset \mathcal{S},$$

hence, $\exists \varepsilon(\cdot) \subset \mathcal{E}$ s.t.

$$\dot{x} = l(x, u_{\delta^*}^*, d) + \varepsilon = f(x, u_{\delta^*}^*, d).$$



Safe Envelopes in the VanderPol System




Figure: For a target \mathcal{T} (black), true reachable sets $\mathcal{R}/\mathcal{R}^-$ (gold) are solved for a *Reach* and *Avoid* game resp. at $t = [0.26]$. The corresponding reachable sets of the safe envelope $\mathcal{R}_{\delta^*}/\mathcal{R}_{\delta^*}^-$ (green) based on the constant Taylor-series linearization error are also solved. The feasible sets $\hat{\mathcal{S}}$ (blue) are over-approximated via DI methods (7) (8).

Limitations

Convexity of the δ^* -Hamiltonian

To ensure the δ^* -minimax solution is not below the true value in a *Reach* game, convexity of the δ^* -Hamiltonian is required.

Convexity of the δ^* -Hamiltonian

To ensure the δ^* -minimax solution is not below the true value in a *Reach* game, convexity of the δ^* -Hamiltonian is required.

Assume the linear Hamiltonians H_l, H_l^- , are convex. These are transformed by the addition of error s.t.,

$$\begin{aligned} H_{l,\mathcal{Z}} &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(t)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}} = H_{l,\mathcal{Z}} - \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p) \\ H_{l,\mathcal{Z}}^- &= \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) - \mathcal{I}_{\mathcal{U}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}}^- = H_{l,\mathcal{Z}}^- + \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p). \end{aligned} \quad (35)$$

Convexity of the δ^* -Hamiltonian

To ensure the δ^* -minimax solution is not below the true value in a *Reach* game, convexity of the δ^* -Hamiltonian is required.

Assume the linear Hamiltonians H_l, H_l^- , are convex. These are transformed by the addition of error s.t.,

$$\begin{aligned} H_{l,\mathcal{Z}} &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(t)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}} = H_{l,\mathcal{Z}} - \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p) \\ H_{l,\mathcal{Z}}^- &= \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) - \mathcal{I}_{\mathcal{U}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}}^- = H_{l,\mathcal{Z}}^- + \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p). \end{aligned} \quad (35)$$

Therefore, in the two-player

- ➊ *Reach* game, H_{δ^*} will not necessarily be convex and must be checked for a specific definition of \mathcal{U}, \mathcal{D} and observed error \mathcal{E} .

Convexity of the δ^* -Hamiltonian

To ensure the δ^* -minimax solution is not below the true value in a *Reach* game, convexity of the δ^* -Hamiltonian is required.

Assume the linear Hamiltonians H_l, H_l^- , are convex. These are transformed by the addition of error s.t.,

$$\begin{aligned} H_{l,\mathcal{Z}} &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(t)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}} = H_{l,\mathcal{Z}} - \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p) \\ H_{l,\mathcal{Z}}^- &= \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) - \mathcal{I}_{\mathcal{U}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}}^- = H_{l,\mathcal{Z}}^- + \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p). \end{aligned} \quad (35)$$

Therefore, in the two-player

- ① *Reach* game, H_{δ^*} will not necessarily be convex and must be checked for a specific definition of \mathcal{U}, \mathcal{D} and observed error \mathcal{E} .
- ② *Avoid* game, $H_{\delta^*}^-$ will be convex for any $\delta^* \geq 0$.

Convexity of the δ^* -Hamiltonian

To ensure the δ^* -minimax solution is not below the true value in a *Reach* game, convexity of the δ^* -Hamiltonian is required.

Assume the linear Hamiltonians H_l, H_l^- , are convex. These are transformed by the addition of error s.t.,

$$\begin{aligned} H_{l,\mathcal{Z}} &= \mathcal{I}_{\mathcal{U}}^*(R_1^\top(t)p) - \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}} = H_{l,\mathcal{Z}} - \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p) \\ H_{l,\mathcal{Z}}^- &= \mathcal{I}_{\mathcal{D}}^*(R_2^\top(t)p) - \mathcal{I}_{\mathcal{U}}^*(R_2^\top(t)p) \Rightarrow H_{\delta^*,\mathcal{Z}}^- = H_{l,\mathcal{Z}}^- + \mathcal{I}_{\mathcal{E}}^*(-\Phi(T-t)^{-\top}p). \end{aligned} \quad (35)$$

Therefore, in the two-player

- ① *Reach* game, H_{δ^*} will not necessarily be convex and must be checked for a specific definition of \mathcal{U}, \mathcal{D} and observed error \mathcal{E} .
- ② *Avoid* game, $H_{\delta^*}^-$ will be convex for any $\delta^* \geq 0$.

Nonlinear \dot{x}_i must have control in 2P Reach

Corollary

Any state x_i , with evolution $\dot{x}_i = f_i(x, u, d)$ s.t. $\delta_i^* = |f - l_i| > 0$ and $\forall u \in \mathcal{U}, f_i(x, u, d) = f_i(x, d)$, will have a non-convex hamiltonian.

Thus, in a two-player Reach game, any state with nonlinear f_i and thus $\delta_i^* > 0$ must have at least some control, or the guarantees of Lemma 3 will not hold.

Nonlinear \dot{x}_i must have control in 2P Reach

Corollary

Any state x_i , with evolution $\dot{x}_i = f_i(x, u, d)$ s.t. $\delta_i^* = |f - l_i| > 0$ and $\forall u \in \mathcal{U}, f_i(x, u, d) = f_i(x, d)$, will have a non-convex hamiltonian.

Thus, in a two-player Reach game, any state with nonlinear f_i and thus $\delta_i^* > 0$ must have at least some control, or the guarantees of Lemma 3 will not hold.

Note, the dubin's car is thus immediately problematic. On the other hand, integrator chains are fine.

Poor Linear Models are Valueless

Regardless of convexity, a large maximum error will be too conservative for meaningful results (empty *Reach* set, infinite *Avoid* set).

Poor Linear Models are Valueless

Regardless of convexity, a large maximum error will be too conservative for meaningful results (empty *Reach* set, infinite *Avoid* set).

Remark. Lem. 3 does not guarantee that $\mathcal{R}_{\delta^*}(\mathcal{T}, t)$ is non-empty. In fact, mapping (26) to the form (19) yields a simple heuristic.

Poor Linear Models are Valueless

Regardless of convexity, a large maximum error will be too conservative for meaningful results (empty *Reach* set, infinite *Avoid* set).

Remark. Lem. 3 does not guarantee that $\mathcal{R}_{\delta^*}(\mathcal{T}, t)$ is non-empty. In fact, mapping (26) to the form (19) yields a simple heuristic.

E.g. if \mathcal{U} is defined by a ball with $Q \triangleq r_{\mathcal{U}} I$ and $\exists \tau$ for $t < \tau$,

$$\delta^* > r_{\mathcal{U}} \|B_1^\top \Phi(t)^{-\top}\| \implies \lim_{t \rightarrow -\infty} \mathcal{R}_{\delta^*}(\mathcal{T}, t) = \emptyset, \quad (36)$$

despite that $\mathcal{R}(\mathcal{T}, t) \neq \emptyset$ in general. The condition ensures a globally-positive Hamiltonian $H_{\delta^*}(x, p, t) \geq 0$.

Methods for Tighter Envelopes

Time-varying Feasibility and Error

To use the state-independent Hopf formula, the error δ^* cannot be a function of x , however, we may define a time-varying error bound $\delta^*(\tau) : [t, T] \mapsto \mathbb{R}$ that implicitly captures a spatial relationship for a tighter bound on the error.

Time-varying Feasibility and Error

To use the state-independent Hopf formula, the error δ^* cannot be a function of x , however, we may define a time-varying error bound $\delta^*(\tau) : [t, T] \mapsto \mathbb{R}$ that implicitly captures a spatial relationship for a tighter bound on the error.

Corollary

(Time-varying Error) Let the maximum antagonistic error be defined by,

$$\delta^*(\tau) \triangleq \max_{x \in \mathcal{S}(\tau)} \|(f - l)(x, u, d)\|, \quad (37)$$

where $\mathcal{S}(\tau) : [t, T] \mapsto \mathbb{K}\mathbb{R}^{n_x}$ maps a time to the compact set containing all trajectories for time interval $[\tau, T]$.

Time-varying Feasibility and Error

To use the state-independent Hopf formula, the error δ^* cannot be a function of x , however, we may define a time-varying error bound $\delta^*(\tau) : [t, T] \mapsto \mathbb{R}$ that implicitly captures a spatial relationship for a tighter bound on the error.

Corollary

(Time-varying Error) Let the maximum antagonistic error be defined by,

$$\delta^*(\tau) \triangleq \max_{x \in \mathcal{S}(\tau)} \|(f - l)(x, u, d)\|, \quad (37)$$

where $\mathcal{S}(\tau) : [t, T] \mapsto \mathbb{K}\mathbb{R}^{n_x}$ maps a time to the compact set containing all trajectories for time interval $[\tau, T]$. Then for reach and avoid games resp.,

$$\begin{aligned} \mathcal{R}_{\delta^*}(\mathcal{T}, t) &\subset \mathcal{R}_{\delta^*(\tau)}(\mathcal{T}, t) \subset \mathcal{R}(\mathcal{T}, t), \\ \mathcal{R}_{\delta^*}^-(\mathcal{T}, t) &\supset \mathcal{R}_{\delta^*(\tau)}^-(\mathcal{T}, t) \supset \mathcal{R}(\mathcal{T}, t). \end{aligned} \quad (38)$$

Time-varying Feasibility and Error

To use the state-independent Hopf formula, the error δ^* cannot be a function of x , however, we may define a time-varying error bound $\delta^*(\tau) : [t, T] \mapsto \mathbb{R}$ that implicitly captures a spatial relationship for a tighter bound on the error.

Corollary

(Time-varying Error) Let the maximum antagonistic error be defined by,

$$\delta^*(\tau) \triangleq \max_{x \in \mathcal{S}(\tau)} \|(f - l)(x, u, d)\|, \quad (37)$$

where $\mathcal{S}(\tau) : [t, T] \mapsto \mathbb{K}\mathbb{R}^{n_x}$ maps a time to the compact set containing all trajectories for time interval $[\tau, T]$. Then for reach and avoid games resp.,

$$\begin{aligned} \mathcal{R}_{\delta^*}(\mathcal{T}, t) &\subset \mathcal{R}_{\delta^*(\tau)}(\mathcal{T}, t) \subset \mathcal{R}(\mathcal{T}, t), \\ \mathcal{R}_{\delta^*}^-(\mathcal{T}, t) &\supset \mathcal{R}_{\delta^*(\tau)}^-(\mathcal{T}, t) \supset \mathcal{R}(\mathcal{T}, t). \end{aligned} \quad (38)$$

If $\mathcal{S}(\tau)$ is evaluated sparsely at $\{s_i\}_{s_1=t}^{s_n=T}$ then the function $\hat{\delta}^*(\tau) \triangleq \{\delta^*(s_i^*) \mid s_i^* \triangleq \max_{s_i \leq \tau} s_i\}$ will yield the same intermediate result.

Time-Varying Error Envelopes in Van der Pol




Figure: For a target \mathcal{T} (black), true reachable sets $\mathcal{R}/\mathcal{R}^-$ (gold) are solved for a *Reach* and *Avoid* game resp. at $t = [0.13, 0.26, 0.39]$. The corresponding reachable sets of the safe envelope $\mathcal{R}_{\delta^*(\tau)}/\mathcal{R}_{\delta^*(\tau)}^-$ (green) based on the TV Taylor-series linearization error are also solved.

Single-Player Error Sets

To assume the set \mathcal{S} contains all trajectories is sufficient but superfluous.
Ultimately, the error need only be bounded on the trajectories evolving from the optimal strategy of Player I for any input of Player II (*in the BRT*).

Single-Player Error Sets

To assume the set \mathcal{S} contains all trajectories is sufficient but superfluous. Ultimately, the error need only be bounded on the trajectories evolving from the optimal strategy of Player I for any input of Player II (*in the BRT*).

Corollary

Suppose $\mathcal{S}_{\mathcal{U}}, \mathcal{S}_{\mathcal{D}} \subset \mathbb{KR}^{n_x}$ are defined as in (9) for control-only and disturbance-only settings resp., and let the maximum errors on these sets be defined as $\delta_{\mathcal{U}}^*$ and $\delta_{\mathcal{D}}^*$. Then for Reach and Avoid games resp.,

$$\begin{aligned}\mathcal{R}_{\delta^*}(\mathcal{T}, t) &\subset \mathcal{R}_{\delta_{\mathcal{U}}^*}(\mathcal{T}, t) \subset \mathcal{R}(\mathcal{T}, t), \\ \mathcal{R}_{\delta^*}^-(\mathcal{T}, t) &\supset \mathcal{R}_{\delta_{\mathcal{D}}^*}^-(\mathcal{T}, t) \supset \mathcal{R}(\mathcal{T}, t).\end{aligned}\tag{39}$$

Single-Player Error Envelopes in Van der Pol




Figure: The reachable sets of the safe envelope $\mathcal{R}_{\delta_{\mathcal{U}}^*(\tau)}/\mathcal{R}_{\delta_{\mathcal{D}}^*(\tau)}^-$ (pink) based on the time-varying Taylor-series linearization error for single-player feasible sets $\mathcal{S}_{\mathcal{U}}, \mathcal{S}_{\mathcal{D}}$ (light blue) are also solved.

Error for Ensemble of Linear Models

Variations of the linear problem may offer variations in the safe envelope, yielding more information.

Error for Ensemble of Linear Models

Variations of the linear problem may offer variations in the safe envelope, yielding more information.

Lemma

Let $\{l_i(x, u, d) = A_i x + B_{1,i}u + B_{2,i}d\}_i$ be a set of linear models and $\delta_i \triangleq \max_{x \in \mathcal{S}} |(f - l_i)|_s|$ be their respective errors. Then for reach and avoid games,

$$\bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}, t) \subset \mathcal{R}(\mathcal{T}, t) \subset \bigcap_i \mathcal{R}_{\delta_i^*}^-(\mathcal{T}, t). \quad (40)$$

Error for Ensemble of Linear Models

Variations of the linear problem may offer variations in the safe envelope, yielding more information.

Lemma

Let $\{l_i(x, u, d) = A_i x + B_{1,i}u + B_{2,i}d\}_i$ be a set of linear models and $\delta_i \triangleq \max_{x \in \mathcal{S}} |(f - l_i)|_s|$ be their respective errors. Then for reach and avoid games,

$$\bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}, t) \subset \mathcal{R}(\mathcal{T}, t) \subset \bigcap_i \mathcal{R}_{\delta_i^*}^-(\mathcal{T}, t). \quad (40)$$

Note, the sub-problems corresponding to each linear model may be solved independently, and hence the load of the total ensemble solution scales in a parallelizable, linear fashion with respect to number of models.

Ensemble Error Envelopes in Van der Pol




Figure: The linear ensemble envelopes $\mathcal{R}_{\delta_{U_i}^*(\tau)}/\mathcal{R}_{D_i}^-(\tau)$ (blues) based on the ensemble trajectories (blue dashed), with time-varying, single-player Taylor-series linearization error are solved. The union and intersection of ensemble envelopes in *Reach* and *Avoid* resp. are also safe envelopes

Target Partitioning Envelopes

One may recall the proposed work hinges on the transformation of the space-dependent error to a *uniformly* applied quantity.

Hence, splitting the target frees some divisions from the maximum error in another.²

²This procedure could be repeated *ad infinitum*, reducing $\varepsilon \rightarrow 0$ for $\mathcal{T}_i = B(x, \varepsilon)$ to converge to a space-optimal formulation of [(9)].

Target Partitioning Envelopes

Lemma

Suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \subseteq \mathcal{T}$. Let S_i be defined as in (9) with $x(T) \in \mathcal{T}_i$, and let δ_i^* be the corresponding error for l . For the Reach game,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, t) \subset \bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (41)$$

Target Partitioning Envelopes

Lemma

Suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \subseteq \mathcal{T}$. Let \mathcal{S}_i be defined as in (9) with $x(T) \in \mathcal{T}_i$, and let δ_i^* be the corresponding error for l . For the Reach game,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, t) \subset \bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (41)$$

For the Avoid game, suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \supset \mathcal{T}$. Let $L(t) \triangleq \max \|x - y\|$ for $x, y \in \mathcal{S}(\mathcal{S}(\mathcal{T}_i, t) \cap \mathcal{S}(\mathcal{T}_j, t), -t)$, the forward feasible set of the intersection of backwards feasible sets of any partition elements.

Target Partitioning Envelopes

Lemma

Suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \subseteq \mathcal{T}$. Let \mathcal{S}_i be defined as in (9) with $x(T) \in \mathcal{T}_i$, and let δ_i^* be the corresponding error for l . For the Reach game,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, t) \subset \bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (41)$$

For the Avoid game, suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \supset \mathcal{T}$. Let $L(t) \triangleq \max \|x - y\|$ for $x, y \in \mathcal{S}(\mathcal{S}(\mathcal{T}_i, t) \cap \mathcal{S}(\mathcal{T}_j, t), -t)$, the forward feasible set of the intersection of backwards feasible sets of any partition elements.

For $x \in \partial \mathcal{T}_i$, if $(\min_{y \in \mathcal{T}} \|x - y\| \geq L(t)) \vee (\exists j, x \in \mathcal{T}_j, \min_{y \in \partial \mathcal{T}_j} \|x - y\| \geq L(t))$, we say the partition “ $L(t)$ -overlaps” and the following property holds,

$$\bigcup_i \mathcal{R}_{\delta_i^*}^-(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (42)$$

Target Partitioning Envelopes

Lemma

Suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \subseteq \mathcal{T}$. Let \mathcal{S}_i be defined as in (9) with $x(T) \in \mathcal{T}_i$, and let δ_i^* be the corresponding error for l . For the Reach game,

$$\mathcal{R}_{\delta^*}(\mathcal{T}, t) \subset \bigcup_i \mathcal{R}_{\delta_i^*}(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (41)$$

For the Avoid game, suppose $\{\mathcal{T}_i\}_i$ is a partition of \mathcal{T} such that $\cup_i \mathcal{T}_i \supset \mathcal{T}$. Let $L(t) \triangleq \max \|x - y\|$ for $x, y \in \mathcal{S}(\mathcal{S}(\mathcal{T}_i, t) \cap \mathcal{S}(\mathcal{T}_j, t), -t)$, the forward feasible set of the intersection of backwards feasible sets of any partition elements.

For $x \in \partial \mathcal{T}_i$, if $(\min_{y \in \mathcal{T}} \|x - y\| \geq L(t)) \vee (\exists j, x \in \mathcal{T}_j, \min_{y \in \partial \mathcal{T}_j} \|x - y\| \geq L(t))$, we say the partition “ $L(t)$ -overlaps” and the following property holds,

$$\bigcup_i \mathcal{R}_{\delta_i^*}^-(\mathcal{T}_i, t) \subset \mathcal{R}(\mathcal{T}, t). \quad (42)$$

Target Partitioning Envelopes in Van der Pol




Figure: The partition envelopes $\mathcal{R}_{\delta_{U_i}^*(\tau)}/\mathcal{R}_{\mathcal{D}_i}^-(\tau)$ (pinks) based on the partitioned target (faint pinks), with time-varying, single-player Taylor-series linearization error are solved. The union of partition envelopes in *Reach* and *Avoid* games is also a safe envelope.

Forward-Backward Feasibility and Error

Finally, there is a substantial improvement in the special case when considering the reachability or avoidability of a local region $\mathcal{X}_g \subset \mathcal{X} \subset \mathbb{R}^{n_x}$, e.g. a single state.

Forward-Backward Feasibility and Error

Finally, there is a substantial improvement in the special case when considering the reachability or avoidability of a local region $\mathcal{X}_g \subset \mathcal{X} \subset \mathbb{R}^{n_x}$, e.g. a single state.

Lemma

Let $\mathcal{S}_g \triangleq \mathcal{S}(\mathcal{X}_g, -t)$ be the feasible set of \mathcal{X}_g at $|t|$ time in the future and δ_g^* be the error. Furthermore, let $\bar{\mathcal{T}} \triangleq \mathcal{T} \cap \mathcal{S}_g$ be its intersection with the target and $\bar{\mathcal{S}} \triangleq \mathcal{S}(\bar{\mathcal{T}}, t)$ be the backwards feasible set of the intersection. Suppose $\bar{\mathcal{S}}_g \triangleq \bar{\mathcal{S}} \cap \mathcal{S}_g$ and $\bar{\delta}_g^*$ to be the error.

Then for Reach and Avoid games resp.,

$$\begin{aligned} V_{\delta^*}(x, t) &\geq V_{\delta_g^*}(x, t) \geq V_{\bar{\delta}_g^*}(x, t) \geq V(x, t), \\ V_{\delta^*}^-(x, t) &\leq V_{\delta_g^*}^-(x, t) \leq V_{\bar{\delta}_g^*}^-(x, t) \leq V(x, t). \end{aligned} \quad x \in \mathcal{X}_g \tag{43}$$

Forward-Backward Feasibility Illustrative Example




Figure: Forward-Backward feasibility sets in the Lorenz 96 Model.

Computation of \mathcal{S} and δ^* *a priori*

There are several developed DI approaches to the problem of safely approximating $\hat{\mathcal{S}} \supset \mathcal{S}$ (10; 11; 7; 12). At simplest, one may consider the “Lipschitz tube” that bounds the flow (12), however, this is unlikely to yield a tight superset.

Computation of \mathcal{S} and δ^* *a priori*

There are several developed DI approaches to the problem of safely approximating $\hat{\mathcal{S}} \supset \mathcal{S}$ (10; 11; 7; 12). At simplest, one may consider the “Lipschitz tube” that bounds the flow (12), however, this is unlikely to yield a tight superset.

Once $\hat{\mathcal{S}} \supset \mathcal{S}$ has been over-approximated, we need to solve δ^* ,

$$\delta^*(\tau) \triangleq \max_{x \in \mathcal{S}(\tau)} \|(f - l)(x, u, d)\|. \quad (44)$$

Computation of \mathcal{S} and δ^* *a priori*

There are several developed DI approaches to the problem of safely approximating $\hat{\mathcal{S}} \supset \mathcal{S}$ (10; 11; 7; 12). At simplest, one may consider the “Lipschitz tube” that bounds the flow (12), however, this is unlikely to yield a tight superset.

Once $\hat{\mathcal{S}} \supset \mathcal{S}$ has been over-approximated, we need to solve δ^* ,

$$\delta^*(\tau) \triangleq \max_{x \in \mathcal{S}(\tau)} \|(f - l)(x, u, d)\|. \quad (44)$$

This may be non-convex. However, if l is the Taylor series linearization, then we may upper-bound with convex objective.

Multi-Agent Dubin's Pursuit-Evasion

Consider N Dubin's cars in the relative frame of an additional car [(13; 14)]. Each agent's relative state, $x_i = [x_{\Delta_i}, y_{\Delta_i}, \theta_{\Delta_i}]$, evolves,

$$\dot{x}_i = \begin{bmatrix} -v_a + v_b \cos(\theta_{\Delta_i}) + ay_{\Delta_i} \\ v_b \sin(\theta_{\Delta_i}) - ax_{\Delta_i} \\ b_i - a \end{bmatrix} \quad (45)$$

where $a \in [-1, 1]$ and b_i are the controls of the central agent and the $i \in [1, N]$ agent resp., and v_a and $v_b (= 3)$ are their velocities resp.

Multi-Agent Dubin's Pursuit-Evasion

Consider N Dubin's cars in the relative frame of an additional car [(13; 14)]. Each agent's relative state, $x_i = [x_{\Delta_i}, y_{\Delta_i}, \theta_{\Delta_i}]$, evolves,

$$\dot{x}_i = \begin{bmatrix} -v_a + v_b \cos(\theta_{\Delta_i}) + ay_{\Delta_i} \\ v_b \sin(\theta_{\Delta_i}) - ax_{\Delta_i} \\ b_i - a \end{bmatrix} \quad (45)$$

where $a \in [-1, 1]$ and b_i are the controls of the central agent and the $i \in [1, N]$ agent resp., and v_a and $v_b (= 3)$ are their velocities resp.

In the game, the central agent seeks to avoid all N other agents which simultaneously aim to reach it. Note, the combined system has $x \in \mathbb{R}^{3N}$.

Multi-Agent Dubin's Pursuit-Evasion

Note, the linearization of this system takes the form (14),

$$\dot{x} = \begin{bmatrix} A_1 & \cdots & 0 \\ & A_2 & \vdots \\ \vdots & \ddots & \\ 0 & \cdots & A_N \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_N \end{bmatrix} + \begin{bmatrix} B_{a_1} \\ B_{a_2} \\ \vdots \\ B_{a_N} \end{bmatrix} a \quad (46)$$

$$+ \begin{bmatrix} B_{b_1} & \cdots & 0 \\ & B_{b_2} & \vdots \\ \vdots & \ddots & \\ 0 & \cdots & B_{b_N} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_N \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_N \end{bmatrix},$$

where $(A, B_a, B_b) \triangleq (D_x f, D_a f, D_b f)|_{\tilde{x}, 0, \tilde{b}}$ and ε is the residual error.

The error is derived by linearizing around the iLQG trajectory $\tilde{x}(\tau)$ without pursuer action, and solving the feasible pursuer set for the initial condition given by $x_i(0) = [10, 0, \pm\pi]$.

Multi-Agent Dubin's Pursuit-Evasion

Capture may occur iff the central agent is close ($r = 1.5$) to one pursuer and near the entire team ($r_f = 5$).

Multi-Agent Dubin's Pursuit-Evasion

Capture may occur iff the central agent is close ($r = 1.5$) to one pursuer and near the entire team ($r_f = 5$).

Hence, the target for each pair (central agent, team member) takes the form of an ellipsoid, given by

$$\Omega_i \triangleq \{x \mid x^\top W_i^{-1} x \leq 1\} \quad (47)$$

where $W_i \in \mathbb{R}^{3N \times 3N}$ is a block-diagonal matrix of diagonal matrices with values of r^2 in the i -th block's first and second diagonals and θ_{max}^2 in the third, and r_f^2 in the j -th block's first and second diagonals θ_{max}^2 in the third for all $j \neq i$.

Multi-Agent Dubin's Pursuit-Evasion

Capture may occur iff the central agent is close ($r = 1.5$) to one pursuer and near the entire team ($r_f = 5$).

Hence, the target for each pair (central agent, team member) takes the form of an ellipsoid, given by

$$\Omega_i \triangleq \{x \mid x^\top W_i^{-1} x \leq 1\} \quad (47)$$

where $W_i \in \mathbb{R}^{3N \times 3N}$ is a block-diagonal matrix of diagonal matrices with values of r^2 in the i -th block's first and second diagonals and θ_{max}^2 in the third, and r_f^2 in the j -th block's first and second diagonals θ_{max}^2 in the third for all $j \neq i$.

The joint system has a target then defined by $\Omega = \cup \Omega_i$, which max-plus algebras certify may be solved by the Hopf formula independently (15).

Multi-Agent Dubin's Pursuit-Evasion

Note, this game may be non-convex, however, in the *Avoid* context a conservative lower-value V_{δ^*} may be solved despite this.

Multi-Agent Dubin's Pursuit-Evasion

Note, this game may be non-convex, however, in the *Avoid* context a conservative lower-value V_{δ^*} may be solved despite this.

We use the Hopf formula with the linearized system and antagonistic error to solve when the central agent may evade capture, given by,

$$\min_{t \in [0,1]} V(x_0, t) = V^*(x_0) > 0. \quad (48)$$

Multi-Agent Dubin's Pursuit-Evasion

Note, this game may be non-convex, however, in the *Avoid* context a conservative lower-value V_{δ^*} may be solved despite this.

We use the Hopf formula with the linearized system and antagonistic error to solve when the central agent may evade capture, given by,

$$\min_{t \in [0,1]} V(x_0, t) = V^*(x_0) > 0. \quad (48)$$

We will solve the surface of $V^*(x_0(\theta_a); v_a, \max |b|) = 0$ i.e. w.r.t. parameters $v_a, \max |b|$ and θ_a .

Multi-Agent Dubin's Pursuit-Evasion

Let x_0 be defined s.t. the agents form a regular N -polygon around the evader, oriented relative to the evader angle θ_a .


Multi-Agent Dubin's Pursuit-Evasion

Let x_0 be defined s.t. the agents form a regular N -polygon around the evader, oriented relative to the evader angle θ_a . Hence,

$$\begin{bmatrix} x_{\Delta_i,0} \\ y_{\Delta_i,0} \end{bmatrix} = R_{\text{rot}} \left((i-1) \frac{2\pi}{N} + \theta_a \right) \begin{bmatrix} 0 \\ r_p \end{bmatrix}$$



$$\theta_{\Delta_i,0} = (i-1) \frac{2\pi}{N} + \theta_a - \pi/2 \quad (49)$$

where $r_p = v_b$ is the initial distance between each agent and the central agent.



Multi-Agent Dubin's Pursuit-Evasion



N – Dubins Capture : $V^* > 0$, 3 Pursuers



Mean time to compute $V^*(x_0; a, b, v_a, v_b)$ for $N = 3$ (9 dimensions) is 5.9 s/agent.

Multi-Agent Dubin's Pursuit-Evasion

N – Dubins Capture : $V^* > 0$, 5 Pursuers



Mean time to compute $V^*(x_0; a, b, v_a, v_b)$ for $N = 5$ (15 dimensions) is 8.0 s/agent.

Thank you!

This research group - people, undergrad and advisor - is truly awesome.

- [1] L. C. Evans and P. E. Souganidis, "Differential games and representation formulas for solutions of Hamilton-Jacobi-isaacs equations," *Indiana University mathematics journal*, vol. 33, no. 5, pp. 773–797, 1984.
- [2] I. Rublev, "Generalized Hopf formulas for the nonautonomous Hamilton-Jacobi equation," *Computational Mathematics and Modeling*, vol. 11, no. 4, pp. 391–400, 2000.
- [3] Q. Wei, "Viscosity solution of the hamilton–jacobi equation by a limiting minimax method," *Nonlinearity*, vol. 27, no. 1, p. 17, 2013.
- [4] Y. Wang, W. Yin, and J. Zeng, "Global convergence of admm in nonconvex nonsmooth optimization," *Journal of Scientific Computing*, vol. 78, pp. 29–63, 2019.
- [5] A. F. Filippov, "Differential equations with discontinuous right-hand side," *Matematicheskii sbornik*, vol. 93, no. 1, pp. 99–128, 1960.
- [6] L. C. Evans, "Envelopes and nonconvex hamilton–jacobi equations," *Calculus of Variations and Partial Differential Equations*, vol. 50, no. 1-2, pp. 257–282, 2014.
- [7] X. Chen, E. Abraham, and S. Sankaranarayanan, "Taylor model flowpipe construction for non-linear hybrid systems," in *2012 IEEE 33rd Real-Time Systems Symposium*. IEEE, 2012, pp. 183–192.



- [8] S. Bogomolov, M. Forets, G. Frehse, K. Potomkin, and C. Schilling, "Juliareach: a toolbox for set-based reachability," in *Proceedings of the 22nd ACM International Conference on Hybrid Systems: Computation and Control*, 2019, pp. 39–44.
- [9] D. Lee and C. J. Tomlin, "Iterative method using the generalized Hopf formula: Avoiding spatial discretization for computing solutions of Hamilton-Jacobi equations for nonlinear systems," in *2019 IEEE 58th Conference on Decision and Control (CDC)*. IEEE, 2019, pp. 1486–1493.
- [10] M. Althoff, O. Stursberg, and M. Buss, "Reachability analysis of nonlinear systems with uncertain parameters using conservative linearization," in *2008 47th IEEE Conference on Decision and Control*. IEEE, 2008, pp. 4042–4048.
- [11] J. K. Scott and P. I. Barton, "Bounds on the reachable sets of nonlinear control systems," *Automatica*, vol. 49, no. 1, pp. 93–100, 2013.
- [12] T. Lew and M. Pavone, "Sampling-based reachability analysis: A random set theory approach with adversarial sampling," in *Conference on robot learning*. PMLR, 2021, pp. 2055–2070.
- [13] I. M. Mitchell, A. M. Bayen, and C. J. Tomlin, "A time-dependent hamilton-jacobi formulation of reachable sets for continuous dynamic games," *IEEE Transactions on automatic control*, vol. 50, no. 7, pp. 947–957, 2005.

- [14] M. R. Kirchner, R. Mar, G. Hewer, J. Darbon, S. Osher, and Y. T. Chow, "Time-optimal collaborative guidance using the generalized Hopf formula," *IEEE Control Systems Letters*, vol. 2, no. 2, pp. 201–206, apr 2018. [Online]. Available: <https://doi.org/10.1109/Flcsys.2017.2785357>
- [15] J. Darbon and S. Osher, "Algorithms for overcoming the curse of dimensionality for certain hamilton–jacobi equations arising in control theory and elsewhere," *Research in the Mathematical Sciences*, vol. 3, no. 1, p. 19, 2016.