

Willy Xiao and Kevin Eskici

STAT 221

Pset 4

Nov 4, 2014

question 1. 1  $p(\lambda, \theta) \propto \lambda^{-1}$

$$\begin{bmatrix} N \\ \theta \end{bmatrix} = f \left( \begin{bmatrix} \lambda \\ \theta \end{bmatrix} \right)$$

$$p(N, \theta) = p(\lambda, \theta) * \left\| \begin{bmatrix} \frac{\partial \lambda}{\partial N} & \frac{\partial \theta}{\partial N} \\ \frac{\partial \lambda}{\partial \theta} & \frac{\partial \theta}{\partial \theta} \end{bmatrix} \right\|$$

$$\text{Our Jacobian is } \begin{vmatrix} \theta & -\lambda N^{-2} \\ N & 1 \end{vmatrix}$$

and its determinant is  $\theta + \frac{\lambda}{N}$

$$\begin{aligned} p(N, \theta) &\propto \frac{1}{\theta N} * \left( \theta + \frac{\lambda}{N} \right) \\ &= \frac{1}{\theta N} * \left( \theta + \frac{\theta N}{N} \right) \\ &= \frac{2}{N} \\ &\propto \frac{1}{N} \end{aligned}$$

2 It is an improper prior:  $\int_0^\infty \int_0^1 \lambda^{-1} d\theta d\lambda \rightarrow \ln(\infty) - \ln(0)$ . This cannot integrate to 1 with any constant factor.

3

4 See R code

5