



UNSW
SYDNEY

MATH1131 Mathematics 1A – Algebra

Lecture 9: Complex Numbers

Lecturer: Sean Gardiner – sean.gardiner@unsw.edu.au

Based on slides by Jonathan Kress

Numbers

The **natural numbers** form the set

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}.$$

These arise naturally from trying to **count** whole objects.

Numbers

The **natural numbers** form the set

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}.$$

These arise naturally from trying to **count** whole objects.

The **integers** form the set

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, 3, \dots\}.$$

These arise naturally from trying to solve equations like $x + 2 = 1$.

Numbers

The **natural numbers** form the set

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}.$$

These arise naturally from trying to **count** whole objects.

The **integers** form the set

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, 3, \dots\}.$$

These arise naturally from trying to solve equations like $x + 2 = 1$.

The **rational numbers** form the set

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p, q \in \mathbb{Z}, q \neq 0 \right\}.$$

These arise naturally from trying to solve equations like $2x = 1$.

Numbers

The **natural numbers** form the set

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}.$$

These arise naturally from trying to **count** whole objects.

The **integers** form the set

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, 3, \dots\}.$$

These arise naturally from trying to solve equations like $x + 2 = 1$.

The **rational numbers** form the set

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p, q \in \mathbb{Z}, q \neq 0 \right\}.$$

These arise naturally from trying to solve equations like $2x = 1$.

The **real numbers** form the set

$$\mathbb{R} = \{\text{all points on the real number line}\}.$$

(This set includes numbers like $\sqrt{2}$, π , and e .)

These arise naturally from trying to solve equations like $x^2 = 2$.

Fields

A **field** \mathbb{F} is a set with two operations, $+$ and \times , that satisfies the following properties for all $x, y, z \in \mathbb{F}$:

$x + y \in \mathbb{F}$ (Closure under addition)

$xy \in \mathbb{F}$ (Closure under multiplication)

$(x + y) + z = x + (y + z)$ (Associativity under addition)

$x(yz) = (xy)z$ (Associativity under multiplication)

$x + y = y + x$ (Commutativity under addition)

$xy = yx$ (Commutativity under multiplication)

$x(y + z) = xy + xz$ and $(x + y)z = xz + yz$ (Distributivity)

There is a $0 \in \mathbb{F}$ such that $x + 0 = x$ for all x . (Existence of 0)

There is a $1 \in \mathbb{F}$ such that $1x = x$ for all x . (Existence of 1)

There is $-x \in \mathbb{F}$ such that $x + (-x) = 0$. (Existence of negatives)

If $x \neq 0$, there is $x^{-1} \in \mathbb{F}$ such that $x^{-1}x = 1$. (Existence of inverses)

Fields – Examples

Example

Which of \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} are fields?

Fields – Examples

Example

Which of \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} are fields?

- \mathbb{N} is **not** a field since it does not satisfy two of the properties: no natural numbers (except 0) have negatives, and no natural numbers (except 1) have inverses.

Example

Which of \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} are fields?

- \mathbb{N} is **not** a field since it does not satisfy two of the properties: no natural numbers (except 0) have negatives, and no natural numbers (except 1) have inverses.
- \mathbb{Z} is **not** a field since it does not satisfy one of the properties: no natural numbers (except ± 1) have inverses.

Example

Which of \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} are fields?

- \mathbb{N} is **not** a field since it does not satisfy two of the properties: no natural numbers (except 0) have negatives, and no natural numbers (except 1) have inverses.
- \mathbb{Z} is **not** a field since it does not satisfy one of the properties: no natural numbers (except ± 1) have inverses.
- \mathbb{Q} **is** a field since all the properties hold: for example, the negative of $\frac{p}{q}$ is $-\frac{p}{q}$, and its inverse is $\frac{q}{p}$. We can also check closure by showing $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd} \in \mathbb{Q}$ and $\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd} \in \mathbb{Q}$.

Example

Which of \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} are fields?

- \mathbb{N} is **not** a field since it does not satisfy two of the properties: no natural numbers (except 0) have negatives, and no natural numbers (except 1) have inverses.
- \mathbb{Z} is **not** a field since it does not satisfy one of the properties: no natural numbers (except ± 1) have inverses.
- \mathbb{Q} **is** a field since all the properties hold: for example, the negative of $\frac{p}{q}$ is $-\frac{p}{q}$, and its inverse is $\frac{q}{p}$. We can also check closure by showing $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd} \in \mathbb{Q}$ and $\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd} \in \mathbb{Q}$.
- \mathbb{R} **is** a field since all the properties hold: for example, the negative of some $r \in \mathbb{R}$ is simply $-r$, and its inverse is $\frac{1}{r}$. To check closure, note that the sum or product of any two numbers on the real number line is also on the real number line.

Complex numbers

Consider the equation

$$x^2 = -1.$$

There is no number in \mathbb{R} that solves this. So we can naturally extend our number system again by including numbers that solve these sorts of equations.

Complex numbers

Consider the equation

$$x^2 = -1.$$

There is no number in \mathbb{R} that solves this. So we can naturally extend our number system again by including numbers that solve these sorts of equations.

First we introduce the **imaginary unit** i which has the property that

$$i^2 = -1.$$

We can think of i as $\sqrt{-1}$.

Complex numbers

Consider the equation

$$x^2 = -1.$$

There is no number in \mathbb{R} that solves this. So we can naturally extend our number system again by including numbers that solve these sorts of equations.

First we introduce the **imaginary unit** i which has the property that

$$i^2 = -1.$$

We can think of i as $\sqrt{-1}$.

The **complex numbers** form the set

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}\}.$$

These arise naturally from trying to solve equations like $x^2 = -1$.

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$(3 + 2i) + (5 - 4i) =$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$(3 + 2i) + (5 - 4i) = (3 + 5) + (2i - 4i)$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$(3 + 2i) - (5 - 4i) =$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$(3 + 2i) - (5 - 4i) = (3 - 5) + (2i + 4i)$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

$$(3 + 2i)(5 - 4i) =$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

$$(3 + 2i)(5 - 4i) = 3 \times 5 + 3 \times (-4i) + 2 \times 5i + 2i \times (-4i)$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

$$\begin{aligned}(3 + 2i)(5 - 4i) &= 3 \times 5 + 3 \times (-4i) + 2 \times 5i + 2i \times (-4i) \\ &= 15 - 12i + 10i - 8i^2\end{aligned}$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

$$\begin{aligned}(3 + 2i)(5 - 4i) &= 3 \times 5 + 3 \times (-4i) + 2 \times 5i + 2i \times (-4i) \\ &= 15 - 12i + 10i - 8i^2 \\ &= 15 - 2i - 8 \times (-1)\end{aligned}$$

Complex numbers – Arithmetic

When performing arithmetic on complex numbers, we treat i like a variable but replace i^2 with -1 .

Example

Find the sum, the difference, and the product of $3 + 2i$ and $5 - 4i$.

$$\begin{aligned}(3 + 2i) + (5 - 4i) &= (3 + 5) + (2i - 4i) \\ &= 8 - 2i\end{aligned}$$

$$\begin{aligned}(3 + 2i) - (5 - 4i) &= (3 - 5) + (2i + 4i) \\ &= -2 + 6i\end{aligned}$$

$$\begin{aligned}(3 + 2i)(5 - 4i) &= 3 \times 5 + 3 \times (-4i) + 2 \times 5i + 2i \times (-4i) \\ &= 15 - 12i + 10i - 8i^2 \\ &= 15 - 2i - 8 \times (-1) \\ &= 23 - 2i\end{aligned}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 9 = 0$.

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 9 = 0$.

If $x^2 = -9$, then

$$x = \pm\sqrt{-9}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 9 = 0$.

If $x^2 = -9$, then

$$\begin{aligned}x &= \pm\sqrt{-9} \\ &= \pm\sqrt{9}\sqrt{-1}\end{aligned}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 9 = 0$.

If $x^2 = -9$, then

$$\begin{aligned}x &= \pm\sqrt{-9} \\&= \pm\sqrt{9}\sqrt{-1} \\&= \pm 3i.\end{aligned}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 9 = 0$.

If $x^2 = -9$, then

$$\begin{aligned}x &= \pm\sqrt{-9} \\&= \pm\sqrt{9}\sqrt{-1} \\&= \pm 3i.\end{aligned}$$

That is, the solutions are $x = 3i$ and $x = -3i$.

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

$$(\text{Recall: } ax^2 + bx + c = 0 \iff x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

$$(\text{Recall: } ax^2 + bx + c = 0 \iff x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times 3}}{2 \times 1}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

$$(\text{Recall: } ax^2 + bx + c = 0 \iff x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

$$\begin{aligned} x &= \frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times 3}}{2 \times 1} \\ &= \frac{-2 \pm \sqrt{-8}}{2} \end{aligned}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

$$(\text{Recall: } ax^2 + bx + c = 0 \iff x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

$$\begin{aligned}x &= \frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times 3}}{2 \times 1} \\&= \frac{-2 \pm \sqrt{-8}}{2} \\&= \frac{-2 \pm 2\sqrt{2}\sqrt{-1}}{2}\end{aligned}$$

Complex numbers – Solving equations

Example

Find the solutions to $x^2 + 2x + 3 = 0$.

Using the quadratic formula:

$$(\text{Recall: } ax^2 + bx + c = 0 \iff x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

$$\begin{aligned}x &= \frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times 3}}{2 \times 1} \\&= \frac{-2 \pm \sqrt{-8}}{2} \\&= \frac{-2 \pm 2\sqrt{2}\sqrt{-1}}{2} \\&= -1 \pm \sqrt{2}i\end{aligned}$$

Complex numbers

Definition

The set \mathbb{C} of complex numbers is given by

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}, i^2 = -1\}$$

Complex numbers

Definition

The set \mathbb{C} of complex numbers is given by

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}, i^2 = -1\}$$

Theorem

The set \mathbb{C} of complex numbers is a **field** with respect to the operations $+$ and \times defined by:

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

and

$$(a + bi)(c + di) = ac - bd + (ad + bc)i.$$

Complex numbers

Definition

The set \mathbb{C} of complex numbers is given by

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}, i^2 = -1\}$$

Theorem

The set \mathbb{C} of complex numbers is a **field** with respect to the operations $+$ and \times defined by:

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

and

$$(a + bi)(c + di) = ac - bd + (ad + bc)i.$$

Exercise

Check that all the field properties do in fact hold.

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$i^3 = i^2 \times i$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \end{aligned}$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$i^4 = i^2 \times i^2$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \end{aligned}$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \\ &= 1. \end{aligned}$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \\ &= 1. \end{aligned}$$

$$\text{So } i^5 = i^4 \times i = i,$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \\ &= 1. \end{aligned}$$

$$\text{So } i^5 = i^4 \times i = i, \quad i^6 = i^4 \times i^2 = -1,$$

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \\ &= 1. \end{aligned}$$

So $i^5 = i^4 \times i = i$, $i^6 = i^4 \times i^2 = -1$, $i^7 = i^4 \times i^3 = -i$, etc.

More examples

Example

Simplify

$$1, i, i^2, i^3, i^4, i^5, i^6, i^7, i^8, i^9, \dots$$

Since $i^2 = -1$, we know

$$\begin{aligned} i^3 &= i^2 \times i \\ &= (-1) \times i \\ &= -i. \end{aligned}$$

Similarly,

$$\begin{aligned} i^4 &= i^2 \times i^2 \\ &= (-1) \times (-1) \\ &= 1. \end{aligned}$$

So $i^5 = i^4 \times i = i$, $i^6 = i^4 \times i^2 = -1$, $i^7 = i^4 \times i^3 = -i$, etc.

In general, we can write $i^{4n+k} = i^k$ for any integer n .

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i)$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i)$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

Substituting in:

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

Substituting in:

$$z^3 - 5z^2 + 9z - 5$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

Substituting in:

$$z^3 - 5z^2 + 9z - 5 = (2 + 11i) - 5(3 + 4i) + 9(2 + i) - 5$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

Substituting in:

$$\begin{aligned} z^3 - 5z^2 + 9z - 5 &= (2 + 11i) - 5(3 + 4i) + 9(2 + i) - 5 \\ &= (2 - 15 + 18 - 5) + (11 - 20 + 9)i \end{aligned}$$

More examples

Example

Show that $z = 2 + i$ is a solution of the cubic equation

$$z^3 - 5z^2 + 9z - 5 = 0.$$

First find z^2 :

$$z^2 = (2 + i)(2 + i) = 4 + 2i + 2i + i^2 = 3 + 4i.$$

Next find z^3 :

$$z^3 = z \times z^2 = (2 + i)(3 + 4i) = 6 + 8i + 3i + 4i^2 = 2 + 11i.$$

Substituting in:

$$\begin{aligned} z^3 - 5z^2 + 9z - 5 &= (2 + 11i) - 5(3 + 4i) + 9(2 + i) - 5 \\ &= (2 - 15 + 18 - 5) + (11 - 20 + 9)i \\ &= 0. \end{aligned}$$